

Conclusions: Considering the prevalence and impact of mental health issues in the perinatal period, it is desirable to structure interventions with a holistic and multidisciplinary approach. Perinatal mental health should be prioritized during the entire process of pregnancy and postnatal period. A network of primary and secondary care systems may allow mitigating and/or overcoming vulnerabilities.

Disclosure: No significant relationships.

Keywords: programme; women; perinatal mental health; intervention

EPV1663

Women's mental health during the COVID 19 pandemic and the problem of domestic violence in a lockdown situation

M. Kachaeva^{1*}, N. Semenova², S. Shport³, V. Vasyanina¹, O. Shishkina¹, N. Skibina⁴ and L. Nazarova⁴

¹V. Serbsky National Medical Research Centre for Psychiatry and Narcology, Forensic Psychiatry, Moscow, Russian Federation;

²Moscow Research Institute of Psychiatry, Clinical Psychology Counseling, Moscow, Russian Federation; ³V. Serbsky National Medical Research Centre for Psychiatry and Narcology, Moscow Research Institute Of Psychiatry, Moscow, Russian Federation and ⁴I. M. Sechenov First Moscow State Medical University, Forensic Psychiatry, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.2237

Introduction: The current situation caused by restrictive measures related to the COVID-19 pandemic provokes the high level of aggressiveness and all forms of domestic violence. These results in mental health problems.

Objectives: The purpose of this study was to find out the consequences of domestic violence against women and to identify psychological problems and mental disorders in women.

Methods: A cohort of 18 females was examined by psychiatrists and psychologists. All women turned to specialists at the Moscow Institute of Psychiatry for help.

Results: All women were victims of violence by their husbands and partners. Lockdown situation associated with COVID 19 pandemic has exacerbated the problem of domestic violence. Clinical assessment has revealed different depressive symptoms, anxiety, fear, suicidal tendencies forming the clinical picture of adjustment disorder (2 cases), acute reaction to stress (3 cases), post-traumatic stress disorder (2 cases), depressive episode (8 cases), eating disorders behavior in the form of bulimia and anorexia (3 cases).

Conclusions: The research has revealed that domestic abuse against women associated with lockdown situation during the COVID 19 pandemic often results in psychological and long-term mental health problems. In these cases, prevention is needed to combat violence against women with the participation of public health specialists, psychologists, psychiatrists, sociologists.

Disclosure: No significant relationships.

Keywords: women; women; mental health,

EPV1665

The impact of sexual hormones on depressive symptoms and alcohol use in a woman with delusional disorder at the perimenopausal stage: case report and review of recent works

G.F. Fucho^{1*}, L. Delgado¹, A. González-Rodríguez², A. Guàrdia², A. Álvarez², S. Crivillés^{1,3}, J.D. Barbero^{1,3}, J.A. Monreal^{2,3,4}, J. Labad^{3,4,5} and D. Palao Vidal^{1,3,4}

¹Parc Taulí University Hospital, Department Of Mental Health, Sabadell, Barcelona, Spain; ²Mutua Terrassa University Hospital, Department Of Mental Health, Terrassa, Spain; ³School of Medicine, Universitat Autònoma de Barcelona, Medicine, Cerdanyola del Vallès, Spain; ⁴Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), Salud Mental, Madrid, Spain and ⁵Hospital of Mataró, Consorci Sanitari Del Maresme, Mataró, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.2238

Introduction: The influence of sexual hormones on mental disorders have been extensively reported. In fact, recent studies suggest that sex hormones may play a relevant role in the pathophysiology of psychosis, may be a precipitant when exogenously administered or even be used as a treatment agent of psychotic disorders.

Objectives: To describe the case of a patient with a recent diagnosis of delusional disorder (DD) with an onset in the perimenopausal period.

Methods: Case report and narrative review focused on the impact of sexual hormones on depressive symptoms and alcohol use comorbidity during perimenopause in DD women by using PubMed database.

Results: Case report: A 48-year-old woman diagnosed with DD. The clinical assessment of climacteric symptoms, as well as the gonadotropins (elevated levels of follicle-stimulating hormone -FSH- and luteinizing hormone -LH-) and estrogen levels monitoring (variations on 17-β-estradiol -E2-) enabled to link the endocrine changes with the onset and course of the psychiatric disorder. During the development of the disease, the patient also presented comorbid depressive symptoms and alcohol use disorder. Review: Estrogen depletion seems to increase the risk of psychosis, while scientific literature is not conclusive in establishing a definitive relationship between depressive symptoms and hormonal imbalance in DD. Alcohol use disorder is a common comorbidity in both perimenopausal women and patients with DD.

Conclusions: Multiaxial management of the case helped clinicians to achieve clinical stabilization. Sex differences as well as hormonal pattern disturbances should receive special attention due to the seminal implications in pharmacotherapy and clinical outcomes.

Disclosure: No significant relationships.

Keywords: Delusional disorder; Depression; Alcohol use; sexual hormones

EPV1666

Self-esteem in Moroccan women: comparison between different age groups and influencing factors

A.K. Rhaouti^{1*}, A. Khallouk², S. Belbachir³, A. Ouanass³ and O. Belakbir⁴

¹Hôpital Ar-Razi, Men's Unit B, Sale, Morocco; ²University Psychiatric Hospital Ar-razi, university hospital center, SALE, Morocco,

University Of Mohammed V, Rabat,, salé, Morocco; ³Hôpital psychiatrique ARRAZI de Salé, Psychiatrie, Salé, Morocco and ⁴Psychiatric University Hospital Ar-razi, Men's Unit B, salé, Morocco
 *Corresponding author.
 doi: 10.1192/j.eurpsy.2022.2239

Introduction: This study revolves around self-esteem which is defined as a basic human characteristic related to self-awareness, emotions, cognitions, behavior, lifestyle, general health and socio-economic factors. This fundamental data of the personality is revealed from one person to another as well as from one period to another. Many studies point out that advancing adults do not necessarily imply a decline in sense of self-worth, although skill losses are very real in many areas of psychological activity. Therefore, it seemed interesting to us to further explore this point in Moroccan women.

Objectives: Evaluate and compare self-esteem among different age groups of Moroccan women. Identify the different influencing factors.

Methods: This is a descriptive cross-sectional study using a questionnaire based on two parts, the first based Rosenberg scale, and the second part to identify the presence of certain factors influencing self-esteem.

Results: Our researches have shown a similar results to those of some previous studies. Indeed, we found out that women experience a significant rise of self-esteem simultaneously with the increase of age. Yet, this self-esteem starts to decline in middle-aged women. Several factors can affect it; we can note on the top, the impact of relationships, education and physical health.

Conclusions: This research contributes to our understanding of Moroccan women's self-esteem and to the identification of factors that influence it. Self-esteem is a core identity issue, essential to personal validation and our ability to experience joy. Previous researches also suggests that self-esteem might influence economic welfare and physical health.

Disclosure: No significant relationships.

Keywords: self-esteem; women

EPV1667

Gender disparities in a psychiatric department in Tunisia

E. Bergaoui*, M. Zrelli, N. Staali, M. Moalla, R. Lansari, A. Larnaout and W. Melki

Razi Hospital, Psychiatry D, Manouba, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.2240

Introduction: Gender disparities exist regarding prevalence, symptomatology and risk factors of mental disorders. In Tunisia, there is only one hospital dedicated entirely to mental health which is Razi hospital.

Objectives: The aim of the present study was to assess gender based mental health disparities in a psychiatric department and its related factors.

Methods: A cross sectional and comparative survey was conducted between March and April 2021 in the department of psychiatry D of Razi Hospital including 70 patients with a sex ratio= 1.

Results: The participants were aged between 17 and 68. Men had higher rate of celibacy: 80% of men against 48.57% of women

($p=0.009$). A total of 11.42% of women were illiterate against 2.85% of men, 48% of men were unemployed against 62.85% of women. There was a significant difference between gender and use of cigarettes, cannabis and alcohol ($p<0.001$). The diagnosis was mood disorders for 35.42% of women and 17.14% of men and schizophrenia for 57.14% of women and 77.14% of men. Gender and modality of hospitalization were significantly associated ($p=0.046$): 14% of women were involuntary hospitalized against 40% of men. Time between symptoms onset and consulting is 3.5 years (± 5.67) for women and 1.77 (± 4.75) for men. The mean number of admissions for women is 1.59 and for men 4.2 ($p=0.009$).

Conclusions: Onset of mental disorders for women is 3 to 4 years later than men. They have better premorbid functioning and better social networks. Gender disparities are not only determined biologically but also socially.

Disclosure: No significant relationships.

Keywords: gender disparities; Tunisia

Others

EPV1668

Simultaneous study of behavioral synchronization of two individuals during a cooperative task

B. Kakuszi*, M. Fullajtar, I. Bitter and P. Czobor

Semmelweis University, Psychiatry And Psychotherapy, Budapest, Hungary

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.2241

Introduction: Interpersonally coordinated behaviors are crucial for social interactions. The "Theory of Mind," or mentalization capacity, of an individual is essential for the establishment of behavioral synchronization. The Reading the Mind in the Eyes Test (RMET) is used to assess mentalization, social cognition and empathy. Previous RMET studies, investigated people in isolation, not in social situations. It is unclear how the RMET predicts functioning during real-life social interactions.

Objectives: To investigate the relationship between the performance measured on the RMET test and the synchronous behavior of two individuals interacting with each other during tasks requiring social collaboration.

Methods: Sample included healthy controls (HC, $n=48$) and patients with ADHD ($n=26$) or schizophrenia (SCH, $n=36$) from an ongoing EEG-hyperscanning study, employing a social coordination condition. We applied a Go/NoGo reaction time (RT) task performed by pairs of participants. Synchronous behavior was characterized by the correlation of participants' RTs. We used the percent (%) correct responses from the RMET to characterize social cognition.

Results: In HC, with better social cognitive performance, the correlation of behavioral responses was significantly ($p<0.05$) higher. In ADHD, better performance on the RMET was also accompanied by better behavioral synchronization, but the association did not reach significance due to the smaller sample size. In SCH, no relationship was detected.

Conclusions: In HC and ADHD, the mentalization ability as measured by RMET is associated with the behavioral synchronization between individuals in social interaction. The lack of