

Research Article

Cite this article: Negrisoli RM and Odero DC (2025) Newly established bermudagrass response to topramezone and photosystem II-inhibitor herbicides. *Weed Technol.* **39**(e39), 1–7. doi: [10.1017/wet.2024.83](https://doi.org/10.1017/wet.2024.83)

Received: 5 August 2024
Revised: 14 October 2024
Accepted: 28 October 2024

Associate Editor:

Sandeep Singh Rana, Bayer Crop Science

Nomenclature:

Ametryn; atrazine; metribuzin; topramezone; bermudagrass, *Cynodon dactylon* (L.) Pers. CYNDA; sugarcane, *Saccharum spp.* hybrid

Keywords:

Perennial grass control; PS II-inhibitor herbicides; mixtures; plant cane

Corresponding author:

D. Calvin Odero; Email: dcodero@ufl.edu

Newly established bermudagrass response to topramezone and photosystem II-inhibitor herbicides

Raphael M. Negrisoli¹  and D. Calvin Odero² 

¹Former Graduate Student, University of Florida, Everglades Research and Education Center, Belle Glade, FL, USA and ²Professor, University of Florida, Everglades Research and Education Center, Belle Glade, FL, USA

Abstract

Bermudagrass is the most troublesome and difficult-to-control perennial grass weed in Florida sugarcane. Once established, it may be effectively controlled only during the sugarcane fallow period using a combination of nonselective herbicides and tillage. Options for selective management of bermudagrass that escape sugarcane fallow period management programs must be evaluated to mitigate its progressive increase as the crop cycle increases from plant cane to ratoon crops. Greenhouse and field studies were conducted in Belle Glade, Florida, from 2017 to 2018, to determine the response of newly established bermudagrass from sprigs with stolons to two or three sequential applications of topramezone (25 and 50 g ha⁻¹) every 14 d, and combinations of topramezone (25 and 50 g ha⁻¹) with herbicides that inhibit photosystem II (PS II) such as atrazine (2,240 g ha⁻¹), ametryn (440 g ha⁻¹), and metribuzin (2,240 g ha⁻¹). Two or three sequential applications of topramezone with a cumulative total of 75 to 100 g ha⁻¹ provided >93% bermudagrass control 42 d after the first sequential application under greenhouse and field conditions. These treatments exhibited 12% chance of survival 70 d after the first sequential application. There was an additive effect of PS II-inhibitor herbicides on bermudagrass control in mixtures with topramezone. The mixture of topramezone (50 g ha⁻¹) with metribuzin and atrazine provided more than 87% and 92% bermudagrass control under greenhouse and field conditions, respectively, 42 d after treatment. Bermudagrass treated with topramezone (50 g ha⁻¹) in a mixture with metribuzin exhibited 23% chance of survival 70 d after treatment. The results show good efficacy of sequential topramezone applications every 14 d or in a mixture with the PS II-inhibitor herbicides atrazine and metribuzin for control of newly established bermudagrass that typically escape control measures during the sugarcane fallow management period.

Introduction

Bermudagrass is one of the world's most serious perennial, grassy weeds associated with many crops, including sugarcane (Holm et al. 1977; Mitich 1989). Sugarcane is an important crop in Florida, where it is grown on approximately 165,000 ha, mainly on organic soils in the Everglades agricultural area in the southern part of the state (USDA-NASS 2024; VanWeelden et al. 2024). Bermudagrass is the most troublesome and difficult-to-control perennial grassy weed in Florida sugarcane fields (Webster 2012). Bermudagrass thrives in the hot and humid summers and cool to mild winters, with infrequent frost in the subtropical climate of southern Florida where sugarcane is cultivated. Because bermudagrass is stoloniferous, it can rapidly spread laterally on the surface by an extensive network of strong, flat stolons and subterraneously by deep, scaly rhizomes (Mitich 1989), making it difficult to control once it becomes established in a sugarcane field. Bermudagrass propagates primarily vegetatively via rhizomes and stolons, although limited spread via seeds can occur (Fernandez 2003; Holm et al. 1977).

Sugarcane is a 3- to 4-yr perennial crop obtained from planting vegetative stalk pieces or setts from late August until early January. In Florida, sugarcane is harvested from mid-October to the end of May. After the first crop in a year, referred to as plant cane, has been harvested, another crop of stalks, called a ratoon, grows from the old plant stubble. One to three ratoon crops may be harvested approximately every year after the first plant cane harvest, depending on cane and sugar yields, prevailing environmental conditions, and occurrences of pests. Interference from established perennial grasses such as bermudagrass can result in shortening the longevity of the sugarcane crop by reducing the number of ratooning years it achieves. During the multiyear sugarcane cropping period, tillage is limited to the row middles, thus allowing bermudagrass to establish over sugarcane rows and become difficult to selectively control (Richard and Dalley 2007). Reduction in sugarcane yield from bermudagrass infestation results in fallowing fields prior to replanting. The fallow period begins after the harvest of the final ratoon crop with

© The Author(s), 2024. Published by Cambridge University Press on behalf of Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



mechanical destruction of the stubble crop and continues until the subsequent planting of the new crop. In Florida, fallow fields can also be planted with crops such as sweet corn (*Zea mays* spp. *saccharata*), snap bean (*Phaseolus vulgaris* L.), lettuce (*Lactuca sativa* L.), radish (*Raphanus sativus* L.), or rice (*Oryza sativa* L.) prior to sugarcane planting (Odero et al. 2023).

Although bermudagrass is a short-stature grass compared to taller grasses such as johnsongrass [*Sorghum halepense* (L.) Pers.] and itchgrass [*Rottboellia cochinchinensis* (Lour.) Claton], which infests sugarcane (Leon and Odero 2018), bermudagrass can produce heavy ground cover that is capable of causing sugarcane yield losses. Bermudagrass interference in sugarcane fields in Louisiana have reduced sugar yields by up to 32% each year in the plant cane crop and the second ratoon crop, with greater yield loss occurring in the plant cane crop (Richard 1992, 1993; Richard and Dalley 2007). Even though no studies have been carried on the effects of bermudagrass interference on sugarcane yield in Florida, yield reduction is certainly higher because of the lack of hard winter seasons that occur in Louisiana, for instance, where bermudagrass becomes dormant, thereby slowing its growth and development. This results in progressively greater bermudagrass infestations in Florida as the crop cycle increases from plant cane into a ratoon crop. Because herbicides do not exist for selective control of bermudagrass in sugarcane fields, its vegetative propagules are usually only destroyed or reduced during the sugarcane fallow period primarily through multiple glyphosate applications and tillage (Ethredge et al. 2009; Miller et al. 1999; Richard 1997). Ethredge et al. (2009) reported that bermudagrass groundcover was reduced by 81% to 95% when treated with multiple applications of glyphosate in combination with tillage. In Florida, where sugarcane is grown in rotation with vegetable crops during the fallow period, bermudagrass vegetative propagules are reduced with graminicides used on those crops (Sandoya-Miranda et al. 2024). However, bermudagrass can still escape into areas of new plant cane that have been established from viable stolons and rhizomes despite the efficacy of fallow management programs (Ethridge et al. 2009). Also, bermudagrass escapes occur when sugarcane is successively replanted immediately after harvest of the final ratoon crop from viable stolons and rhizomes (Fernandez 2003). Approximately 30% of the Florida sugarcane acreage is plant cane with 66% and 34% from the fallow and successive planting systems, respectively (VanWeelden et al. 2024).

It is important that management programs be developed that target bermudagrass that escapes fallow management programs to mitigate any increases over time during the multiyear sugarcane cropping period. Topramezone is a postemergence herbicide that was recently approved for use on sugarcane to control broadleaf and grassy weeds among the crop (Anonymous 2018; Negrisoli et al. 2020). Topramezone belongs to the pyrazole family, which inhibits 4-hydroxyphenylpyruvate dioxygenase (HPPD), a key enzyme in the biosynthesis of carotenoids (Grossmann and Ehrhardt 2007; Jhala et al. 2023; Shaner 2014). Topramezone elicits bleaching or whitening in susceptible plant species followed by death (Grossmann and Ehrhardt 2007; Jhala et al. 2023; Shaner 2014). It has been shown to significantly bleach bermudagrass leaf tissue and reduce total chlorophyll, lutein, and total xanthophyll cycle pigment concentrations (Brosnan et al. 2011; Elmore et al. 2011). Topramezone and mixtures with herbicides that inhibit photosystem II (PS II) provide more efficacious weed control compared with topramezone applied alone (Fluttert et al. 2022; Negrisoli et al. 2020). Triazine herbicides that inhibit PS II, including atrazine, ametryn, and metribuzin, are widely used in

Florida sugarcane for weed control (Odero and Dusky 2021). Therefore, it is important to determine the efficacy of topramezone in mixtures with PS II-inhibitor herbicides used in Florida sugarcane to control newly established bermudagrass plants that escape fallow management programs. Thus, the objective of this study was to evaluate the level of control of newly established bermudagrass from sprigs with stolons using 1) sequential applications of topramezone, and 2) topramezone applied alone or in combination with PS II-inhibitor herbicides atrazine, ametryn, and metribuzin.

Materials and Methods

Sequential Topramezone Application Study

Greenhouse Experiment

Two greenhouse experiments were conducted at the University of Florida Everglades Research and Education Center (EREC) in Belle Glade, Florida, from 2017 to 2018. Bermudagrass was established using sprigs collected from sugarcane fields at the EREC. Sprigs with stolons averaging 15 cm long with two to four rooting nodes were planted into round 7.6-L pots (16-cm top diameter and 17 cm height) filled with a commercial potting medium (Fafard®; Sun Gro Horticulture, Agawam, MA) on September 18, 2017, and December 13, 2017, for the first and second experimental runs, respectively. The basal part of sprigs with stolons that had rooting nodes were planted approximately 8 cm below the potting medium surface. A slow-release 14-14-14 fertilizer (Osmocote®; The Scotts Company, Marysville, OH) was mixed with the potting medium, and the plants were kept in a greenhouse with a maximum temperature of 30 °C under natural light for the entire study. The plants were watered as needed to ensure that moisture was not a limiting factor. Plants averaged 16 cm in length, ascending to decumbent stems 21 d after establishment. Herbicide treatments were applied on October 9, 2017, for the first experimental run and on January 3, 2018, for the second experimental run after plant sizes were recorded 21 d after establishment.

The experiment was a completely randomized design with five replications of each treatment. Topramezone (Armezon®, BASF Corporation, Research Triangle Park, NC) was applied at 25 and 50 g ha⁻¹ once or sequentially every 14 d for up to three applications for a total of nine treatments (Table 1). All herbicide treatments included methylated seed oil (99% ai; Dyne-Amic®; Helena Chemical Co., Collierville, TN) at 10 mL L⁻¹. A nontreated control was included for comparison. Herbicide treatments were applied using a moving-nozzle spray chamber (Generation II Spray Booth; Devries Manufacturing Hollandale, MN) equipped with a TeeJet 8002E nozzle tip (TeeJet Technologies, Glendale Heights, IL) calibrated to deliver 187 L ha⁻¹ at 172 kPa. Plants were returned to the greenhouse immediately after herbicide application in the spray chamber.

Bermudagrass control was visually estimated using a scale of 0 to 100, with 0 being no control and 100 being complete plant death, estimated 14 d after the final or third sequential topramezone treatment (equivalent to 42 d after the first sequential or single application). Plants were harvested at the soil surface level 42 d after the first sequential or single application and dried in an oven for 48 h at 60 °C to determine aboveground dry weight.

Field Experiment

Two field experiments were conducted at the EREC in Belle Glade, Florida (26.6583°N, 80.6260°W) in 2018 and 2019. The soil was a

Table 1. Newly established bermudagrass control and aboveground dry weight in response to sequential applications of topramezone 42 d after the first sequential treatment (equivalent to 14 d after the final third sequential application) in greenhouse and field experiments combined over two experimental runs in 2017 to 2018 in Belle Glade, FL.^{a,b}

Treatment program ^c	Rate	Greenhouse		Field	
		Control	Dry weight	Control	P(survival) ^d
	g ha ⁻¹	%	g	%	%
Nontreated control ^e	–	–	9.19 b	–	100
Topramezone	25	78 c	1.21 a	61 b	100
Topramezone	50	89 b	1.34 a	84 a	88
Topramezone	25 fb 25	100 a	1.26 a	87 a	75
Topramezone	50 fb 50	100 a	0.95 a	99 a	12
Topramezone	25 fb 25 fb 25	100 a	0.94 a	98 a	25
Topramezone	50 fb 25	100 a	0.84 a	93 a	37
Topramezone	25 fb 50	99 a	1.10 a	93 a	37
Topramezone	50 fb 25 fb 25	100 a	1.24 a	98 a	12
Topramezone	25 fb 25 fb 50	100 c	0.93 a	100 b	12

^aAbbreviation: fb, followed by.^bMeans followed by the same letter within a column are not significantly different according to Tukey's test ($P < 0.05$).^cSequential applications were made every 14 d.^dP(survival) is the probability of bermudagrass survival (the presence or absence of bermudagrass green tissue) 70 d after the first sequential or single topramezone application (equivalent to 28 d after the final third sequential application).^eNontreated control data were not included in the analysis because there was no variance.

Dania muck (Euic, hyperthermic, shallow Lithic Haplosaprists), pH 7.3, and with 85% organic matter. Soil pH and organic matter content were determined using the method described by Fernandez et al. (2019). Bermudagrass was established from sprigs with stolons as described for the greenhouse experiment but using muck soil as the potting medium. The plants were kept in the greenhouse for 14 d before transplanting in the field when plants averaged 12 cm in length with ascending to decumbent stems. The fields were conventionally prepared and treated with atrazine (4.5 kg ha⁻¹) and pendimethalin at (4.3 kg ha⁻¹) applied preemergence prior to bermudagrass transplantation to control other weeds. Plots were subsequently kept free of any other weeds by hand-pulling. The first experimental run was transplanted in the field on March 6, 2018, and the second experimental run on April 2, 2019. The field was irrigated with sprinklers as needed to ensure establishment of the transplanted bermudagrass. The experiment was set up as described to simulate the occurrence of bermudagrass escapes in a plant cane field because it was difficult to obtain a homogenous infestation of new escapes in a field large enough to set up the experiment. Once established, herbicides were applied 21 d after transplanting on April 10, 2018, for the first experimental run, and on May 7, 2019, for the second experimental run. Plants averaged 36 cm in length from the longest ends with spreading mat-forming stolons.

The experiment was a randomized complete block design with four replications. Plots were 2 m by 2 m. One pot was transplanted at the center of each plot. Topramezone was applied at 25 and 50 g ha⁻¹ once or sequentially every 14 d for up to three applications for nine treatments (Table 1). All treatments included methylated seed oil at 10 mL L⁻¹. A nontreated control was included for comparison. Herbicides were applied using a CO₂-pressurized backpack sprayer with a swath width of 1.8 m calibrated to deliver 187 L ha⁻¹ at 276 kPa using TeeJet XR11002VS nozzle tips at a walking speed of 4.8 km h⁻¹.

Visual estimation of bermudagrass control was made using a scale of 0 to 100, with 0 being no control and 100 being complete plant death at 14 d after the final or third sequential topramezone application (equivalent to 42 d after the first sequential or single topramezone application). The binomial response of the presence

or absence of bermudagrass green tissue was recorded 28 d after the final control rating from each plot (equivalent to 70 d after the first sequential or single topramezone application) as 1 or 0, respectively, to determine chances of bermudagrass surviving topramezone sequential treatments.

Topramezone and Mixtures with PS II-Inhibitor Herbicides Greenhouse Experiment

Two greenhouse experiments were conducted at the EREC in 2017 to 2018. Bermudagrass was established from sprigs with stolons as described for the sequential greenhouse experiment using the same potting medium on September 18, 2017, and December 13, 2017, for the first and second experimental runs, respectively. Plants of the same size as described in the sequential greenhouse experiment were treated with herbicides 21 d after planting on October 9, 2017, for the first experimental run, and on January 3, 2018, for the second experimental run.

The experiment was a completely randomized design with five replications of each treatment. Topramezone was applied at 25 and 50 g ha⁻¹, or in mixtures with atrazine at 2,240 g ha⁻¹, ametryn at 440 g ha⁻¹, or metribuzin at 2,240 g ha⁻¹ for a total of eight treatments (Table 2). These are commonly used rates of the herbicides that inhibit PS II used for weed control in Florida sugarcane production (Odero and Dusky 2021). Methylated seed oil at 10 mL L⁻¹ was included in all treatments. A nontreated control was included for comparison. Herbicide treatments were applied as described for the sequential greenhouse experiment.

Visual estimation of bermudagrass control was made using a scale of 0 to 100, with 0 being no control and 100 being complete plant death at 42 d after treatment. Plants were harvested at the soil surface level 42 d after treatment, and aboveground dry weight was determined as described for the sequential greenhouse experiment.

Field Experiment

Two field experiments were conducted at the EREC in 2018 and 2019 at the same location as the sequential field experiment. Bermudagrass was established from sprigs with stolons, transplanted, and maintained in the field as described for the sequential

Table 2. Newly established bermudagrass control and aboveground dry weight in response to topramezone alone or in mixtures with photosystem II inhibitor herbicides (atrazine, metribuzin, ametryn) 42 d after treatment in greenhouse and field experiments combined over two experimental runs in 2017 and 2018 in Belle Glade, FL.^a

Treatment program	Rate	Greenhouse experiment		Field experiment	
		Control	Dry weight	Control	P(survival) ^b
	g ha ⁻¹	%	g	%	%
Nontreated control ^c	–	–	7.00 c	–	100
Topramezone	25	72 b	2.60 b	63 b	100
Topramezone	50	86 ab	0.74 a	79 ab	89
Topramezone + atrazine	25 + 2,240	87 ab	0.89 a	84 a	76
Topramezone + metribuzin	25 + 2,240	90 ab	0.47 a	87 a	63
Topramezone + ametryn	25 + 440	89 ab	0.59 a	79 ab	76
Topramezone + atrazine	50 + 2,240	87 ab	0.57 a	92 a	36
Topramezone + metribuzin	50 + 2,240	100 a	0.18 a	95 a	23
Topramezone + ametryn	50 + 440	97 a	0.26 a	88 a	63

^aMeans followed by the same letter within a column are not significantly different according to Tukey's test ($P < 0.05$).

^bP(survival) is the probability of bermudagrass survival (the presence or absence of bermudagrass green tissue) 70 d after treatment.

^cNontreated control data were not included in the analysis because there was no variance.

field experiment. The first and second experimental runs were transplanted in the field on March 6, 2018, and April 2, 2019, respectively. Herbicides were applied on plants of similar sizes as we did for the sequential field experiment 21 d after transplanting on April 10, 2018, for the first experimental run, and on May 7, 2019, for the second experimental run.

The experiment was a randomized complete block design with four replications. Plots were 2-m by 2-m with one pot transplanted at the center of each plot. Topramezone was applied alone and in mixtures at rates as we did for the greenhouse experiment (Table 2). A nontreated control was included for comparison. Herbicides were applied as described for the sequential field experiment.

Bermudagrass control was visually estimated using a scale of 0 to 100, with 0 being no control and 100 being complete plant death at 42 d after treatment application. The binomial response of presence or absence of bermudagrass green tissue was recorded as 1 or 0, respectively, from each plot 70 d after treatment to determine chances of bermudagrass surviving topramezone and mixture treatments.

Statistical Analysis

Bermudagrass control and aboveground dry weight data for the topramezone sequential and mixtures with PS II-inhibitor herbicides studies were subjected to ANOVA using the LME4 package (Bates et al. 2024) with the R statistical language (v. 4.2.2; R Core Team 2022). The control and aboveground dry weight data were analyzed using a mixed-effects model with the *lmer* function in the LME4 package for both greenhouse and field experiments. Treatment was considered a fixed effect, while experimental run and replication nested within experimental run were considered as random effects. Estimated marginal means were calculated where significant effects were detected, and the post hoc Tukey test was performed for all pairwise comparisons ($P < 0.05$) using the EMMEANS package of R (Lenth 2024). A generalized linear model was used to conduct analysis of deviance for the binomial response data of presence or absence of bermudagrass green tissue for the topramezone sequential and mixtures with PS II-inhibitor herbicides field experiments. Analysis of deviance, analogous to ANOVA, is appropriate for binomial data (Venables and Ripley 2002). The binary outcome of presence or absence of bermudagrass green tissue was analyzed using the *glmer* function with a binomial

family in the LME4 package for both topramezone sequential and mixture field experiments. The probability of having green bermudagrass tissue 70 d after treatment was determined for each observation using the *predict* function in R followed by aggregation of the predicted probabilities by treatment and replication nested within run using the *aggregate* function in R. The mean probabilities of each treatment across replication nested within run was predicted using the *aggregate* function for both topramezone sequential and mixture field experiments.

Results and Discussion

Sequential Topramezone Application Study

The main effect of herbicide treatment was significant for newly established bermudagrass control and aboveground dry weight accumulation in response to sequential topramezone applications for the greenhouse experiment ($P < 0.05$). Bermudagrass control ranged from 78% to complete control 42 d after the first sequential or single application of topramezone (equivalent to 14 d after the final sequential application) (Table 1). A single application of topramezone at 25 g ha⁻¹ controlled bermudagrass less (78%) effectively than topramezone at 50 g ha⁻¹ (89%) and the sequential applications of topramezone (>99%). All sequential applications of topramezone controlled bermudagrass by more than 99% and were more effective than single applications. All topramezone applications reduced bermudagrass aboveground biomass accumulation 42 d after the first sequential or single application compared with that of the nontreated control even though no significant differences between the treatments were observed (Table 1). Similar to the greenhouse experiment, the effect of herbicide treatment was important for controlling newly established bermudagrass in the field ($P < 0.05$). Single or sequential topramezone applications controlled bermudagrass by 61% to 100% 42 d after the first sequential or single application in the field (Table 1). A single application of topramezone at 25 g ha⁻¹ controlled bermudagrass less effectively than the single topramezone application of 50 g ha⁻¹ and sequential applications in the field. Sequential applications of topramezone in the field controlled bermudagrass by 87% to 100%, with no differences between the treatments. Sequential applications of topramezone with a cumulative total of 75 to 100 g ha⁻¹ controlled bermudagrass by >93% in the field. Brewer et al. (2017) predicted that a single

application of topramezone at 36.8 g ha^{-1} would cause a maximum bermudagrass injury of 81%. In comparison, a sequential application of the same rate, applied at a 21-d interval, resulted in a higher injury level of 93%. Topramezone can be applied at 25 to 50 g ha^{-1} on sugarcane with not more than 50 g ha^{-1} per application, and not more than a cumulative total of 100 g ha^{-1} per year (Anonymous 2018). These results indicate that sequential applications of topramezone at 14-d intervals provided greater control of newly established bermudagrass than single applications in greenhouse and field experiments. Topramezone injury to bermudagrass was manifested as bleaching of leaf tissue with most of the bleaching observed 14 d after treatment. Brosnan et al. (2011) reported that sequential applications of topramezone for bermudagrass control should occur at 14- to 21-d intervals because the stress imposed by the herbicide was greatest at 14 and 21 d after treatment when peak visual bleaching occurred. These intervals were ideal because bermudagrass increased its photoprotective xanthophyll cycle pigments, a mechanism that allows it to recover from HPPD-inhibiting herbicide injury 14 and 21 d after treatment, and allowing it to recover by 35 d after treatment (Brosnan et al. 2011). Similarly, Elmore et al. (2011) suggested that sequential applications of topramezone on hybrid bermudagrass should occur at 14- to 21-d intervals based on occurrence of peak visual bleaching and maximum pigment reductions at these intervals.

The effect of single or sequential topramezone applications with respect to bermudagrass probability of survival in the field based on the presence of green tissue 28 d after the final control rating (equivalent to 70 d after the first sequential or single topramezone application) was significant ($P < 0.05$). The probability of bermudagrass survival was higher for single topramezone application treatments than for sequential treatments (Table 1). Bermudagrass treated with single application of topramezone at 25 g ha^{-1} was estimated to have 100% chance of survival 70 d after the first sequential or single application compared with 88% for topramezone at 50 g ha^{-1} and 12% to 75% for sequential applications. Bermudagrass treated with two and three sequential applications of topramezone with a cumulative total of 100 g ha^{-1} had 12% chance of survival compared with two applications with a cumulative total of 75 g ha^{-1} , which had 37% chance of survival. The results show that topramezone sequential applications every 14 d with a cumulative total of 100 g ha^{-1} is necessary to reduce chances of newly established bermudagrass surviving topramezone treatment application as the crop season progresses. The present study was conducted under full sunlight with no shading on bermudagrass, which is atypical in a sugarcane field. Sugarcane canopy closure can be slow, intermediate, or fast depending on the sugarcane cultivar. For example, sugarcane varieties CP 80-1762 and CP 88-1743 have intermediate and fast canopy closures, respectively (Sandhu et al. 2022a, 2022b). Weed growth and development are influenced by the speed of sugarcane canopy closure. Bermudagrass foliar growth and ground cover are negatively affected by shading (Bittencourt 2009; Juraimi et al. 2004). Because shading reduces bermudagrass growth and development, it is likely that the effect of topramezone in combination with shading as sugarcane canopy closes can be important for its long-term control. For fields likely to have bermudagrass escaping fallow management programs, planting sugarcane cultivars that emerge early will generate a high stalk population with fewer upright leaves, and produce shading, which can be important in ensuring increased efficacy of topramezone for management of bermudagrass.

Topramezone and Mixtures with PS II-Inhibitor Herbicides

The main effect of herbicide treatment was significant for newly established bermudagrass control and aboveground dry weight accumulation in response to application of topramezone and mixtures with PS II-inhibitor herbicides for the greenhouse experiment ($P < 0.05$). Bermudagrass control ranged from 72% to 100% at 42 d after treatment (Table 2). Topramezone applied alone at 25 g ha^{-1} provided 72% control at 42 d after treatment compared with 86% control with the highest rate, although these differences were not significant. Mixtures of topramezone at 25 g ha^{-1} with atrazine, metribuzin, and ametryn provided 87%, 90%, and 89% control, respectively, compared with 97%, 100%, and 87% control with topramezone at 50 g ha^{-1} in mixtures with atrazine, metribuzin, and ametryn, respectively. Bermudagrass control was improved by 15% to 18% when mixtures of topramezone at the lower rate were applied with the PS II-inhibitor herbicides compared with 1% to 14% improvement with mixtures of topramezone at the highest rate.

The combination of topramezone (50 g ai ha^{-1}) with metribuzin provided the greatest control of bermudagrass (100%), whereas topramezone (25 g ha^{-1}) plus ametryn provided the least control (87%), although these differences were not significant. All treatments significantly reduced bermudagrass aboveground biomass compared to the nontreated control (Table 2). However, topramezone at 25 g ai ha^{-1} produced significantly greater aboveground biomass accumulation compared with the higher rate and all mixtures. The effect of herbicide treatment was significant for control of newly established bermudagrass in the field ($P < 0.05$). Bermudagrass control ranged from 63% to 79% with topramezone-only treatments compared with 84% to 95% control provided by topramezone mixtures with PS II-inhibitor herbicides 42 d after treatment in the field (Table 2). Mixtures of topramezone at 25 g ha^{-1} with PS II-inhibitor herbicides controlled bermudagrass by 79% to 84% compared with 88% to 95% control provided by mixtures with topramezone at 50 g ha^{-1} .

Bermudagrass control was improved by 16% to 21% with mixtures of topramezone at the lower rate with the triazine herbicides compared with 7% to 13% improvement when mixtures of topramezone were applied at the highest rate in the field. Similar to what occurred in the greenhouse experiment, the improvement in bermudagrass control with PS II-inhibitor herbicides was much more profound when mixtures with topramezone at 25 g ha^{-1} were applied compared with 50 g ha^{-1} . The mixture of 50 g ai ha^{-1} topramezone plus metribuzin provided the greatest bermudagrass control (95%), whereas topramezone (25 g ha^{-1}) plus ametryn offered the least control (79%) even though these differences were not significantly different.

Results from this study indicate that the level of bermudagrass control improved with combinations of topramezone with atrazine, ametryn, and metribuzin. Enhancement of control of several weed species by combining herbicides that inhibit HPPD and PS II has been reported (Armel et al. 2007; Flutterm et al. 2022; Negrisoli et al. 2020). The interactive effects of HPPD-inhibitor and PS II-inhibitor herbicides have mainly been reported to involve atrazine (Flutterm et al. 2022). In this study, bermudagrass control improved with the combination of topramezone with metribuzin and ametryn, with the highest level of control occurring with the combination with metribuzin. Similarly, Negrisoli et al. (2020) reported improved control of fall panicum (*Panicum dichotomiflorum* Michx.) in sugarcane with the mixture of topramezone and metribuzin.

The effect of topramezone and mixtures with PS II-inhibitor herbicides regarding probability of survival of newly established bermudagrass in the field based on the presence of green tissue at 70 d after treatment application was significant ($P < 0.05$). Like the topramezone sequential study, the probability of bermudagrass survival was higher with single topramezone applications compared to mixtures with PS II-inhibitor herbicides (Table 2). Bermudagrass treated with topramezone alone at 25 g ha⁻¹ was estimated to have 100% chance of survival compared with 89% for topramezone at 50 g ha⁻¹, and 23% to 76% for mixtures with PS II-inhibitor herbicides. For mixtures of the PS II-inhibitor herbicides with topramezone at 25 g ha⁻¹, bermudagrass had 63% to 76% chance of survival compared with 23% to 63% with the mixtures with topramezone at 50 g ha⁻¹. Bermudagrass treated with topramezone at 50 g ha⁻¹ in mixtures with metribuzin and atrazine had the lowest chances of survival at 23% and 36%, respectively. Like the sequential study, the effect of topramezone and mixtures with PS II-inhibitor herbicides might probably have been more profound under shading because of the negative effect of shading on growth and development of bermudagrass (Bittencourt 2009; Juraimi et al. 2004).

Results from this study indicated that topramezone applied sequentially or in mixtures with PS II-inhibitor herbicides atrazine, ametryn, and metribuzin provided acceptable control of newly established bermudagrass from sprigs with stolons. Sequential applications with a cumulative total of 75 to 100 g ha⁻¹ provided acceptable bermudagrass control in the field, and applications with a cumulative total of 100 g ha⁻¹ had a 12% chance of survival at 70 d after the first sequential treatment. The mixture of topramezone (50 g ai ha⁻¹) plus metribuzin provided the greatest bermudagrass control, with a 23% chance of survival 70 d after treatment. The use of sugarcane weed management programs involving sequential topramezone applications or mixtures with PS II-inhibiting herbicides in plant cane can provide growers with options for reducing the progressive increment of newly established bermudagrass in the sugarcane crop cycle. Such bermudagrass infestations escape control programs during sugarcane the fallow management period or in successively planted sugarcane.

Practical Implications

Bermudagrass is a difficult-to-control perennial grass weed in Florida sugarcane fields. Sugarcane is a multiyear crop in which tillage is limited to the row middles, thus allowing bermudagrass to establish over the rows and become difficult to control during the entire crop cycle because of lack of selective postemergence herbicides. Bermudagrass is only effectively managed after the final ratoon crop of sugarcane has been harvested using a combination of multiple glyphosate applications and tillage during the fallow period. However, bermudagrass, which escapes fallow management programs or when sugarcane is successively replanted, can re-establish from viable stolons and rhizomes. Topramezone is a postemergence herbicide that can be safely used on sugarcane for control of broadleaf and grass weeds. It can be used to manage newly established bermudagrass escapes in plant cane. Two or three sequential applications of topramezone every 14 d provide acceptable control of newly established bermudagrass. Mixtures of topramezone with PS II-inhibitor herbicides, particularly metribuzin and atrazine, commonly used in sugarcane production, improved the control of newly established bermudagrass. These findings illustrate that sugarcane growers can mitigate the adverse effects of newly established bermudagrass in plant cane using

sequential applications of topramezone and in mixtures with metribuzin and atrazine. Topramezone can be applied to achieve acceptable management of bermudagrass, and in mixtures with other herbicides it can be used to manage other problematic broadleaf and annual grass weeds in sugarcane.

Acknowledgments. We thank Leonard Fox from the University of Florida Everglades Research and Education Center for his help with the project.

Funding. BASF Corporation provided partial funding the project.

Competing Interests. The authors declare they have no competing interests.

References

- Anonymous (2018) Armezon herbicide for broadleaf and grass control in sugarcane. APN# 1809002 Armezon-Sugarcane-FL-2018. Florham Park, NJ: BASF Corporation
- Armell GR, Rardon PL, McComrick MC, Ferry NM (2007) Differential response of several carotenoid biosynthesis inhibitors in mixtures with atrazine. *Weed Technol* 21:947–953
- Bates D, Maechler M, Bolker B, Walker S (2024) lme4: Linear Mixed-Effects Models using 'Eigen' and S4. Version 1.1-35.3. <https://cran.r-project.org/web/packages/lme4/>. Accessed: May 25, 2024
- Bittencourt MA (2009) Seasonal growth and shading potential of sugarcane (*Saccharum* spp. hybrids) and shade response of perennial weeds [master's thesis]. Baton Rouge: Louisiana State University. 48 p
- Brewer JR, Willis J, Rana SS, Askew SD (2017) Response of six turfgrass species and four weeds to three HPPD-inhibiting herbicides. *Agron J* 109:1777–1784
- Brosnan JT, Kopsell D A, Elmore MT, Breeden GK (2011) Changes in 'Riviera' bermudagrass [*Cynodon dactylon* (L.) Pers.] carotenoid pigments after treatment with three p-hydroxyphenylpyruvate dioxygenase-inhibiting herbicides. *HortScience* 46:493–498
- Elmore MT, Brosnan JT, Kopsell DA, Mueller TC (2011) Response of hybrid bermudagrass (*Cynodon dactylon* × *C. transvaalensis*) to three HPPD inhibitors. *Weed Sci* 59:458–463
- Etheredge LM Jr, Griffin JL, Salassi ME (2009) Efficacy and economics of summer fallow conventional and reduced-tillage programs for sugarcane. *Weed Technol* 23:274–279
- Fernandez JV, Odero DC, MacDonald GE, Ferrell JA, Sellers BA, Wilson PC (2019) Field dissipation of S-metolachlor in organic and mineral soils used for sugarcane production in Florida. *Weed Technol* 34:362–370
- Fernandez ON (2003) Establishment of *Cynodon dactylon* from stolon and rhizome fragments. *Weed Res* 43:130–138
- Flutter JC, Soltani N, Galla M, Hooker DC, Robinson DE, Sikkema PH (2022) Additive and synergistic interactions of 4-hydroxyphenylpyruvate dioxygenase (HPPD) and photosystem II (PSII) inhibitors for the control of glyphosate-resistant horseweed (*Conyza canadensis*) in corn. *Weed Sci* 70:319–327
- Grossmann K, Ehrhardt T (2007) On the mechanism and selectivity of the corn herbicide topramezone: a new inhibitor of 4-hydroxyphenylpyruvate dioxygenase. *Pest Manag Sci* 63:429–439
- Holm LG, Plucknett DL, Pancho JV, Herberger JP (1977) *The World's Worst Weeds: Distribution and Biology*. Honolulu: University Press of Hawaii. 610 p
- Jhala AJ, Kumar V, Yadav R, Jha P, Jugulam M, Williams MM II, Hausman NE, Dayan FE, Burton PM, Dale RP, Norsworthy JK (2023) 4-hydroxyphenylpyruvate dioxygenase (HPPD)-inhibiting herbicides: past, present, and future. *Weed Technol* 37:1–14
- Juraimi AS, Drennan DSH, Anuar N (2004) The effects of shading on the growth, development and partitioning of biomass in bermudagrass (*Cynodon dactylon* (L.) Pers.). *J Biol Sci* 4:756–762
- Lenth RV (2024) emmeans: Estimated marginal means, aka least-squares means. Version 1.10.2. <https://cran.r-project.org/web/packages/emmeans/index.html>. Accessed: May 25, 2024
- Miller DK, Griffin JL, Richard EP Jr (1999) Summer fallow and after-planting bermudagrass (*Cynodon dactylon*) control programs for sugarcane (*Saccharum* spp. hybrids). *Weed Technol* 13:127–131
- Mitich LW (1989) Bermudagrass. *Weed Technol* 3:433–435

- Negrisola RM, Otero DC, MacDonald GE, Sellers BA, Laughinghouse HD (2020) Sugarcane response and fall panicum (*Panicum dichotomiflorum*) control with topramezone and triazine herbicides. *Weed Technol* 34: 241–249
- Leon RG, Otero DC (2018) Weed management in sugarcane. Pages 455–469 in Korres NE, Burgos NR, Duke SO, eds. *Weed Control: Sustainability, Hazards, and Risks in Cropping Systems Worldwide*. 1st ed. Boca Raton, FL: CRC Press/Taylor & Francis Group
- Otero DC, Beuzelin JM, VanWeelden M, Kammerer CL, Raid RN, Swanson S, Rott P, Mossler M (2023) Florida crop/pest profile: sugarcane. *Electronic Data Information Sources PI-171*. Gainesville: University of Florida Institute of Food and Agricultural Science, Florida Cooperative Extension.
- Otero DC, Dusky JA (2021) Weed management in sugarcane. *Electronic Data Information Sources SS-AGR-09*. Gainesville: University of Florida Institute of Food and Agricultural Science, Florida Cooperative Extension
- R Core Team (2022) R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. <https://www.Rproject.org/>. Accessed: April 23, 2024
- Richard EP Jr (1992) Bermudagrass interference during a three-year sugarcane crop cycle. *Proc Int Soc Sugar Cane Technol* 21:31–41
- Richard EP Jr (1993) Preemergence herbicide effects on bermudagrass (*Cynodon dactylon*) interference in sugarcane (*Saccharum* spp. hybrids). *Weed Technol* 7:578–584
- Richard EP Jr (1997) Effects of fallow bermudagrass (*Cynodon dactylon*) control programs on newly planted sugarcane (*Saccharum* spp. hybrids). *Weed Technol* 11:677–682
- Richard EP Jr, Dalley CD (2007) Sugarcane response to bermudagrass interference. *Weed Technol* 21:941–946
- Sandhu HS, Singh MP, Gilber RA (2022a) Sugarcane cultivar CP 88-1743. *Electronic Data Information Sources SS-AGR-118*. Gainesville: University of Florida Institute of Food and Agricultural Science, Florida Cooperative Extension
- Sandhu HS, Singh MP, Gilber RA (2022b) Sugarcane cultivar CP 88-1762. *Electronic Data Information Sources SS-AGR-121*. Gainesville: University of Florida Institute of Food and Agricultural Science, Florida Cooperative Extension.
- Sandoya-Miranda G, Kanissery R, Dufault NS, Desaegeer J, Meszaros A, Beuzelin J, Xavier KT (2024) Leavy vegetable production. Pages 187–232 in Dittmar P, Agehara S, Dufault NS, eds. *Vegetable Production Handbook of Florida*. Gainesville: University of Florida Institute of Food and Agricultural Science, Florida Cooperative Extension
- Shaner DL, ed (2014) Pages 449–450 in *Herbicide Handbook*. 10th ed. Lawrence, KS: Weed Science Society of America
- VanWeelden M, Kammerer C, Davidson W, Baltazar M, Tice R (2024) Sugarcane variety census: Florida 2023. *Sugar J* 87:6–11
- Venables WN, Ripley BD (2002) *Modern Applied Statistics with S*. 4th ed. New York: Springer. 495 p
- Webster TM (2012) Weed survey - southern states: grass crops subsection. *Proc South Weed Sci Soc* 65:267–288
- [USDA-NASS] U.S. Department of Agriculture–National Agricultural Statistics Service. (2024) Acreage. <https://usda.library.cornell.edu/concern/publications/j098zb09z>. Accessed May 9, 2024