- odeficiency virus transmission in household settings—United States. *MMWR* 1994;43:347,353-356.
- Centers for Disease Control. Recommendations for prevention of HIV transmission in health-care settings. MMWR 1987;36(suppl 2S):1S-8S.
- Hussain SA, Latid ABA, Choudhary AAAA. Risk of surgeons: a survey of accidental injuries during operations. Br J Surg 1988;75:314-316.
- Gerberding JL, Littell C, Tarkington A, Brown A, Schecter WP. Risk of exposure of surgical personnel to patients' blood during surgery at San Francisco General Hospital. N Engl J Med 1990;322:1788-1793.
- 8. Quebbeman EJ, Telford GL, Hubbard S, et al. Risk of blood contamination and injury to operating room personnel. *Ann Surg* 1991;214:614-620.
- 9. Panlilio AL, Foy DR, Edwards JR, et al. Blood contacts during surgical procedures. *JAMA* 1991;265(12):1533-1537.
- Popejoy SL, Fry DE. Blood contact and exposure in the operating room. Surg Gynecol Obstet 1991;172:480-483.
- White MC, Lynch P. Blood contact and exposures among operating room personnel: a multicenter study. Am J Infect Control 1993;21:243-248.
- Tokars JI, Bell DM, Marcus R, et al. Percutaneous injuries during surgical procedures. JAMA 1992;267:2899-2904.
- Panlilio AL, Welch BA, Bell DM, et al. Blood and amniotic fluid contact sustained by obstetric personnel during deliveries. Am J Obstet Gynecol 1992;167:703-708.
- Tokars JI, Chamberland ME, Schable CA, et al. A survey of occupational blood contact and HIV infection among orthopedic surgeons. *IAMA* 1992;268(4):489-494.
- Cohn GM, Seifer DB. Blood exposure in single versus double gloving during pelvic surgery. Am J Obstet Gynecol 1990;162:715-717.

- 16. Telford GL, Quebbeman EJ. Assessing the risk of blood exposure in the operating room. Am J Infect Control 1993;21:351-356.
- 17. Quebbeman EJ, Telford GJ, Wadsworth K, Hubbard S, Goodman H, Gottlieb MS. Double gloving: protecting surgeons from blood contamination in the operating room. *Arch Surg* 1992;127:213-217.
- Matta H, Thompson AM, Rainey JB. Does wearing two pairs of gloves protect operating theatre staff from skin contamination? Br Med J 1988;297:597-598.
- McLeod GG. Needlestick injuries at operations for trauma. J Bone Joint Surg Br 1989;71-B(3):489-491.
- Cole RP, Gault DT. Glove perforation during plastic surgery. Br J Plast Surg 1989;42:481-483.
- Dodds RDA, Barker SGE, Morgan NH, Donaldson DR, Thomas MH. Self protection in surgery: the use of double gloves. Br J Surg 1990;77:219-220.
- Gani JS, Anseline PF, Bissett RL. Efficacy of double versus single gloving in protecting the operating team. Aust N Z J Surg 1990;60:171-175.
- 23. Bennett B, Duff P. The effect of double gloving on frequency of glove perforations. *Obstet Gynecol* 1991;78:1019-1022.
- Doyle PM, Alvi S, Johanson R. The effectiveness of double gloving in obstetrics and gynaecology. Br J Obstet Gynaecol 1992;99:83-84.
- Chapman S, Duff P. Frequency of glove perforations and subsequent blood contact in association with selected obstetric surgical procedures. Am J Obstet Gynecol 1993;168:1354-1357.
- Rose DA, Ramiro NZ, Perlman JL, Schecter WP, Gerberding JL.
   Two gloves or not two gloves? Intraoperative glove utilization at San Francisco General Hospital. *Infect Control Hosp Epidemiol* 1994;15:(4,suppl)P37.
- 27. Webb JM, Pentlow BD. Double gloving and surgical technique. *Ann R Coll Surg Engl* 1993;75:291-292.

## Microbe Names—Are You Keeping Up?

## by Gina Pugliese, RN, MS Medical News Editor

In a recent article in *Infectious Disease Alert*, Dr. Stan Deresinski highlighted some of the changes in nomenclature, taxonomy, and classification of pathogenic organisms. By now, most know that enterococci are no longer Group D streptococci and that the former *Neisseria catarrhalis* briefly was known as *Branhamella* 

catarrhalis and now has become Moraxella catarrhalis. Other pathogenic organisms, he notes, that recently have changed names include Mycobacterium mucogenicum (formerly Mycobacterium chelonae-like organism or MCLO); Prevotella melaninogenica (formerly Bacteroides melaninogenicus); Burkholderia cepacia (formerly Pseudomonas cepacia); Chrysobacterium meningosepticum (formerly Flavobacterium meningo-

septicum); and Stenotrophomonas maltophilia (formerly Xanthomonas maltophilia). Dr. Deresinski points out that although these changes may drive us crazy, they may help to maintain our value as knowledge brokers.

From: Deresinski S. Are you keeping up with the name game? *Infectious Disease Alert*. October 15, 1995:14-15.