

# High spatial resolution observations of OH 231.8+4.2

M. Matsuura<sup>1,2,3</sup>, O. Chesneau<sup>4</sup>, A. A. Zijlstra<sup>1</sup>, W. Jaffe<sup>5</sup>, L. B. F. M. Waters<sup>6,7</sup>, J. A. Yates<sup>8</sup>, E. Lagadec<sup>1</sup>, and T. M. Gledhill<sup>9</sup>

<sup>1</sup>University of Manchester, Sackville Street, P.O. Box 88, Manchester M60 1QD, UK

<sup>2</sup>Queen's University of Belfast, Belfast BT7 1NN, Northern Ireland, UK

<sup>3</sup>National Astronomical Observatory of Japan, Tokyo 181-8588, Japan  
email: mikako@optik.mtk.nao.ac.jp

<sup>4</sup>Observatoire de la Côte d'Azur, Department of Gemini-CNRS-UMR 6203,  
F-06130, Grasse, France

<sup>5</sup>Leiden Observatory, P.B. 9513, Leiden 2300 RA, The Netherlands

<sup>6</sup>University of Amsterdam, Kruislaan 403, 1098 SJ, Amsterdam, The Netherlands

<sup>7</sup>Katholieke Universiteit Leuven, Celestijnenlaan 200B, 3001 Heverlee, Belgium

<sup>8</sup>University College London, Gower street, London, WC1E 6BT, United Kingdom

<sup>9</sup>Centre for Astrophysics Research, University of Hertfordshire, Hertfordshire AL10 9AB, UK

**Abstract.** We have observed the bipolar post-AGB candidate OH 231.8+4.2, using the mid-infrared interferometer MIDI and the infrared camera with the adaptive optics system NACO on the Very Large Telescope. The NACO images at 2.12 and 3.8  $\mu\text{m}$  show a bipolar outflow and a flared disk or torus. An unresolved core ( $<200$  mas in FWHM) is found at the centre of OH 231.8+4.2 in the 3.8  $\mu\text{m}$  image. This compact source is resolved with the interferometer. The fringes from the four baselines consistently show the presence of a compact circumstellar material with an inner radius of 30–40 mas, which is equivalent to 40–50 AU at 1.3 kpc. This clearly shows that the mid-infrared compact source is not the central star (3 AU) but circumstellar material.


**Keywords.** stars: AGB and post-AGB, (stars:) circumstellar matter, infrared: stars, infrared: ISM, stars: imaging stars: mass loss

---


Among post-AGB stars and planetary nebulae, a high fraction of stars show asymmetric circumstellar envelopes, while typical AGB winds are spherical symmetric. The most extreme shape of a nebula is a bipolar one. One of the hypotheses about the formation of the bipolar shape invokes a binary-disk scenario (Balick & Adams 2002). Part of the material lost during the intensive AGB wind is trapped in the binary system, and a disk is formed around the companion or around the binary system. The disk restricts the (low density but high velocity) post-AGB and PN wind in the equatorial plane, and focusses the wind towards the two poles.

OH 231.8+4.2 (hereafter OH 231) is the best-known bipolar post-AGB candidate. It is probably associated with the open cluster, M 46 (Jura & Morris 1985), so that its distance is relatively well determined (1.3 kpc), and also the initial mass (about  $3 M_{\odot}$ ).

To search for evidence of a compact, constraining disk, we observed OH 231 with the Very Large Telescope (VLT), using both the infrared adaptive optics cameras NACO and the VLTI mid-infrared interferometre MIDI. MIDI was used on 2005 March 2nd, using the telescopes Melipal (UT3) and Yepun (UT4). We obtained four baselines with a difference of 21 degree position angle, and with a length range of 47–62 meters. NACO observations were obtained on 2004 March 6th. Narrow band (NB) 2.12 and L'-band filters are used, both measuring continuum emission. Details of the observations are described in Matsuura *et al.* (2006).



**Figure 1.** NACO  $16.2 \times 23.5$  arcsec<sup>2</sup> images (a: NB-band and b: L'-band). Lines in image (b) show the approximate baseline position angle for MIDI observations (OB1 and OB4). Insert (c) shows the brightest region of L-band image in a different colour scale so as to clearly show the unresolved compact object inside. North is up and east is to the left.



**Figure 2.** Left: correlated flux of the mid-infrared ‘core’ (unresolved source in NACO image). The vertical axis gives the radius needed to reproduce the visibilities.

Fig. 1 shows the NACO NB2.12 and L'-band images of OH 231. The central region is still obscured in the NB2.12 image and the bipolar outflow is brighter. On the other hand, in the L'-band the central region is brighter than the outflow. The central region has a very red color, indicating obscuration by a very high optical depth. There is an unresolved core in the centre of the L'-band image with a size  $< 200$  mas (FWHM).

This compact core produced fringes with the interferometer. The left panel in Fig. 2 shows the correlated flux obtained with the MIDI. The visibility data are interpreted assuming a smooth Gaussian profile, as described by Leinert *et al.* (2004). Fig. 2 shows the radii (HWHM) of the Gaussians which reproduce the observed visibilities. These are typically 30–40 mas. At the distance of 1.3 kpc, the radii are about 40–50 AU, as displayed on the right side of the y-axis in the right panel Fig. 2. The mid-infrared compact source is not the central star (3 AU) but is circumstellar. In Matsuura *et al.* (2006) we discuss the possibility that this represents the confining disk.

## References

- Balick, B., & Frank, A. 2002, *ARA&A* 40, 439  
 Jura, M., & Morris, M. 1985, *ApJ*, 292, 487  
 Leinert Ch., van Boekel R., Waters B.F.M.M. *et al.* 2004, *A&A* 423, 537  
 Matsuura, M., Chesneau, O., Zijlstra, A.A. *et al.* 2006, *ApJ*, 646, L123