




## Regular Article

# Let's face it! The role of social anxiety and executive functions in recognizing others' emotions from faces: Evidence from autism and specific learning disorders

Rachele Lievore<sup>1</sup> , Ramona Cardillo<sup>1,2</sup>  and Irene C. Mammarella<sup>1</sup> 

<sup>1</sup>Department of Developmental and Social Psychology, University of Padua, Padua, Italy and <sup>2</sup>Department of Women's and Children's Health, University of Padua, Padua, Italy

### Abstract

Youth with different developmental disorders might experience challenges when dealing with facial emotion recognition (FER). By comparing FER and related emotional and cognitive factors across developmental disorders, researchers can gain a better understanding of challenges and strengths associated with each condition. The aim of the present study was to investigate how social anxiety and executive functioning might underlie FER in youth with and without autism spectrum disorders (ASD) and specific learning disorders (SLD). The study involved 263 children and adolescents between 8 and 16 years old divided into three groups matched for age, sex, and IQ: 60 (52 M) with ASD without intellectual disability, 63 (44 M) with SLD, and 140 (105 M) non-diagnosed. Participants completed an FER test, three executive functions' tasks (inhibition, updating, and set-shifting), and parents filled in a questionnaire reporting their children's social anxiety. Our results suggest that better FER was consistent with higher social anxiety and better updating skills in ASD, while with lower social anxiety in SLD. Clinical practice should focus on coping strategies in autistic youth who could feel anxiety when facing social cues, and on self-efficacy and social worries in SLD. Executive functioning should also be addressed to support social learning in autism.

**Keywords:** autism spectrum disorders; executive functions; facial emotion recognition; social anxiety; specific learning disorders

(Received 31 July 2023; revised 12 November 2023; accepted 5 January 2024; First Published online 8 February 2024)

### Introduction

Recognizing others' emotional states from their facial expressions is a crucial ability in social functioning. Individuals can derive a considerable amount of information through nonverbal facial cues conveying signals to others about thoughts, feelings, attitudes, and intentions (Ekman & Friesen, 2003; Ekman, 1993). Indeed, facial emotion recognition (FER) is necessary for adapting one's own social actions and feedbacks, with the aim of creating worthwhile social interactions (Fischer et al., 2019). In this sense, understanding emotional manifestations is a prerequisite for responding appropriately to other people's assumed internal state, whether it means offering help, or simply an acknowledgement, or even a disapproval (Van Kleef, 2009). However, youth with different developmental disorders, such as autism spectrum disorder (ASD) and specific learning disorders (SLD), might experience challenges when dealing with emotion recognition, with specific emotional and cognitive factors associated with such difficulties. By comparing FER and associated factors across various neurodevelopmental disorders, researchers can gain a better understanding of the specific challenges and strengths associated with each condition.

Specifically, evidence suggests that social difficulties experienced by autistic people might be partly due to a reduced social reciprocity, mainly conveyed by their weakness to detect other people's emotional and mental expressions (Baron-Cohen et al., 2009; Chiu et al., 2022). Challenges in FER are consistently associated with autism spectrum disorder (ASD) (for a meta-analysis see Lozier et al., 2014), an early onset neurodevelopmental condition characterized by deficits in social communication and social interaction alongside stereotypic, repetitive, restricted behaviors and interests (American Psychiatric Association, 2013). However, the overall picture of FER in autism is inconsistent, and complicated by the substantial variability between the huge amount of studies in task features, and participants' characteristics, which could count as potential moderating factors contributing to the mixed findings in the literature (for systematic reviews and meta-analyses, see Black et al., 2017; Harms et al., 2010; Leung et al., 2022; Lozier et al., 2014; Uljarevic & Hamilton, 2013; Yeung, 2022). Some studies have identified a generalized face-emotion recognition deficit across all emotional facial expressions in autistic children (Lindner & Rosén, 2006; Lozier et al., 2014; Rump et al., 2009), whereas others revealed a lower performance for one or a subset of expressions, highlighting emotion-specific challenges mainly for negative emotions, which have variously included anger, fear, disgust, sadness, and surprise (see for example, Ashwin et al., 2007; Economides et al., 2020; Humphreys et al., 2007). Moreover,

**Corresponding author:** R. Lievore; Email: [rachele.lievore@studenti.unipd.it](mailto:rachele.lievore@studenti.unipd.it)

**Cite this article:** Lievore, R., Cardillo, R., & Mammarella, I. C. (2025). Let's face it! The role of social anxiety and executive functions in recognizing others' emotions from faces: Evidence from autism and specific learning disorders. *Development and Psychopathology* 37: 451–463, <https://doi.org/10.1017/S0954579424000038>



challenges with emotion recognition in ASD might also vary based on the type of measure (i.e., accuracy vs reaction times), task (i.e., verbal vs non-verbal), emotion complexity (i.e., basic vs complex emotions), processing (i.e., holistic vs featural), and stimuli (i.e., faces vs speech prosody) (for reviews and meta-analyses see, Leung *et al.*, 2022; Lievore *et al.*, 2023; Yeung, 2022). Overall, the large variability of findings on the topic denotes a literature gap which is incapable of establishing a clear framework; for this reason, it is essential to increase the evidence-based results on the study of FER in autism, by also embracing possible associated mechanisms.

While it appears that FER is an area of significant challenge for children with ASD, questions have arisen encompassing those with SLD. People with SLD show difficulties in reading, written expression, and/or mathematics, which have to be substantially and quantifiably below the levels expected for the individual's chronological age and may affect academic achievement or daily functioning (American Psychiatric Association, 2013). Academic achievement has been discovered to be intimately linked to social adaptive functioning (Brigman *et al.*, 2007; Sung & Chang, 2010). In particular, the model of social-emotional learning emphasizes that learning in school is a profoundly social process (Denham & Brown, 2010). About that, longitudinal research confirmed that social skills are important predictors of future school and work success (Whiting & Robinson, 2001). An extensive body of literature has linked social behavior to school achievement, highlighting how early adaptive social skills of children might provide the foundation for both positive attitudes toward learning and a constructive classroom atmosphere (Brigman *et al.*, 2007; Sung & Chang, 2010).

Some authors have explored whether children with SLD might face challenges with recognizing others' emotions. Despite the limited number of research in this area, most studies revealed that individuals with SLD were more inaccurate and slower in detecting facial emotions as compared to non-diagnosed participants (Albayrak *et al.*, 2022; Bloom & Heath, 2010; Ciray & Turan, 2022; Economides *et al.*, 2020; Operto *et al.*, 2020). Specific-emotion difficulties (e.g., for angry faces; Ciray & Turan, 2022) emerge as well as a generalized challenge, with the tendency to assign some emotional states to neutral expressions (Operto *et al.*, 2020). Some authors concluded that problems with FER might be specifically related to high-level visual deficit (Economides *et al.*, 2020); people with dyslexia, a subtype of SLD that impacts reading ability, matched faces more slowly, especially when the faces differed in viewpoint, and discriminated more poorly between similar faces (but not cars) (Gabay *et al.*, 2017; Sigurdardottir *et al.*, 2015; Tarkiainen *et al.*, 2003). However, research is still needed to disentangle the possible mechanisms that underlie challenges with others' emotion detection from facial expressions in children with SLD.

### Facial emotion recognition and social anxiety

When facing others' emotions within social interactions, some people might feel inadequate or uncomfortable, resulting in tension or anxiety (Heimberg *et al.*, 2014). From this point of view, anxiety is not necessarily negatively conceptualized, and under certain circumstances, may serve as useful for navigating social situations (McNeil & Randall, 2014). Sometimes instead, anxiety might reach the disabling extremity of the continuum, preventing people to maintain close social relationships. This feeling is usually named social anxiety, which is characterized by cognitive, emotional, behavioral, and physiological symptoms before or during social or performance situations, concerning the fear to be

exposed to possible scrutiny by others, to be negatively evaluated and a tendency for avoiding interactions (American Psychiatric Association, 2013; Lievore *et al.*, 2022). A recent social information processing model of childhood social anxiety development (Nikolić, 2020), postulates cognitive biases when encoding social stimuli, such as facial emotional expressions. Feelings of rejection can derive from a facial expression perceived as hostile or when misinterpreted, and might consequently provoke anxiety in people for whom approval is meaningful (for a meta-analysis see O'Toole *et al.*, 2013). Findings are inconsistent on the types of other people's emotions that can trigger the development of social anxiety. Some do support the idea that effects are not emotion-specific (Bell *et al.*, 2011; Button *et al.*, 2013; Silvia *et al.*, 2006), whereas others have proven that people with social anxiety recognize less accurately or are more sensitive to certain emotions (Battaglia *et al.*, 2004; Richards *et al.*, 2002; Simonian *et al.*, 2001; Tseng *et al.*, 2017). In addition, an attentional bias toward threatening facial expressions has been proven in socially anxious children and adults (Bantini *et al.*, 2016; Dudeney *et al.*, 2015; Waters *et al.*, 2011). However, the relationship between face emotion recognition and social anxiety focusing on neurodevelopmental disorders has not been investigated yet.

The experience of negative emotions, such as fear or anxiety, within the social contexts is common in most children and adolescents diagnosed with different neurodevelopmental disorders, such as ASD and SLD (Harms *et al.*, 2010; Lievore *et al.*, 2022; Rostami *et al.*, 2014; Terras *et al.*, 2009). Some researchers have proposed a bidirectional relationship between social challenges and social anxiety in ASD (Bellini, 2006; White *et al.*, 2010). The presence of social anxiety, together with previous negative social experiences, could prevent people from engaging in social situations, reducing the opportunity to practice social skills and to empower social efficacy. Therefore, autistic individuals might have increased social worries as a consequence of the self-awareness of their social problems (internal sources of threat, for example, making a self-identified mistake in social situations) and of rejections or negative judgment from others (external sources of threat) (Rosen & Lerner, 2018; Spain *et al.*, 2018). In this regard, for some autistic individuals, faces may not be perceived as emotionally interesting causing indifference toward social interactions (Ashwin *et al.*, 2007), while for others, faces may be overstimulating, and possibly associated with higher levels of social anxiety (Kleinmans *et al.*, 2010). However, the relationship between the ability to recognize facial emotions and levels of social anxiety in children and adolescents with ASD has not been extensively investigated yet. The few studies on the topic revealed that social anxiety might be related to poorer FER in young adults and adolescents with ASD (Antezana *et al.*, 2023), and to poorer fear recognition in autistic adults (Corden *et al.*, 2008). Research on social attention in autism revealed that self-reported fears of negative evaluation predict greater gaze duration to social threat stimuli (White *et al.*, 2015), and that autonomic reactivity to eye gaze negatively correlates with face recognition performance in autistic children (Bal *et al.*, 2010; Joseph *et al.*, 2008). Other studies found no association between behavioral ratings of social anxiety and emotion recognition performance across 7–13 years old autistic children (Wong *et al.*, 2012) and adults (Spain *et al.*, 2016).

In respect of children with SLD, social anxiety might be driven by specific challenges associated with their academic impairments. Children with SLD might face a unique set of socioemotional challenges as a result of their academic difficulties (Livingston *et al.*, 2018). Specifically, to social anxiety, different studies have

revealed a higher rate in children and adults with SLD compared to non-diagnosed (Carroll & Iles, 2006; Mammarella et al., 2016; Rostami et al., 2014; Terras et al., 2009). Consequently, to school challenges and negative experiences, children with SLD might experience feelings of anxiety, shame, and inadequacy (Elgendi et al., 2021), and any negative feedback from school may affect their socioemotional functioning furthering internalizing problems (Donolato et al., 2022; Sahoo et al., 2015; Thaler et al., 2010). Multiple experiences of these feelings can result in lowering self-esteem, social skills deficits with peers, causing loneliness and troubles finding friends or relationships, and resulting in higher levels of social anxiety (Kohli et al., 2005). Though, the association between social skills, such as FER, and social anxiety in children and adolescents with SLD, has not been systematically examined yet.

### Facial emotion recognition and executive functions

Besides underpinning emotional mechanisms, there is accumulating evidence that social development and emotion knowledge, might be supported by higher-order cognitive skills, such as executive functions (EFs). EFs refer to an umbrella term that includes a set of top-down mental processes, crucial for controlling, monitoring and planning thoughts and behaviors, as well as problem-solving, self-controlling, considering alternatives and flexibly adjusting to new settings (Diamond, 2013; Miyake et al., 2000). According to Miyake et al. (2000), the three core processes of EFs are inhibitory control (i.e., the ability to deliberately inhibit automatic responses when required), updating (i.e., the ability to update and monitor information in the working memory), and cognitive flexibility (i.e., set shifting, the ability to switch between mental sets to adjust to changed priorities). Altogether EFs help us to reach goal-directed behaviors, and are therefore essential for success in school and work situations, as well as everyday living. A widely cited biopsychosocial model (SOCIAL; Beauchamp & Anderson, 2010) combines the biological underpinnings and socio-cognitive skills that underlie the social functioning (EFs, communication, socio-emotional skills), as well as the internal and external aspects that mediate these skills. A developing body of literature has indeed underlined the relation between EFs and social abilities (Denham et al., 2012; Moriguchi, 2014; Spikman et al., 2013). EFs might serve as basis for the development of emotional understanding, since attentional and inhibitory control may be required for children to correctly behave in social contexts, and practising with social skills (Carlson & Wang, 2007; Lecce et al., 2017). In particular, cognitive flexibility showed a consistent positive correlation with FER in various populations, given that the ability to handle with complex emotional scripts might be supported by EFs (David et al., 2014; Lee et al., 2009; Martins et al., 2016). Moreover, working memory might play a role in the recognition of other persons' emotions, because of the need to process the emotional cue while temporarily holding it in mind (Neumann et al., 2021; Phillips et al., 2008).

Impairments in executive functioning have been described in different neurodevelopmental disorders, including autism and SLD (Brosnan et al., 2002; Hill, 2004). In autism, the "executive dysfunction" theory has been put forward to try to describe autistic symptoms (Hill, 2004; Hughes, 2001; Ozonoff, 1998). Impairments in EFs have been proposed as a cause of not only the rigid and repetitive interests and behaviors in ASD, but also of the primary challenges with reciprocal social interactions. However, there is a multitude of conflicting evidence on executive dysfunction in ASD,

indicating the inconsistency of EF deficit across subdomains, thus supporting heterogeneity in EFs in this clinical population (Dajani et al., 2016). In this regard, sometimes autistic people display prominent challenges with EFs in everyday life, while amply performing on structured neuropsychological tests (Kenworthy et al., 2008). The association between the ability to recognize others' emotions and EFs has been quite investigated in children with ASD (Joseph & Tager-Flusberg, 2004; Sivaratnam et al., 2018; Torske et al., 2018), postulating a link between the domains.

Concerning children with SLD, challenges with EFs have been proven in many studies, in particular vis-à-vis working memory (Meltzer, 2011; Swanson & Sachse-Lee, 2001). Due to the evident difficulties of children with SLD in working memory, research has focused at most on this area of EFs. Verbal working memory seems to be linked to reading competences (Schuchardt et al., 2008), whereas visuospatial working memory to math achievement (Caviola et al., 2020; Giofrè et al., 2014; Mammarella et al., 2018). In a recent study, however, children with dyslexia were able to similarly perform an updating task to the control group (Artuso et al., 2021). Regarding other EFs' components, findings are inconsistent. Some supported the idea of an executive impairment when assessing set-shifting and response inhibition in SLD (Crisci et al., 2021; Singh et al., 2021; Szűcs et al., 2013), whereas others did not confirm it when assessing planning, sequencing, or inhibiting responses, reporting comparable outcomes to non-diagnosed participants (Brosnan et al., 2002; Desoete & De Weerd, 2013). Contrary to ASD, the association between the ability to recognize others' emotions and EFs has never been studied in children with SLD.

### The present study

To the best of our knowledge, no previous studies investigated the abilities to recognize facial emotion expressions and associated internalizing (social anxiety) and cognitive (EFs) factors, by taking into account (and comparing) groups of children with and without ASD and SLD. By comparing FER and associated factors across various neurodevelopmental disorders, researchers can gain a better understanding of the specific challenges and strengths associated with each condition. In this regard, a great amount of studies has been conducted in the field of autism with huge variability across findings, but little attention has been given to the maladaptive effects of experiencing learning difficulties.

Thus, the first aim of the present study was to evaluate FER skills in children with different neurodevelopmental disorders (ASD, SLD) as compared to non-diagnosed (ND) participants, by also considering how different emotions are perceived. The second aim was to understand which emotional and cognitive mechanisms might be associated with FER abilities, and how these relationships might differ among groups (ASD, SLD, and ND). To do so, we investigated the role of social anxiety and executive functioning (inhibition, updating, and set-shifting) in predicting performance on an FER task.

Based on previous findings (Black et al., 2017; Bloom & Heath, 2010; Operto et al., 2020; Uljarevic & Hamilton, 2013), it is worth hypothesizing a worse general performance on the FER task in children with ASD and SLD, as compared to those without these conditions. As regards the recognition of single emotions, we expect that the clinical groups will find more difficulties in detecting negative facial expressions, such as fear and anger (Ashwin et al., 2007; Ciray & Turan, 2022; Humphreys et al., 2007).

Concerning our second aim, we expect a statistically significant association between FER ability and the reported social anxiety in



both clinical groups (ASD, SLD). In children with ASD, higher levels of social anxiety might be consistent with a poorer recognition of other people's emotional expressions because of their widely recognized social challenges and discomfort within the social contexts (Antezana et al., 2023; Corden et al., 2008; Kleinhans et al., 2010). Similarly, children and adolescents with SLD with higher levels of social anxiety might display worse FER ability, as a result of avoidance toward others' emotions which they find difficulties to interpret (Bögels & Mansell, 2004; Mansell et al., 1999). The fear of being negatively evaluated and diminished self-efficacy could play a role as well in the association between social anxiety and others' emotion recognition (Livingston et al., 2018; Rostami et al., 2014; Terras et al., 2009). Nonetheless, the hypothesis on the link between social anxiety and FER in SLD is solely exploratory due to the lack of literature on the topic.

Moreover, we predict that EFs might support the performance on the FER task in all considered groups (Martins et al., 2016; Neumann et al., 2021; Phillips et al., 2008; Sivaratnam et al., 2018; Torske et al., 2018).

## Method

### Participants

The study involved 263 children and adolescents aged between 8 and 16 years old divided into three groups: 60 (52 M) with autism spectrum disorders (ASD), 63 (44 M) with SLD, and 140 (105 M) matched non-diagnosed (ND) peers. The three groups did not statistically differ in chronological age [ $F(2, 260) = 1.04, p = .35, Adj R^2 = .001$ ; ASD = 12.05 (8.33), SLD = 12.53 (1.94), ND = 12.07 (2.06)], sex distribution [ $\chi^2 = .86, df = 2, p = .65$ ], or total IQ [ $F(2, 260) = 1.14, p = .32, Adj R^2 = .002$ ; ASD = 106.63 (15.68), SLD = 106.17 (10.47), ND = 108.66 (11.22)]. All children and adolescents were native Italian speakers, and none had any visual or hearing impairments.

The ND group comprised healthy children with typical development with no psychiatric, neurological, or neurodevelopmental disorders. They were engaged and examined individually at school. The clinical groups (ASD, SLD) were assessed at the child and adolescent psychiatric services, where they referred. All participants in the clinical groups had been previously diagnosed with ASD or SLD, according to the DSM-IV-TR or the DSM-5 (American Psychiatric Association, 2000, 2013) or ICD-10 (World Health Organization, 1992) criteria. Specifically, the SLD group was composed as follows: 44 % of the participants had major difficulties in mathematics, reading and/or writing, 35 % in reading, 13 % in writing, and 8 % in mathematics. To confirm their diagnosis, an experimenter, blind to the group membership of the participants, was responsible for testing all the children using specific clinical assessment materials. Diagnosis of ASD was confirmed using the Autism Diagnostic Interview – Revised (ADI-R; Rutter et al., 2005). The groups statistically differed in all subscales of ADI-R (Reciprocal social interaction:  $F(2, 260) = 122.78$ ; Language/communication:  $F(2, 260) = 87.81$ ; Repetitive behaviors/interests:  $F(2, 260) = 73.94$ ), with autistic participants having higher scores than those with SLD and ND ( $p_s < .001$ ). Diagnoses of SLD were also confirmed by implementing some age-appropriate subtests which evaluated reading and math competencies (AC-MT-3 6-14, Cornoldi et al., 2020; DDE-2, Sartori et al., 2007; MT-Avanzate-3-Clinica, Cornoldi et al., 2017). In both reading (words - errors:  $F(2, 260) = 72.93$ , words - time  $F(2, 260) = 45.68$ ; pseudo-words - errors:  $F(2, 260) = 39.82$ , pseudo-words - time  $F(2, 260) = 28.56$ ) and math (mental calculation - accuracy:  $F(2, 260) = 23.67$ , mental calculation - time

$F(2, 260) = 13.34$ ) subtests, the SLD group was more impaired than the ASD and ND groups ( $p_s < .001$ ).

Inclusion criterion for the current study was a standard score of 80 or more for total IQ as assessed by the Wechsler Intelligence Scales (WISC IV; Wechsler, 2003). Participants who were taking medication, or with other known genetic conditions, a history of neurological diseases, comorbid psychopathologies, or certified physical and intellectual disabilities were excluded.

### Materials

#### Facial emotion recognition

An ad hoc computerized task was created to assess FER abilities. It was developed by using the EU-Emotion Stimulus Set, which was created as part of the ASC-Inclusion project within the European Community's Seventh Framework Programme (FP7/2007-2013; O'Reilly et al., 2016).

Stimuli were static images of actors, taken from dynamic video clips with a neutral white background, that expressed a facial emotion. The frame for the static images was captured at the peak of the expressed emotion, so that the emotion appeared as close as possible to a real-life situation. A direct frontal view of each actor, featuring only their shoulders and head within the frame, was depicted. The participant was required to choose whether two actors expressed the same or different emotions by pressing distinct keys on the keyboard ("z" if "same emotion", "m" if "different emotions"). There were no time restrictions to respond. The task comprised 144 trials divided into two blocks of 72 with a small break after the first 36 trials of each block. In half of the trials the two actors showed the same emotion ( $n = 72$ ), in half different emotions ( $n = 72$ ). The six basic emotions were assessed: happiness, surprise, sadness, fear, anger, and disgust. Each of the six emotions was the correct answer 12 times during the entire task. Actors were of different ethnicity and age range 10–70 years old (ten females and nine males). The trials were counterbalanced for actors' age (adult–adult, child–child, and adult–child) and emotions' intensity (high and low). Raw scores were calculated for total accuracy. In addition, correct responses for the single emotions (happiness, surprise, sadness, fear, anger, and disgust) were considered.

For the current sample, Cronbach's  $\alpha$  (total accuracy) = .83 (C.I. = .79–.86).

#### Social anxiety

The parent-report form of the Italian version (Paloscia et al., 2017) of the Multidimensional Anxiety Scale for Children (MASC-2; March, 2012) was administered to evaluate the presence of social anxiety symptoms. Specifically, the Social Anxiety subscale (composed by Humiliation/Rejection, and Performance Fears) was administered. This dimension includes 9 items rated on a 4-point scale from 0 (*never*) to 3 (*often*). Raw scores were converted into T scores by using normative data that considers the child's age and sex.

Cronbach's  $\alpha$  (Social Anxiety scale) = .88.

#### Executive functions

The tripartite Miyake's model of EFs (Miyake et al., 2000) was applied to assess executive functioning in our sample.

#### Inhibition

Inhibitory control was assessed with a computerized go/no-go task (adapted from Christ et al., 2007). The task included 120 trials, composed by two blocks (60 + 60) with a break between them. On

each trial, one of four stimuli (blue, red, yellow, and green dots) was centrally displayed in a computer monitor. In the first block, children were asked to press the spacebar as quickly as possible when a blue dot appeared (target; go trials) and to give no response when a dot in any other color showed (non-target; no-go trials). In the second block, children were asked to press the spacebar as quickly as possible when a dot of any color except for blue appeared (target; go trials) and to deliver no response when a blue dot appeared (non-target; no-go trials). Performance on the go trials corresponds to a measure of attention, whereas errors on the no-go trials to a measure of inhibition. Within each block, stimuli were produced in a random order, and targets were presented on a minority of trials (25%). After an intertrial interval of 2,000 ms, a new trial was presented. At the beginning of the task, after giving the instructions, eight practise trials were presented, in which children were provided a feedback on the response ("correct", "incorrect", and "too slow" when the response was not given within 2,000 ms). Errors on the no-go trials (number of reactions to the non-target stimuli) were considered as a measure of inhibition.

Cronbach's  $\alpha$  (for both go and no-go trials) = .94.

### Updating

This computerized task, adapted from Cardillo et al. (2021), was used to assess the ability to visually and spatially update information into the working memory system. Participants were shown a  $4 \times 4$  blank matrix on the computer screen. A total of eight sequences of various shapes (i.e., triangle, circle, star, square, rhombus, and pentagon) appeared in the matrix. Participants were asked to recall the last position of certain shapes shown on the screen. The number of shapes to recall ranges from 2 to 5, depending on the level's difficulty. Sequences of 6, 8, 10, or 12 shapes were used, and there were two trials for each sequence length. Before starting the experiment, participants were presented with three practice trials. Accuracy was calculated as a proportion of items' last positions correctly recalled out of the total positions to remember (Giofrè & Mammarella, 2014).

Cronbach's  $\alpha$  = .80 (C.I. = .75–.83).

### Set-shifting

A computerized version of the Wisconsin Card Sort Test (WCST; Heaton et al., 1993) has been designed from the original edition. Each trial comprised four stimulus cards and a response card, and the participant's goal was to correctly match the response card to one of the stimulus cards according to one of three grouping principles. After each trial, participants received feedback on the answer's correctness. The stimulus cards appeared in a row at the top of the computer screen, and the response cards showed one at a time at the bottom of the screen. The four stimulus cards displayed the following symbols: (a) one red triangle, (b) two green stars, (c) three yellow crosses, and (d) four blue circles. There were two decks of 64 response cards, each containing from one to four identical symbols (triangles, stars, crosses, or circles) of a particular color (red, green, yellow, or blue). The first correct sorting principle was color, followed by form and then number. The participant had to correctly match the stimulus and the response cards 10 times consecutively within each criterion. The test finished when the participant was able to match the response cards following six criteria (twice each sorting principle), or when he/she used two identical decks of 64 response cards (128 cards total). The total number of perseverative errors, calculated as proportions to the number of administered trials, was used in the analysis.

A perseverative error is defined as a failure to shift category after receiving negative feedback from the previous trial.

Generalizability coefficient = .60 (from .37 to .72) (Heaton et al., 1993).

### Procedure

The study was approved by the ethics review board of the Authors' institution. After obtaining the written consent of children's parents to their participation in the study, participants were tested individually in a quiet room at specialized centers (ASD, SLD) or at school (ND) during three sessions (a screening and two experimental phases) lasting approximately 40 minutes each. In the screening phase, the IQ was calculated and only participants who scored above 80 were included; reading and math competences were also evaluated in this initial phase. Meanwhile, parents were administered the ADI-R for child autistic symptoms. The first experimental phase included the FER test (first block), go-no go, and updating tasks. The second one comprised the FER task (second block) and the WCST. The tasks in the experimental sessions were presented in a counterbalanced order. At the same time, parents completed the MASC-2 for assessing the child social anxiety. The experimenter provided instructions on each task, letting the participant practice before starting the experiment. The computerized tasks (FER and EFs' tasks) were created and administered using PsychoPy3 (Peirce et al., 2019) and a laptop computer with a 15-inch LCD screen.

### Statistical approach

First, a series of univariate ANOVAs were performed to estimate differences between the three groups in the measures of interest. Post hoc comparisons between groups were assessed through the Tukey's correction test. Effect sizes were also computed, using both adjusted R-squared ( $Adj R^2$ ), which represents the effect size of the comparisons between the three groups for the factors considered, and Cohen's  $d$ , which instead expresses the effect size of the pairwise comparisons between the groups (for details see Supplementary results).

Second, a series of regression models were developed to assess the effect of the measured variables on the dependent variable. To do this, we created a full model with FER (total accuracy) as dependent variable, and group, social anxiety, inhibition, updating and set-shifting as predictors. Age of the child was included as a covariate. Starting from the full model, we built the various models by removing one variable at time, to evaluate the principal effects of the single predictors on the dependent variable. In addition, the interaction effects between each variable and the group were tested, starting from the full model we added one interaction effect at time. We compared each model with the full model, using the Akaike Information Criterion (AIC) and the  $Adj R^2$ . A lower AIC indicates a better model. The  $Adj R^2$  represents the proportion of variance in the outcome variable which is explained by the predictor variables in the sample. The  $Adj R^2$  is a modified version of  $R^2$  that adjusts for the number of predictors in a regression model and allows for comparing models with a different number of predictors (Miles, 2005). Cohen (1988) has defined the effects as follows: a very weak effect  $< 0.02$ , a weak effect between  $0.02 \leq R^2 < 0.13$ , a moderate effect between  $0.13 \leq R^2 < 0.26$ , and a large effect  $\geq 0.26$ , whereas a recent paper establishes that an  $R^2$  of at least 0.10 is acceptable in social science empirical modeling (Ozili, 2023).

Data were analyzed using R version 1.3.1093 (R Core Team, 2022). The "lme4" package was used to run the regression models

**Table 1.** Descriptive statistics and statistical comparisons by group for correct responses on the facial emotion recognition task, in children and adolescents with Autism spectrum disorders (ASD), specific learning disorders (SLD), and non-diagnosed (ND)

Measures	ASD ( <i>n</i> = 60)	SLD ( <i>n</i> = 63)	ND ( <i>n</i> = 140)	<i>F</i> (2, 260)	<i>p</i>	<i>Adj R</i> <sup>2</sup>	Post-hoc comparisons
Total accuracy	112.13 (10.35)	114.62 (7.93)	119.49 (8.36)	17.03	<.001	.11	ASD < ND, SLD < ND
Happiness	11.05 (1.39)	11.09 (1.17)	11.47 (.97)	4.10	.02	.02	ASD < ND
Surprise	8.70 (2.59)	8.76 (2.21)	9.72 (1.79)	7.29	<.001	.04	ASD < ND, SLD < ND
Sadness	10.25 (1.96)	9.93 (1.97)	10.62 (1.50)	3.57	.03	.02	SLD < ND
Fear	6.71 (3.12)	7.03 (2.53)	7.85 (2.15)	5.23	.006	.03	ASD < ND
Anger	6.91 (2.64)	7.02 (2.77)	7.81 (2.31)	3.76	.02	.02	ASD < ND
Disgust	7.98 (2.05)	8.33 (2.05)	9.45 (1.99)	13.72	<.001	.09	ASD < ND, SLD < ND

**Table 2.** Descriptive statistics and statistical comparisons by group for social anxiety and executive functions (inhibition, updating, and set-shifting), in children and adolescents with Autism spectrum disorders (ASD), specific learning disorders (SLD), and non-diagnosed (ND)

Measures	Variables	ASD ( <i>n</i> = 60)	SLD ( <i>n</i> = 63)	ND ( <i>n</i> = 140)	<i>F</i> (2, 260)	<i>p</i>	<i>Adj R</i> <sup>2</sup>	Post-hoc comparisons
Social anxiety	T score	66.35 (14.79)	67.16 (11.16)	58.19 (12.03)	15.43	<.001	.10	ASD > ND, SLD > ND
Inhibition	No-go (errors)	4.23 (3.57)	4.08 (3.43)	3.75 (3.42)	.46	.63	-.004	–
Updating	Total accuracy (%)	63.60 (18.58)	70.80 (16.23)	72.87 (13.63)	7.56	<.001	.05	ASD < ND, ASD < SLD
Set-shifting	Perseverative errors (%)	13.72 (5.45)	14.04 (5.96)	13.59 (6.89)	.10	.88	-.006	–

and the AIC (Bates et al., 2015), and the “ggplot2” package to obtain the graphical effects (Wickham, 2016).

## Results

### Preliminary analyses

Tables 1 and 2 display descriptive statistics and statistical comparisons between the three groups in all considered measures. Effect sizes of the pairwise comparisons between the groups (Cohen’s *d*), are reported in Table S1 (Supplementary results).

### Facial emotion recognition

Table 1 concerns to the performance in the FER task with respect to total accuracy, and single emotions (happiness, surprise, sadness, fear, anger, and disgust). The three groups statistically differed in the scores of all the considered variables in this test, with a worse performance for clinical groups as compared to ND. Both the ASD (*Tukey’s p* < .001, *Cohen’s d* = −.82) and SLD (*Tukey’s p* < .001, *Cohen’s d* = −.59) groups achieved an overall lower performance in recognizing emotions as compared to the ND group, *F* (2, 260) = 17.03, *p* < .001, *Adj R*<sup>2</sup> = .11. Specifically, the ASD group scored worse than the ND group in recognizing happiness (*Tukey’s p* = .04, *Cohen’s d* = −.38), surprise (*Tukey’s p* = .005, *Cohen’s d* = −.49), fear (*Tukey’s p* = .01, *Cohen’s d* = −.46), anger (*Tukey’s p* = .04, *Cohen’s d* = −.37), and disgust (*Tukey’s p* < .001, *Cohen’s d* = −.73). Instead, the SLD group obtained lower scores as compared to the ND group in detecting surprise (*Tukey’s p* = .008, *Cohen’s d* = −.49), sadness (*Tukey’s p* = .03, *Cohen’s d* = −.41), and disgust (*Tukey’s p* < .001, *Cohen’s d* = −.55).

### Social anxiety

Concerning social anxiety (see Table 2), a statistically significant effect of the group emerged, *F* (2, 260) = 15.43, *p* < .001, *Adj R*<sup>2</sup> = .10. Parents of children and adolescents with ASD (*Tukey’s p* < .001, *Cohen’s d* = .63) and SLD (*Tukey’s p* < .001, *Cohen’s d* = .76) reported greater levels of social anxiety as compared to the ND group.

**Table 3.** Results of the comparisons between regression models with facial emotion recognition (total accuracy) as dependent variable, and all the other measured variables and their interactions with group as predictors. Age of the child has been included as a covariate

Model	<i>F</i>	<i>p</i>	AIC	<i>R</i> <sup>2</sup> adjusted
Full model – Group	15.46	<.001	1872.64	.18
Full model – Social anxiety	.15	.69	1844.69	.28
Full model – Inhibition	13.04	<.001	1857.65	.23
Full model – Updating	4.27	.04	1848.91	.25
Full model – Set shifting	1.19	.28	1845.75	.27
Full model + Group*Social anxiety	3.13	.04	1844.11	.28
Full model + Group*Inhibition	1.39	.25	1847.65	.26
Full model + Group*Updating	3.42	.03	1843.50	.28
Full model + Group*Set shifting	.89	.41	1848.67	.26

Note. AIC, Akaike Information Criterion. Full model = Age + Group + Social anxiety + Inhibition + Updating + Set-shifting. [AIC = 1846.53, *R*<sup>2</sup> adjusted = .26].


### Executive functions

As regards the EFs’ tasks (see Table 2), the ASD group performed significantly worse than the ND (*Tukey’s p* < .001, *Cohen’s d* = −.61) and SLD (*Tukey’s p* = .03, *Cohen’s d* = −.41) groups in the updating task, *F* (2, 260) = 7.56, *p* < .001, *Adj R*<sup>2</sup> = .05.


No statistically significant differences between groups emerged as regard to the other measures.

### Regression models

A series of linear regressions were tested in order to investigate the association between the FER and the hypothesized predictors (i.e., group, social anxiety, inhibition, updating, and set-shifting). Age of the child was included as a covariate. The interactive effect of group was also considered (see Table 3).



**Figure 1.** Significant interaction effect of the group with social anxiety (T score), and facial emotion recognition (total accuracy) as a dependent variable. Error bands represent 95% confidence intervals.



**Figure 2.** Significant interaction effect of the group with updating (% accuracy), and facial emotion recognition (total accuracy) as a dependent variable. Error bands represent 95% confidence intervals.

A significant main effect of group was found (full model with group:  $AIC = 1846.53$ ,  $R^2_{adj} = .26$ , model without group:  $AIC = 1872.64$ ,  $R^2_{adj} = .18$ ;  $F = 15.46$ ,  $p < .001$ ). The model coefficients showed that the ASD and SLD groups were less accurate than the ND group in the FER task ( $p_s < .001$ ).

The main effects of inhibition (full model with inhibition:  $AIC = 1846.53$ ,  $R^2_{adj} = .26$ , model without inhibition:  $AIC = 1857.65$ ,  $R^2_{adj} = .23$ ;  $F = 13.04$ ,  $p < .001$ ) and updating (full model with updating:  $AIC = 1846.53$ ,  $R^2_{adj} = .26$ , model without updating:  $AIC = 1848.91$ ,  $R^2_{adj} = .25$ ;  $F = 4.27$ ,  $p = .04$ ) were also significant. The model coefficients showed that participants who made more

errors in the inhibition task ( $p = .001$ ) and were more accurate and in the updating task ( $p < .001$ ), had higher performance in the FER task. Differently, the effects of social anxiety and set-shifting were not statistically significant ( $p = .69$  and  $p = .28$  respectively).

As regards the interaction effects between measures and group, the analysis revealed two statistically significant interaction effects. The interaction between group and social anxiety was found to significantly predict the performance on the FER task (model with interaction:  $AIC = 1844.11$ ,  $R^2_{adj} = .28$ , model without interaction:  $AIC = 1846.53$ ,  $R^2_{adj} = .26$ ;  $F = 3.13$ ,  $p = .04$ ). More specifically, as Figure 1 shows, higher levels of social anxiety are



consistent with greater scores on the FER task in ASD group, whereas higher levels of social anxiety are related to lower scores on the FER task in SLD group, and no relation emerged for ND group.

Also the interaction between group and updating was found to be statistically significant (model with interaction:  $AIC = 1843.50$ ,  $R^2_{adj} = .28$ , model without interaction:  $AIC = 1846.53$ ,  $R^2_{adj} = .26$ ;  $F = 3.42$ ,  $p = .03$ ). As Figure 2 shows, a better performance in the updating task is significantly associated to higher scores in the FER task only in the ASD group.

No other statistically significant main or interaction effects occurred.

## Discussion

The present study sought to investigate the abilities to recognize facial emotion expressions in children and adolescents with and without ASD and SLD, by also considering how different emotions are identified. A second aim was to study how social anxiety and executive functioning (inhibition, updating, and set-shifting) might underlie the performance on a FER task in these three groups (ASD, SLD, and ND).

In line with previous data in literature (Black *et al.*, 2017; Bloom & Heath, 2010; Operto *et al.*, 2020; Uljarevic & Hamilton, 2013), our findings revealed that performance on the FER task was statistically worse in both clinical groups (ASD, SLD) as compared to the ND group. Recognizing other people's emotions from facial expression represents a foundation for adaptive social functioning (Fischer *et al.*, 2019; Van Kleef, 2009), so youth with ASD and SLD might find challenging to navigate the social world because of their difficulties to interpret emotional dispositions, matched with requests, intentions, and feedbacks in dyadic social exchanges. However, differences between ASD and SLD emerged as regards the type of conveyed emotion. Autistic participants seem to experience difficulties in detecting all emotions included in our task (except for sadness) as compared to the ND group, giving support to the hypothesis of a generalized face-emotion recognition deficit in ASD (Leung *et al.*, 2022; Lindner & Rosén, 2006; Lozier *et al.*, 2014; Rump *et al.*, 2009). It is worth noting that a lack of global expertise in basic social processes, like FER, may be associated to profound deficits in social reciprocity (Høyland *et al.*, 2017; Trevisan & Birmingham, 2016), particularly when individuals fail to recognize primary emotions (e.g., happiness). Interestingly, children and adolescents with SLD demonstrated an overall worse performance in the FER test, but more remarkably specific challenges with certain emotions (Ciray & Turan, 2022), not experiencing any problems with happiness, fear, and anger. Apart from happiness which is the first learnt (and more common) emotion, fear, and anger are adaptive emotions with a long evolutionary history, representing reactions to threatening stimuli (Marsh *et al.*, 2005), thus it may be easier to recognize and interpret them. Instead, participants with SLD demonstrated problems with interpreting facial expressions such as those expressing disgust, surprise, and sadness. However, future studies should verify if such findings can be replicated because of the lack of literature on the recognition of specific emotions in SLD.

### Facial emotion recognition and social anxiety

To further explore the ability of recognizing others' emotions from their facial expressions, it is worth investigating whether internalizing symptoms might play a role. Our findings revealed that both parents of children and adolescents with ASD and SLD reported higher levels of social anxiety in their sons and daughters

as compared to ND participants, consistent with data in literature (Bellini, 2006; Livingston *et al.*, 2018; Terras *et al.*, 2009; White *et al.*, 2010). Therefore, youth with ASD and SLD might experience concerns and emotional activation when acting in front of other people, and this could prevent them from being involved in social situations (Rostami *et al.*, 2014; Spain *et al.*, 2018). This might trigger a cascade effect which could lower the possibility of these children of practising with social skills and with interpreting others' emotions and intentions to appropriately socially behave.

Indeed, what is noteworthy is the relationship between social anxiety and FER abilities depending on the group of membership. In fact, a statistically significant interaction emerged between social anxiety and group in predicting FER ability.

In contrast with our initial hypotheses, for children with ASD a better recognition of other people's facial emotional expressions was consistent with higher levels of social anxiety reported by parents. Better FER skills might trigger social anxiety in autistic children and adolescents, because of the distress with social interactions, which may increase when they have more advanced sociocognitive abilities and are more aware of the social context, as revealed by research on socially anxious individuals (Hunter *et al.*, 2009; Nikolic *et al.*, 2019; Tibi-Elhanany & Shamay-Tsoory, 2011). Alternatively, autistic children with higher levels of social anxiety may have enhanced attentional resources toward external stimuli (Schultz & Heimberg, 2008), as a consequence of hypervigilance when executing a socially based test. Indeed, emotional faces might be perceived as threatening for people with ASD because of the awareness of their own social challenges (Brewer *et al.*, 2022), causing a biased attentive processing (Koster *et al.*, 2006). Higher attentional levels could have resulted in a better performance, because of the increased focus on the FER task: anxiety may not weaken the performance on a task when it enables the use of compensatory strategies (Eysenck *et al.*, 2007). An interesting study conducted by Livingston *et al.* (2019) stated that autistic adolescents who show better outcome (e.g., good social skills) than others, are usually the ones with higher IQ and EFs but also higher levels of anxiety. In this sense, compensation may come at a cost, promoting exhaustion, and anxiety for trying to meet social demands.

In agreement with our initial hypotheses instead, a worse FER ability was associated with higher levels of social anxiety in children with SLD. One potential explanation for this result is that youth with SLD are more in need to receive social feedbacks, especially in school, as a consequence of their awareness of their academic difficulties, lower self-efficacy, and fear of being negatively evaluated (Carroll & Iles, 2006; Elgendi *et al.*, 2021; Rostami *et al.*, 2014; Terras *et al.*, 2009). This necessity for feedbacks from others might negatively affect emotional regulation and boost social anxiety when other people's emotions are not well understood. Moreover, higher levels of social anxiety, worries, and emotional activation in children with SLD might be consistent with avoidance and safety behaviors (Bögels & Mansell, 2004; Mansell *et al.*, 1999), which could have prevented them from accurately perform on the FER task.

### Facial emotion recognition and executive functions

According to the literature, not only internalizing symptoms such as social anxiety but also cognitive processes are involved in the processing of others' emotions. As regards the role of EFs, the main effect of inhibition emerges in predicting FER in the whole sample. The bidirectional relation between executive functioning and social abilities has been widely explored: EFs are required to correctly



behave in social contexts, and practising with social skills may support the development of cognitive processes associated with behavioral regulation (Denham et al., 2012; Moriguchi, 2014; Spikman et al., 2013). In this sense, the ability to inhibit impulsive behaviors is significantly correlated with the capacity to detect others' emotions, but also with the ability to perform a computerized task that requires to press the correct answer as quickly as possible (Martins et al., 2016; Neumann et al., 2021). Besides, a statistically significant interaction emerged between updating and group membership in predicting FER ability. Updating skills seem to be predictive of social abilities in autistic children and adolescents (Phillips et al., 2008; Sivaratnam et al., 2018; Torske et al., 2018). As a fact, higher order EFs may support social engagement, acting as an important compensatory mechanism (Johnson et al., 2015; Livingston et al., 2019). In line with these previous studies, our findings revealed that ASD-diagnosed children who exhibited the highest levels of updating skills performed at a level equivalent to non-diagnosed participants on the FER task. It appears as though this underlying executive function has a compensatory impact on social skills in individuals with autism.

### Limitations and potential implications

Despite the novelty and significance of our findings, some limitations of the present study need to be considered. The first limitation might be intrinsic to the FER task, which is composed by static images of facial emotional expressions, and though previous research stated the reliability of this type of stimuli, also other tasks could be useful to study emotion recognition. For example, dynamic cues (e.g., morphing from neutral to angry) and videos of social interactions might be implemented to examine emotion recognition from an ecological point of view, which resembles more the real world (Cardillo et al., 2023; Ekman & Friesen, 2003). The absence of time constraints in the FER task may have allowed the use of compensatory strategies. For instance, autistic children without intellectual disability could potentially use their cognitive resources to effectively recognize the emotions of others (Harms et al., 2010; Leung et al., 2022). Future studies should investigate this aspect, which may provide valuable insights into the strategies employed by children with different neurodevelopmental conditions in the recognition of emotions. Moreover, only the six basic emotions have been investigated (Ekman, 1993), thus it could be interesting to explore how children and adolescents diagnosed with ASD and SLD perceive and interpret more complex emotions, especially the "social" emotions (e.g., shame, guilt), which could be more related to the development of social anxiety and internalizing symptoms. Future research should further study the underlying mechanisms of social challenges in children with SLD and the association between academic competences, social learning, and executive functioning.

Potential implications of the present research should target clinical practise when handling social skills. From our findings, higher awareness of others' emotions is consistent with higher social anxiety levels in ASD, thus it could be possible that children who do not dispose of social strategies could feel tension when they face social demands with the growing awareness of their own social difficulties. From this point of view, social motivation and social skills should be empowered to prevent negative feelings associated with emotional reciprocity. Instead, a key factor for preventing the rising of social problems in children with SLD might be enhancing self-efficacy and lowering worries concerning others' negative

judgment, allowing the possibility of practising social skills and interacting with peers. Lastly, executive functioning should also be addressed to support social learning in the case of ASD-diagnosed children and adolescents.

Overall, despite similar weaknesses in FER in ASD and SLD, different underlying mechanisms may lie behind such performances in these two different neurodevelopmental conditions. Understanding these differences can guide practitioners in designing individualized interventions, tailored to the specific challenges of the examined clinical profile. Our findings provide new insights into the relationship between FER, social anxiety, and executive functioning in children and adolescents with different neurodevelopmental disorders. Noteworthy, specific patterns distinctly emerged in children with ASD and SLD in the association between social anxiety and FER, whereas executive functioning seems to boost social skills in autism.

**Supplementary material.** The supplementary material for this article can be found at <https://doi.org/10.1017/S0954579424000038>.

**Author contribution.** All authors contributed to the study's conception and design. RL conceptualized the study, RL and RC performed the literature search and data analysis, and RL drafted the work. All authors (RL, RC, and ICM) discussed versions of the draft. ICM supervised and critically revised the work and approved the final manuscript.

**Funding statement.** None.

**Competing interests.** None.

### References

- Albayrak, Z. S., Kadak, M. T., Gürbüz, H. G. A., & Doğançün, B. (2022). Emotion recognition skill in specific learning disorder and attention-deficit hyperactivity disorder. *Alpha Psychiatry*, 23(6), 268–273. <https://doi.org/10.5152/alphapsychiatry.2022.22219>
- Antezana, L., Valdespino, A., Wieckowski, A. T., Coffman, M. C., Carlton, C. N., Garcia, K. M., Gracanin, D., White, S. W., & Richey, J. A. (2023). Social anxiety symptoms predict poorer facial emotion recognition in autistic male adolescents and young adults without intellectual disability. *Journal of Autism and Developmental Disorders*, <https://doi.org/10.1007/s10803-023-05998-5>
- Artuso, C., Borgatti, R., & Palladino, P. (2021). Phonological memory updating and developmental dyslexia: The role of long-term knowledge. *Child Neuropsychology*, 27(6), 718–733. <https://doi.org/10.1080/09297049.2021.1888907>
- Ashwin, E. T., Baron-Cohen, S., Wheelwright, S., O'Riordan, M., & Bullmore, E. T. (2007). Differential activation of the amygdala and the social brain during fearful face-processing in asperger syndrome. *Neuropsychologia*, 45(1), 2–14. <https://doi.org/10.1016/j.neuropsychologia.2006.04.014>
- Bal, E., Harden, E., Lamb, D., Van Hecke, A. V., Denver, J. W., & Porges, S. W. (2010). Emotion recognition in children with autism spectrum disorders: Relations to eye gaze and autonomic state. *Journal of Autism and Developmental Disorders*, 40(3), 358–370. <https://doi.org/10.1007/s10803-009-0884-3>
- Bantin, T., Stevens, S., Gerlach, A. L., & Hermann, C. (2016). What does the facial dot-probe task tell us about attentional processes in social anxiety? A systematic review. *Journal of Behavior Therapy and Experimental Psychiatry*, 50, 40–51. <https://doi.org/10.1016/j.jbtep.2015.04.009>
- Baron-Cohen, S., Golan, O., & Ashwin, E. (2009). Can emotion recognition be taught to children with autism spectrum conditions? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364(1535), 3567–3574. <https://doi.org/10.1098/rstb.2009.0191>
- Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67(1), 1–48. <https://doi.org/10.18637/jss.v067.i01>
- Battaglia, M., Ogliari, A., Zanoni, A., Villa, F., Citterio, A., Binaghi, F., Fossati, A., & Maffei, C. (2004). Children's discrimination of expressions of

- emotions: Relationship with indices of social anxiety and shyness. *Journal of the American Academy of Child & Adolescent Psychiatry*, 43(3), 358–365. <https://doi.org/10.1097/00004583-200403000-00019>
- Beauchamp, M. H., & Anderson, V. (2010). SOCIAL: An integrative framework for the development of social skills. *Psychological Bulletin*, 136(1), 39–64. <https://doi.org/10.1037/a0017768>
- Bell, C., Bourke, C., Colhoun, H., Carter, F., Frampton, C., & Porter, R. (2011). The misclassification of facial expressions in generalised social phobia. *Journal of Anxiety Disorders*, 25(2), 278–283. <https://doi.org/10.1016/j.janxdis.2010.10.001>
- Bellini, S. (2006). The development of social anxiety in adolescents with autism spectrum disorders. *Focus On Autism and Other Developmental Disabilities*, 21(3), 138–145. <https://doi.org/10.1177/10883576060210030201>
- Black, M. H., Chen, N. T. M., Iyer, K. K., Lipp, O. V., Bölte, S., Falkner, M., Tan, T., & Girdler, S. (2017). Mechanisms of facial emotion recognition in autism spectrum disorders: Insights from eye tracking and electroencephalography. *Neuroscience & Biobehavioral Reviews*, 80, 488–515. <https://doi.org/10.1016/j.neubiorev.2017.06.016>
- Bloom, E., & Heath, N. (2010). Recognition, expression, and understanding facial expressions of emotion in adolescents with nonverbal and general learning disabilities. *Journal of Learning Disabilities*, 43(2), 180–192. <https://doi.org/10.1177/0022219409345014>
- Bögels, S. M., & Mansell, W. (2004). Attention processes in the maintenance and treatment of social phobia: Hypervigilance, avoidance and self-focused attention. *Clinical Psychology Review*, 24(7), 827–856. <https://doi.org/10.1016/j.cpr.2004.06.005>
- Brewer, N., Lucas, C. A., Georgopoulos, M. A., & Young, R. L. (2022). Facing up to others' emotions: No evidence of autism-related deficits in metacognitive awareness of emotion recognition. *Autism Research*, 15(8), 1508–1521. <https://doi.org/10.1002/aur.2781>
- Brigman, G. A., Webb, L. D., & Campbell, C. (2007). Building skills for school success: Improving the academic and social competence of students. *Professional School Counseling*, 10(3), 2156759X0701000310. <https://doi.org/10.1177/2156759X0701000310>
- Brosnan, M., Demetre, J., Hamill, S., Robson, K., Shepherd, H., & Cody, G. (2002). Executive functioning in adults and children with developmental dyslexia. *Neuropsychologia*, 40(12), 2144–2155. [https://doi.org/10.1016/S0028-3932\(02\)00046-5](https://doi.org/10.1016/S0028-3932(02)00046-5)
- Button, K., Lewis, G., Penton-Voak, I., & Munafò, M. (2013). Social anxiety is associated with general but not specific biases in emotion recognition. *Psychiatry Research*, 210(1), 199–207. <https://doi.org/10.1016/j.psychres.2013.06.005>
- Cardillo, R., Crisci, G., Seregni, S., & Mammarella, I. C. (2023). Social perception in children and adolescents with ADHD: The role of higher-order cognitive skills. *Research in Developmental Disabilities*, 135, 104440. <https://doi.org/10.1016/j.ridd.2023.104440>
- Cardillo, R., Mammarella, I. C., Demurie, E., Giofrè, D., & Roeyers, H. (2021). Pragmatic language in children and adolescents with Autism spectrum disorder: Do theory of mind and executive functions have a mediating role? *Autism Research*, 14(5), 932–945. <https://doi.org/10.1002/aur.2423>
- Carlson, S. M., & Wang, T. S. (2007). Inhibitory control and emotion regulation in preschool children. *Cognitive Development*, 22(4), 489–510. <https://doi.org/10.1016/j.cogdev.2007.08.002>
- Carroll, J. M., & Iles, J. E. (2006). An assessment of anxiety levels in dyslexic students in higher education. *British Journal of Educational Psychology*, 76(3), 651–662. <https://doi.org/10.1348/000709905X66233>
- Caviola, S., Colling, L. J., Mammarella, I. C., & Szűcs, D. (2020). Predictors of mathematics in primary school: Magnitude comparison, verbal and spatial working memory measures. *Developmental Science*, 23(6), e12957. <https://doi.org/10.1111/desc.12957>
- Chiu, H.-M., Chen, C.-T., Tsai, C.-H., Li, H.-J., Wu, C.-C., Huang, C.-Y., & Chen, K.-L. (2022). Theory of mind predicts social interaction in children with Autism spectrum disorder: A two-year follow-up study. *Journal of Autism and Developmental Disorders*, 53(9), 3659–3669. <https://doi.org/10.1007/s10803-022-05662-4>
- Christ, S. E., Holt, D. D., White, D. A., & Green, L. (2007). Inhibitory control in children with Autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 37(6), 1155–1165. <https://doi.org/10.1007/s10803-006-0259-y>
- Ciray, R. O., & Turan, S. (2022). Searching the underlying mechanisms of specific learning disorder: An emotion recognition and social cognition aspect for Turkish clinical youth population. *Applied Neuropsychology: Child*, 0(0), 1–7. <https://doi.org/10.1080/21622965.2022.2156290>
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Lawrence Erlbaum.
- Corden, B., Chilvers, R., & Skuse, D. (2008). Avoidance of emotionally arousing stimuli predicts social-perceptual impairment in Asperger's syndrome. *Neuropsychologia*, 46(1), 137–147. <https://doi.org/10.1016/j.neuropsychologia.2007.08.005>
- Cornoldi, C., Mammarella, I. C., & Caviola, S. (2020). AC-MT-3 6-14 anni. *Prove per la clinica*. Erickson.
- Cornoldi, C., Pra Baldi, A., & Giofrè, D. (2017). *Prove MT avanzate-3-clinica*. Giunti-Edu.
- Crisci, G., Caviola, S., Cardillo, R., & Mammarella, I. C. (2021). Executive functions in neurodevelopmental disorders: Comorbidity overlaps between attention deficit and hyperactivity disorder and specific learning disorders. *Frontiers in Human Neuroscience*, 15, 594234.
- Dajani, D. R., Llabre, M. M., Nebel, M. B., Mostofsky, S. H., & Uddin, L. Q. (2016). Heterogeneity of executive functions among comorbid neurodevelopmental disorders. *Scientific Reports*, 6(1), 36566. <https://doi.org/10.1038/srep36566>
- David, D. P., Soeiro-de-Souza, M. G., Moreno, R. A., & Bio, D. S. (2014). Facial emotion recognition and its correlation with executive functions in bipolar I patients and healthy controls. *Journal of Affective Disorders*, 152–154, 288–294. <https://doi.org/10.1016/j.jad.2013.09.027>
- Denham, S. A., Bassett, H. H., Way, E., Mincic, M., Zinsser, K., & Graling, K. (2012). Preschoolers' emotion knowledge: Self-regulatory foundations, and predictions of early school success. *Cognition and Emotion*, 26(4), 667–679. <https://doi.org/10.1080/02699931.2011.602049>
- Denham, S. A., & Brown, C. (2010). "Plays nice with others": Social-emotional learning and academic success. *Early Education and Development*, 21(5), 652–680. <https://doi.org/10.1080/10409289.2010.497450>
- Desoete, A., & De Weert, F. (2013). Can executive functions help to understand children with mathematical learning disorders and to improve instruction? *Learning Disabilities – A Contemporary Journal*, 11(2), 27–39.
- American Psychiatric Association (2000). *Diagnostic and statistical manual of mental disorders* (4th ed.). Author.
- Diamond, A. (2013). Executive functions. *Annual Review of Psychology*, 64(1), 135–168. <https://doi.org/10.1146/annurev-psych-113011-143750>
- Donolato, E., Cardillo, R., Mammarella, I. C., & Melby-Lervåg, M. (2022). Research review: Language and specific learning disorders in children and their co-occurrence with internalizing and externalizing problems: A systematic review and meta-analysis. *Journal of Child Psychology and Psychiatry*, 63(5), 507–518. <https://doi.org/10.1111/jcpp.13536>
- American Psychiatric Association (2013). *DSM-5. Diagnostic and statistical manual of mental disorders* (5th ed.). American Psychiatric Association.
- Dudeney, J., Sharpe, L., & Hunt, C. (2015). Attentional bias towards threatening stimuli in children with anxiety: A meta-analysis. *Clinical Psychology Review*, 40, 66–75. <https://doi.org/10.1016/j.cpr.2015.05.007>
- Economides, A., Baiano, C., Zappullo, I., Conson, M., Kalli-Laouri, J., Laouris, Y., & Esposito, A. (2020). Is Autism, attention deficit hyperactivity disorder (ADHD) and specific learning disorder linked to impaired emotion recognition in primary school aged children? *2020 IEEE International Conference On Human-Machine Systems (ICHMS)*, 1–6. <https://doi.org/10.1109/ICHMS49158.2020.9209413>
- Ekman, P. (1993). Facial expression and emotion. *American Psychologist*, 48(4), 384–392. <https://doi.org/10.1037/0003-066X.48.4.384>
- Ekman, P., & Friesen, W. V. (2003). *Unmasking the face: A guide to recognizing emotions from facial clues*. ISHK.
- Elgendy, M. M., Stewart, S. H., MacKay, E. J., & Deacon, S. H. (2021). Two aspects of psychological functioning in undergraduates with a history of

- reading difficulties: Anxiety and self-efficacy. *Annals of Dyslexia*, 71(1), 84–102. <https://doi.org/10.1007/s11881-021-00223-3>
- Eysenck, M. W., Derakshan, N., Santos, R., & Calvo, M. G. (2007). Anxiety and cognitive performance: Attentional control theory. *Emotion*, 7(2), 336–353. <https://doi.org/10.1037/1528-3542.7.2.336>
- Fischer, A. H., Pauw, L. S., & Manstead, A. S. R. (2019). Emotion recognition as a social act: The role of the expresser-observer relationship in recognizing emotions. In U. Hess, & S. Hareli (Eds.), *The social nature of emotion expression: What emotions can tell us about the world* (pp. 7–24). Springer International Publishing. [https://doi.org/10.1007/978-3-030-32968-6\\_2](https://doi.org/10.1007/978-3-030-32968-6_2)
- Gabay, Y., Dundas, E., Plaut, D., & Behrmann, M. (2017). Atypical perceptual processing of faces in developmental dyslexia. *Brain and Language*, 173, 41–51. <https://doi.org/10.1016/j.bandl.2017.06.004>
- Giofrè, D., & Mammarella, I. C. (2014). The relationship between working memory and intelligence in children: Is the scoring procedure important? *Intelligence*, 46, 300–310. <https://doi.org/10.1016/j.intell.2014.08.001>
- Giofrè, D., Mammarella, I. C., & Cornoldi, C. (2014). The relationship among geometry, working memory, and intelligence in children. *Journal of Experimental Child Psychology*, 123, 112–128. <https://doi.org/10.1016/j.jecp.2014.01.002>
- Harms, M. B., Martin, A., & Wallace, G. L. (2010). Facial emotion recognition in autism spectrum disorders: A review of behavioral and neuroimaging studies. *Neuropsychology Review*, 20(3), 290–322. <https://doi.org/10.1007/s11065-010-9138-6>
- Heaton, R. K., Chelune, G. J., Talley, J. L., Kay, G. G., & Curtiss, G. (1993). *Wisconsin card sorting test manual*. Psychological Assessment Resources.
- Heimberg, R. G., Hofmann, S. G., Liebowitz, M. R., Schneier, F. R., Smits, J. A. J., Stein, M. B., Hinton, D. E., & Craske, M. G. (2014). Social anxiety disorder in Dsm-5. *Depression and Anxiety*, 31(6), 472–479. <https://doi.org/10.1002/da.22231>
- Hill, E. L. (2004). Executive dysfunction in autism. *Trends in Cognitive Sciences*, 8(1), 26–32. <https://doi.org/10.1016/j.tics.2003.11.003>
- Høyland, A. L., Nærland, T., Engstrøm, M., Lydersen, S., & Andreassen, O. A. (2017). The relation between face-emotion recognition and social function in adolescents with autism spectrum disorders: A case control study. *PLOS ONE*, 12(10), e0186124. <https://doi.org/10.1371/journal.pone.0186124>
- Hughes, C. (2001). Executive dysfunction in autism: Its nature and implications for the everyday problems experienced by individuals with autism. In *The development of Autism: Perspectives from theory and research* (pp. 255–275). Lawrence Erlbaum Associates Publishers.
- Humphreys, K., Minshew, N., Leonard, G. L., & Behrmann, M. (2007). A fine-grained analysis of facial expression processing in high-functioning adults with autism. *Neuropsychologia*, 45(4), 685–695. <https://doi.org/10.1016/j.neuropsychologia.2006.08.003>
- Hunter, L. R., Buckner, J. D., & Schmidt, N. B. (2009). Interpreting facial expressions: The influence of social anxiety, emotional valence, and race. *Journal of Anxiety Disorders*, 23(4), 482–488. <https://doi.org/10.1016/j.janxdis.2008.10.004>
- Johnson, M. H., Jones, E. J. H., & Gliga, T. (2015). Brain adaptation and alternative developmental trajectories. *Development and Psychopathology*, 27(2), 425–442. <https://doi.org/10.1017/S0954579415000073>
- Joseph, R. M., Ehrman, K., McNally, R., & Keehn, B. (2008). Affective response to eye contact and face recognition ability in children with ASD. *Journal of the International Neuropsychological Society*, 14(6), 947–955. <https://doi.org/10.1017/S1355617708081344>
- Joseph, R. M., & Tager-Flusberg, H. (2004). The relationship of theory of mind and executive functions to symptom type and severity in children with autism. *Development and Psychopathology*, 16(1), 137–155. <https://doi.org/10.1017/S095457940404444X>
- Kenworthy, L., Yerys, B. E., Anthony, L. G., & Wallace, G. L. (2008). Understanding executive control in Autism spectrum disorders in the lab and in the real world. *Neuropsychology Review*, 18(4), 320–338. <https://doi.org/10.1007/s11065-008-9077-7>
- Kleinmans, N. M., Richards, T., Weaver, K., Johnson, L. C., Greenson, J., Dawson, G., & Aylward, E. (2010). Association between amygdala response to emotional faces and social anxiety in autism spectrum disorders. *Neuropsychologia*, 48(12), 3665–3670. <https://doi.org/10.1016/j.neuropsychologia.2010.07.022>
- Kohli, A., Malhotra, S., Mohanty, M., Khehra, N., & Kaur, M. (2005). Specific learning disabilities in children: Deficits and neuropsychological profile. *International Journal of Rehabilitation Research*, 28(2), 165–169.
- Koster, E. H. W., Crombez, G., Verschuere, B., & De Houwer, J. (2006). Attention to threat in anxiety-prone individuals: Mechanisms underlying attentional bias. *Cognitive Therapy and Research*, 30(5), 635–643. <https://doi.org/10.1007/s10608-006-9042-9>
- Lecce, S., Bianco, F., Devine, R. T., & Hughes, C. (2017). Relations between theory of mind and executive function in middle childhood: A short-term longitudinal study. *Journal of Experimental Child Psychology*, 163, 69–86. <https://doi.org/10.1016/j.jecp.2017.06.011>
- Lee, S. J., Lee, H.-K., Kwon, Y.-S., Lee, C. T., & Lee, K.-U. (2009). The impact of executive function on emotion recognition and emotion experience in patients with Schizophrenia. *Psychiatry Investigation*, 6(3), 156. <https://doi.org/10.4306/pi.2009.6.3.156>
- Leung, F. Y. N., Sin, J., Dawson, C., Ong, J. H., Zhao, C., Veić, A., & Liu, F. (2022). Emotion recognition across visual and auditory modalities in autism spectrum disorder: A systematic review and meta-analysis. *Developmental Review*, 63, 101000. <https://doi.org/10.1016/j.dr.2021.101000>
- Lievore, R., Cardillo, R., Lanfranchi, S., & Mammarella, I. C. (2022). Chapter four - social anxiety in children and adolescents with autism spectrum disorders. In R. M. Hodapp, D. J. Fidler, & S. Lanfranchi (Eds.), *International review of research in developmental disabilities*. (vol. 63, p. 131–186). Academic Press. <https://doi.org/10.1016/bs.irdd.2022.09.003>
- Lievore, R., Crisci, G., & Mammarella, I. C. (2023). Emotion recognition in children and adolescents with ASD and ADHD: A systematic review. *Review Journal of Autism and Developmental Disorders*. <https://doi.org/10.1007/s40489-023-00407-0>
- Lindner, J. L., & Rosén, L. A. (2006). Decoding of emotion through facial expression, prosody and verbal content in children and adolescents with Asperger's syndrome. *Journal of Autism and Developmental Disorders*, 36(6), 769–777. <https://doi.org/10.1007/s10803-006-0105-2>
- Livingston, E. M., Siegel, L. S., & Ribary, U. (2018). Developmental dyslexia: Emotional impact and consequences. *Australian Journal of Learning Difficulties*, 23(2), 107–135. <https://doi.org/10.1080/19404158.2018.1479975>
- Livingston, L. A., Colvert, E., & Happé, F. (2019). Good social skills despite poor theory of mind: Exploring compensation in autism spectrum disorder. *Journal of Child Psychology and Psychiatry*, 60(1), 102–110. <https://doi.org/10.1111/jcpp.12886>
- Lozier, L. M., Vanmeter, J. W., & Marsh, A. A. (2014). Impairments in facial affect recognition associated with autism spectrum disorders: A meta-analysis. *Development and Psychopathology*, 26(4pt1), 933–945. <https://doi.org/10.1017/S0954579414000479>
- Mammarella, I. C., Caviola, S., Giofrè, D., & Szűcs, D. (2018). The underlying structure of visuospatial working memory in children with mathematical learning disability. *British Journal of Developmental Psychology*, 36(2), 220–235. <https://doi.org/10.1111/bjdp.12202>
- Mammarella, I. C., Ghisi, M., Bomba, M., Bottesi, G., Caviola, S., Broggi, F., & Nacinovich, R. (2016). Anxiety and depression in children with nonverbal learning disabilities, reading disabilities, or typical development. *Journal of Learning Disabilities*, 49(2), 130–139. <https://doi.org/10.1177/0022219414529336>
- Mansell, W., Clark, D. M., Ehlers, A., & Chen, Y.-P. (1999). Social anxiety and attention away from emotional faces. *Cognition and Emotion*, 13(6), 673–690. <https://doi.org/10.1080/026999399379032>
- March, J. S. (2012). *Multidimensional anxiety scale for children* (2nd ed.). Multi-Health Systems.
- Marsh, A. A., Ambady, N., & Kleck, R. E. (2005). The effects of fear and anger facial expressions on approach- and avoidance-related behaviors. *Emotion*, 5(1), 119–124. <https://doi.org/10.1037/1528-3542.5.1.119>
- Martins, E. C., Osório, A., Veríssimo, M., & Martins, C. (2016). Emotion understanding in preschool children: The role of executive functions. *International Journal of Behavioral Development*, 40(1), 1–10. <https://doi.org/10.1177/0165025414556096>
- McNeil, D. W., & Randall, C. L. (2014). Conceptualizing and describing social anxiety and its disorders. In S. G. Hofmann, & P. M. DiBartolo (Eds.), *Social*



- anxiety: *Clinical, developmental, and social perspectives* (3rd ed. pp. 3–26). Academic Press.
- Meltzer, L. (2011). *Executive function in education, first edition: From theory to practice*. Guilford Press.
- Miles, J. (2005). R-squared, adjusted R-squared. In *Encyclopedia of statistics in behavioral science*. John Wiley & Sons, Ltd, <https://doi.org/10.1002/0470013192.bsa526>
- Miyake, A., Friedman, N. P., Emerson, M. J., Witzki, A. H., Howerter, A., & Wager, T. D. (2000). The unity and diversity of executive functions and their contributions to complex, frontal lobe, tasks: A latent variable analysis. *Cognitive Psychology*, 41(1), 49–100. <https://doi.org/10.1006/cogp.1999.0734>
- Moriguchi, Y. (2014). The early development of executive function and its relation to social interaction: A brief review. *Frontiers in Psychology*, 5, 388.
- Neumann, R., Völker, J., Hajba, Z., & Seiler, S. (2021). Lesions and reduced working memory impair emotion recognition in self and others. *Cognition and Emotion*, 35(8), 1527–1542. <https://doi.org/10.1080/02699931.2021.1983521>
- Nikolić, M. (2020). Disturbed social information processing as a mechanism in the development of social anxiety disorder. *Child Development Perspectives*, 14(4), 258–264. <https://doi.org/10.1111/cdep.12390>
- Nikolić, M., van der Storm, L., Colonesi, C., Brummelman, E., Kan, K. J., & Bögels, S. (2019). Are socially anxious children poor or advanced mindreaders? *Child Development*, 90(4), 1424–1441. <https://doi.org/10.1111/cdev.13248>
- O'Reilly, H., Pigat, D., Fridenson, S., Berggren, S., Tal, S., Golan, O., Bölte, S., Baron-Cohen, S., & Lundqvist, D. (2016). The EU-emotion stimulus set: A validation study. *Behavior Research Methods*, 48(2), 567–576. <https://doi.org/10.3758/s13428-015-0601-4>
- O'Toole, M. S., Hougaard, E., & Mennin, D. S. (2013). Social anxiety and emotion knowledge: A meta-analysis. *Journal of Anxiety Disorders*, 27(1), 98–108. <https://doi.org/10.1016/j.janxdis.2012.09.005>
- Operto, F. F., Pastorino, G. M. G., Stellato, M., Morcaldi, L., Vetri, L., Carotenuto, M., Viggiano, A., & Coppola, G. (2020). Facial emotion recognition in children and adolescents with specific learning disorder. *Brain Sciences*, 10(8), 473. <https://doi.org/10.3390/brainsci10080473>
- Ozili, P. K. (2023). The acceptable R-square in empirical modelling for social science research. In *Social research methodology and publishing results: A guide to non-native english speakers* (pp. 134–143). IGI Global. <https://doi.org/10.4018/978-1-6684-6859-3.ch009>
- Ozonoff, S. (1998). Assessment and remediation of executive dysfunction in autism and Asperger syndrome. In E. Schopler, G. B. Mesibov, & L. J. Kunce (Eds.), *Asperger syndrome or high-functioning Autism?* (pp. 263–289). Springer US. [https://doi.org/10.1007/978-1-4615-5369-4\\_12](https://doi.org/10.1007/978-1-4615-5369-4_12)
- Paloscia, C., Giangregorio, A., Guerini, R., & Melchiori, F. M. (2017). *MASC 2—multidimensional anxiety scale for children—manuale versione Italiana*. Hogrefe.
- Peirce, J., Gray, J. R., Simpson, S., MacAskill, M., Höchenberger, R., Sogo, H., Kastman, E., & Lindeløv, J. K. (2019). PsychoPy2: Experiments in behavior made easy. *Behavior Research Methods*, 51(1), 195–203. <https://doi.org/10.3758/s13428-018-01193-y>
- Phillips, L. H., Channon, S., Tunstall, M., Hedenstrom, A., & Lyons, K. (2008). The role of working memory in decoding emotions. *Emotion*, 8(2), 184–191. <https://doi.org/10.1037/1528-3542.8.2.184>
- R Core Team (2022). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.R-project.org>
- Richards, A., French, C. C., Calder, A. J., Webb, B., Fox, R., & Young, A. W. (2002). Anxiety-related bias in the classification of emotionally ambiguous facial expressions. *Emotion*, 2(3), 273–287. <https://doi.org/10.1037/1528-3542.2.3.273>
- Rosen, T. E., & Lerner, M. D. (2018). Error-related brain activity and anxiety symptoms in youth with autism spectrum disorder. *Autism Research*, 11(2), 342–354. <https://doi.org/10.1002/aur.1898>
- Rostami, M., Veisi, N., Jafarian Dehkordi, F., & Alkasir, E. (2014). Social anxiety in students with learning disability: Benefits of acceptance and commitment therapy. *Practice in Clinical Psychology*, 2(4), 277–284. <http://jpcp.uswr.ac.ir/article-1-201-en.html>
- Rump, K. M., Giovannelli, J. L., Minshew, N. J., & Strauss, M. S. (2009). The development of emotion recognition in individuals with Autism. *Child Development*, 80(5), 1434–1447. <https://doi.org/10.1111/j.1467-8624.2009.01343.x>
- Rutter, M., Le Couteur, A., & Lord, C. (2005). *Autism diagnostic interview – revised*. Giunti OS Special Organization.
- Sahoo, M. K., Biswas, H., & Padhy, S. K. (2015). Psychological co-morbidity in children with specific learning disorders. *Journal of Family Medicine and Primary Care*, 4(1), 21–25. <https://doi.org/10.4103/2249-4863.152243>
- Sartori, G., Job, R., & Tressoldi, P. E. (2007). *DDE-2. Batteria per la valutazione della dislessia e della disortografia evolutiva [battery for the assessment of developmental dyslexia and dysorthography]*. Giunti OS.
- Schuchardt, K., Maehler, C., & Hasselhorn, M. (2008). Working memory deficits in children with specific learning disorders. *Journal of Learning Disabilities*, 41(6), 514–523. <https://doi.org/10.1177/0022219408317856>
- Schultz, L. T., & Heimberg, R. G. (2008). Attentional focus in social anxiety disorder: Potential for interactive processes. *Clinical Psychology Review*, 28(7), 1206–1221. <https://doi.org/10.1016/j.cpr.2008.04.003>
- Sigurdardottir, H. M., Ívarsson, E., Kristinsdóttir, K., & Kristjánsson, Á. (2015). Impaired recognition of faces and objects in dyslexia: Evidence for ventral stream dysfunction? *Neuropsychology*, 29(5), 739–750. <https://doi.org/10.1037/neu0000188>
- Silvia, P. J., Allan, W. D., Beauchamp, D. L., Maschauer, E. L., & Workman, J. O. (2006). Biased recognition of happy facial expressions in social anxiety. *Journal of Social and Clinical Psychology*, 25(6), 585–602. <https://doi.org/10.1521/jscp.2006.25.6.585>
- Simonian, S. J., Beidel, D. C., Turner, S. M., Berkes, J. L., & Long, J. H. (2001). Recognition of facial affect by children and adolescents diagnosed with social phobia. *Child Psychiatry and Human Development*, 32(2), 137–145. <https://doi.org/10.1023/A:1012298707253>
- Singh, J., Arun, P., & Bajaj, M. K. (2021). Theory of mind and executive functions in children with attention deficit hyperactivity disorder and specific learning disorder. *Indian Journal of Psychological Medicine*, 43(5), 392–398. <https://doi.org/10.1177/0253717621999807>
- Sivaratnam, C., Newman, L., & Rinehart, N. (2018). Emotion-recognition and theory of mind in high-functioning children with ASD: Relationships with attachment security and executive functioning. *Research in Autism Spectrum Disorders*, 53, 31–40. <https://doi.org/10.1016/j.rasd.2018.05.005>
- Spain, D., Happé, F., Johnston, P., Campbell, M., Sin, J., Daly, E., Ecker, C., Anson, M., Chaplin, E., Glaser, K., Mendez, A., Lovell, K., & Murphy, D. G. (2016). Social anxiety in adult males with autism spectrum disorders. *Research in Autism Spectrum Disorders*, 32, 13–23. <https://doi.org/10.1016/j.rasd.2016.08.002>
- Spain, D., Sin, J., Linder, K. B., McMahon, J., & Happé, F. (2018). Social anxiety in autism spectrum disorder: A systematic review. *Research in Autism Spectrum Disorders*, 52, 51–68. <https://doi.org/10.1016/j.rasd.2018.04.007>
- Spikman, J. M., Boelen, D. H. E., Pijnenborg, G. H. M., Timmerman, M. E., van der Naalt, J., & Fasotti, L. (2013). Who benefits from treatment for executive dysfunction after brain injury? Negative effects of emotion recognition deficits. *Neuropsychological Rehabilitation*, 23(6), 824–845. <https://doi.org/10.1080/09602011.2013.826138>
- Sung, Y. Y., & Chang, M. (2010). Which social skills predict academic performance of elementary school students. *Journal On Educational Psychology*, 3(3), 23–34.
- Swanson, H. L., & Sachse-Lee, C. (2001). A subgroup analysis of working memory in children with reading disabilities: Domain-general or domain-specific deficiency? *Journal of Learning Disabilities*, 34(3), 249–263. <https://doi.org/10.1177/002221940103400305>
- Szűcs, D., Devine, A., Soltesz, F., Nobes, A., & Gabriel, F. (2013). Developmental dyscalculia is related to visuo-spatial memory and inhibition impairment. *Cortex*, 49(10), 2674–2688. <https://doi.org/10.1016/j.cortex.2013.06.007>
- Tarkiainen, A., Helenius, P., & Salmelin, R. (2003). Category-specific occipitotemporal activation during face perception in dyslexic individuals: An MEG study. *NeuroImage*, 19(3), 1194–1204. [https://doi.org/10.1016/S1053-8119\(03\)00161-7](https://doi.org/10.1016/S1053-8119(03)00161-7)



- Terras, M. M., Thompson, L. C., & Minnis, H. (2009). Dyslexia and psycho-social functioning: An exploratory study of the role of self-esteem and understanding. *Dyslexia*, 15(4), 304–327. <https://doi.org/10.1002/dys.386>
- Thaler, N. S., Kazemi, E., & Wood, J. J. (2010). Measuring anxiety in youth with learning disabilities: Reliability and validity of the multidimensional anxiety scale for children (MASC). *Child Psychiatry & Human Development*, 41(5), 501–514. <https://doi.org/10.1007/s10578-010-0182-5>
- Tibi-Elhanany, Y., & Shamay-Tsoory, S. G. (2011). Social cognition in social anxiety: First evidence for increased empathic abilities. *The Israel Journal of Psychiatry and Related Sciences*, 48(2), 98–106.
- Torske, T., Nærland, T., Øie, M. G., Stenberg, N., & Andreassen, O. A. (2018). Metacognitive aspects of executive function are highly associated with social functioning on parent-rated measures in children with Autism spectrum disorder. *Frontiers in Behavioral Neuroscience*, 11, 258.
- Trevisan, D. A., & Birmingham, E. (2016). Are emotion recognition abilities related to everyday social functioning in ASD? A meta-analysis. *Research in Autism Spectrum Disorders*, 32, 24–42. <https://doi.org/10.1016/j.rasd.2016.08.004>
- Tseng, H.-H., Huang, Y.-L., Chen, J.-T., Liang, K.-Y., Lin, C.-C., & Chen, S.-H. (2017). Facial and prosodic emotion recognition in social anxiety disorder. *Cognitive Neuropsychiatry*, 22(4), 331–345. <https://doi.org/10.1080/13546805.2017.1330190>
- Uljarevic, M., & Hamilton, A. (2013). Recognition of emotions in autism: A formal meta-analysis. *Journal of Autism and Developmental Disorders*, 43(7), 1517–1526. <https://doi.org/10.1007/s10803-012-1695-5>
- Van Kleef, G. A. (2009). How emotions regulate social life: The emotions as social information (EASI) model. *Current Directions in Psychological Science*, 18(3), 184–188. <https://doi.org/10.1111/j.1467-8721.2009.01633.x>
- Waters, A. M., Mogg, K., Bradley, B. P., & Pine, D. S. (2011). Attention bias for angry faces in children with social phobia. *Journal of Experimental Psychopathology*, 2(4), 475–489. <https://doi.org/10.5127/jep.018111>
- Wechsler, D. (2003). *Wechsler intelligence scale for children—fourth edition (WISC-IV)*. The Psychological Corporation.
- White, S. W., Albano, A. M., Johnson, C. R., Kasari, C., Ollendick, T., Klin, A., Oswald, D., & Scahill, L. (2010). Development of a cognitive-behavioral intervention program to treat anxiety 55 social anxiety iand social deficits in teens with high-functioning autism. *Clinical Child and Family Psychology Review*, 13(1), 77–90. <https://doi.org/10.1007/s10567-009-0062-3>
- White, S. W., Maddox, B. B., & Panneton, R. K. (2015). Fear of negative evaluation influences eye gaze in adolescents with autism spectrum disorder: A pilot study. *Journal of Autism and Developmental Disorders*, 45(11), 3446–3457. <https://doi.org/10.1007/s10803-014-2349-6>
- Whiting, P. R., & Robinson, G. L. (2001). The interpretation of emotion from facial expressions for children with a visual sub-type of dyslexia. *Australian Journal of Learning Disabilities*, 6(4), 6–14. <https://doi.org/10.1080/19404150109546682>
- Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. Springer-Verlag. <https://ggplot2.tidyverse.org>
- Wong, N., Beidel, D. C., Sarver, D. E., & Sims, V. (2012). Facial emotion recognition in children with high functioning Autism and children with social phobia. *Child Psychiatry & Human Development*, 43(5), 775–794. <https://doi.org/10.1007/s10578-012-0296-z>
- World Health Organization (1992). *The ICD-10 classification of mental and behavioural disorders: Clinical descriptions and diagnostic guidelines*. Author.
- Yeung, M. K. (2022). A systematic review and meta-analysis of facial emotion recognition in autism spectrum disorder: The specificity of deficits and the role of task characteristics. *Neuroscience & Biobehavioral Reviews*, 133, 104518. <https://doi.org/10.1016/j.neubiorev.2021.104518>