

COMPLETENESS OF TWO THEORIES ON ORDERED ABELIAN GROUPS AND EMBEDDING RELATIONS

YUICHI KOMORI

§0. Introduction

The first order language \mathcal{L} that we consider has two nullary function symbols $0, 1$, a unary function symbol $-$, a binary function symbol $+$, a unary relation symbol $0 <$, and the binary relation symbol $=$ (equality). Let \mathcal{L}' be the language obtained from \mathcal{L} , by adding, for each integer $n > 0$, the unary relation symbol $n|$ (read " n divides"). The terms $1 + \cdots + 1$ and $t + \cdots + t$ (1 and t repeated n times) will be written as n and nt , the term $t + (-s)$ as $t - s$, the atomic formula $0 < t - s$ as $t < s$, and the formulas $u < t \wedge t < s$ and $t = s \vee t < s$ as $u < t < s$ and $t \leq s$, respectively. We now give some axiom systems for abelian groups with a semidiscrete total ordering.

(a) The axioms for abelian groups:

$$(x + y) + z = x + (y + z)$$

$$x + y = y + x$$

$$x + 0 = x$$

$$x - x = 0.$$

(b) The axioms for a total ordering compatible with group structures:

$$0 < x \wedge 0 < y \rightarrow 0 < x + y$$

$$\neg(0 < x \wedge 0 < -x)$$

$$x = 0 \vee 0 < x \vee 0 < -x.$$

(c) The axioms for a semi-discrete ordering:

$$0 < 1$$

$$2x < 1 \vee 1 < 2x.$$

(d) The axioms for infinitesimals:

$$2x < 1 \rightarrow nx < 1 \text{ for each } n > 2.$$

The axioms for $n|$ are

Received October 26, 1978.

- (e) $n|x \leftrightarrow \exists y \exists z (-1 < 2z < 1 \wedge x = ny + z)$ for each $n > 0$, and
- (f) $n|x \vee n|x + 1 \vee n|x + 2 \vee \dots \vee n|x + n - 1$ for each $n > 1$.
- (g) The axioms for divisible infinitesimals:
 $-1 < 2x < 1 \rightarrow \exists y (x = ny)$ for each $n > 1$.
- (h) The axiom for discrete orderings:
 $\neg (0 < x < 1)$.
- (i) The axiom for existence of infinitesimals:
 $\exists x (0 < x < 1)$.

The language of the theory \mathcal{S} is \mathcal{L} . The set of axioms of \mathcal{S} is $(a) \cup (b) \cup (c) \cup (d)$ (We will write (a, b, c, d) for this set in future.). The language of the theory \mathcal{D} is \mathcal{L} as well. The set of axioms of \mathcal{D} is $\mathcal{S} \cup (h)$ (equivalent to (a, b, c, h)), which is equivalent to $(a, b) \cup \{0 < x \leftrightarrow x = 1 \vee 1 < x\}$. We call a model of \mathcal{S} (or \mathcal{D}) an *abelian group with a semi-discrete (or discrete) total ordering*. The languages of the theories \mathcal{SS} , \mathcal{SC} and \mathcal{DC} are all \mathcal{L}' . The sets of axioms of \mathcal{SS} , \mathcal{SC} and \mathcal{DC} are $\mathcal{S} \cup (e, f)$, $\mathcal{SS} \cup (g, i)$ and $\mathcal{SS} \cup (g, h)$ (equivalent to $\mathcal{D} \cup (e, f)$), respectively.

Let Z and Q be the set of integers and the set of rationals, respectively. Consider the group $ZQ = Z \times Q$ ordered as follows: $0 < (x, y)$ if and only if either $0 < x$ or $x = 0$ and $0 < y$. ZQ is a model of \mathcal{S} which contains Z as a submodel (identifying $(n, 0)$ with n). Of course, Z is a model of \mathcal{D} , but ZQ is not a model of \mathcal{D} . It is clear that each model of \mathcal{S} is also a model of (e), or, more precisely, given a model of \mathcal{S} , there is a unique value of $n|$ so that (e) is satisfied. ZQ is a model of \mathcal{SC} as well.

It is known that \mathcal{DC} allows elimination of quantifiers and that it is complete (cf. Kreisel and Krivine [2] p. 54. There exists an error in the proof, but it is easy to correct the error.). In § 1, we shall show that \mathcal{SC} allows elimination of quantifiers and that it is complete and decidable.

In § 2, we shall show that any model of \mathcal{S} (\mathcal{SS} , \mathcal{D}) can be embedded in some model of \mathcal{SS} (\mathcal{SC} , \mathcal{DC} , respectively). Embedding relations will be used to show some results on the first order semantics for \mathcal{L} and \mathcal{L}' . One of them is that for any universal formula F of \mathcal{L}' and any model A of $\mathcal{SS} \cup (i)$, F is valid in ZQ if and only if F is valid in A . In the paper [1], we shall make use of it for giving a complete description of super-Lukasiewicz propositional logics.

Z is a model of $\mathcal{SS} \cup (g)$, but not a model of \mathcal{SC} . Therefore, (i) is not a consequence of the set (a, b, c, d, e, f, g) . Consider the group $ZZ = Z \times Z$ ordered as follows: $0 < (x, y)$ if and only if either $0 < x$ or $x = 0$

and $0 < y$. This group (identifying $(n, 0)$ with n) is a model of $SS \cup (i)$, but not a model of SC . Hence, (g) is not a consequence of the set (a, b, c, d, e, f, i).

§1. Elimination of quantifiers

To show that SC admits elimination of quantifiers, we consider a formula F of the form $\exists x(\alpha_1 \wedge \dots \wedge \alpha_n)$ where each α_i is an atomic formula of \mathcal{L}' or the negation of an atomic formula of \mathcal{L}' . Thus α_i is of one of the forms $t = s, t \neq s, 0 < t, \neg(0 < t), n|t$ or $\neg(n|t)$.

In this section, the derivations from SS are done without notice. As we shall prove by the way that DC admits elimination of quantifiers, we notice the use when we use (g) or (i). $t = s, t \neq s, \neg(0 < t)$ and $\neg(n|t)$ are equivalent to $t - s = 0, 0 < t - s \vee 0 < s - t, t = 0 \vee 0 < -t$ and $n|t + 1 \vee \dots \vee n|t + n - 1$, respectively. Hence we can suppose that each α_i is of one of the forms $t = 0, 0 < t$ or $n|t$.

Each term t can be written in the form $px + s$ with $p \in \mathbb{Z}$ and s a term which does not contain x . If $p = 0$, the atomic formula can be taken out of the scope of $\exists x$.

Thus the formula F can be written in the form

$$(*) \quad \exists x(p_1x < t_1 \wedge \dots \wedge p_jx < t_j \wedge u_1 < q_1x \wedge \dots \wedge u_k < q_kx \wedge r_1x = v_1 \wedge \dots \wedge r_lx = v_l \wedge n_1|s_1x - w_1 \wedge \dots \wedge n_m|s_mx - w_m)$$

where the p, q, r, s are in N (the set of natural numbers which does not contain 0) and t, u, v, w are terms which do not contain x .

For any $k \in N, t = u, t < u$ and $n|t$ are equivalent to $kt = ku, kt < ku$ and $kn|kt$, respectively. Hence, taking the least common multiple (l.c.m.) of $p_1, \dots, p_j, q_1, \dots, q_k, r_1, \dots, r_l, s_1, \dots, s_m$ we can suppose that $p_1 = \dots = p_j = q_1 = \dots = q_k = r_1 = \dots = r_l = s_1 = \dots = s_m = p$ in the formula (*). Then the formula (*) is equivalent to

$$\exists x(x < t_1 \wedge \dots \wedge x < t_j \wedge u_1 < x \wedge \dots \wedge u_k < x \wedge x = v_1 \wedge \dots \wedge x = v_l \wedge n_1|x - w_1 \wedge \dots \wedge n_m|x - w_m \wedge \exists y(x = py)).$$

It follows from $SS \cup (g)$ that $\exists y(x = py)$ is equivalent to $p|x$. If $l \geq 1$, then the above formula equivalent to

$$v_1 < t_1 \wedge \dots \wedge v_1 < t_j \wedge u_1 < v_1 \wedge \dots \wedge u_k < v_1 \wedge v_1 = v_2 \wedge \dots \wedge v_1 = v_l \wedge n_1|v_1 - w_1 \wedge \dots \wedge n_m|v_1 - w_m \wedge p|v_1$$

which is quantifier free. Hence we can assume that $l = 0$. If $j \geq 2$, the formula F is equivalent to

$$(t_1 < t_2 \wedge \exists x(x < t_1 \wedge x < t_3 \wedge \dots)) \vee (t_2 < t_1 \wedge \exists x(x < t_2 \wedge x < t_3 \wedge \dots))$$

and we are reduced to the case of a formula with $j - 1$ atomic formulas of the form $x < t$. Hence we can assume that $j \leq 1$. Similarly, we can assume that $k \leq 1$.

Suppose that $j = k = 1$. Let n be l.c.m. of n_1, n_2, \dots, n_m, p . Let C_q be the formula

$$\exists y \exists z (0 < ny + z + q < t_1 - u_1 \wedge -1 < 2z < 1) \wedge n_1 | q + u_1 - w_1 \wedge \dots \wedge n_m | q + u_1 - w_m \wedge p | q + u_1 \quad (q = 0, 1, 2, \dots, n - 1).$$

Then, F is equivalent to $C_0 \vee C_1 \vee \dots \vee C_{n-1}$.

It suffices to show that $\exists y \exists z (0 < ny + z + q < t \wedge -1 < 2z < 1)$ is equivalent to some quantifier free formula. It follows from $SS \cup (g)$ that it is equivalent to $\exists y (0 < ny + q < t)$. If $q = 0$, then it is equivalent to $0 < t$ in SC (equivalent to $n < t$ in DC). If $0 < q \leq n - 1$, then it is equivalent to $-1 < 2(t - q)$ in SC (equivalent to $q < t$ in DC).

When $j = 0$ or $k = 0$, F is equivalent to $E_0 \vee E_1 \vee \dots \vee E_{n-1}$ where E_q is the formula $n_1 | q + u_1 - w_1 \wedge \dots \wedge n_m | q + u_1 - w_m \wedge p | q + u_1$.

This completes our proof that SC (and DC) allows elimination of quantifiers.

Because any atomic formula without variables is equivalent to $0 = 0$ or $0 \neq 0$, any quantifier free formula without variables is equivalent to $0 = 0$ or $0 \neq 0$. Hence any closed formula is equivalent to $0 = 0$ or $0 \neq 0$. Therefore SC (and DC) is complete.

THEOREM 1.1. *Both theories SC and DC allow elimination of quantifiers, and they are complete and decidable.*

§2. Embedding relations

THEOREM 2.1. *Any model of SS can be embedded in some model of SC .*

Proof. Let A be a model of SS . When A satisfies (h), we consider $A \times Q$ ordered as $0 < (x, y)$ if and only if either $0 < x$ or $x = 0$ and $0 < y$. Then $A \times Q$ is a model of SC and the mapping $f: A \rightarrow A \times Q$ such that $f(x) = (x, 0)$ is an embedding of A in $A \times Q$. Suppose A does not satisfy (h), that is, satisfies (i). Let B be the set $\{(x, n) | n | x \text{ and } n > 0 \text{ and } x \in A\}$.

We define functions $-$, $+$ and a relation $0 <$ on B as follows: $-(x, n) = (-x, n)$, $(x, m) + (y, n) = (nx + my, mn)$, $0 < (x, n)$ if and only if $0 < x$. The relation \sim on B defined by $(x, m) \sim (y, n)$ if and only if $nx = my$ is a congruence relation. B/\sim is a model of SC and the mapping $g: A \rightarrow B/\sim$ such that $g(x) = [(x, 1)]$ (equivalence class containing $(x, 1)$) is an embedding of A in B/\sim . Q.E.D.

By the Embedding Theorem (cf. [2] p. 40) and the fact that ZQ is a model of complete theory SC , we have

COROLLARY 2.2. *For any universal formula F of \mathcal{L}' , F is valid in ZQ if and only if F is valid in every model of SS .*

The following corollary is used for giving a complete description of super-Lukasiewicz propositional logics in a subsequent paper [1].

COROLLARY 2.3. *For any universal formula F of \mathcal{L}' and any model A of $SS \cup (i)$, F is valid in ZQ if and only if F is valid in A .*

Proof. By Corollary 2.2, F is valid in A if F is valid in ZQ . Conversely, suppose that F is valid in A . ZZ can be embedded in any model of $SS \cup (i)$. Hence F is valid in ZZ . Any finitely generated submodel of ZQ is isomorphic to Z or ZZ . Hence it can be embedded in ZZ . Therefore, F is valid in ZQ . Q.E.D.

LEMMA 2.4. *For any model A of S and any elements x_1, x_2, \dots, x_q of A , if $m + n_1x_1 + \dots + n_qx_q = 0$, then $m = n_1 = \dots = n_q = 0$ or there exist elements y_1, y_2, \dots, y_{q-1} of A and integers k_{ij} ($1 \leq i \leq q, 0 \leq j \leq q - 1$) such that $x_i = k_{i0} + \sum_{j=1}^{q-1} k_{ij}y_j$ for any i ($1 \leq i \leq q$).*

Proof. Choosing the signs of x_1, x_2, \dots, x_q suitably, we can assume that n_1, n_2, \dots, n_q are positive or zero. We prove this lemma by induction on $n_1 + n_2 + \dots + n_q$.

Case 1. Suppose that at least two of n_1, n_2, \dots, n_q are positive. We can assume that $0 < n_1 \leq n_2$. Then we have

$$m + n_1(x_1 + x_2) + (n_2 - n_1)x_2 + n_3x_3 + \dots + n_qx_q = 0.$$

By the hypothesis of induction, there exist elements y_1, \dots, y_{q-1} of A and integers p_{ij} such that

$$x_1 + x_2 = p_{10} + \sum_{j=1}^{q-1} p_{1j} y_j \quad \text{and} \quad x_i = p_{i0} + \sum_{j=1}^{q-1} p_{ij} y_j \quad (i = 2, 3, \dots, q).$$

Then we have $x_1 = (p_{10} - p_{20}) + \sum_{k=1}^{q-1} (p_{1k} - p_{2k}) y_k$. It completes our proof of Case 1 that we put $k_{1j} = p_{1j} - p_{2j}$ ($0 \leq j \leq q - 1$) and $k_{ij} = p_{ij}$ ($2 \leq i \leq q, 0 \leq j \leq q - 1$).

Case 2. We can assume that $n_1 \neq 0$ and $n_2 = n_3 = \dots = n_q = 0$. Since it follows from S that $\forall x(mx \neq n)$ for every mutually prime integers m, n , n_1 is a factor of m . Hence, we have $x_1 = -m/n_1$. Put $y_i = x_{i+1}$ ($i = 1, 2, \dots, q - 1$). Q.E.D.

THEOREM 2.5. *Any model of S can be embedded in some model of SS .*

Proof. It suffices to show that any model of S generated by a finite set can be embedded in some model of SS (cf. Theorem 13 in [1] p. 41). Let A be a model of S which have n generators a_1, a_2, \dots, a_n but can not be generated by $n - 1$ generators. We define a relation $0 <$ on $Z \times Q^n$ as follows: $0 < (m, q_1, \dots, q_n)$ if and only if $0 < pm + (pq_1)a_1 + \dots + (pq_n)a_n$ where p is l.c.m. of denominators of q_1, \dots, q_n ,

In order to prove that $Z \times Q^n$ is a model of SS , it suffices to show that $m = k_1 = \dots = k_n = 0$ if $m + k_1 a_1 + \dots + k_n a_n = 0$. By Lemma 2.4, a_1, a_2, \dots, a_n can be generated by $n - 1$ generators if $m + k_1 a_1 + \dots + k_n a_n = 0$, and $m \neq 0$ or $k_i \neq 0$ for some i . Hence, $Z \times Q^n$ is a model of SS .

Let f be a function from A into $Z \times Q^n$ such that $f(m + k_1 a_1 + \dots + k_n a_n) = (m, k_1, \dots, k_n)$. Then f is an embedding of A in $Z \times Q^n$. Q.E.D.

By Theorem 2.1 and Theorem 2.5, we have

THEOREM 2.6. *Any model of S can be embedded in some model of SC .*

The following corollary can be proved similarly to Corollary 2.3.

COROLLARY 2.7. *For any universal formula F of \mathcal{L} and any model A of $S \cup (i)$, F is valid in ZQ if and only if F is valid in A .*

We can prove the following theorem quite similarly to Theorem 2.5.

THEOREM 2.8. *Any model of D can be embedded in some model of DC .*

Since any model of D has a submodel Z which is a model of DC , we have the following corollary.

COROLLARY 2.9. *For any universal formula F of \mathcal{L} and any model A of D , F is valid in Z if and only if F is valid in A .*

Any model of $(a, b) \cup \{0 < 1\}$, that is, any totally ordered abelian group with 1 contains Z as a submodel. Hence, for any model A of $(a, b) \cup \{0 < 1\}$, the set of open formulas valid in A is included in the set of open formulas valid in Z which equals to the set of open theorems of D . The theory D is an open theory, that is, axiomatizable by only open formulas. Therefore, we have

THEOREM 2.10. *The theory D is the greatest element of the class of open consistent theories in \mathcal{L} containing $(a, b) \cup \{0 < 1\}$.*

REFERENCES

- [1] Y. Komori, Super-Lukasiewics propositional logics, to appear.
- [2] G. Kreisel and J. L. Krivine, Elements of Mathematical Logic, North-Holland, Amsterdam, 1967.

Shizuoka University