OCTOBER 1987 VOLUME 8, NUMBER 10



EDITORIAL

The Relationship Between APIC and SHEA: "Closely Watched Trains"
George W. Counts, MD

ORIGINAL ARTICLES

Survey on Testing Criteria and Reporting Methods for Human Immunodeficiency Virus Serologic Tests in Veterans Administration Medical Centers Susan Mottice, PhD; Sego Matsumiya, RN; Larry Reimer, MD

Inactivation of Human Immunodeficiency Virus by BetadineJoan C. Kaplan, PhD; David C. Crawford, BS; Amy G. Durno, BS;
Robert T. Schooley, MD


Diffusion and Adoption of CDC Guidelines for the Prevention and Control of Nosocomial Infections in US Hospitals
David D. Celentano, ScD; Laura L. Morlock, PhD; Faye E. Malitz, BS

The Detection of Blood on Nonporous Environmental Surfaces: An Approach for Assessing Factors Contributing to the Risk of Occupational Exposure to Blood in the Autopsy Suite Lorraine R. Beaumont, DrPH

Brief Report: An Outbreak of Scabies Among Employees in a Hospital-associated Commercial Laundry
Michelle C. Thomas; Dale H. Giedinghagen; Gerald L. Hoff, PhD

SPECIAL SECTIONS

Topics in Clinical Microbiology: Vibrio vulnificus Hilda Ratner, BA, MA



Protection Solution to High-Risk Situations



EPA ACCEPTED: For complete disinfection (including T.B. activity)

"SPORICIDIN, DILUTED 1:16, IS TUBERCULOCIDAL AND CAN BE USED AND RE-USED FOR 30 DAYS FOR 100% HOSPITAL LEVEL DISINFECTION IN 10 MINUTES AT TRUE ROOM TEMPERATURE, **68°F AND ABOVE•** (REG. #8383-5)."

Sporicidin is the **ONLY** dilutable glutaraldehyde-based 10 minute Important Product Information: disinfectant that is registered for tuberculocidal re-use without heating. Respiratory and other equipment were used in the studies supporting the re-use claims. It is recommended for respiratory therapy and operating room instruments and equipment.

Sporicidin is recommended by all major scope manufacturers.

**Cidex now must be heated to 77°F / 25°C for 45 minutes for tuberculocidal activity. Cidex, and Cidex type products, require a much higher glutaraldehyde concentration (2%) for disinfection and are less effective than Sporicidin, a glutaraldehyde (0.13%)-phenate complex.

Under the most stringent and scientific guidelines, Sporicidin is the best sterilizing/disinfecting product available.



utio l. hm.h, no. Curtis L. Lynch, M.D., M.P.H. Medical Director (800) 424-3733



INFECTION CONTROL®

Table of Contents

EDITORIAL

The Relationship Between APIC and SHEA: 'Closely Watched Trains" George W. Counts, MD			404
ORIGINAL ARTICLES			
Survey on Testing Criteria and Reporting Methods for Human Immunodeficiency Virus Serologic Tests in Veterans Administration Medical Centers Susan Mottice, PhD; Sego Matsumiya, RN; Larry Reimer, MD			407
Inactivation of Human Immunodeficiency Virus by Betadine Joan C. Kaplan, PhD; David C. Crawford, BS; Amy G. Durno, BS; Robert T. Schooley, MD			412
Diffusion and Adoption of CDC Guidelines for the Prevention and Control of Nosocomial Infections in US Hospitals David D. Celentano, ScD; Laura L. Morlock, PhD; Faye E. Malitz, BS			415
The Detection of Blood on Non An Approach for Assessing Fac Occupational Exposure to Bloo Lorraine R. Beaumont, DrPH	tors Co	ntributing to the Risk of	424
Brief Report: An Outbreak of Scabies Among Employees in a Hospital-associated Commercial Laundry Michelle C. Thomas; Dale H. Giedinghagen; Gerald L. Hoff, PhD			427
SPECIAL SECTIONS			
Topics in Clinical Microbiology: <i>Vibrio vulnificus</i> Hilda Ratner, BA, MA			430
DEPARTMENTS			
Information for Authors	399	Book Reviews	434
Letters to the Editor	400	Calendar of Events	436

The ideas and opinions expressed by contributing authors do not necessarily reflect those of the editors or publisher.

Publisher: Infection Control (ISSN-0195-9417) is published monthly by SLACK Incorporated, 6900 Grove Road, Thorofare, New Jersey 08086. Telephone: (609) 848-1000.

Copyright 1987: All rights reserved. No part of this publication may be reproduced without written permission from the publisher.

Subscriptions. Requests should be addressed to the publisher (except Japan). In Japan, contact Woodbell Scope Incorporated, Mansui Bldg., 9-18, Kanda Surugadai 2-chome, Chiyoda-ku, Tokyo 101, Japan. Subscription rates in the US and possessions: Individual: One year—\$40.00; Two years—\$66.00; Three years—\$94.00. Institutional: One year—\$55.00; Two years—\$92.00; Three years—\$128.00; Canada: \$15.00 additional each year; all other countries: \$25.00 additional each year. Single copies of current issues may be obtained for \$7.00, United States and possessions; \$13.00 all other countries.

Change of address: Notice should be sent to the publisher six weeks in advance of effective date. Include old and new addresses with zip codes. The publisher cannot accept responsibility for undelivered copies. Second-class postage is paid at Thorofare, New Jersey 08086. Postmaster: Send address changes to SLACK Incorporated, 6900 Grove Road, Thorofare, NJ 08086.

As of Volume 1, Number 1, INFECTION CONTROL is listed in Index Medicus, Current Contents—Clinical Practice, Hospital Literature Index, and Cumulative Index to Nursing and Allied Health Literature.

INFECTION CONTROL® / Editorial Board

EDITOR

Richard P. Wenzel, MD, MSc Iowa City, Iowa

SENIOR ASSOCIATE EDITOR

William Schaffner, MD Nashville, Tennessee

ASSOCIATE EDITORS

Sue Crow, MSN, RN Shreveport, Louisiana

John E. McGowan, Jr., MD Atlanta, Georgia

Dennis G. Maki, MD Madison, Wisconsin

SLACK

SLACK Incorporated 6900 Grove Road Thorofare, New Jersey 08086

Publisher Richard N. Roash

Editorial Director Donna Carpenter

Senior Editor/Development John C. Carter

Editorial Coordinator Sandra L. Patterson

Assistant Editor Dawn A. Zwick

Editorial Assistant Mary Lou Jerrell

Production Manager

Jim Griffin

Production Coordinator Lester Robeson

Circulation Manager Kevin J. Fenton

Advertising Manager Randall Roash

Advertising Sales Representative Wayne M. McCourt

Advertising Sales Coordinator Betty Martz

Classified Advertising Representative

EDITORIAL OFFICES

C41 General Hospital University of Iowa Hospitals and Clinics Iowa City, Iowa 52242 (319) 356-0464

William J. Hausler, Jr., PhD R. Michael Massanari, MD, MS

Michael Pfaller, MD Robert F. Woolson, PhD

Managing Editor Marikay Klein

EDITORIAL ADVISORY BOARD

Robert C. Aber, MD Hershey, Pennsylvania

Paul Arnow, MD Chicago, Illinois

Charles S. Bryan, MD Columbia, South Carolina

John P. Burke, MD Salt Lake City, Utah

Mary Castle, MPH, RN Berkeley, California

Marie B. Coyle, PhD Seattle, Washington

William Craig, MD Madison, Wisconsin

Burke A. Cunha, MD Mineola, New York

Richard E. Dixon, MD Trenton, New Jersey

Leigh G. Donowitz, MD Charlottesville, Virginia

Mark Eggleston, PharmD Washington, DC

Harvey A. Elder, MD Loma Linda, California

Bruce Farber, MD Pittsburgh, Pennsylvania

Martin S. Favero, PhD Atlanta, Georgia

Peter C. Fuchs, MD, PhD Portland, Oregon

Richard A. Garibaldi, MD Farmington, Connecticut

Angella Goetz, MNEd, RN Pittsburgh, Pennsylvania

Donald A. Goldmann, MD Boston, Massachusetts

Dieter H.M. Gröschel, MD Charlottesville, Virginia Peter A. Gross, MD Hackensack, New Jersey

Karen Hadley, MPH, RN New Orleans, Louisiana

David K. Henderson, MD Bethesda, Maryland

Peter N.R. Heseltine, MD Los Angeles, California

Cyrus C. Hopkins, MD Boston, Massachusetts

Allen B. Kaiser, MD Nashville, Tennessee

Donald L. Kaiser, DrPH Charlottesville, Virginia

Elaine Larson, PhD Cabin John, Maryland

Harold Laufman, MD, PhD New York, New York

Barbara McArthur, PhD, RN Detroit, Michigan

Rob Roy MacGregor, MD Philadelphia, Pennsylvania

C. Glen Mayhall, MD Richmond, Virginia

Patti J. Miller, MS Charlottesville, Virginia

Harry C. Nottebart, Jr., JD, MD London, England Richmond, Virginia

Michael T. Osterholm, PhD, MPH Minneapolis, Minnesota

James E. Peacock, Jr., MD Winston-Salem, North Carolina

Frank S. Rhame, MD Minneapolis, Minnesota

William A. Rutala, PhD, MPH Chapel Hill, North Carolina

William E. Scheckler, MD Madison, Wisconsin

Walter E. Stamm, MD Seattle, Washington

Charles W. Stratton, MD Nashville, Tennessee

Richard L. Sweet, MD San Francisco, California

George H. Talbot, MD Philadelphia, Pennsylvania

Timothy R. Townsend, MD Baltimore, Maryland

James Veazey, MD Albany, New York

FOREIGN ADVISORY BOARD

Graham Ayliffe, MD, FRCPath. Birmingham, England

Professor G. Berencsi Szeged, Hungary

David Birnbaum, MPH Sidney, British Columbia, Canada

Professor Jaap Dankert Groningen, Netherlands

Professor Dr. F. Daschner Freiburg, West Germany

Lars O. Kallings, MD Stockholm, Sweden

Professor W.B. Kędzia Sieroca, Poland

Professor A.P. Krasilnikow Minsk, USSR

Professor Dr. W. Marget Munich, West Germany

Jonathan L. Meakins, MD, FRCSC, FACS Montreal, Canada

Bertil Nyström, MD Huddinge, Sweden

Ian Phillips, MA, MD, MRCPath. London, England

Samuel Ponce de Leon, MD Mexico City, Mexico

Hans Reber, MD Basel, Switzerland

Professor Gerald Reybrouck Leuven, Belgium

Manfred L. Rotter, MD, DipBact Vienna, Austria

Theodore Sacks, MD Jerusalem, Israel

Dr. Bernhard M. Thimm Federal Republic of Germany

Professor Dr. med. H.P. Werner Mainz, West Germany

Professor Dr. W. Weuffen Greifswald, German Democratic Republic

https://doi.org/10.7017/50195941700066492 Published online by Cambridge University Press

TOUGH ON STAPH

Solve 69% of Tomorrow's Problems.

The need for an effective, cost efficient method of infection control of catheter-associated UTI has been answered. Introducing the CURITY® Infection Control Foley Catheterization System. No other system has been clinically proven to be as effective in reducing the incidence of UTI.

Clinically Proven to **Substantially Reduce** UTI!

The Curity infection control system has been clinically proven to reduce the incidence of UTI by an astounding 69%!

In vivo studies conducted at the University of Cincinnati Medical

https://doi.org/10.1017/S0195941700066492 Published online by Ca

Center, and presented at the 86th Annual Meeting of the American Society for Microbiology, show the Curity infection control system to effectively reduce the incidence of UTI by 69% in patients catheterized from 2-30 days. In addition, the Curity test group had more total days free of infection than the control group.1

Why the System is So Effective.

The reason the Curity® system works like no other system is because it is like no other system. Three major preventive measures against infection are incorporated into the unique system design:

1. A Retrograde Contamination Guard which prevents bacteria from entering the drain port.

2. The CURITY® Ultramer™ catheter, which offers greater lubricity than other coated or uncoated catheters.2


3. A Tamper Discouraging Seal which discourages disconnection at the sterile catheter-tubing iunction.

Together they present an imposing set of

barriers against contamination of the urinary drainage system.

What It All Means to You.

What this means to you is now, for the first time, you have a new weapon in the fight against UTI.








The Retrograde Contamination Guard, the CURITY $^{\oplus}$ Ultramer $^{\leftarrow}$ catheter and the Tamper Discouraging Seal are three major preventive measures against infection. Together they present an imposing set of barriers against contamination of the urinary drainage system.

A weapon that could mean substantial savings to both hospital and patient. For example, a 69% reduction in the incidence of UTI could save the average 250 bed hospital up to \$127,660 in treatment costs annually.3 Much of which might not be reimbursible under DRG's.

To find out how much your hospital can save while improving patient care, call or write:

> The Kendall Company Hospital Products One Federal Street Boston, MA 02101 1-800-225-2600 (in MA: 1-800-882-2000).



REFERENCES:

- 1. Cicmanec, J. and Al-Juburi, A.Z.: "Evaluation of the CURITY Retrograde Contamination Guard Urinary Drainage System in Preventing Catheter Associated Urinary Tract Infections. Presented at the 86th Annual Meeting of the American Society for Microbiology, Washington, D.C., March 1986.
- 2. Supportive material available on request.
- 3. Supportive material available on request.

