



The restricted quantum double of the Yangian

Curtis Wendlandt

Abstract. Let \mathfrak{g} be a complex semisimple Lie algebra with associated Yangian $Y_{\hbar}\mathfrak{g}$. In the mid-1990s, Khoroshkin and Tolstoy formulated a conjecture which asserts that the algebra $DY_{\hbar}\mathfrak{g}$ obtained by doubling the generators of $Y_{\hbar}\mathfrak{g}$, called the Yangian double, provides a realization of the quantum double of the Yangian. We provide a uniform proof of this conjecture over $\mathbb{C}[[\hbar]]$ which is compatible with the theory of quantized enveloping algebras. As a by-product, we identify the universal R -matrix of the Yangian with the canonical element defined by the pairing between the Yangian and its restricted dual.

1 Introduction

This article is a continuation of [41], which studied the Yangian double $DY_{\hbar}\mathfrak{g}$ associated with an arbitrary symmetrizable Kac–Moody algebra \mathfrak{g} through the lens of a \mathbb{Z} -graded algebra homomorphism

$$\Phi_z : DY_{\hbar}\mathfrak{g} \rightarrow LY_{\hbar}\widehat{\mathfrak{g}}_z \subset Y_{\hbar}\mathfrak{g}[[z^{\pm 1}]].$$

Here, $LY_{\hbar}\widehat{\mathfrak{g}}_z$ is a naturally defined \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -algebra, described explicitly in Lemma 4.5, and $Y_{\hbar}\mathfrak{g}$ is the Yangian of \mathfrak{g} , defined over $\mathbb{C}[[\hbar]]$. This homomorphism, called the *formal shift operator*, naturally extends the so-called shift homomorphism τ_z on the Yangian, and has a number of remarkable properties. For instance, it induces a family of isomorphisms between completions of $DY_{\hbar}\mathfrak{g}$ and $Y_{\hbar}\mathfrak{g}$, realizes $Y_{\hbar}\mathfrak{g}$ as a degeneration of $DY_{\hbar}\mathfrak{g}$, and is injective provided \mathfrak{g} is of finite type or of simply laced affine type. In addition, it was applied in [18] to characterize the category of finite-dimensional representations of $DY_{\hbar}\mathfrak{g}$, for $\hbar \in \mathbb{C}^{\times}$ and \mathfrak{g} of finite type, as the tensor-closed Serre subcategory of that of the Yangian consisting of those representations which have no *poles* at zero.

In this article, we narrow our focus to the case where \mathfrak{g} is a finite-dimensional simple Lie algebra, and apply these results in conjunction with those of the recent article [17] to prove one of the main conjectures from the work [32] of Khoroshkin and Tolstoy. Namely, we establish that $DY_{\hbar}\mathfrak{g}$, which is defined by doubling the generators of $Y_{\hbar}\mathfrak{g}$ (see Definition 4.1), is isomorphic to the *restricted* quantum double of the Yangian $Y_{\hbar}\mathfrak{g}$, where the prefix “restricted” indicates that all duality operations are taken so as to respect the underlying gradings. As a consequence of this result and its proof, we find

Received by the editors May 16, 2023; accepted February 5, 2024.

Published online on Cambridge Core February 16, 2024.

AMS subject classification: 17B37, 81R10.

Keywords: Yangians, affine quantum groups, R -matrices, Yang–Baxter equation, quantization.



that Φ_z identifies the universal R -matrix of $DY_{\hbar}\mathfrak{g}$, which arises from the quantum double construction, with Drinfeld’s universal R -matrix $\mathcal{R}(z) \in (Y_{\hbar}\mathfrak{g} \otimes Y_{\hbar}\mathfrak{g})[[z^{-1}]]$. Our argument makes essential use of the constructive proof of the existence of $\mathcal{R}(z)$ given in [17], which is independent from Drinfeld’s cohomological construction of $\mathcal{R}(z)$ from the foundational article [6].

1.1 Main results

Let us now sketch our main results in detail. The two results alluded to above form Parts (1) and (2) of the following theorem.

Theorem I *There is a unique \mathbb{Z} -graded Hopf algebra structure on $DY_{\hbar}\mathfrak{g}$ over $\mathbb{C}[[\hbar]]$ such that the formal shift operator*

$$\Phi_z : DY_{\hbar}\mathfrak{g} \hookrightarrow \widehat{LY_{\hbar}\mathfrak{g}}_z \subset Y_{\hbar}\mathfrak{g}[[z^{\pm 1}]]$$

intertwines the Hopf structures on $DY_{\hbar}\mathfrak{g}$ and $Y_{\hbar}\mathfrak{g}$. Moreover:

- (1) $DY_{\hbar}\mathfrak{g}$ is isomorphic, as a \mathbb{Z} -graded Hopf algebra, to the restricted quantum double of the Yangian $Y_{\hbar}\mathfrak{g}$.
- (2) Under the above identifications, the universal R -matrix \mathcal{R} of $DY_{\hbar}\mathfrak{g}$ satisfies

$$(\Phi_w \otimes \Phi_z)(\mathcal{R}) = \mathcal{R}(w - z) \in Y_{\hbar}\mathfrak{g}^{\otimes 2}[[w]][[z^{-1}]].$$

This is a combination of the three main results of this article: Theorems 7.5, 8.4, and 9.6. Part (1) is the statement of our second main result – Theorem 8.4 – and is precisely the variant of the conjecture from [32, Section 2], which we establish in the present article. Our approach to proving it is, in a certain sense, dual to the strategy outlined in [32], which was brought to fruition for $\mathfrak{g} = \mathfrak{sl}_2$. In more detail, our argument hinges on the fact, proven in Proposition 7.1, that the universal R -matrix $\mathcal{R}(z)$ of the Yangian gives rise to a $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$\Phi_z^{\vee} : \dot{Y}_{\hbar}\mathfrak{g}^{\star} \rightarrow Y_{\hbar}\mathfrak{g}[[z^{-1}]]$$

which is compatible with the Hopf algebra structure on $Y_{\hbar}\mathfrak{g}$ and the co-opposite Hopf structure on the dual $\dot{Y}_{\hbar}\mathfrak{g}^{\star}$ of the Yangian $Y_{\hbar}\mathfrak{g}$ taken in the category of \mathbb{Z} -graded quantized enveloping algebras. That is, $\dot{Y}_{\hbar}\mathfrak{g}^{\star}$ is the restricted (or graded) dual of the Drinfeld–Gavarini [7, 19] subalgebra $\dot{Y}_{\hbar}\mathfrak{g} \subset Y_{\hbar}\mathfrak{g}$, defined in Section 5, and provides a homogeneous quantization of the restricted dual $t^{-1}\mathfrak{g}[t^{-1}]$ to the \mathbb{N} -graded Lie bialgebra $\mathfrak{g}[t]$, as we prove in detail in Section 6 (see Theorem 6.7).

Using the construction of $\mathcal{R}(z)$ given in [17] and properties of Φ_z established in [41], we deduce that the image of $\dot{Y}_{\hbar}\mathfrak{g}^{\star}$ under Φ_z^{\vee} is contained in the image of Φ_z . We may thus compose Φ_z^{\vee} with Φ_z^{-1} to obtain a $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$\dot{\iota} := \Phi_z^{-1} \circ \Phi_z^{\vee} : Y_{\hbar}\mathfrak{g}^{\vee} \rightarrow DY_{\hbar}\mathfrak{g},$$

where $Y_{\hbar}\mathfrak{g}^{\vee} := (\dot{Y}_{\hbar}\mathfrak{g}^{\star})^{\text{cop}}$. In our first main result – Theorem 7.5 – we show that there is a unique \mathbb{Z} -graded Hopf algebra structure on $DY_{\hbar}\mathfrak{g}$ for which both $\dot{\iota}$ and the natural inclusion $\iota : Y_{\hbar}\mathfrak{g} \rightarrow DY_{\hbar}\mathfrak{g}$ are injective homomorphisms of graded Hopf algebras. This is exactly the Hopf structure alluded to in the statement of Theorem I, and is such that $DY_{\hbar}\mathfrak{g}$ provides a homogeneous quantization of the restricted Drinfeld double

$\mathfrak{g}[t^{\pm 1}]$ of $\mathfrak{g}[t]$. Using Theorem 7.5, it is then not difficult to establish Part (1) above (*i.e.*, Theorem 8.4) using the double cross-product realization of the restricted quantum double (see Section 8.2).

Our third and final main result, Theorem 9.6, is a strengthening of Part (2) above. Indeed, it outputs the Gauss decomposition for \mathcal{R} while identifying each factor appearing in this decomposition with the factors $\mathcal{R}^{\pm}(z)$ and $\mathcal{R}^0(z)$ of $\mathcal{R}(z)$, which were studied in detail in [16, 17].

1.2 Motivation

Part (2) of Theorem I implies that $\mathcal{R}(z)$ can be recovered from the canonical element defined by the pairing between $Y_{\hbar\mathfrak{g}}$ and its restricted dual $Y_{\hbar\mathfrak{g}}^* \subset DY_{\hbar\mathfrak{g}}$ by applying the injection Φ_{-z} to its second tensor factor:

$$(\mathbf{1} \otimes \Phi_{-z})(\mathcal{R}) = \mathcal{R}(z) \in Y_{\hbar\mathfrak{g}}^{\otimes 2}[[z^{-1}]].$$

Here, we refer the reader to Theorem 9.6 for further details, which takes into account the topological subtleties surrounding this statement. Obtaining this interpretation of $\mathcal{R}(z)$ is in fact our original motivation for addressing the conjecture of [32, Section 2], and brings the theory surrounding the universal R -matrix of the Yangian to a more equal footing with that of the (extended, untwisted) quantum affine algebra $U_q(\widehat{\mathfrak{g}})$ and the quantum loop algebra $U_q(L\mathfrak{g})$. The differences are, however, still quite pronounced. Indeed, $U_q(\widehat{\mathfrak{g}})$ is itself nearly the quantum double of its (quantum Kac–Moody) Borel subalgebra $U_q(\mathfrak{b}^+)$, and its universal R -matrix \mathcal{R} lies in a completion of $U_q(\mathfrak{b}^+) \otimes U_q(\mathfrak{b}^-)$. One then recovers the universal R -matrix $\mathcal{R}_{L\mathfrak{g}}$ of $U_q(L\mathfrak{g})$ as a truncation of \mathcal{R} , and its z -dependent analogue is

$$\mathcal{R}_{L\mathfrak{g}}(z) := (\mathbf{1} \otimes D_{z^{-1}})(\mathcal{R}_{L\mathfrak{g}}) \in U_q(L\mathfrak{g})^{\otimes 2}[[z]],$$

where D_z is given by the \mathbb{Z} -grading on $U_q(L\mathfrak{g})$ (see [10, Section 9.4] or [14, Section 4], for instance). Crucially, \mathcal{R} can be constructed by computing dual bases with respect to the pairing between $U_q(\mathfrak{b}^+)$ and $U_q(\mathfrak{b}^-)$, and was done explicitly by Damiani in [5]. In contrast, the Yangian $Y_{\hbar\mathfrak{g}}$ is not of Kac–Moody type and does not arise as a Hopf algebra from the quantum double construction applied to any analogue of $U_q(\mathfrak{b}^+)$.¹ In addition, $\mathcal{R}(z)$ and $\mathcal{R}_{L\mathfrak{g}}(z)$ exhibit significantly different analytic behaviour when evaluated on finite-dimensional representations [17]. Nonetheless, the results of this article further cement that there are very strong parallels to be drawn between the two pictures. Indeed, one obtains the $Y_{\hbar\mathfrak{g}}$ -analogue of the above story by replacing $U_q(\mathfrak{b}^+)$ by $Y_{\hbar\mathfrak{g}}$, $U_q(L\mathfrak{g})$ by the Yangian double $DY_{\hbar\mathfrak{g}}$, and D_z by the formal shift operator Φ_z .

It should be noted that it appears that this realization of $\mathcal{R}(z)$ has been anticipated for some time in the mathematical physics community; see, for instance, [40, Section 5], which considers its super-analogue. The direction taken therein is, however, based on both the conjecture from [32, Section 2] at the heart of the present article, and on the infinite product formulas for the factors \mathcal{R}^{\pm} of \mathcal{R} given

¹We refer the reader to [42, Section 4] for a related construction of $Y_{\hbar\mathfrak{g}}$ with respect to its deformed Drinfeld coproduct, which does not endow $Y_{\hbar\mathfrak{g}}$ with the structure of a Hopf algebra.

[32, Section 5], which remain conjectural. Some more discussion on this point is given in Section 9.4.

1.3 Remarks

Let us now give a few brief remarks. First, it is essential that the Yangian double $DY_{\hbar}\mathfrak{g}$ is defined as a topological algebra over $\mathbb{C}[[\hbar]]$ for the above results to hold true. To expand on this, $DY_{\hbar}\mathfrak{g}$ can be realized as the \hbar -adic completion of a \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -algebra $\mathbb{D}Y_{\hbar}\mathfrak{g}^j$ defined by generators and relations (see Remark 4.2). One can further specialize \hbar to any nonzero complex number ζ to obtain a \mathbb{C} -algebra $\mathbb{D}Y_{\zeta}\mathfrak{g} = \mathbb{D}Y_{\hbar}\mathfrak{g}^j / (\hbar - \zeta)\mathbb{D}Y_{\hbar}\mathfrak{g}^j$, whose category of finite-dimensional representations was characterized in terms of that of the corresponding Yangian $Y_{\zeta}(\mathfrak{g})$ in [18]. Though this category has a tensor structure which corresponds to the Hopf structure on $DY_{\hbar}\mathfrak{g}$, it is important to note that $\mathbb{D}Y_{\zeta}\mathfrak{g}$ is *not* a Hopf algebra over \mathbb{C} , and in particular it does not coincide with the (restricted) quantum double of $Y_{\zeta}(\mathfrak{g})$ defined in any reasonable sense.

That being said, $\mathbb{D}Y_{\zeta}\mathfrak{g}$ admits a natural \mathbb{Z} -filtration corresponding to the \mathbb{Z} -grading on $DY_{\hbar}\mathfrak{g}$, and the expectation is that the formal completion of $\mathbb{D}Y_{\zeta}\mathfrak{g}$ with respect to this filtration coincides with the (restricted) quantum double of $Y_{\zeta}(\mathfrak{g})$ taken in the appropriate category of \mathbb{Z} -filtered, complete topological Hopf algebras. This is in fact the version of Part (2) of Theorem I conjectured in [32], and is consistent with the situation that transpires in type A for the R -matrix realization of the Yangian, which has been developed in great detail in the recent article [38]. For our purposes, it is more natural to work over $\mathbb{C}[[\hbar]]$ within the framework of quantized enveloping algebras first developed by Drinfeld [7], where we may study $DY_{\hbar}\mathfrak{g}$ from the point of view of quantization of Lie bialgebras. At the same time, many of our results are “global” (in the sense of [20]) and admit an interpretation over both $\mathbb{C}[[\hbar]]$ and \mathbb{C} , including the realization of $\mathcal{R}(z)$ provided by Theorem 9.6 (see Appendix A).

1.4 Outline

The article is written so as to provide a complete picture, accessible to non-experts, where possible. For this reason, we take great care to lay the foundation needed to state and prove the results outlined in Section 1.1. The first three sections – Sections 2–4 – are intended to serve a preliminary role: Section 2 surveys the theory of \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -modules, algebras, and Hopf algebras, including homogeneous quantizations of graded Lie bialgebras. This theory plays a prominent role throughout the article. In Section 3, we review the definition and main properties of the Yangian $Y_{\hbar}\mathfrak{g}$, defined both over $\mathbb{C}[[\hbar]]$ and $\mathbb{C}[[\hbar]]$. Notably, this includes a review of the construction of the universal R -matrix $\mathcal{R}(z)$ carried out in [17]. Section 4 is focused on the Yangian double $DY_{\hbar}\mathfrak{g}$ and, in particular, on reviewing the main results of [41] (see Theorems 4.6 and 4.8).

In Sections 5 and 6, we study the Drinfeld–Gavarini subalgebra $\check{Y}_{\hbar}\mathfrak{g}$ of the Yangian, its $\mathbb{C}[[\hbar]]$ -form, and its restricted dual $\check{Y}_{\hbar}\mathfrak{g}^*$ in detail. This includes a detailed proof that $\check{Y}_{\hbar}\mathfrak{g}^*$ provides a homogeneous quantization of $t^{-1}\mathfrak{g}[t^{-1}]$, equipped with the Yangian Lie bialgebra structure (see Definition 6.1 and Theorem 6.7).

The last three sections of the article contain its three main results: Theorems 7.5, 8.4, and 9.6. We refer the reader to Section 1.1 above, where these are outlined in detail. Finally, in Appendix Appendix A, we explain how to translate the construction of the universal R -matrix given in [17] for $\hbar \in \mathbb{C}^\times$ to the setting of the present article, in which \hbar is a formal variable (see Proposition A.1, which appears in Section 3.6 as Theorem 3.8).

2 Homogeneous quantizations

2.1 Topological modules

Recall that a $\mathbb{C}[[\hbar]]$ -module M is *separated* if the intersection of the family of submodules $\hbar^n M$ is trivial, and it is *complete* if the natural $\mathbb{C}[[\hbar]]$ -linear map

$$M \rightarrow \varprojlim_n (M/\hbar^n M)$$

is surjective, where the inverse limit is taken over the set \mathbb{N} of nonnegative integers. In particular, M is both separated and complete if and only if the above map is an isomorphism. If M is separated, complete, and torsion-free as a $\mathbb{C}[[\hbar]]$ -module, then it is said to be *topologically free*. This is equivalent to the existence of a $\mathbb{C}[[\hbar]]$ -module isomorphism $M \cong V[[\hbar]]$ for a complex vector space V . Such an isomorphism is specified by a choice of complement $V \subset M$ to $\hbar M$:

$$M = V \oplus \hbar M.$$

More generally, if M is any $\mathbb{C}[[\hbar]]$ -module, then the space $V = M/\hbar M$ is called the *semiclassical limit* of M . Similarly, the semiclassical limit of a $\mathbb{C}[[\hbar]]$ -linear map $\tau : M \rightarrow N$ is the \mathbb{C} -linear map $\bar{\tau} : M/\hbar M \rightarrow N/\hbar N$ uniquely determined by the commutativity of the diagram

$$\begin{array}{ccc} M & \xrightarrow{\tau} & N \\ \downarrow & & \downarrow \\ M/\hbar M & \xrightarrow{\bar{\tau}} & N/\hbar N. \end{array}$$

As the following elementary result illustrates, the semiclassical limit of a $\mathbb{C}[[\hbar]]$ -module homomorphism encodes important information about the original map.

Lemma 2.1 *Let M, N, τ , and $\bar{\tau}$ be as above.*

- (1) *Suppose that M is separated, N is torsion-free, and $\bar{\tau}$ is injective. Then τ is injective.*
- (2) *Suppose that M is complete, N is separated, and $\bar{\tau}$ is surjective. Then τ is surjective.*

The topological tensor product $M \widehat{\otimes} N$ of two $\mathbb{C}[[\hbar]]$ -modules M and N is the \hbar -adic completion of the algebraic tensor product $M \otimes_{\mathbb{C}[[\hbar]]} N$:

$$M \widehat{\otimes} N = \varprojlim_n (M \otimes_{\mathbb{C}[[\hbar]]} N) / \hbar^n (M \otimes_{\mathbb{C}[[\hbar]]} N).$$

If M and N are topologically free with $M \cong V[[\hbar]]$ and $N \cong W[[\hbar]]$, then $M \widehat{\otimes} N$ is topologically free and isomorphic to $(V \otimes_{\mathbb{C}} W)[[\hbar]]$.

In this article, we shall say that M is a *topological module* over $\mathbb{C}[[\hbar]]$ if M is a $\mathbb{C}[[\hbar]]$ -module which is both separated and complete. For any such module, we have

$$\mathbb{C}[[\hbar]] \widehat{\otimes} M \cong M \cong M \widehat{\otimes} \mathbb{C}[[\hbar]].$$

Similarly, by a *topological algebra* A over $\mathbb{C}[[\hbar]]$, we shall always mean that A is a $\mathbb{C}[[\hbar]]$ -algebra which is both separated and complete as a module over $\mathbb{C}[[\hbar]]$. In particular, the multiplication m can be viewed as a $\mathbb{C}[[\hbar]]$ -linear map

$$m : A \widehat{\otimes} A \rightarrow A.$$

A *topological Hopf algebra* H over $\mathbb{C}[[\hbar]]$ is a topological $\mathbb{C}[[\hbar]]$ -algebra equipped with a counit $\varepsilon : H \rightarrow \mathbb{C}[[\hbar]]$, a coproduct $\Delta : H \rightarrow H \widehat{\otimes} H$, and an antipode $S : H \rightarrow H$ which collectively satisfy the axioms of a Hopf algebra with all tensor products given by the topological tensor product $\widehat{\otimes}$. By modifying these definitions in the expected way, one obtains the notion of a topological coalgebra and bialgebra over $\mathbb{C}[[\hbar]]$.

If M and N are topological $\mathbb{C}[[\hbar]]$ -modules and $N \cong W[[\hbar]]$ is topologically free, then the space of $\mathbb{C}[[\hbar]]$ -module homomorphisms $\text{Hom}_{\mathbb{C}[[\hbar]]}(M, N)$ is separated, complete, and torsion-free. If, in addition, M is topologically free with $M \cong V[[\hbar]]$, then one has

$$\text{Hom}_{\mathbb{C}[[\hbar]]}(M, N) \cong \text{Hom}_{\mathbb{C}}(V, W)[[\hbar]].$$

In particular, the $\mathbb{C}[[\hbar]]$ -linear dual $M^* := \text{Hom}_{\mathbb{C}[[\hbar]]}(M, \mathbb{C}[[\hbar]])$ of a topologically free $\mathbb{C}[[\hbar]]$ -module $M \cong V[[\hbar]]$ satisfies $M^* \cong V^*[[\hbar]]$.

2.2 Graded topological modules

Let us now turn toward the \mathbb{Z} -graded analogues of the above definitions. Henceforth, we view $\mathbb{C}[[\hbar]] = \bigoplus_{k \in \mathbb{N}} \mathbb{C}\hbar^k$ as an \mathbb{N} -graded ring. For brevity, we shall denote its \mathbb{N} -graded quotient $\mathbb{C}[[\hbar]]/\hbar^n \mathbb{C}[[\hbar]]$ by K_n , for each $n \in \mathbb{N}$.

Definition 2.2 We say that a topological $\mathbb{C}[[\hbar]]$ -module M is *\mathbb{Z} -graded* if, for each $n \in \mathbb{N}$, $M/\hbar^n M = \bigoplus_{k \in \mathbb{Z}} (M/\hbar^n M)_k$ is a \mathbb{Z} -graded K_n -module and the natural homomorphism

$$M/\hbar^{n+1} M \rightarrow M/\hbar^n M$$

is \mathbb{Z} -graded. If $(M/\hbar^n M)_k$ is trivial for $k < 0$, we say that M is \mathbb{N} -graded.

A $\mathbb{C}[[\hbar]]$ -module homomorphism $M \rightarrow N$ between \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -modules M and N is said to be *\mathbb{Z} -graded* if the induced morphisms

$$M/\hbar^n M \rightarrow N/\hbar^n N$$

are all \mathbb{Z} -graded. More generally, it is \mathbb{Z} -graded of degree $a \in \mathbb{Z}$ if each of these induced morphisms is homogeneous of degree a .

The category of \mathbb{Z} -graded topological modules is closed under the tensor product $\widehat{\otimes}$. Indeed, this follows from the elementary observation that, given two $\mathbb{C}[[\hbar]]$ -modules M and N , one has

$$(M \widehat{\otimes} N)/\hbar^n (M \widehat{\otimes} N) \cong (M \otimes_{\mathbb{C}[[\hbar]]} N)/\hbar^n (M \otimes_{\mathbb{C}[[\hbar]]} N) \cong M/\hbar^n M \otimes_{K_n} N/\hbar^n N,$$

which can be equipped with the standard tensor product grading, provided $M/\hbar^n M$ and $N/\hbar^n N$ are both \mathbb{Z} -graded.

Definition 2.3 A topological algebra A is said to be \mathbb{Z} -graded if it is graded as a topological $\mathbb{C}[[\hbar]]$ -module and the multiplication map

$$m : A \widehat{\otimes} A \rightarrow A$$

is a \mathbb{Z} -graded homomorphism. Similarly, a topological Hopf algebra H is \mathbb{Z} -graded if it is \mathbb{Z} -graded as a topological algebra and the structure maps

$$\Delta : H \rightarrow H \widehat{\otimes} H, \quad \varepsilon : H \rightarrow \mathbb{C}[[\hbar]], \quad S : H \rightarrow H$$

are all \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -module homomorphisms. Equivalently, a topological algebra or Hopf algebra H is \mathbb{Z} -graded if the conditions of Definition 2.2 hold and each $H/\hbar^n H$ is a \mathbb{Z} -graded algebra or Hopf algebra over K_n , respectively.

Of course, one also has the notion of a \mathbb{Z} -graded topological coalgebra and bialgebra, which are defined by making the obvious modifications to the above definition.

The prototypical example of a \mathbb{Z} -graded topological module over $\mathbb{C}[[\hbar]]$ is $M = V[[\hbar]]$ where $V = \bigoplus_{k \in \mathbb{Z}} V_k$ is a \mathbb{Z} -graded complex vector space. In this case, one has

$$M/\hbar^n M \cong V[[\hbar]]/\hbar^n V[[\hbar]],$$

which is naturally graded, as $V[[\hbar]]$ is graded with $V[[\hbar]]_k = \bigoplus_{n \geq 0} \hbar^n V_{k-n}$, and $\hbar^n V[[\hbar]]$ is a graded submodule. The assertion that M is \mathbb{Z} -graded may be recaptured as follows. For each $k \in \mathbb{Z}$, set

$$M_k := \varprojlim_n V[[\hbar]]_k/\hbar^n V[[\hbar]]_{k-n} \cong \prod_{n \in \mathbb{N}} \hbar^n V_{k-n} \subset V[[\hbar]].$$

Then each M_k is a closed subspace of M satisfying $M_k/\hbar^n M_{k-n} \cong (M/\hbar^n M)_k$, and M contains the \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -module $\bigoplus_{k \in \mathbb{Z}} M_k$ as a dense $\mathbb{C}[[\hbar]]$ -submodule. Moreover, the \hbar -adic topology on this submodule coincides with the subspace topology, so M is the \hbar -adic completion of $\bigoplus_{k \in \mathbb{Z}} M_k$. If, in addition, V is \mathbb{N} -graded, then $\bigoplus_{k \in \mathbb{Z}} M_k$ coincides with the polynomial space $V[[\hbar]] \subset V[[\hbar]]$.

The below lemma provides an equivalent characterization of the definition of a \mathbb{Z} -graded topological module and algebra which generalizes this picture.

Lemma 2.4 A topological $\mathbb{C}[[\hbar]]$ -module M is \mathbb{Z} -graded if and only if it admits a dense, \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -submodule

$$M_{\mathbb{Z}} = \bigoplus_{k \in \mathbb{Z}} M_k \subset M$$

with each M_k a closed subspace of M and $\hbar^n M \cap M_{\mathbb{Z}} = \hbar^n M_{\mathbb{Z}}$ for all $n \in \mathbb{N}$.

If, in addition, M has the structure of a topological algebra, then it is \mathbb{Z} -graded if and only if the above conditions hold and $M_{\mathbb{Z}}$ is a \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -subalgebra of M .

If M is a \mathbb{Z} -graded topological module, then the k th component M_k of $M_{\mathbb{Z}}$ from Lemma 2.4 is uniquely determined and recovered as the inverse limit

$$(2.1) \quad M_k := \varprojlim_n (M/\hbar^n M)_k \subset \varprojlim_n M/\hbar^n M = M.$$

Moreover, one has $(M/\hbar^n M)_k \cong M_k/\hbar^n M_{k-n}$ for all $n \in \mathbb{N}$ and $k \in \mathbb{Z}$.

We further observe that, for each $k \in \mathbb{Z}$, the system of linear projections $\pi_{n,k} : M/\hbar^n M \rightarrow (M/\hbar^n M)_k$ gives rise to a projection

$$\pi_k := \varprojlim_n \pi_{n,k} : M \rightarrow M_k$$

which restricts to the projection of $M_{\mathbb{Z}}$ onto its k th homogeneous component. In particular, a $\mathbb{C}[[\hbar]]$ -linear map $\tau : M \rightarrow N$ between \mathbb{Z} -graded topological modules is graded if and only if $\pi_k^N \circ \tau = \tau \circ \pi_k^M$ for each $k \in \mathbb{Z}$.

We conclude this preliminary subsection with two corollaries of the above discussion. The first shows that any topologically free \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -module is of the form described above Lemma 2.4.

Corollary 2.5 *Suppose M is a \mathbb{Z} -graded topologically free module over $\mathbb{C}[[\hbar]]$, and let V denote the \mathbb{Z} -graded complex vector space $M/\hbar M = \bigoplus_{k \in \mathbb{Z}} M_k/\hbar M_{k-1}$. Then*

$$M \cong V[[\hbar]]$$

as a \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -module. In particular, one has

$$M_{\mathbb{Z}} = \bigoplus_{k \in \mathbb{Z}} M_k \cong \bigoplus_{k \in \mathbb{Z}} V[[\hbar]]_k \subset V[[\hbar]], \quad \text{where } V[[\hbar]]_k = \prod_{n \in \mathbb{N}} \hbar^n V_{k-n}.$$

Proof This is a refinement of the elementary result, alluded to at the beginning of the section, that $M \cong V[[\hbar]]$ as a $\mathbb{C}[[\hbar]]$ -module (see [26, Proposition XVI.2.4], for instance). In more detail, an isomorphism of \mathbb{Z} -graded topological modules $M \cong V[[\hbar]]$ is specified by choosing, for each $k \in \mathbb{Z}$, a complement $V_k \subset M_k$ to $\hbar M_{k-1}$ in M_k :

$$M_k = V_k \oplus \hbar M_{k-1}.$$

Setting $V := \bigoplus_{k \in \mathbb{Z}} V_k \subset M_{\mathbb{Z}}$, we then have

$$\begin{aligned} V &\cong M_{\mathbb{Z}}/\hbar M_{\mathbb{Z}} = M/\hbar M, \\ M_k/\hbar^n M_{k-n} &\cong V_k \oplus \hbar V_{k-1} \oplus \cdots \oplus \hbar^{n-1} V_{k-n+1} \cong V[[\hbar]]_k/\hbar^n V[[\hbar]]_{k-n}, \\ M/\hbar^n M &\cong \bigoplus_{k \in \mathbb{Z}} M_k/\hbar^n M_{k-n} \cong \bigoplus_{k \in \mathbb{Z}} V[[\hbar]]_k/\hbar^n V[[\hbar]]_{k-n} = V[[\hbar]]/\hbar^n V[[\hbar]], \end{aligned}$$

where the third line is an identification of \mathbb{Z} -graded modules. Here, we note that the second line follows from the definition of V_k and that M is a torsion-free $\mathbb{C}[[\hbar]]$ -module. Taking inverse limits, one finds that $M \cong V[[\hbar]]$ as \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -modules. ■

Let us now shift our attention to the case where $\dot{M} = \bigoplus_{k \in \mathbb{N}} \dot{M}_k$ is an \mathbb{N} -graded $\mathbb{C}[[\hbar]]$ -module. Any such module is automatically separated, and so embeds into its \hbar -adic completion

$$M = \varprojlim_n (\dot{M}/\hbar^n \dot{M}),$$

which is an \mathbb{N} -graded topological $\mathbb{C}[[\hbar]]$ -module. Moreover, if \dot{M} is a torsion-free $\mathbb{C}[[\hbar]]$ -module, then M is topologically free as a $\mathbb{C}[[\hbar]]$ -module. Since \dot{M}_{k-n} is trivial for $n > k$, the submodule M_k of M (see (2.1)) coincides with \dot{M}_k and so, in the notation of Lemma 2.4, one has $M = M_{\mathbb{N}}$. These observations, coupled with Corollary 2.5 and that $V[[\hbar]]_k = V[\hbar]_k$ when V is \mathbb{N} -graded, yield the following.

Corollary 2.6 *Let \dot{M} be an \mathbb{N} -graded, torsion-free $\mathbb{C}[[\hbar]]$ -module. Then M is a topologically free \mathbb{N} -graded $\mathbb{C}[[\hbar]]$ -module. Moreover, we have*

$$\dot{M}_k = \varprojlim_n (\dot{M}_k / \hbar^n \dot{M}_{k-n}) = M_k \quad \text{for all } k \in \mathbb{N}.$$

Consequently, \dot{M} coincides with $M_{\mathbb{N}}$ and there is an isomorphism of \mathbb{N} -graded $\mathbb{C}[[\hbar]]$ -modules

$$\dot{M} \cong V[[\hbar]] = \bigoplus_{k \in \mathbb{N}} V[\hbar]_k, \quad \text{where } V := \dot{M} / \hbar \dot{M}.$$

Note that if \dot{M} is an \mathbb{N} -graded $\mathbb{C}[[\hbar]]$ -algebra or Hopf algebra, then M is automatically an \mathbb{N} -graded topological algebra or Hopf algebra, respectively.

2.3 The restricted dual

For a given \mathbb{Z} -graded complex vector space $V = \bigoplus_n V_n$, we let $V^* = \bigoplus_n (V^*)_n \subset V^*$ denote the restricted, or graded, dual of V , where

$$(V^*)_n = \{f \in V^* : f(V_m) \subset \mathbb{C}_{m+n}\} \cong (V_{-n})^*$$

and \mathbb{C} is given the trivial grading with $\mathbb{C}_0 = \mathbb{C}$ and $\mathbb{C}_m = \{0\}$ for $m \neq 0$. One can similarly define the restricted dual $M^* \subset M^*$ in the category of \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -modules. In this subsection, we will recall some properties of this duality operation in the \mathbb{N} -graded setting which will be applied to construct the dual Yangian in Section 6.

Suppose that M is an \mathbb{N} -graded, topological $\mathbb{C}[[\hbar]]$ -module with \mathbb{N} -graded $\mathbb{C}[[\hbar]]$ -submodule $M_{\mathbb{N}} = \bigoplus_{k \in \mathbb{N}} M_k$ as in Lemma 2.4. For each $n \in \mathbb{N}$, let J_k denote the \hbar -adic completion of the ideal $\bigoplus_{k \geq n} M_k$ of $M_{\mathbb{N}}$. The gradation topology on M is the topology associated with the descending filtration

$$M = J_0 \supset J_1 \supset \dots \supset J_n \supset \dots$$

Equipped with this terminology, we may make the following definition.

Definition 2.7 The restricted dual M^* is defined to be the $\mathbb{C}[[\hbar]]$ -submodule of M^* consisting of those f which are continuous with respect to the gradation topology:

$$M^* := \{f \in M^* : f(J_k) \subset \hbar^n \mathbb{C}[[\hbar]] \quad \forall n \in \mathbb{N} \text{ and } k \gg 0\}.$$

The restricted dual of any \mathbb{N} -graded topological $\mathbb{C}[[\hbar]]$ -module M is easily seen to be separated, complete, and torsion-free. Let us now see that it admits a \mathbb{Z} -graded structure.

For each $a \in \mathbb{Z}$, let $\text{Hom}_{\mathbb{C}[[\hbar]]}^a(M, \mathbb{C}[[\hbar]]) \subset M^*$ denote the (closed) subspace consisting of \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -module homomorphisms $f : M \rightarrow \mathbb{C}[[\hbar]]$ of degree a . Equivalently,

$$\text{Hom}_{\mathbb{C}[[\hbar]]}^a(M, \mathbb{C}[[\hbar]]) = \{f \in M^* : f(M_k) \subset \mathbb{C}[[\hbar]]_{k+a} \quad \forall k \in \mathbb{N}\}.$$

Then the sum $\sum_{a \in \mathbb{Z}} \text{Hom}_{\mathbb{C}[[\hbar]]}^a(M, \mathbb{C}[[\hbar]])$ is direct and the space

$$(M^*)_{\mathbb{Z}} := \bigoplus_{a \in \mathbb{Z}} \text{Hom}_{\mathbb{C}[[\hbar]]}^a(M, \mathbb{C}[[\hbar]]) \subset M^*$$

is a \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -submodule of M^* .

Remark 2.8 Under the natural identification of M^* with $\text{Hom}_{\mathbb{C}[[\hbar]]}(M_{\mathbb{N}}, \mathbb{C}[[\hbar]])$, one has $\text{Hom}_{\mathbb{C}[[\hbar]]}^a(M, \mathbb{C}[[\hbar]]) \cong \text{Hom}_{\mathbb{C}[[\hbar]]}^a(M_{\mathbb{N}}, \mathbb{C}[[\hbar]])$ and $(M^*)_{\mathbb{Z}}$ coincides with the graded dual $(M_{\mathbb{N}})^* \subset \text{Hom}_{\mathbb{C}[[\hbar]]}(M_{\mathbb{N}}, \mathbb{C}[[\hbar]])$ of $M_{\mathbb{N}}$ taken in the category of \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -modules.

It is not difficult to prove that, for each $n \in \mathbb{N}$, one has

$$\hbar^n M^* \cap (M^*)_{\mathbb{Z}} = \hbar^n (M^*)_{\mathbb{Z}} \quad \text{and} \quad (M^*)_{\mathbb{Z}} / \hbar^n (M^*)_{\mathbb{Z}} \cong M^* / \hbar^n M^*.$$

Consequently, M^* coincides with the \hbar -adic completion of $(M^*)_{\mathbb{Z}}$ and, by Lemma 2.4, is a \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -module. We note that, although Definition 2.7 is strictly for an \mathbb{N} -graded $\mathbb{C}[[\hbar]]$ -module M , one can define the restricted dual in the \mathbb{Z} -graded setting precisely as the \hbar -adic completion of the space $(M^*)_{\mathbb{Z}}$.

If M is itself topologically free with $M \cong V[[\hbar]]$ for a graded vector space $V = \bigoplus_{k \in \mathbb{N}} V_k$, then the natural homomorphism $M^* / \hbar M^* \rightarrow V^*$ is an isomorphism of graded vector spaces. As M^* is topologically free, Corollary 2.5 yields the following.

Corollary 2.9 Suppose that M is a topologically free \mathbb{N} -graded $\mathbb{C}[[\hbar]]$ -module with $M \cong V[[\hbar]]$ for a graded vector space $V = \bigoplus_{k \in \mathbb{N}} V_k$. Then M^* is isomorphic to $V^*[[\hbar]]$ as a \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -module.

We shall say that a topologically free \mathbb{N} -graded $\mathbb{C}[[\hbar]]$ -module is of *finite type* if the graded components V_k of $V \cong M/\hbar M$ from the above corollary are all finite-dimensional complex vector spaces.

Now suppose that H is an \mathbb{N} -graded topological Hopf algebra with coproduct Δ , counit ε , antipode S , product m , and unit ι . Since these are all \mathbb{N} -graded $\mathbb{C}[[\hbar]]$ -module homomorphisms and $\mathbb{C}[[\hbar]]^* \cong \mathbb{C}[[\hbar]]$, taking transposes yields \mathbb{Z} -graded maps

$$\begin{aligned} \Delta^t : (H \widehat{\otimes} H)^* &\rightarrow H^*, & \varepsilon^t : \mathbb{C}[[\hbar]] &\rightarrow H^*, & S^t : H^* &\rightarrow H^* \\ m^t : H^* &\rightarrow (H \widehat{\otimes} H)^*, & \iota^t : H^* &\rightarrow \mathbb{C}[[\hbar]] \end{aligned}$$

which formally satisfy the axioms of a Hopf algebra. In particular, H^* is a topological $\mathbb{C}[[\hbar]]$ -algebra with unit ε^t and product given by restricting Δ^t . It is not, in general, a topological coalgebra (or Hopf algebra) as m^t does not necessarily have image in $H^* \widehat{\otimes} H^*$. However, this is the case when H is of finite type, as we now explain.

In general, for any two \mathbb{N} -graded topological $\mathbb{C}[[\hbar]]$ -modules M and N , there is a canonical injective homomorphism of \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -modules

$$\gamma : M^* \widehat{\otimes} N^* \hookrightarrow (M \widehat{\otimes} N)^*.$$

If M and N are topologically free with $M \cong V[[\hbar]]$ and $N \cong W[[\hbar]]$, then the semiclassical limit of γ is the natural inclusion $V^* \otimes_{\mathbb{C}} W^* \hookrightarrow (V \otimes_{\mathbb{C}} W)^*$ which is an isomorphism provided the graded components of V or W are all finite-dimensional. This observation, together with Lemma 2.1, implies the following proposition.

Proposition 2.10 *Let M and N be topologically free, \mathbb{N} -graded $\mathbb{C}[[\hbar]]$ -modules and suppose that either M or N is of finite type. Then γ is an isomorphism of \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -modules*

$$\gamma : M^* \widehat{\otimes} N^* \xrightarrow{\sim} (M \widehat{\otimes} N)^*.$$

Consequently, if H is a topologically free \mathbb{N} -graded Hopf algebra of finite type, then H^* is a \mathbb{Z} -graded topological Hopf algebra over $\mathbb{C}[[\hbar]]$.

Remark 2.11 Henceforth, we shall simply write \otimes for the topological tensor product $\widehat{\otimes}$. More generally, the use of the symbol \otimes will always be clear from context and will be clarified should any ambiguity arise.

2.4 Homogeneous quantizations

Let us now recall some basic constructions from the theory of quantum groups, adapted to the graded setting.

A topological Hopf algebra H over $\mathbb{C}[[\hbar]]$ is called a *quantized enveloping algebra* if it is a flat deformation of the universal enveloping algebra $U(\mathfrak{b})$ of a complex Lie algebra \mathfrak{b} as a Hopf algebra. Equivalently:

- The semiclassical limit $H/\hbar H$ of H is isomorphic to $U(\mathfrak{b})$ as a Hopf algebra.
- H is topologically free, and thus isomorphic to $U(\mathfrak{b})[[\hbar]]$ as a $\mathbb{C}[[\hbar]]$ -module.

If $H = U_{\hbar} \mathfrak{b}$ is a quantized enveloping algebra with semiclassical limit $U(\mathfrak{b})$, then \mathfrak{b} inherits from $U_{\hbar} \mathfrak{b}$ the structure of a Lie bialgebra with cocommutator $\delta_{\mathfrak{b}} : \mathfrak{b} \rightarrow \mathfrak{b} \wedge \mathfrak{b} \subset U(\mathfrak{b})^{\otimes 2}$ given by the formula

$$(2.2) \quad \delta_{\mathfrak{b}}(x) := \frac{\Delta(\dot{x}) - \Delta^{op}(\dot{x})}{\hbar} \pmod{\hbar U_{\hbar} \mathfrak{b} \otimes U_{\hbar} \mathfrak{b}} \quad \forall x \in \mathfrak{b},$$

where $\dot{x} \in U_{\hbar} \mathfrak{b}$ is any lift of x . We refer the reader to Propositions 6.2.3 and 6.2.7 of [4] for a detailed discussion of this point.

Conversely, if $(\mathfrak{b}, \delta_{\mathfrak{b}})$ is a Lie bialgebra, then a *quantization* of $(\mathfrak{b}, \delta_{\mathfrak{b}})$ is a quantized enveloping algebra $U_{\hbar} \mathfrak{b}$ with semiclassical limit $U(\mathfrak{b})$, such that $\delta_{\mathfrak{b}}$ coincides with the cocommutator (2.2).

Now let us shift our attention to the graded setting. In what follows, we will say that a Lie bialgebra $(\mathfrak{b}, \delta_{\mathfrak{b}})$ is \mathbb{Z} -graded if $\mathfrak{b} = \bigoplus_{k \in \mathbb{Z}} \mathfrak{b}_k$ is \mathbb{Z} -graded as a Lie algebra, and the cocommutator $\delta_{\mathfrak{b}}$ is a graded linear map of degree d , for some $d \in \mathbb{Z}$. That is, $\delta_{\mathfrak{b}} \in \text{Hom}_{\mathbb{C}}^d(\mathfrak{b}, \mathfrak{b}^{\otimes 2})$.

Definition 2.12 Let \mathfrak{b} be a \mathbb{Z} -graded complex Lie bialgebra with cocommutator $\delta_{\mathfrak{b}}$. Then a *homogeneous quantization* of $(\mathfrak{b}, \delta_{\mathfrak{b}})$ is a topological Hopf algebra $U_{\hbar}\mathfrak{b}$ satisfying:

- (1) $U_{\hbar}\mathfrak{b}$ is a quantization of $(\mathfrak{b}, \delta_{\mathfrak{b}})$.
- (2) $U_{\hbar}\mathfrak{b}$ is \mathbb{Z} -graded as a topological Hopf algebra, and the natural inclusion

$$\mathfrak{b} \hookrightarrow U(\mathfrak{b}) \cong U_{\hbar}\mathfrak{b}/\hbar U_{\hbar}\mathfrak{b}$$

is a \mathbb{Z} -graded linear map.

Note that the last condition guarantees that the grading on $U(\mathfrak{b})$ inherited from $U_{\hbar}\mathfrak{b}$ coincides with that induced by the Lie algebra grading on \mathfrak{b} . In addition, since the coproduct Δ is homogeneous of degree zero, (2.2) implies that the cocommutator $\delta_{\mathfrak{b}}$ must be of degree $d = -1$.

We shall employ similar terminology to the above in the \mathbb{N} -graded setting over $\mathbb{C}[[\hbar]]$. Namely, if $(\mathfrak{b}, \delta_{\mathfrak{b}})$ is an \mathbb{N} -graded Lie bialgebra, then a *homogeneous quantization* of $(\mathfrak{b}, \delta_{\mathfrak{b}})$ over $\mathbb{C}[[\hbar]]$ is a space $\mathbb{U}_{\hbar}\mathfrak{b}$ such that:

- (1) $\mathbb{U}_{\hbar}\mathfrak{b}$ is an \mathbb{N} -graded torsion-free Hopf algebra over $\mathbb{C}[[\hbar]]$.
- (2) The semiclassical limit $\mathbb{U}_{\hbar}\mathfrak{b}/\hbar\mathbb{U}_{\hbar}\mathfrak{b}$ is isomorphic to $U(\mathfrak{b})$ as a graded Hopf algebra, with the cocommutator $\delta_{\mathfrak{b}}$ given by (2.2).

Note that, by Corollary 2.6, such a quantization $\mathbb{U}_{\hbar}\mathfrak{b}$ is isomorphic to $U(\mathfrak{b})[[\hbar]]$ as an \mathbb{N} -graded $\mathbb{C}[[\hbar]]$ -module, and its \hbar -adic completion is a homogeneous quantization of \mathfrak{b} over $\mathbb{C}[[\hbar]]$.

2.5 The Yangian Manin triple

The most well-known, nontrivial, example of a homogeneous quantization is the Yangian $Y_{\hbar}\mathfrak{g}$ associated with an arbitrary simple Lie algebra \mathfrak{g} over the complex numbers. In this article, we shall encounter two other, closely related, examples: the dual Yangian $Y_{\hbar}\mathfrak{g}^*$ and the Yangian double $DY_{\hbar}\mathfrak{g}$. Collectively, these three quantum groups arise as a quantization of a restricted Manin triple structure on $(\mathfrak{t}, \mathfrak{t}_+, \mathfrak{t}_-)$, where

$$\mathfrak{t} := \mathfrak{g}[[t^{\pm 1}]], \quad \mathfrak{t}_+ := \mathfrak{g}[[t]], \quad \text{and} \quad \mathfrak{t}_- := t^{-1}\mathfrak{g}[[t^{-1}]].$$

In this section, we briefly recall how this structure is defined.

The Lie algebra $\mathfrak{t} = \mathfrak{g}[[t^{\pm 1}]]$ comes equipped with a nondegenerate, invariant bilinear form $\langle \cdot, \cdot \rangle : \mathfrak{t} \otimes \mathfrak{t} \rightarrow \mathbb{C}$ given by

$$(2.3) \quad \langle f(t), g(s) \rangle := -\text{Res}_t(f(t), g(t)),$$

where (\cdot, \cdot) is a fixed symmetric, invariant, and nondegenerate bilinear form on \mathfrak{g} , which has been extended to a $\mathbb{C}[[t^{\pm 1}]]$ -valued bilinear form on \mathfrak{t} by $\mathbb{C}[[t^{\pm 1}]]$ -linearity. The above form is a degree 1 element of the restricted dual $(\mathfrak{t} \otimes \mathfrak{t})^*$, as defined in the beginning of Section 2.3. Namely, it vanishes on

$$(\mathfrak{t} \otimes \mathfrak{t})_k = \bigoplus_{a+b=k} \mathfrak{t}_a \otimes \mathfrak{t}_b$$

for any $k \neq -1$, and restricts to a nondegenerate pairing $\mathfrak{t}_a \otimes \mathfrak{t}_{-a-1} \rightarrow \mathbb{C}$ for any $a \in \mathbb{Z}$. Moreover, one has the polarization

$$\mathfrak{t} = \mathfrak{t}_+ \oplus \mathfrak{t}_- \quad \text{for} \quad \mathfrak{t}_+ = \mathfrak{g}[t], \quad \mathfrak{t}_- = t^{-1}\mathfrak{g}[t^{-1}]$$

with \mathfrak{t}_+ and \mathfrak{t}_- isotropic, graded Lie subalgebras of \mathfrak{t} , with gradings concentrated in nonnegative and nonpositive degrees, respectively. Said in fewer words, this collection of data gives rise to a *restricted Manin triple* $(\mathfrak{t}, \mathfrak{t}_+, \mathfrak{t}_-)$ (see Sections 5.2 and 5.3 of [2]).

Since each homogeneous component \mathfrak{t}_k of \mathfrak{t} is finite-dimensional, there are dual Lie bialgebra structures on \mathfrak{t}_+ and \mathfrak{t}_- , obtained as follows. The residue form (2.3) yields isomorphisms of graded vector spaces

$$\text{Res}_\pm : \mathfrak{t}_\pm \xrightarrow{\sim} \mathfrak{t}_\mp^*$$

which are homogeneous of degree 1: $\text{Res}_\pm(\mathfrak{t}_{\pm, n}) = (\mathfrak{t}_\mp^*)_{n+1}$ for all $n \in \mathbb{Z}$. Dualizing Lie brackets then gives rise to honest, degree -1 , Lie bialgebra cobrackets

$$\delta_\pm = [\cdot, \cdot]_{\mathfrak{t}_\mp^*}^t : \mathfrak{t}_\pm \rightarrow \mathfrak{t}_\pm \wedge \mathfrak{t}_\pm.$$

Observe that the Casimir tensor $\Omega_{\mathfrak{g}} \in (\mathfrak{g} \otimes \mathfrak{g})^{\mathfrak{g}}$ satisfies

$$([x \otimes 1, \Omega_{\mathfrak{g}}], y \otimes z)_{\mathfrak{g} \otimes \mathfrak{g}} = -(x, [y, z]) \quad \forall x, y, z \in \mathfrak{g},$$

where $(\cdot, \cdot)_{\mathfrak{g} \otimes \mathfrak{g}} = (\cdot, \cdot) \otimes (\cdot, \cdot) \circ (23) : \mathfrak{g} \otimes \mathfrak{g} \otimes \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathbb{C}$. It follows readily from this observation and the definition of Res_\pm that δ_+ and δ_- are given explicitly on each graded component by

$$\begin{aligned} \delta_+(xt^k) &= \sum_{a+b=k-1} [x \otimes 1, \Omega_{\mathfrak{g}}] t^a s^b \in \mathfrak{g}[t] \otimes \mathfrak{g}[s] = \mathfrak{t}_+ \otimes \mathfrak{t}_+, \\ \delta_-(xt^{-k-1}) &= \sum_{a+b=k} [x \otimes 1, \Omega_{\mathfrak{g}}] t^{-a-1} s^{-b-1} \in t^{-1}\mathfrak{g}[t^{-1}] \otimes s^{-1}\mathfrak{g}[s^{-1}] = \mathfrak{t}_- \otimes \mathfrak{t}_-, \end{aligned}$$

where we have used the natural identification of $(\mathfrak{g} \otimes \mathfrak{g})[t^{\pm 1}, s^{\pm 1}]$ with $\mathfrak{g}[t^{\pm 1}] \otimes \mathfrak{g}[s^{\pm 1}]$, and a, b take values in \mathbb{N} . Since

$$(z - w) \sum_{a=0}^k z^a w^{k-a} = z^{k+1} - w^{k+1},$$

the linear map $\delta := \delta_+ \oplus (-\delta_-) : \mathfrak{t} \rightarrow \mathfrak{t} \wedge \mathfrak{t}$ is given by the formula

$$\delta(f)(t, s) = \left[f(t) \otimes 1 + 1 \otimes f(s), \frac{\Omega_{\mathfrak{g}}}{t-s} \right] \in \mathfrak{g}[t^{\pm 1}] \otimes \mathfrak{g}[s^{\pm 1}] \quad \forall f(t) \in \mathfrak{g}[t^{\pm 1}]$$

and defines a Lie bialgebra structure on the Lie algebra \mathfrak{t} such that $(\mathfrak{t}_+, \delta_+)$ and $(\mathfrak{t}_-, -\delta_-)$ are Lie sub-bialgebras. This construction identifies \mathfrak{t} with the *restricted Drinfeld double* $D(\mathfrak{t}_+)$ of the \mathbb{N} -graded Lie bialgebra \mathfrak{t}_+ , as defined in [2, Section 5.4], for instance.

3 The Yangian revisited

3.1 The Lie algebra \mathfrak{g}

We henceforth fix \mathfrak{g} to be a finite-dimensional simple Lie algebra over the complex numbers, with invariant form $(,)$ as in Section 2.5. Let $\mathfrak{h} \subset \mathfrak{g}$ be a Cartan subalgebra, $\{\alpha_i\}_{i \in I} \subset \mathfrak{h}^*$ a basis of simple roots, and $\{\alpha_i^\vee\}_{i \in I}$ the set of simple coroots, so that $\alpha_j(\alpha_i^\vee) = a_{ij} = 2(\alpha_i, \alpha_j)/(\alpha_i, \alpha_i)$ are the entries of the Cartan matrix $\mathbf{A} = (a_{ij})_{i,j \in I}$ of \mathfrak{g} . Let $\Delta^+ \subset \mathfrak{h}^*$ be the associated set of positive roots, and let $Q = \bigoplus_{i \in I} \mathbb{Z}\alpha_i$ and $Q_+ = \bigoplus_{i \in I} \mathbb{N}\alpha_i$ denote the root lattice and its positive cone, respectively, where we recall that \mathbb{N} denotes the set of nonnegative integers. Set

$$d_{ij} = \frac{(\alpha_i, \alpha_j)}{2} \text{ and } d_i = d_{ii} \quad \forall i, j \in I.$$

We normalize $(,)$, if necessary, so that the square length of a short root is 2. In particular, we then have $\{d_i\}_{i \in I} \subset \{1, 2, 3\}$. Let $\{e_i, f_i\}_{i \in I}$ denote the Chevalley generators of \mathfrak{g} , as in [24, Section 1.3], and set

$$h_i = d_i \alpha_i^\vee, \quad x_i^+ = \sqrt{d_i} e_i, \quad x_i^- = \sqrt{d_i} f_i \quad \forall i \in I.$$

These normalized generators satisfy $(x_i^+, x_i^-) = 1$ and $h_i = [x_i^+, x_i^-]$ for all $i \in I$.

3.2 The Yangian

We now recall the definition of the Yangian $Y_{\hbar}(\mathfrak{g})$. Let S_m denote the symmetric group on $\{1, \dots, m\}$.

Definition 3.1 The Yangian $Y_{\hbar}(\mathfrak{g})$ is the unital associative $\mathbb{C}[\hbar]$ -algebra generated by $\{x_{ir}^\pm, h_{ir}\}_{i \in I, r \in \mathbb{N}}$, subject to the following relations for $i, j \in I$ and $r, s \in \mathbb{N}$:

$$(3.1) \quad [h_{ir}, h_{js}] = 0,$$

$$(3.2) \quad [h_{i0}, x_{js}^\pm] = \pm 2d_{ij} x_{js}^\pm,$$

$$(3.3) \quad [x_{ir}^+, x_{js}^-] = \delta_{ij} h_{i,r+s},$$

$$(3.4) \quad [h_{i,r+1}, x_{js}^\pm] - [h_{ir}, x_{j,s+1}^\pm] = \pm \hbar d_{ij} (h_{ir} x_{js}^\pm + x_{js}^\pm h_{ir}),$$

$$(3.5) \quad [x_{i,r+1}^\pm, x_{js}^\pm] - [x_{ir}^\pm, x_{j,s+1}^\pm] = \pm \hbar d_{ij} (x_{ir}^\pm x_{js}^\pm + x_{js}^\pm x_{ir}^\pm),$$

$$(3.6) \quad \sum_{\pi \in S_m} \left[x_{i,r_{\pi(1)}}^\pm, \left[x_{i,r_{\pi(2)}}^\pm, \dots, \left[x_{i,r_{\pi(m)}}^\pm, x_{js}^\pm \right] \dots \right] \right] = 0,$$

where in the last relation $i \neq j$, $m = 1 - a_{ij}$, and $r_1, \dots, r_m \in \mathbb{N}$.

The Yangian $Y_{\hbar}(\mathfrak{g})$ is an \mathbb{N} -graded $\mathbb{C}[\hbar]$ -algebra, with grading

$$Y_{\hbar}(\mathfrak{g}) = \bigoplus_{k \in \mathbb{N}} Y_{\hbar}(\mathfrak{g})_k$$

determined by $\deg x_{ir}^\pm = \deg h_{ir} = r$ for all $i \in I$ and $r \in \mathbb{N}$. Moreover, Definition 6.1 is such that $Y_{\hbar}(\mathfrak{g})$ provides an \mathbb{N} -graded $\mathbb{C}[\hbar]$ -algebra deformation of the enveloping algebra $U(\mathfrak{t}_+)$, where we recall that $\mathfrak{t}_+ = \mathfrak{g}[t]$. Indeed, the identification

$Y_{\hbar}(\mathfrak{g})/\hbar Y_{\hbar}(\mathfrak{g}) \cong U(\mathfrak{t}_+)$ is induced by the graded algebra epimorphism $q : Y_{\hbar}(\mathfrak{g}) \rightarrow U(\mathfrak{t}_+)$ given on generators by

$$q : x_{ir}^{\pm} \mapsto x_i^{\pm} t^r, \quad h_{ir} \mapsto h_i t^r \quad \forall i \in \mathbf{I} \text{ and } r \in \mathbb{N}.$$

In addition, the relations (3.1)–(3.6) imply that the assignment

$$x_i^{\pm} \mapsto x_{i0}^{\pm}, \quad h_i \mapsto h_{i0} \quad \forall i \in \mathbf{I}$$

determines a \mathbb{C} -algebra homomorphism $U(\mathfrak{g}) \rightarrow Y_{\hbar}(\mathfrak{g})$, which is injective as its composition with q is the identity map $1_{U(\mathfrak{g})}$ on $U(\mathfrak{g})$. Henceforth, we shall identify \mathfrak{g} with its image in $Y_{\hbar}(\mathfrak{g})$ without further comment.

To specify the standard Hopf algebra structure on $Y_{\hbar}(\mathfrak{g})$, we first note that $Y_{\hbar}(\mathfrak{g})$ is generated as a $\mathbb{C}[\hbar]$ -algebra by the set $\mathfrak{g} \cup \{t_{i1}\}_{i \in \mathbf{I}} \subset Y_{\hbar}(\mathfrak{g})$, where

$$t_{i1} := h_{i1} - \frac{\hbar}{2} h_{i0}^2 \quad \forall i \in \mathbf{I}.$$

More precisely, for each $s > 0$, x_{is}^{\pm} and $h_{i,s+1}$ are determined by the recursive formulas

$$x_{is}^{\pm} = \pm \frac{1}{2d_i} [t_{i1}, x_{i,s-1}^{\pm}] \quad \text{and} \quad h_{i,s+1} = [x_{is}^+, x_{i1}^-].$$

Now let $r \in \mathfrak{n}_- \otimes \mathfrak{n}_+$ denote the canonical tensor associated with the pairing $(,)|_{\mathfrak{n}_- \times \mathfrak{n}_+}$, where $\mathfrak{n}_{\pm} = \bigoplus_{\alpha \in \Delta^{\pm}} \mathfrak{g}_{\pm\alpha}$ is the Lie subalgebra of \mathfrak{g} generated by $\{x_i^{\pm}\}_{i \in \mathbf{I}}$. Equivalently, r is the unique preimage of the identity map under the natural isomorphism $\mathfrak{n}_- \otimes \mathfrak{n}_+ \xrightarrow{\sim} \text{End}_{\mathbb{C}}(\mathfrak{n}_-)$, determined by $(,)|_{\mathfrak{n}_- \times \mathfrak{n}_+}$. In addition, we set

$$r_i := [h_i \otimes 1, r] \quad \forall i \in \mathbf{I}.$$

If $x_{\alpha}^{\pm} \in \mathfrak{g}_{\pm\alpha}$ are root vectors satisfying $(x_{\alpha}^+, x_{\alpha}^-) = 1$, then one has the formulae

$$r = \sum_{\alpha \in \Delta^+} x_{\alpha}^- \otimes x_{\alpha}^+ \quad \text{and} \quad r_i = - \sum_{\alpha \in \Delta^+} \alpha(h_i) x_{\alpha}^- \otimes x_{\alpha}^+.$$

The following proposition describes the Hopf algebra structure on $Y_{\hbar}(\mathfrak{g})$, where $m : Y_{\hbar}(\mathfrak{g})^{\otimes 2} \rightarrow Y_{\hbar}(\mathfrak{g})$ denote the multiplication map.

Proposition 3.2 *The Yangian $Y_{\hbar}(\mathfrak{g})$ is an \mathbb{N} -graded Hopf algebra with counit ε , coproduct Δ , and antipode S uniquely determined by the requirement that \mathfrak{g} is primitive and that, for each $i \in \mathbf{I}$, one has*

$$\varepsilon(t_{i1}) = 0, \quad \Delta(t_{i1}) = t_{i1} \otimes 1 + 1 \otimes t_{i1} + \hbar r_i, \quad S(t_{i1}) = -t_{i1} + m(\hbar r_i).$$

In particular, $Y_{\hbar}(\mathfrak{g})$ is an \mathbb{N} -graded Hopf algebra deformation of $U(\mathfrak{t}_+)$ over $\mathbb{C}[\hbar]$.

The crux of the proof of this proposition lies in showing that Δ is an algebra homomorphism. Though this is a consequence of [6, Theorem 2] and [8, Theorem 1] (see also [22, Theorem 2.6]), a complete proof has only recently appeared in [21] (see Theorem 4.9 therein).

The Yangian $Y_{\hbar}(\mathfrak{g})$ also admits a Q -grading compatible with the above \mathbb{N} -grading; that is to say, it is $\mathbb{N} \times Q$ -graded as a Hopf algebra. This Q -grading arises from the

adjoint action of the Cartan subalgebra $\mathfrak{h} \subset \mathfrak{g}$ on $Y_{\hbar}(\mathfrak{g})$. Namely, one has $Y_{\hbar}(\mathfrak{g}) = \bigoplus_{\beta \in Q} Y_{\hbar}(\mathfrak{g})_{\beta}$, where $Y_{\hbar}(\mathfrak{g})_{\beta}$ is just the β -weight space

$$Y_{\hbar}(\mathfrak{g})_{\beta} = \{x \in Y_{\hbar}(\mathfrak{g}) : [h, x] = \beta(h)x \quad \forall h \in \mathfrak{h}\} \quad \forall \beta \in Q.$$

3.3 Automorphisms

There are two families of (anti)automorphisms of $Y_{\hbar}(\mathfrak{g})$ which will play an especially pronounced role in this article: the *shift* automorphisms and the *Chevalley involution*. The former are a family $\{\tau_c\}_{c \in \mathbb{C}} \subset \text{Aut}(Y_{\hbar}(\mathfrak{g}))$ which give rise to an action of the additive group \mathbb{C} on $Y_{\hbar}(\mathfrak{g})$ by Hopf algebra automorphisms. In more detail, τ_c is defined explicitly by

$$(3.7) \quad \tau_c(x_i^{\pm}(u)) = x_i^{\pm}(u - c) \quad \text{and} \quad \tau_c(h_i(u)) = h_i(u - c) \quad \forall i \in \mathbf{I},$$

where we have introduced the generating series $x_i^{\pm}(u)$ and $h_i(u)$ in $Y_{\hbar}(\mathfrak{g})[[u^{-1}]]$ by

$$x_i^{\pm}(u) = \sum_{r \in \mathbb{N}} x_{i,r}^{\pm} u^{-r-1} \quad \text{and} \quad h_i(u) = \sum_{r \in \mathbb{N}} h_{i,r} u^{-r-1}.$$

Replacing c by a formal variable z in (3.7), one obtains an \mathbb{N} -graded embedding

$$(3.8) \quad \tau_z : Y_{\hbar}(\mathfrak{g}) \hookrightarrow Y_{\hbar}(\mathfrak{g})[z]$$

called the *formal shift homomorphism*, where $\deg z = 1$. Let us now turn to defining the Chevalley involution, beginning with the following lemma.

Lemma 3.3 *The assignments ω and ζ defined by*

$$\begin{aligned} \omega(x_i^{\pm}(u)) &= x_i^{\mp}(u), & \omega(h_i(u)) &= h_i(u), \\ \zeta(x_i^{\pm}(u)) &= x_i^{\pm}(-u), & \zeta(h_i(u)) &= h_i(-u), \end{aligned}$$

extend to commuting anti-involutions ω and ζ of $Y_{\hbar}(\mathfrak{g})$. Moreover, ω and ζ satisfy

$$\begin{aligned} \tau_c \circ \omega &= \omega \circ \tau_c, & \tau_{-c} \circ \zeta &= \zeta \circ \tau_c \quad \forall c \in \mathbb{C}, \\ \varepsilon \circ \omega &= \varepsilon, & (\omega \otimes \omega) \circ \Delta &= \Delta^{\text{op}} \circ \omega, & \omega \circ S &= S \circ \omega, \\ \varepsilon \circ \zeta &= \varepsilon, & (\zeta \otimes \zeta) \circ \Delta &= \Delta \circ \zeta, & \zeta \circ S^{-1} &= S \circ \zeta. \end{aligned}$$

This result, which has appeared in various forms in the literature (for instance, [3, Proposition 2.9]), is readily established using Definition 6.1 and the relations of Proposition 3.2. By the lemma, ω is an involutive Hopf algebra anti-automorphism of $Y_{\hbar}(\mathfrak{g})$, which we call the *Chevalley involution* of $Y_{\hbar}(\mathfrak{g})$. On $\mathfrak{g} \subset Y_{\hbar}(\mathfrak{g})$, this recovers the standard Chevalley involution, given by

$$\omega(x_i^{\pm}) = x_i^{\mp} \quad \text{and} \quad \omega(h_i) = h_i \quad \forall i \in \mathbf{I}.$$

Similarly, under the identification $Y_{\hbar}(\mathfrak{g})/\hbar Y_{\hbar}(\mathfrak{g}) \cong U(\mathfrak{t}_+)$, the semiclassical limit $\bar{\omega} : U(\mathfrak{t}_+) \rightarrow U(\mathfrak{t}_+)$ of ω coincides with the anti-involution of $U(\mathfrak{t}_+)$ uniquely extending the Lie algebra anti-automorphism

$$(3.9) \quad \bar{\omega}(xt^r) = \omega(x)t^r \quad \forall x \in \mathfrak{g} \text{ and } r \in \mathbb{N}.$$

In addition, we note that the composite $\kappa := \omega \circ \zeta$ is an involutive algebra automorphism of $Y_{\hbar}(\mathfrak{g})$, given explicitly by

$$(3.10) \quad \kappa(x_i^\pm(u)) = x_i^\mp(-u) \quad \text{and} \quad \kappa(h_i(u)) = h_i(-u) \quad \forall i \in I.$$

This automorphism is itself often called the Chevalley or Cartan involution of $Y_{\hbar}(\mathfrak{g})$, though here we shall reserve the former terminology for ω .

3.4 Poincaré–Birkhoff–Witt theorem

An important foundational result in the theory of Yangians is the Poincaré–Birkhoff–Witt theorem, which asserts the flatness of $Y_{\hbar}(\mathfrak{g})$ as an \mathbb{N} -graded Hopf algebra deformation of $U(\mathfrak{t}_+)$. It can be stated concisely as follows.

Theorem 3.4 *The Yangian $Y_{\hbar}(\mathfrak{g})$ is a torsion-free $\mathbb{C}[\hbar]$ -module, and thus provides a flat deformation of $Y_{\hbar}(\mathfrak{g})/\hbar Y_{\hbar}(\mathfrak{g}) \cong U(\mathfrak{t}_+)$ as a graded Hopf algebra over $\mathbb{C}[\hbar]$. In particular, $Y_{\hbar}(\mathfrak{g})$ is isomorphic to $U(\mathfrak{t}_+)[\hbar]$ as an \mathbb{N} -graded $\mathbb{C}[\hbar]$ -module.*

As $Y_{\hbar}(\mathfrak{g})$ is an \mathbb{N} -graded algebra deformation of $U(\mathfrak{t}_+)$, an isomorphism $Y_{\hbar}(\mathfrak{g}) \cong U(\mathfrak{t}_+)[\hbar]$ can be obtained by specifying an ordered, homogeneous, lift $\mathbb{G} \subset Y_{\hbar}(\mathfrak{g})$ of any fixed homogeneous basis of the Lie algebra \mathfrak{t}_+ . For our purposes, it will be useful to specify a class of isomorphisms of this type with a number of useful properties.

For each $\beta \in \Delta^+$, we may choose $i(\beta) \in I$ and $\mathbf{X}^\beta \in U(\mathfrak{n}_+)_{\beta - \alpha_{i(\beta)}} \subset U(\mathfrak{g})$ such that

$$(3.11) \quad x_\beta^+ := \mathbf{X}^\beta \cdot x_{i(\beta)}^+ \in \mathfrak{g}_\beta \quad \text{and} \quad x_\beta^- := \omega(x_\beta^+) \in \mathfrak{g}_{-\beta}$$

satisfy the duality condition $(x_\beta^+, x_\beta^-) = 1$, where \mathbf{X}^β acts on $x_{i(\beta)}^+$ via the adjoint action of \mathfrak{g} on $U(\mathfrak{g})$. In particular, we can (and shall) take $\mathbf{X}^{\alpha_i} = 1$ for all $i \in I$, so that $x_{\alpha_i}^\pm = x_i^\pm$. For each $k \in \mathbb{N}$, we then set

$$x_{\beta,k}^+ := \mathbf{X}^\beta \cdot x_{i(\beta),k}^+ \in Y_{\hbar}(\mathfrak{g})_\beta \quad \text{and} \quad x_{\beta,k}^- := \omega(x_{\beta,k}^+) \in Y_{\hbar}(\mathfrak{g})_{-\beta},$$

where \mathfrak{g} now operates on $Y_{\hbar}(\mathfrak{g})$ via the adjoint action. This definition is such that $q(x_{\beta,k}^\pm) = x_\beta^\pm t^k$ for all $\beta \in \Delta^+$ and $k \in \mathbb{N}$, and hence the set of elements

$$\mathbb{G} := \bigcup_{k \in \mathbb{N}} \{h_{ik}, x_{\beta,k}^\pm\}_{i \in I, \beta \in \Delta^+}$$

reduces modulo \hbar to the basis of \mathfrak{t}_+ consisting of all Cartan elements $h_i t^k$ and root vector $x_\beta^\pm t^k$. For each choice of total order \leq on \mathbb{G} , the corresponding set of ordered monomials

$$B(\mathbb{G}) = \{x_1 x_2 \cdots x_n : n \in \mathbb{N}, x_i \in \mathbb{G} \text{ and } x_i \leq x_j \ \forall i < j\}$$

is therefore a homogeneous basis of the $\mathbb{C}[\hbar]$ -module $Y_{\hbar}(\mathfrak{g})$, and so defines an isomorphism \mathbb{N} -graded modules

$$(3.12) \quad v_{\mathbb{G}} : Y_{\hbar}(\mathfrak{g}) \xrightarrow{\sim} U(\mathfrak{t}_+)[\hbar]$$

uniquely determined by the property that $v_{\mathbb{G}}|_{B(\mathbb{G})}$ coincides with the restriction of the quotient map q to $B(\mathbb{G})$. We note that $v_{\mathbb{G}}$ is automatically an isomorphism of \mathfrak{h} -modules, and is thus Q -graded.

We shall single out a subclass of isomorphisms of this type which are compatible with Chevalley involutions and satisfy a triangularity condition. To make this precise, we must first recall that $Y_{\hbar}(\mathfrak{g})$ admits a triangular decomposition, compatible with the decomposition

$$\mathfrak{g} = \mathfrak{n}_+ \oplus \mathfrak{h} \oplus \mathfrak{n}_-$$

Let us define $Y_{\hbar}^0(\mathfrak{g})$ and $Y_{\hbar}^{\pm}(\mathfrak{g})$ to be the unital associative subalgebras of $Y_{\hbar}(\mathfrak{g})$ generated by $\{h_{ir}\}_{i \in I, r \in \mathbb{N}}$ and $\{x_{ir}^{\pm}\}_{i \in I, r \in \mathbb{N}}$, respectively. These are $\mathbb{N} \times Q$ -graded subalgebras of $Y_{\hbar}(\mathfrak{g})$. The triangular decomposition of $Y_{\hbar}(\mathfrak{g})$ is then encoded by the following proposition, which is a well-known consequence of Theorem 3.4.

Proposition 3.5

- (1) $Y_{\hbar}^{\pm}(\mathfrak{g})$ is isomorphic to the unital, associative $\mathbb{C}[\hbar]$ -algebra generated by the set $\{x_{ir}^{\pm}\}_{i \in I, r \in \mathbb{N}}$, subject to relations (3.5) and (3.6) of Definition 6.1:

$$\begin{aligned} [x_{i,r+1}^{\pm}, x_{js}^{\pm}] - [x_{ir}^{\pm}, x_{j,s+1}^{\pm}] &= \pm \hbar d_{ij}(x_{ir}^{\pm} x_{js}^{\pm} + x_{js}^{\pm} x_{ir}^{\pm}), \\ \sum_{\pi \in S_m} [x_{i,r_{\pi(1)}}^{\pm}, [x_{i,r_{\pi(2)}}^{\pm}, \dots, [x_{i,r_{\pi(m)}}^{\pm}, x_{js}^{\pm}] \dots]] &= 0, \end{aligned}$$

where all indices are constrained as in Definition 6.1. In particular, $Y_{\hbar}^{\pm}(\mathfrak{g})$ is an \mathbb{N} -graded, torsion-free $\mathbb{C}[\hbar]$ -algebra deformation of $U(\mathfrak{n}_{\pm}[t])$.

- (2) The assignment $h_{ik} \mapsto h_i t^k$, for all $i \in I$ and $k \in \mathbb{N}$, extends to an isomorphism of \mathbb{N} -graded, commutative $\mathbb{C}[\hbar]$ -algebras

$$\xi : Y_{\hbar}^0(\mathfrak{g}) \xrightarrow{\sim} U(\mathfrak{h}[t])[\hbar] = S(\mathfrak{h}[t])[\hbar].$$

- (3) The multiplication map

$$m : Y_{\hbar}^+(\mathfrak{g}) \otimes Y_{\hbar}^0(\mathfrak{g}) \otimes Y_{\hbar}^-(\mathfrak{g}) \rightarrow Y_{\hbar}(\mathfrak{g})$$

is an isomorphism of graded $\mathbb{C}[\hbar]$ -modules.

As a consequence of Part (1) of Proposition 3.5 and Corollary 2.6, one has $Y_{\hbar}^{\pm}(\mathfrak{g}) \cong U(\mathfrak{n}_{\pm}[t])[\hbar]$ as \mathbb{N} -graded $\mathbb{C}[\hbar]$ -modules. Following the procedure outlined at the beginning of the section, let us fix an arbitrary total order \leq_+ on the union

$$\mathbb{G}_+ = \bigcup_{k \in \mathbb{N}} \{x_{\beta,k}^+\}_{\beta \in \Delta^+} = \mathbb{G} \cap Y_{\hbar}^+(\mathfrak{g}).$$

The set of ordered monomials $B(\mathbb{G}_+)$ in \mathbb{G}_+ is a basis of $Y_{\hbar}^+(\mathfrak{g})$, and thus gives rise to an isomorphism of $\mathbb{N} \times Q$ -graded $\mathbb{C}[\hbar]$ -modules

$$v_+ : Y_{\hbar}^+(\mathfrak{g}) \xrightarrow{\sim} U(\mathfrak{n}_+[t])[\hbar],$$

sending each ordered monomial in \mathbb{G}_+ to its image in $Y_{\hbar}^+(\mathfrak{g})/\hbar Y_{\hbar}^+(\mathfrak{g}) \cong U(\mathfrak{n}_+[t])$. Using the Chevalley involution ω and its semiclassical limit $\tilde{\omega}$ (see (3.9)), we then obtain an isomorphism

$$v_- := \tilde{\omega} \circ v_+ \circ \omega : Y_{\hbar}^-(\mathfrak{g}) \xrightarrow{\sim} U(\mathfrak{n}_-[t])[\hbar].$$

Combining v_{\pm} with ξ from Part (2) of Proposition 3.5 outputs an isomorphism of $\mathbb{N} \times Q$ -graded $\mathbb{C}[[\hbar]]$ -modules

$$(3.13) \quad v := \tilde{m} \circ (v_+ \otimes \xi \otimes v_-) \circ m^{-1} : Y_{\hbar}(\mathfrak{g}) \xrightarrow{\sim} U(\mathfrak{t}_+)[[\hbar]],$$

where $\tilde{m} : U(\mathfrak{n}_+[t]) \otimes_{\mathbb{C}} U(\mathfrak{h}[t]) \otimes_{\mathbb{C}} U(\mathfrak{n}_-[t]) \xrightarrow{\sim} U(\mathfrak{t}_+)$ is the multiplication map, which we extend trivially by $\mathbb{C}[[\hbar]]$ -linearity. By construction, v is compatible with the underlying triangular decompositions on $Y_{\hbar}(\mathfrak{g})$ and $U(\mathfrak{t}_+)$ and satisfies

$$v \circ \omega = \tilde{\omega} \circ v.$$

The definition (3.13) is such that $v = v_{\mathbb{G}}$ for any total order \leq on \mathbb{G} which restricts to \leq_+ , satisfies $x^+ \leq h \leq x^-$ for all $x^{\pm} \in \mathbb{G} \cap Y_{\hbar}^{\pm}(\mathfrak{g})$ and $h \in Y_{\hbar}^0(\mathfrak{g})$, and for which ω is a decreasing function on \mathbb{G} . We will denote the inverse of v by μ :

$$\mu := v^{-1} : U(\mathfrak{t}_+)[[\hbar]] \rightarrow Y_{\hbar}(\mathfrak{g}).$$

Note that, for any total order on \mathbb{G} , one has $\mu(x) = v_{\mathbb{G}}^{-1}(x)$ for all $x \in \mathbb{G}$.

3.5 Quantization

As a consequence of Proposition 3.2 and Theorem 3.4, the Yangian $Y_{\hbar}(\mathfrak{g})$ provides a homogeneous quantization of an \mathbb{N} -graded Lie bialgebra structure on the Lie algebra \mathfrak{t}_+ over $\mathbb{C}[[\hbar]]$, with cocommutator δ determined by the formula (2.2). By Proposition 3.2, δ is uniquely determined by $\delta(\mathfrak{g}) = 0$ and

$$\delta(h_i t) = r_i - r_i^{21} = [h_i \otimes 1, \Omega_{\mathfrak{g}}] = \left[h_i t \otimes 1 + 1 \otimes h_i s, \frac{\Omega_{\mathfrak{g}}}{t-s} \right] = \delta_+(h_i t) \quad \forall i \in I,$$

and thus coincides with δ_+ from Section 2.5. This recovers the following well-known result, originally due to Drinfeld [6, Theorem 2]:

Theorem 3.6 *$Y_{\hbar}(\mathfrak{g})$ is a homogeneous quantization of $(\mathfrak{t}_+, \delta_+)$ over $\mathbb{C}[[\hbar]]$.*

As explained in Section 2.4, it follows immediately that the \hbar -adic completion

$$(3.14) \quad Y_{\hbar} \mathfrak{g} := \varprojlim_n (Y_{\hbar}(\mathfrak{g}) / \hbar^n Y_{\hbar}(\mathfrak{g}))$$

is a homogeneous quantization of $(\mathfrak{t}_+, \delta_+)$ over $\mathbb{C}[[\hbar]]$. We refer the reader to Definition 2.12 and Corollary 2.6 for a detailed discussion of this point.

Remark 3.7 Let $Y_{\hbar}^{\pm} \mathfrak{g}$ and $Y_{\hbar}^0 \mathfrak{g}$ denote the topological $\mathbb{C}[[\hbar]]$ -algebras

$$Y_{\hbar}^{\pm} \mathfrak{g} := \varprojlim_n Y_{\hbar}^{\pm}(\mathfrak{g}) / \hbar^n Y_{\hbar}^{\pm}(\mathfrak{g}) \quad \text{and} \quad Y_{\hbar}^0 \mathfrak{g} := \varprojlim_n Y_{\hbar}^0(\mathfrak{g}) / \hbar^n Y_{\hbar}^0(\mathfrak{g}).$$

It follows from Corollary 2.6 and Proposition 3.5 that these are subalgebras of $Y_{\hbar} \mathfrak{g}$, with $Y_{\hbar}^0 \mathfrak{g}$ isomorphic to $U(\mathfrak{h}[t])[[\hbar]] \cong S(\mathfrak{h}[t])[[\hbar]]$ as an \mathbb{N} -graded topological $\mathbb{C}[[\hbar]]$ -algebra, and $Y_{\hbar}^{\pm} \mathfrak{g}$ a topologically free \mathbb{N} -graded $\mathbb{C}[[\hbar]]$ -algebra with semiclassical limit equal to $U(\mathfrak{n}_{\pm}[t])$. By Part (3) of Proposition 3.5, the product m on $Y_{\hbar} \mathfrak{g}$ gives rise to an isomorphism of \mathbb{N} -graded topological $\mathbb{C}[[\hbar]]$ -modules

$$(3.15) \quad m : Y_{\hbar}^+ \mathfrak{g} \otimes Y_{\hbar}^0 \mathfrak{g} \otimes Y_{\hbar}^- \mathfrak{g} \xrightarrow{\sim} Y_{\hbar} \mathfrak{g},$$

where, following Remark 2.11, \otimes should now be understood to be the topological tensor product $\widehat{\otimes}$ of $\mathbb{C}[[\hbar]]$ -modules. For later purposes, we note that the product on $Y_{\hbar}\mathfrak{g}$ also gives rise to an isomorphism

$$Y_{\hbar}^{-}\mathfrak{g} \otimes Y_{\hbar}^0\mathfrak{g} \otimes Y_{\hbar}^{+}\mathfrak{g} \xrightarrow{\sim} Y_{\hbar}\mathfrak{g}$$

which can be realized as $\kappa \circ m \circ (\kappa_{-} \otimes \kappa_0 \otimes \kappa_{+})$, where m is the isomorphism (3.15), κ is the involutive automorphism of $Y_{\hbar}\mathfrak{g}$ defined in (3.10) (extended by continuity), and $\kappa_{\chi} := \kappa|_{Y_{\hbar}^{\chi}\mathfrak{g}} : Y_{\hbar}^{\chi}\mathfrak{g} \xrightarrow{\sim} Y_{\hbar}^{-\chi}\mathfrak{g}$.

3.6 The universal R-matrix

We complete our survey of $Y_{\hbar}(\mathfrak{g})$ by reviewing the construction of the universal R-matrix $\mathcal{R}(z)$ of the Yangian, whose existence and uniqueness was first established by Drinfeld in [6, Theorem 3]. We shall, however, need a refined version of Drinfeld’s theorem only recently proven in [17, Section 7.4], which reconstructs $\mathcal{R}(z)$ from the factors in its Gauss decomposition

$$\mathcal{R}(z) = \mathcal{R}^{+}(z)\mathcal{R}^0(z)\mathcal{R}^{-}(z).$$

Let us begin with a few preliminaries. For each positive integer n , let

$$Y_{\hbar}(\mathfrak{g})^{\otimes n}[z; z^{-1}] = \bigcup_{k \in \mathbb{N}} z^k Y_{\hbar}(\mathfrak{g})^{\otimes n}[[z^{-1}]] \subset Y_{\hbar}(\mathfrak{g})^{\otimes n}[[z^{\pm 1}]]$$

denote the algebra of formal Laurent series in z^{-1} with coefficients in $Y_{\hbar}(\mathfrak{g})^{\otimes n}$. Following [41, Section 4.2], we then introduce the subspace

$$\widehat{Y_{\hbar}\mathfrak{g}}_z^{(n)} := \prod_{k \in \mathbb{N}} (Y_{\hbar}(\mathfrak{g})^{\otimes n})_k z^{-k} \subset Y_{\hbar}(\mathfrak{g})^{\otimes n}[[z^{-1}]],$$

where $(Y_{\hbar}(\mathfrak{g})^{\otimes n})_k$ is the k th graded component of the \mathbb{N} -graded algebra $Y_{\hbar}(\mathfrak{g})^{\otimes n}$. This is a \mathbb{C} -algebra isomorphic to the completion of $Y_{\hbar}(\mathfrak{g})^{\otimes n}$ with respect to its grading. The $\mathbb{C}[z, z^{-1}]$ -submodule of $Y_{\hbar}(\mathfrak{g})^{\otimes n}[z; z^{-1}]$ that it generates is a \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -algebra

$$(3.16) \quad \mathbb{L}(\widehat{Y_{\hbar}\mathfrak{g}}_z^{(n)}) := \bigoplus_{k \in \mathbb{Z}} z^k \widehat{Y_{\hbar}\mathfrak{g}}_z^{(n)} \subset Y_{\hbar}(\mathfrak{g})^{\otimes n}[z; z^{-1}].$$

Though for the moment we shall only be interested in the case where $n = 2$, such formal series spaces will reappear in later sections. In addition to the above, we shall make use of two functions $Q_{+} \rightarrow \mathbb{N}$. First, we have the standard additive height function ht given by

$$\text{ht}(\beta) = \sum_{i \in I} n_i \quad \text{for each } \beta = \sum_{i \in I} n_i \alpha_i \in Q_{+}.$$

Second, we have an auxiliary function $\nu : Q_{+} \rightarrow \mathbb{N}$ defined by

$$(3.17) \quad \nu(\beta) = \min\{k \in \mathbb{N} : \exists \beta_1, \dots, \beta_k \in \Delta^{+} \text{ with } \beta = \beta_1 + \dots + \beta_k\},$$

where it is understood that $\nu(0) = 0$.

Let us now recall the construction of the factor $\mathcal{R}^-(z)$. Fix a Cartan element

$$\zeta \in \mathfrak{h} \setminus \bigcup_{\beta \neq 0} \text{Ker}(\beta),$$

where the union runs over all nonzero $\beta \in Q_+$. We then introduce

$$\mathcal{R}_\beta^-(z) \in (Y_{\hbar}^-(\mathfrak{g})_{-\beta} \otimes Y_{\hbar}^+(\mathfrak{g})_{\beta})[[z^{-1}]] \quad \forall \beta \in Q_+$$

by setting $\mathcal{R}_0^-(z) = 1$ and defining $\mathcal{R}_\beta^-(z)$ inductively in $\text{ht}(\beta)$ using the formula

$$(3.18) \quad \mathcal{R}_\beta^-(z) = \hbar \sum_{p \geq 0} \frac{\mathbf{T}(\zeta)^p}{(z\beta(\zeta))^{p+1}} \sum_{\alpha \in \Delta^+} \alpha(\zeta) \mathcal{R}_{\beta-\alpha}^-(z) (x_\alpha^- \otimes x_\alpha^+),$$

where $\mathcal{R}_\gamma^-(z) = 0$ whenever $\gamma \notin Q_+$ and $\mathbf{T}(\zeta) = \text{ad}(\mathbf{T}(\zeta) \otimes 1 + 1 \otimes \mathbf{T}(\zeta))$, with $\mathbf{T} : \mathfrak{h} \rightarrow Y_{\hbar}^0(\mathfrak{g})$ the embedding determined by

$$\mathbf{T}(h_i) = t_{i1} \quad \forall i \in \mathbf{I}.$$

Using the fact that, for each $p \in \mathbb{N}$, $\mathbf{T}(\zeta)^p z^{-p-1}$ is a homogeneous operator on $\mathbb{L}(\widehat{Y_{\hbar} \mathfrak{g}_z^{(2)}})$ of degree -1 , one deduces from the recursive formula (3.18) that

$$\mathcal{R}_\beta^-(z) \in \hbar^{v(\beta)} \mathbb{L}(\widehat{Y_{\hbar} \mathfrak{g}_z^{(2)}})_{-v(\beta)} \subset z^{-v(\beta)} Y_{\hbar}(\mathfrak{g})^{\otimes 2}[[z^{-1}]] \quad \forall \beta \in Q_+.$$

As the set $\{\beta \in Q_+ : v(\beta) \leq k\}$ is finite for any $k \in \mathbb{N}$, we obtain a well-defined formal series

$$\mathcal{R}^-(z) = \sum_{\beta \in Q_+} \mathcal{R}_\beta^-(z) \in (Y_{\hbar}^-(\mathfrak{g}) \otimes Y_{\hbar}^+(\mathfrak{g}))[[z^{-1}]]$$

which by construction satisfies $\mathcal{R}^-(z) \in 1 + \hbar \mathbb{L}(\widehat{Y_{\hbar} \mathfrak{g}_z^{(2)}})_{-1} \subset \widehat{Y_{\hbar} \mathfrak{g}_z^{(2)}}$.

By Theorem 4.1 of [17], $\mathcal{R}^-(z)$ is independent of the choice of $\zeta \in \mathfrak{h}$ made above and satisfies a number of remarkable properties. Notably, it intertwines $\tau_z \otimes 1 \circ \Delta$ and the formal, deformed Drinfeld coproduct Δ_z^D on $Y_{\hbar}(\mathfrak{g})$, as defined in [17, Section 3.4]. We will not make direct use of these properties here, and refer the reader to [17] for a detailed treatment of $\mathcal{R}^-(z)$.

Let us now recall the definition of the abelian R -matrix $\mathcal{R}^0(z) \in Y_{\hbar}^0(\mathfrak{g})^{\otimes 2}[[z^{-1}]]$ from [17, Section 6]. Let $\mathbf{B} = (d_i a_{ij})$ denote the symmetrization of the Cartan matrix \mathbf{A} . Given an indeterminate v , we let $\mathbf{B}(v) = ([d_i a_{ij}]_v) \subset \text{GL}_1(\mathbb{Q}(v))$ be the associated matrix of v -numbers, where

$$[m]_v = \frac{v^m - v^{-m}}{v - v^{-1}}.$$

Then it is known [16, Theorem A.1] that the auxiliary matrix

$$\mathbf{C}(v) = (c_{ij}(v)) = [2\kappa]_v \mathbf{B}(v)^{-1}$$

has entries $c_{ij}(v)$ in $\mathbb{N}[v, v^{-1}]$, where 4κ is the eigenvalue of the Casimir element of \mathfrak{g} in the adjoint representation.

Next, for each index $i \in \mathbf{I}$, we introduce the series $t_i(u) \in \hbar Y_{\hbar}^0(\mathfrak{g})[[u^{-1}]]$ and its inverse Borel transform $B_i(u) \in \hbar Y_{\hbar}^0(\mathfrak{g})[[u]]$ by

$$t_i(u) = \hbar \sum_{r \geq 0} t_{ir} u^{-r-1} = \log(1 + \hbar h_i(u)) \quad \text{and} \quad B_i(u) = \hbar \sum_{r \geq 0} \frac{t_{ir}}{r!} u^r.$$

Note that t_{i1} coincides with the element of the same name introduced in Section 3.2. From these data, we obtain an element $\mathcal{L}(z) \in (\hbar/z)^2 Y_{\hbar}^0(\mathfrak{g})^{\otimes 2}[[z^{-1}]]$ defined by

$$\mathcal{L}(z) = T^{2\kappa} \sum_{i,j \in \mathbf{I}} c_{ij}(T) B_i(\partial_z) \otimes B_j(-\partial_z)(-z^{-2}),$$

where T is the shift operator $T(f(z)) = f(z + \frac{\hbar}{2})$ on $Y_{\hbar}(\mathfrak{g})^{\otimes 2}[[z^{-1}]]$. Equivalently,

$$T = \exp\left(\frac{\hbar}{2} \partial_z\right) : Y_{\hbar}(\mathfrak{g})^{\otimes 2}[[z^{-1}]] \rightarrow Y_{\hbar}(\mathfrak{g})^{\otimes 2}[[z^{-1}]].$$

As $\mathcal{L}(z) \in \hbar z^{-2} Y_{\hbar}^0(\mathfrak{g})^{\otimes 2}[[z^{-1}]]$ and $Y_{\hbar}^0(\mathfrak{g})^{\otimes 2}$ is torsion-free, there is a unique solution $\mathcal{S}(z)$ to the formal difference equation

$$\mathcal{L}(z) = \mathcal{S}(z + 2\kappa\hbar) - \mathcal{S}(z) \quad \text{with} \quad \mathcal{S}(z) \in z^{-1} Y_{\hbar}^0(\mathfrak{g})^{\otimes 2}[[z^{-1}]].$$

If $g(z) \in z^{-1}\mathbb{C}[[z^{-1}]]$ is the unique solution of $-z^{-2} = g(z+1) - g(z)$, then by Proposition 6.6 of [17], we have

$$(3.19) \quad \mathcal{S}(z) = \frac{T^{2\kappa}}{(2\kappa\hbar)^2} \sum_{i,j \in \mathbf{I}} c_{ij}(T) B_i(\partial_z) \otimes B_j(-\partial_z) \left(g\left(\frac{z}{2\kappa\hbar}\right) \right),$$

where $g(z/2\kappa\hbar)$ is viewed as an element of $\mathbb{L}(\widehat{Y_{\hbar}\mathfrak{g}_z^{(2)}})$. The abelian R -matrix $\mathcal{R}^0(z)$ is defined to be the formal series exponential of this solution:

$$\mathcal{R}^0(z) = \exp(\mathcal{S}(z)) \in 1 + z^{-1} Y_{\hbar}^0(\mathfrak{g})^{\otimes 2}[[z^{-1}]].$$

Equivalently, it is the unique formal solution in $1 + z^{-1} Y_{\hbar}^0(\mathfrak{g})^{\otimes 2}[[z^{-1}]]$ of the equation

$$\mathcal{R}^0(z + 2\kappa\hbar) = \mathcal{A}(z)\mathcal{R}^0(z),$$

where $\mathcal{A}(z) = \exp(\mathcal{L}(z))$. As T and $\hbar^{-2} B_i(\partial_z) \otimes B_j(-\partial_z)$ are homogeneous operators of degree zero on $\mathbb{L}(\widehat{Y_{\hbar}\mathfrak{g}_z^{(2)}})$, it follows from (3.19) that

$$\mathcal{R}^0(z) \in 1 + \hbar \mathbb{L}(\widehat{Y_{\hbar}\mathfrak{g}_z^{(2)}})_{-1} \subset \widehat{Y_{\hbar}\mathfrak{g}_z^{(2)}}.$$

We are now in a position to introduce the universal R -matrix of the Yangian. Set $\mathcal{R}^+(z) = \mathcal{R}_{21}^{-1}(-z)^{-1}$ and define

$$\mathcal{R}(z) := \mathcal{R}^+(z)\mathcal{R}^0(z)\mathcal{R}^-(z) \in 1 + z^{-1} Y_{\hbar}(\mathfrak{g})^{\otimes 2}[[z^{-1}]].$$

The following result is the content of Theorem 7.4 of [17].

Theorem 3.8 $\mathcal{R}(z)$ is the unique formal series in $1 + z^{-1} Y_{\hbar}(\mathfrak{g})^{\otimes 2}[[z^{-1}]]$ satisfying the intertwiner equation

$$(3.20) \quad \tau_z \otimes \mathbf{I} \circ \Delta^{\text{op}}(x) = \mathcal{R}(z) \cdot \tau_z \otimes \mathbf{I} \circ \Delta(x) \cdot \mathcal{R}(z)^{-1} \quad \forall x \in Y_{\hbar}(\mathfrak{g})$$

in $Y_{\hbar}(\mathfrak{g})^{\otimes 2}[[z; z^{-1}]]$, in addition to the cabling identities

$$\begin{aligned} \Delta \otimes \mathbf{1}(\mathcal{R}(z)) &= \mathcal{R}_{13}(z)\mathcal{R}_{23}(z), \\ \mathbf{1} \otimes \Delta(\mathcal{R}(z)) &= \mathcal{R}_{13}(z)\mathcal{R}_{12}(z) \end{aligned}$$

in $Y_{\hbar}(\mathfrak{g})^{\otimes 3}[[z^{-1}]]$. Moreover, $\mathcal{R}(z)$ has the following properties:

- (1) It is unitary: $\mathcal{R}(z)^{-1} = \mathcal{R}_{21}(-z)$.
- (2) For any $a, b \in \mathbb{C}$, one has

$$(\tau_a \otimes \tau_b)\mathcal{R}(z) = \mathcal{R}(z + a - b).$$

- (3) $\mathcal{R}(z)$ is a homogeneous, degree zero, element of $\mathbb{L}(\widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)})$, with

$$\mathcal{R}(z) - 1 \in \hbar \mathbb{L}(\widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)})_{-1} = \hbar z^{-1} \widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)}$$

and semiclassical limit given by

$$q^{\otimes 2} \hbar^{-1}(\mathcal{R}(z) - 1) = \frac{\Omega_{\mathfrak{g}}}{z + t - w} \in (U(\mathfrak{g}[t]) \otimes U(\mathfrak{g}[w]))[[z^{-1}]].$$

The series $\mathcal{R}(z)$ is called the *universal R-matrix* of the Yangian, and is related to the element $\mathcal{R}^D(z) \in Y_{\hbar}(\mathfrak{g})^{\otimes 2}[[z^{-1}]]$ introduced by Drinfeld in Theorem 3 of [6] by $\mathcal{R}(z) = \mathcal{R}^D(-z)^{-1}$ (see Section 1.1 and Corollary 7.4 of [17]).

Remark 3.9 Strictly speaking, the results of [17] are stated with \hbar replaced by an arbitrary nonzero complex number. However, it is easy to translate between the numerical and formal \hbar settings via a homogenization procedure, and for the sake of completeness we make this rigorous in Appendix Appendix A (see Proposition A.1).

The final result of this section shows that, in particular, $\mathcal{R}(z)$ is invariant under the Chevalley involution ω of Section 3.3.

Corollary 3.10 *The universal R-matrix $\mathcal{R}(z)$ satisfies*

$$(\omega \otimes \omega)\mathcal{R}(z) = \mathcal{R}(z) \quad \text{and} \quad (\varsigma \otimes \varsigma)\mathcal{R}(z) = \mathcal{R}_{21}(z) = (\kappa \otimes \kappa)\mathcal{R}(z).$$

Proof Set $\mathcal{R}^{\omega}(z) := (\omega \otimes \omega)\mathcal{R}(z)$ and $\mathcal{R}^{\varsigma}(z) := (\varsigma \otimes \varsigma)\mathcal{R}(z)$. Applying the anti-automorphisms $\omega \otimes \omega$ and $\varsigma \otimes \varsigma$ to the intertwiner equation (8.3), while making use of the relations of Lemma 3.3, we find that

$$\begin{aligned} \tau_z \otimes \mathbf{1} \circ \Delta(x) &= \mathcal{R}^{\omega}(z)^{-1} \cdot \tau_z \otimes \mathbf{1} \circ \Delta^{\text{op}}(x) \cdot \mathcal{R}^{\omega}(z), \\ \tau_{-z} \otimes \mathbf{1} \circ \Delta^{\text{op}}(x) &= \mathcal{R}^{\varsigma}(z)^{-1} \cdot \tau_{-z} \otimes \mathbf{1} \circ \Delta(x) \cdot \mathcal{R}^{\varsigma}(z) \end{aligned}$$

for all $x \in Y_{\hbar}(\mathfrak{g})$. Hence, $\mathcal{R}^{\omega}(z)$ and $\mathcal{R}^{\varsigma}(-z)^{-1} = \mathcal{R}_{21}^{\varsigma}(z)$ are both solutions of (8.3). One verifies similarly that these both satisfy the cabling identities, and hence coincide with $\mathcal{R}(z)$ by the uniqueness statement of Theorem 3.8. Since $\kappa = \varsigma \circ \omega$, this completes the proof of the proposition. ■

4 The Yangian double

We now recall the definition and main properties of the Yangian double $DY_{\hbar}\mathfrak{g}$, including a review of some of the results of [41]. These results, summarized in

Theorems 4.6 and 4.8, will play an integral role in establishing in Sections 7 and 8 that $DY_{\hbar}\mathfrak{g}$ is a homogeneous quantization of the Lie bialgebra $\mathfrak{t} = \mathfrak{g}[t^{\pm 1}]$ isomorphic to the restricted quantum double of the Yangian.

4.1 The Yangian double

The definition of the Yangian double $DY_{\hbar}\mathfrak{g}$ is obtained by allowing the second index of the generators in Definition 6.1 to take values in \mathbb{Z} , while working in the category of topological $\mathbb{C}[[\hbar]]$ -algebras:

Definition 4.1 The Yangian double $DY_{\hbar}\mathfrak{g}$ is the unital, associative $\mathbb{C}[[\hbar]]$ -algebra topologically generated by $\{x_{ir}^{\pm}, h_{ir}\}_{i \in \mathbf{I}, r \in \mathbb{Z}}$, subject to the relations (3.1)–(3.6) of Definition 6.1. In terms of generating series

$$\mathcal{X}_i^{\pm}(u) = \sum_{r \in \mathbb{Z}} x_{ir}^{\pm} u^{-r-1} \quad \text{and} \quad \mathcal{H}_i(u) = \sum_{r \in \mathbb{Z}} h_{ir} u^{-r-1},$$

these defining relations can be expressed as follows, for $i, j \in \mathbf{I}$:

$$\begin{aligned} [\mathcal{H}_i(u), \mathcal{H}_j(v)] &= 0, \\ [h_{i0}, \mathcal{X}_j^{\pm}(u)] &= \pm 2d_{ij} \mathcal{X}_j^{\pm}(u), \\ (u - v \mp \hbar d_{ij}) \mathcal{H}_i(u) \mathcal{X}_j^{\pm}(v) &= (u - v \pm \hbar d_{ij}) \mathcal{X}_j^{\pm}(v) \mathcal{H}_i(u), \\ (u - v \mp \hbar d_{ij}) \mathcal{X}_i^{\pm}(u) \mathcal{X}_j^{\pm}(v) &= (u - v \pm \hbar d_{ij}) \mathcal{X}_j^{\pm}(v) \mathcal{X}_i^{\pm}(u), \\ [\mathcal{X}_i^+(u), \mathcal{X}_j^-(v)] &= \delta_{ij} u^{-1} \delta(v/u) \mathcal{H}_i(v), \\ \sum_{\pi \in S_m} [\mathcal{X}_i^{\pm}(u_{\pi(1)}), [\mathcal{X}_i^{\pm}(u_{\pi(2)}), \dots, [\mathcal{X}_i^{\pm}(u_{\pi(m)}), \mathcal{X}_j^{\pm}(v)]] \dots]] &= 0, \end{aligned}$$

where $\delta(u) = \sum_{r \in \mathbb{Z}} u^r \in \mathbb{C}[[u^{\pm 1}]]$ is the formal delta function and in the last relation $i \neq j$ and $m = 1 - a_{ij}$.

$DY_{\hbar}\mathfrak{g}$ is \mathbb{Z} -graded as a topological $\mathbb{C}[[\hbar]]$ -algebra, with grading induced by the degree assignment $\deg x_{ir}^{\pm} = \deg h_{ir} = r$ for all $i \in \mathbf{I}$ and $r \in \mathbb{Z}$. That is, if $DY_{\hbar}\mathfrak{g}_k$ denotes the closure of the subspace of $DY_{\hbar}\mathfrak{g}$ spanned over the complex numbers by monomials in x_{ir}^{\pm}, h_{ir} and \hbar of total degree k , then

$$DY_{\hbar}\mathfrak{g} := \bigoplus_{k \in \mathbb{Z}} DY_{\hbar}\mathfrak{g}_k$$

is a dense, \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -subalgebra of $DY_{\hbar}\mathfrak{g}$ satisfying the conditions of Lemma 2.4. In particular, in the notation of Section 2.2, one has $DY_{\hbar}\mathfrak{g} = DY_{\hbar}\mathfrak{g}_{\mathbb{Z}}$.

Remark 4.2 Let $DY_{\hbar}\mathfrak{g}^j$ denote the $\mathbb{C}[[\hbar]]$ -algebra generated by $\{x_{ir}^{\pm}, h_{ir}\}_{i \in \mathbf{I}, r \in \mathbb{Z}}$, subject to the defining relations (3.1)–(3.6). Then $DY_{\hbar}\mathfrak{g}^j$ is a \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -algebra, and there is a natural algebra homomorphism

$$j : DY_{\hbar}\mathfrak{g}^j \rightarrow DY_{\hbar}\mathfrak{g} \subset DY_{\hbar}\mathfrak{g}.$$

The kernel of j is the graded ideal $\bigcap_{n \in \mathbb{N}} \hbar^n DY_{\hbar}\mathfrak{g}^j$, and $DY_{\hbar}\mathfrak{g}$ can be recovered as the \hbar -adic completion of $DY_{\hbar}\mathfrak{g}^j$ (see [41, Proposition 2.7]). Thus, $j(DY_{\hbar}\mathfrak{g}^j)$ is a dense,

\mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -subalgebra of $DY_{\hbar}\mathfrak{g}$. It is, however, a proper subalgebra of $\mathbb{D}Y_{\hbar}\mathfrak{g}$ as the graded components $j(\mathbb{D}Y_{\hbar}\mathfrak{g}_k^j)$ are not closed in $DY_{\hbar}\mathfrak{g}$. Rather, one has

$$DY_{\hbar}\mathfrak{g}_k = \varprojlim_n (\mathbb{D}Y_{\hbar}\mathfrak{g}_k^j / \hbar^n \mathbb{D}Y_{\hbar}\mathfrak{g}_{k-n}^j) \quad \forall k \in \mathbb{Z}.$$

The above definition implies that $DY_{\hbar}\mathfrak{g}$ is a \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -algebra deformation of the enveloping algebra $U(\mathfrak{t})$, where we recall that $\mathfrak{t} = \mathfrak{g}[[\hbar^{\pm 1}]]$. Analogously to the Yangian case recalled in Section 3.2, the identification $DY_{\hbar}\mathfrak{g} / \hbar DY_{\hbar}\mathfrak{g} \cong U(\mathfrak{t})$ is induced by the graded $\mathbb{C}[[\hbar]]$ -algebra epimorphism $DY_{\hbar}\mathfrak{g} \rightarrow U(\mathfrak{t})$ given by

$$x_{ir}^{\pm} \mapsto x_i^{\pm} t^r, \quad h_{ir} \mapsto h_i t^r \quad \forall i \in I \text{ and } r \in \mathbb{Z}.$$

The Poincaré–Birkhoff–Witt theorem for $DY_{\hbar}\mathfrak{g}$, established in Theorem 6.2 of [41], asserts that $DY_{\hbar}\mathfrak{g}$ is a topologically free $\mathbb{C}[[\hbar]]$ -module, and thus a flat deformation of $U(\mathfrak{t})$:

Theorem 4.3 *$DY_{\hbar}\mathfrak{g}$ is a flat deformation of the \mathbb{Z} -graded algebra $DY_{\hbar}\mathfrak{g} / \hbar DY_{\hbar}\mathfrak{g} \cong U(\mathfrak{t})$ over $\mathbb{C}[[\hbar]]$. In particular, $DY_{\hbar}\mathfrak{g} \cong U(\mathfrak{t})[[\hbar]]$ as a \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -module.*

The notation for the generators of $DY_{\hbar}\mathfrak{g}$ may seem, on the surface, to conflict with the notation used for generators in the Yangian associated with \mathfrak{g} . However, there is a natural \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$(4.1) \quad \iota : Y_{\hbar}\mathfrak{g} \rightarrow DY_{\hbar}\mathfrak{g}$$

sending each generator of $Y_{\hbar}(\mathfrak{g}) \subset Y_{\hbar}\mathfrak{g}$ to the corresponding element of $DY_{\hbar}\mathfrak{g}$, denoted with the same symbol. By Corollary 4.4 of [41], ι is injective, and we shall henceforth identify $Y_{\hbar}\mathfrak{g}$ with $\iota(Y_{\hbar}\mathfrak{g})$.

4.2 Automorphisms and root vectors

To each $i \in I$, we may associate series $\dot{x}_i^{\pm}(u)$ and $\dot{h}_i(u)$ in $DY_{\hbar}\mathfrak{g}[[u]]$ by setting

$$\dot{x}_i^{\pm}(u) := x_i^{\pm}(u) - \mathcal{X}_i^{\pm}(u) \quad \text{and} \quad \dot{h}_i(u) := h_i^{\pm}(u) - \mathcal{H}_i^{\pm}(u).$$

The following lemma is then a straightforward consequence of the defining relations of $DY_{\hbar}\mathfrak{g}$, where ω and ς are as in Lemma 3.3.

Lemma 4.4 *There are unique extensions of the anti-automorphisms ω and ς of $Y_{\hbar}(\mathfrak{g})$ to anti-automorphisms of the $\mathbb{C}[[\hbar]]$ -algebra $DY_{\hbar}\mathfrak{g}$ such that, for each $i \in I$,*

$$\begin{aligned} \omega(\dot{x}_i^{\pm}(u)) &= \dot{x}_i^{\mp}(u), & \omega(\dot{h}_i(u)) &= \dot{h}_i(u), \\ \varsigma(\dot{x}_i^{\pm}(u)) &= \dot{x}_i^{\pm}(-u), & \varsigma(\dot{h}_i(u)) &= \dot{h}_i(-u). \end{aligned}$$

Following the terminology from Section 3.3, we shall refer to the involution ω as the Chevalley involution of $DY_{\hbar}\mathfrak{g}$.

The adjoint action of $\mathfrak{h} \subset DY_{\hbar}\mathfrak{g}$ on $DY_{\hbar}\mathfrak{g}$ gives rise to a topological Q -grading on the $\mathbb{C}[[\hbar]]$ -algebra $DY_{\hbar}\mathfrak{g}$ (cf. Corollary 6.5 and [41, Section 3.1]) with graded components given by the weight spaces

$$DY_{\hbar}\mathfrak{g}_{\beta} := \{x \in DY_{\hbar}\mathfrak{g} : [h, x] = \beta(h)x \quad \forall h \in \mathfrak{h}\} \quad \forall \beta \in Q.$$

That is to say, each of these subspaces is a closed $\mathbb{C}[[\hbar]]$ -submodule of $DY_{\hbar}\mathfrak{g}$, and the direct sum

$$DY_{\hbar}\mathfrak{g}_Q := \bigoplus_{\beta \in Q} DY_{\hbar}\mathfrak{g}_{\beta}$$

is a Q -graded dense $\mathbb{C}[[\hbar]]$ -subalgebra of $DY_{\hbar}\mathfrak{g}$ whose subspace topology coincides with its \hbar -adic topology. Here, we have borrowed, and modified appropriately, the terminology of Section 2.2. It should be emphasized that the word *topological* is key in this statement, as the Q -graded algebra $DY_{\hbar}\mathfrak{g}_Q$ is a proper subset of $DY_{\hbar}\mathfrak{g}$.

We now introduce root vectors in $DY_{\hbar}\mathfrak{g}$ of arbitrary degree, following the procedure used in Section 3.4. Recall from (3.11) that to each positive root $\beta \in \Delta^+$, we attached an index $i(\beta) \in I$ and an element $X^{\beta} \in U(\mathfrak{n}_+)_{\beta - \alpha_{i(\beta)}}$. For each $k \in \mathbb{Z}$, we then set

$$x_{\beta,k}^+ := X^{\beta} \cdot x_{i(\beta),k}^+ \in DY_{\hbar}\mathfrak{g}_{\beta} \quad \text{and} \quad x_{\beta,k}^- := \omega(x_{\beta,k}^+) \in DY_{\hbar}\mathfrak{g}_{-\beta},$$

where \mathfrak{g} operates on $DY_{\hbar}\mathfrak{g}$ via the adjoint action. For $k \in \mathbb{N}$, these elements are identical to those introduced below (3.11). Moreover, we have

$$x_{\beta}^{\pm} t^r = x_{\beta,r}^{\pm} \pmod{\hbar} \quad \forall \beta \in \Delta^+, r \in \mathbb{Z}.$$

It shall be convenient for us to organize the above elements into generating series $x_{\beta}^{\pm}(u) \in DY_{\hbar}\mathfrak{g}_{\pm\beta}[[u^{-1}]]$ and $\dot{x}_{\beta}^{\pm}(u) \in DY_{\hbar}\mathfrak{g}_{\pm\beta}[[u]]$ by setting

$$x_{\beta}^{\pm}(u) := \sum_{r \in \mathbb{N}} x_{\beta,r}^{\pm} u^{-r-1} \quad \text{and} \quad \dot{x}_{\beta}^{\pm}(u) := - \sum_{r \in \mathbb{N}} x_{\beta,-r-1}^{\pm} u^r \quad \forall \beta \in \Delta^+.$$

4.3 The formal shift operator

We now shift our attention to recalling some of the main constructions of [41], subject to our standing assumption that \mathfrak{g} is a finite-dimensional simple Lie algebra. To begin, we introduce a number of relevant spaces built from the Yangian $Y_{\hbar}(\mathfrak{g})$, following Sections 4.1 and 4.2 of [41] and Section 3.6 above. First, let

$$\widehat{Y_{\hbar}\mathfrak{g}} = \prod_{k \in \mathbb{N}} Y_{\hbar}(\mathfrak{g})_k$$

denote the formal completion of $Y_{\hbar}(\mathfrak{g})$ with respect to its \mathbb{N} -grading. This is a topologically free $\mathbb{C}[[\hbar]]$ -algebra containing $Y_{\hbar}\mathfrak{g}$ as a subalgebra (see [15, Proposition 6.3] or [41, Lemma 4.1]).

Next, let $\mathbb{L}\widehat{Y_{\hbar}\mathfrak{g}}_z$ and $\widehat{Y_{\hbar}\mathfrak{g}}_z^{(1)}$ denote the subspaces $\mathbb{L}(\widehat{Y_{\hbar}\mathfrak{g}}_z^{(1)})$ and $\widehat{Y_{\hbar}\mathfrak{g}}_z^{(1)}$ of the space of Laurent series $Y_{\hbar}(\mathfrak{g})[[z; z^{-1}]]$ introduced in Section 3.6. That is,

$$\widehat{Y_{\hbar}\mathfrak{g}}_z = \prod_{k \in \mathbb{N}} Y_{\hbar}(\mathfrak{g})_k z^{-k} \subset Y_{\hbar}(\mathfrak{g})[[z^{-1}]],$$

and $\mathbb{L}\widehat{Y_{\hbar}\mathfrak{g}}_z$ is the \mathbb{Z} -graded subalgebra of $Y_{\hbar}(\mathfrak{g})[[z; z^{-1}]]$ over $\mathbb{C}[[\hbar]]$ defined by

$$\mathbb{L}\widehat{Y_{\hbar}\mathfrak{g}}_z = \bigoplus_{n \in \mathbb{Z}} z^n \widehat{Y_{\hbar}\mathfrak{g}}_z.$$

The following lemma, established in [41, Proposition 4.2], provides a characterization of the \hbar -adic completion of $\mathbb{L}\widehat{Y_{\hbar}\mathfrak{g}}_z$.

Lemma 4.5 *The \hbar -adic completion $\widehat{\mathbb{L}Y_{\hbar}\mathfrak{g}_z}$ of $\mathbb{L}Y_{\hbar}\mathfrak{g}_z$ is the subspace of $Y_{\hbar}\mathfrak{g}[[z^{\pm 1}]]$ consisting of formal series*

$$\sum_{k \in \mathbb{Z}} z^k f_k(z), \quad f_k(z) \in \widehat{Y_{\hbar}\mathfrak{g}_z}$$

with the property that, for each $n \in \mathbb{N}$, $\exists N_n \in \mathbb{N}$ such that

$$f_k(z) \in (\hbar/z)^n \widehat{Y_{\hbar}\mathfrak{g}_z} \quad \forall |k| \geq N_n.$$

Moreover, $\widehat{\mathbb{L}Y_{\hbar}\mathfrak{g}_z}$ is a topologically free \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -algebra with

$$(\widehat{\mathbb{L}Y_{\hbar}\mathfrak{g}_z})_{\mathbb{Z}} = \widehat{\mathbb{L}Y_{\hbar}\mathfrak{g}_z}.$$

The last statement of the lemma employs the notation from Lemma 2.4 and follows from the fact that $\widehat{\mathbb{L}Y_{\hbar}\mathfrak{g}_z}$ is a torsion-free \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -algebra and that each subspace $z^k \widehat{Y_{\hbar}\mathfrak{g}_z}$ is closed in $Y_{\hbar}\mathfrak{g}[[z^{\pm 1}]]$, equipped with the \hbar -adic topology, and thus in $\widehat{\mathbb{L}Y_{\hbar}\mathfrak{g}_z}$.

Next, recall that τ_z is the formal shift homomorphism of the Yangian introduced in (3.8), which we may view as a $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$\tau_z : Y_{\hbar}\mathfrak{g} \rightarrow Y_{\hbar}\mathfrak{g}[[z]].$$

In addition, we set $\partial_z^{(n)} = \frac{1}{n!} \partial_z^n$ for each $n \in \mathbb{N}$, where ∂_z is the partial derivative operator with respect to z . The following theorem, which is a combination of a special case of Theorems 4.3 and 6.2 of [41], introduces the so-called *formal shift operator* Φ_z on $DY_{\hbar}\mathfrak{g}$.

Theorem 4.6 *There is a unique homomorphism of $\mathbb{C}[[\hbar]]$ -algebras*

$$\Phi_z : DY_{\hbar}\mathfrak{g} \rightarrow \widehat{\mathbb{L}Y_{\hbar}\mathfrak{g}_z}$$

with the property that $\Phi_z \circ \iota = \tau_z$. Moreover:

(1) Φ_z is injective, and satisfies

$$\Phi_z(\dot{x}_{\beta}^{\pm}(u)) = \sum_{n \in \mathbb{N}} (-1)^n u^n \partial_z^{(n)} x_{\beta}^{\pm}(-z) \quad \forall \beta \in \Delta^+,$$

$$\Phi_z(\dot{h}_i(u)) = \sum_{n \in \mathbb{N}} (-1)^n u^n \partial_z^{(n)} h_i(-z) \quad \forall i \in I.$$

(2) The restriction of Φ_z to $DY_{\hbar}\mathfrak{g}$ is a \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$\Phi_z|_{DY_{\hbar}\mathfrak{g}} : DY_{\hbar}\mathfrak{g} \rightarrow \widehat{\mathbb{L}Y_{\hbar}\mathfrak{g}_z} = \bigoplus_{n \in \mathbb{Z}} z^n \widehat{Y_{\hbar}\mathfrak{g}_z} \subset Y_{\hbar}(\mathfrak{g})[z; z^{-1}].$$

Remark 4.7 In the terminology of Section 2.2, Part (2) is equivalent to the assertion that Φ_z is a \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -algebra homomorphism. This is implied by Part (2) of Theorem 4.3 in [41], which asserts that the composition $\Phi_z \circ \jmath$ is a \mathbb{Z} -graded algebra homomorphism, where \jmath is as in Remark 4.2.

By [41, Proposition 4.2(4)], the evaluation map

$$(4.2) \quad \mathcal{E}_v : \widehat{\mathbb{L}Y_{\hbar}\mathfrak{g}_z} \rightarrow \widehat{Y_{\hbar}\mathfrak{g}}, \quad f(z) \mapsto f(1),$$

is an epimorphism of $\mathbb{C}[[\hbar]]$ -algebras. We may thus compose Φ_z with $\mathcal{E}\mathcal{U}$ to obtain a $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$\Phi := \mathcal{E}\mathcal{U} \circ \Phi_z : DY_{\hbar}\mathfrak{g} \rightarrow \widehat{Y_{\hbar}\mathfrak{g}}.$$

By Theorem 6.2 of [41], this homomorphism is injective. One of the main results of [41] is that Φ induces an isomorphism between completions of $DY_{\hbar}\mathfrak{g}$ of $Y_{\hbar}(\mathfrak{g})$. To make this precise, let $\mathcal{J} \subset DY_{\hbar}\mathfrak{g}$ denote the kernel of the composition

$$DY_{\hbar}\mathfrak{g} \xrightarrow{\hbar \rightarrow 0} U(\mathfrak{g}[t^{\pm 1}]) \xrightarrow{t \rightarrow 1} U(\mathfrak{g}),$$

and define $\widehat{DY_{\hbar}\mathfrak{g}}$ to be the completion of $DY_{\hbar}\mathfrak{g}$ with respect to the \mathcal{J} -adic filtration

$$DY_{\hbar}\mathfrak{g} = \mathcal{J}^0 \supset \mathcal{J} \supset \mathcal{J}^2 \supset \dots \supset \mathcal{J}^n \supset \dots.$$

We then have the following analogue of [15, Theorem 6.2], which is a special case of Theorem 5.5 in [41].

Theorem 4.8 Φ is injective and induces an isomorphism of $\mathbb{C}[[\hbar]]$ -algebras

$$\widehat{\Phi} : \widehat{DY_{\hbar}\mathfrak{g}} \xrightarrow{\sim} \widehat{Y_{\hbar}\mathfrak{g}}$$

with inverse Γ uniquely extending the embedding $\iota \circ \tau_{-1} : Y_{\hbar}(\mathfrak{g}) \rightarrow DY_{\hbar}\mathfrak{g}$.

Remark 4.9 One subtle consequence of this result is that the natural homomorphism

$$DY_{\hbar}\mathfrak{g} \rightarrow \widehat{DY_{\hbar}\mathfrak{g}}$$

is injective. Indeed, its composition with the isomorphism $\widehat{\Phi}$ recovers the injection Φ . Henceforth, we shall freely make use of this fact and view $DY_{\hbar}\mathfrak{g}$ as a subalgebra of $\widehat{DY_{\hbar}\mathfrak{g}}$. We further note that the subspace topology on $DY_{\hbar}\mathfrak{g}$, with respect to the \hbar -adic topology on $\widehat{DY_{\hbar}\mathfrak{g}}$, coincides with the \hbar -adic topology on $DY_{\hbar}\mathfrak{g}$. Indeed, as $\widehat{DY_{\hbar}\mathfrak{g}} \cong \widehat{Y_{\hbar}\mathfrak{g}}$ is torsion-free, to see this it suffices to show that

$$\hbar \widehat{DY_{\hbar}\mathfrak{g}} \cap DY_{\hbar}\mathfrak{g} = \hbar DY_{\hbar}\mathfrak{g}.$$

This, however, follows immediately from the injectivity of the semiclassical limit of Φ , established in [41, Theorem 6.2]. In particular, this discussion implies that $DY_{\hbar}\mathfrak{g}$ is a closed subspace of the topological $\mathbb{C}[[\hbar]]$ -module $\widehat{DY_{\hbar}\mathfrak{g}}$. Similarly, one deduces that $\Phi_z(DY_{\hbar}\mathfrak{g})$ is a closed subspace of $\widehat{LY_{\hbar}\mathfrak{g}_z}$.

To conclude this preliminary section on $DY_{\hbar}\mathfrak{g}$, we introduce the auxiliary $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$(4.3) \quad \Gamma_z := \Gamma \circ \mathcal{E}\mathcal{U} : \widehat{LY_{\hbar}\mathfrak{g}_z} \rightarrow \widehat{DY_{\hbar}\mathfrak{g}}$$

which has the property that $\Gamma_z|_{\text{Im}(\Phi_z)} = \Phi_z^{-1}$. This homomorphism shall play a prominent role in the main result of Section 7 and its proof (see Theorem 7.5).

5 The Drinfeld–Gavarini Yangian

In this section and Section 6, we give a self-contained exposition to the dual Yangian $\check{Y}_{\hbar}\mathfrak{g}^*$, which provides a homogeneous quantization of the graded dual $\mathfrak{t}_- = t^{-1}\mathfrak{g}[t^{-1}]$

to the Lie bialgebra $\mathfrak{t}_+ = \mathfrak{g}[t]$, as will be explained in detail in Section 6. The definition of $Y_{\hbar}\mathfrak{g}^*$ takes as input the so-called Drinfeld–Gavarini subalgebra of the Yangian. The goal of the present section is to introduce this subalgebra and survey some of its key properties.

5.1 Quantum duality

To provide context, let us first briefly recall the general construction of the dual of a quantized enveloping algebra, following [7, Section 7] and [19, Section 4.4] (see also [11] and [1, Section 2.19], for example).

Suppose that $U_{\hbar}\mathfrak{b}$ is a quantization of a finite-dimensional Lie bialgebra $(\mathfrak{b}, \delta_{\mathfrak{b}})$, where we follow the terminology and notation from Section 2.4. One would then like to introduce a notion of duality which sends $U_{\hbar}\mathfrak{b}$ to a quantization of the Lie bialgebra dual $(\mathfrak{b}^*, \delta_{\mathfrak{b}^*} = [\cdot, \cdot]_{\mathfrak{b}}^*)$ to $(\mathfrak{b}, \delta_{\mathfrak{b}})$. The first crucial observation is that $\mathbb{C}[[\hbar]]$ -linear dual $U_{\hbar}\mathfrak{b}^* = \text{Hom}_{\mathbb{C}[[\hbar]]}(U_{\hbar}\mathfrak{b}, \mathbb{C}[[\hbar]])$ of $U_{\hbar}\mathfrak{b}$ is not itself a quantized enveloping algebra (see Lemma 2.1 of [19], in addition to [7, Section 7] and [1, Section 2.19]). The correct notion of duality within the category of quantized enveloping algebras was introduced in [7, Section 7]. One considers the $\mathbb{C}[[\hbar]]$ -submodule

$$U_{\hbar}\mathfrak{b}' := \{x \in U_{\hbar}\mathfrak{b} : (1 - \varepsilon)^{\otimes n} \Delta^n(x) \in \hbar^n U_{\hbar}\mathfrak{b}^{\otimes n} \ \forall n \in \mathbb{N}\} \subset U_{\hbar}\mathfrak{b},$$

where ε and Δ are the counit and coproduct, respectively, on the topological Hopf algebra $U_{\hbar}\mathfrak{b}$, and all notation is as in Section 5.2 below. Then, by [19, Proposition 3.6], $U_{\hbar}\mathfrak{b}'$ is a quantized formal series Hopf algebra, with semiclassical limit isomorphic as an algebra to the completion of the symmetric algebra $S(\mathfrak{b}) = \bigoplus_{n \in \mathbb{N}} S^n(\mathfrak{b})$ with respect to its standard grading. In particular, this means that although $U_{\hbar}\mathfrak{b}'$ is not in general a topological Hopf algebra over $\mathbb{C}[[\hbar]]$ in the sense of Section 2.1, it is a topological Hopf algebra with respect to the $J_{\mathfrak{b}}$ -adic topology, where

$$J_{\mathfrak{b}} = \hbar U_{\hbar}\mathfrak{b} \cap U_{\hbar}\mathfrak{b}' = \varepsilon|_{U_{\hbar}\mathfrak{b}'}^{-1}(\hbar \mathbb{C}[[\hbar]]).$$

The subspace $U_{\hbar}\mathfrak{b}^{\circ} \subset (U_{\hbar}\mathfrak{b}')^*$ consisting of continuous linear forms with respect to this topology is then a quantization of $(\mathfrak{b}^*, \delta_{\mathfrak{b}^*})$. This is the *quantized enveloping algebra dual* of $U_{\hbar}\mathfrak{b}$.

Remark 5.1 Here, we note that $U_{\hbar}\mathfrak{b}^{\circ}$ can be equivalently defined as the \hbar -adic completion of the $\mathbb{C}[[\hbar]]$ -module

$$(U_{\hbar}\mathfrak{b}^*)^{\times} = \sum_{n \in \mathbb{N}} \hbar^{-n} \mathfrak{m}_{\mathfrak{b}}^n \subset \mathbb{C}((\hbar)) \otimes_{\mathbb{C}[[\hbar]]} U_{\hbar}\mathfrak{b}^*,$$

where $\mathfrak{m}_{\mathfrak{b}} := \{f \in U_{\hbar}\mathfrak{b}^* : f(1) \in \hbar \mathbb{C}[[\hbar]]\}$. That this produces a topological Hopf algebra which can be identified with $U_{\hbar}\mathfrak{b}^{\circ}$ is a nontrivial result, which is part of the *quantum duality principle*. This was first announced in [7, Section 7], and proven in detail in [19] (see Theorem 1.6 therein). We will not, however, need this equivalent formulation in the present article.

In our setting, $\mathfrak{b} = \mathfrak{t}_+ = \mathfrak{g}[t]$ is *not* finite-dimensional, but rather an \mathbb{N} -graded Lie bialgebra $\mathfrak{b} = \bigoplus_{n \in \mathbb{N}} \mathfrak{b}_n$ with finite-dimensional graded components. As $Y_{\hbar}\mathfrak{g} = U_{\hbar}\mathfrak{b}$ is a homogeneous quantization of \mathfrak{t}_+ , the above construction remains valid, provided the notion of duality is adjusted so as to respect the underlying gradings. In fact, one can

replace $U_{\hbar} \mathfrak{b}'$ with an \mathbb{N} -graded topological Hopf algebra $\check{Y}_{\hbar} \mathfrak{g}$ over $\mathbb{C}[[\hbar]]$ of finite type, and $U_{\hbar} \mathfrak{b}^{\circ}$ with the restricted dual of $\check{Y}_{\hbar} \mathfrak{g}$, as defined in Section 2.3. The topological Hopf algebra $\check{Y}_{\hbar} \mathfrak{g}$ and its $\mathbb{C}[[\hbar]]$ -form $\check{Y}_{\hbar}(\mathfrak{g}) = (\check{Y}_{\hbar} \mathfrak{g})_{\mathbb{N}}$ (see Section 2.2) are the focus of the present section.

5.2 The Drinfeld–Gavarini subalgebra

Let us define Δ^n for any $n \in \mathbb{N}$ by setting $\Delta^0 = \varepsilon$, $\Delta^1 = 1 = 1_{Y_{\hbar}(\mathfrak{g})}$ and

$$\Delta^n := (\Delta \otimes \mathbf{1}^{\otimes(n-2)}) \circ \Delta^{n-1} : Y_{\hbar}(\mathfrak{g}) \rightarrow Y_{\hbar}(\mathfrak{g})^{\otimes n}$$

for all $n \geq 2$. We then define the $\mathbb{C}[[\hbar]]$ -submodule $\check{Y}_{\hbar}(\mathfrak{g}) \subset Y_{\hbar}(\mathfrak{g})$ by

$$\check{Y}_{\hbar}(\mathfrak{g}) := \{x \in Y_{\hbar}(\mathfrak{g}) : (1 - \varepsilon)^{\otimes n} \Delta^n(x) \in \hbar^n Y_{\hbar}(\mathfrak{g})^{\otimes n} \ \forall n \in \mathbb{N}\}.$$

By Lemma 3.2 and Proposition 3.5 of [27] (see also [19, Proposition 2.6], [20, Theorem 3.5], and [13, Lemma A.1]), $\check{Y}_{\hbar}(\mathfrak{g})$ is a subalgebra of $Y_{\hbar}(\mathfrak{g})$ which is commutative modulo \hbar :

$$(5.1) \quad [x, y] \in \hbar \check{Y}_{\hbar}(\mathfrak{g}) \quad \forall x, y \in \check{Y}_{\hbar}(\mathfrak{g}).$$

As the structure maps $\mathbf{1}$, ε , and Δ^n are graded, $\check{Y}_{\hbar}(\mathfrak{g})$ inherits from $Y_{\hbar}(\mathfrak{g})$ the structure of an \mathbb{N} -graded algebra. We shall call $\check{Y}_{\hbar}(\mathfrak{g})$ the *Drinfeld–Gavarini subalgebra* of the Yangian $Y_{\hbar}(\mathfrak{g})$. Its algebraic structure has been described in detail by Tsybaliuk and Weekes in Appendix Appendix A of [13], following the general results obtained in the works [19, 20] of Gavarini. In this subsection, we review, and partially extend, this description.

Let $R_{\hbar}(U(\mathfrak{t}_+))$ denote the Rees algebra associated with the standard enveloping algebra filtration \mathbf{F}_{\bullet} on $U(\mathfrak{t}_+)$:

$$R_{\hbar}(U(\mathfrak{t}_+)) = \bigoplus_{n \in \mathbb{N}} \hbar^n \mathbf{F}_n(U(\mathfrak{t}_+)) \subset U(\mathfrak{t}_+)[[\hbar]].$$

Consider now the symmetric algebra $S(\hbar \mathfrak{t}_+) \subset S(\mathfrak{t}_+)[[\hbar]]$ on $\hbar \mathfrak{t}_+$. Here, \hbar can be viewed as a gradation parameter associated with the standard \mathbb{N} -grading on the symmetric algebra $S(\mathfrak{t}_+)$. Namely, $S(\hbar \mathfrak{t}_+) \cap \hbar^n S(\mathfrak{t}_+)$ is precisely the n th symmetric power $S^n(\hbar \mathfrak{t}_+) = \hbar^n S^n(\mathfrak{t}_+)$, and $S(\hbar \mathfrak{t}_+) \cong S(\mathfrak{t}_+)$ as an \mathbb{N} -graded algebra. As $U(\mathfrak{t}_+)$ is a filtered deformation of $S(\hbar \mathfrak{t}_+)$ (that is, one has $\text{gr } U(\mathfrak{t}_+) \cong S(\hbar \mathfrak{t}_+)$), $R_{\hbar}(U(\mathfrak{t}_+))$ is a flat deformation of $S(\hbar \mathfrak{t}_+)$ over $\mathbb{C}[[\hbar]]$. Let

$$q : R_{\hbar}(U(\mathfrak{t}_+)) \twoheadrightarrow S(\hbar \mathfrak{t}_+)$$

be the natural quotient map, under the identification of $R_{\hbar}(U(\mathfrak{t}_+))/\hbar R_{\hbar}(U(\mathfrak{t}_+))$ with $\text{gr } U(\mathfrak{t}_+) \cong S(\hbar \mathfrak{t}_+)$. In what follows, we shall be primarily interested in the *loop gradings* on $R_{\hbar}(U(\mathfrak{t}_+))$ and $S(\hbar \mathfrak{t}_+)$, inherited from the natural grading on $U(\mathfrak{t}_+)[[\hbar]]$ compatible with the \mathbb{N} -grading on \mathfrak{t}_+ . Namely, one has

$$\text{deg}(\hbar \mathfrak{t}_{+,k}) = \text{deg}(\hbar \mathfrak{g} t^k) = k + 1 \quad \forall k \in \mathbb{N}.$$

We shall denote the n th graded component of $S(\hbar \mathfrak{t}_+)$ by $S_n(\hbar \mathfrak{t}_+)$ so that

$$S(\hbar \mathfrak{t}_+) = \bigoplus_{n \in \mathbb{N}} S_n(\hbar \mathfrak{t}_+).$$

Recall from Section 3.4 that v_G denotes the graded $\mathbb{C}[\hbar]$ -module isomorphism

$$v_G : Y_{\hbar}(\mathfrak{g}) \xrightarrow{\sim} U(\mathfrak{t}_+)[\hbar]$$

defined in (3.12), which depends on a fixed total order on the set G . We equip $\hbar G$ with the induced ordering, for which multiplication by \hbar defines an isomorphism of ordered sets $G \xrightarrow{\sim} \hbar G$, and we let $B(\hbar G) \subset Y_{\hbar}(\mathfrak{g})$ denote the corresponding set of ordered monomials in $\hbar G$. We further recall that $\mu : U(\mathfrak{t}_+)[\hbar] \rightarrow Y_{\hbar}(\mathfrak{g})$ is the inverse of the specific choice $v = v_G$ defined in (3.13).

The following proposition is a consequence of Proposition 3.3 of [20] in addition to Proposition A.2 and Theorem A.7 of [13] (see also [19, Section 3.5]).

Proposition 5.2 *Let \dot{v}_G denote the restriction of v_G to $\dot{Y}_{\hbar}(\mathfrak{g})$. Then:*

- (1) $\dot{Y}_{\hbar}(\mathfrak{g})$ is an \mathbb{N} -graded Hopf subalgebra of $Y_{\hbar}(\mathfrak{g})$.
- (2) \dot{v}_G is an isomorphism of \mathbb{N} -graded $\mathbb{C}[\hbar]$ -modules

$$\dot{v}_G : \dot{Y}_{\hbar}(\mathfrak{g}) \xrightarrow{\sim} R_{\hbar}(U(\mathfrak{t}_+)).$$

- (3) $\dot{Y}_{\hbar}(\mathfrak{g})$ is generated as a $\mathbb{C}[\hbar]$ -algebra by $\hbar\mu(\mathfrak{t}_+)$ and has basis $B(\hbar G)$.
- (4) The composition $\dot{q} := q \circ \dot{v}_G$ is an epimorphism of \mathbb{N} -graded algebras which descends to an isomorphism

$$\dot{Y}_{\hbar}(\mathfrak{g})/\hbar\dot{Y}_{\hbar}(\mathfrak{g}) \xrightarrow{\sim} S(\hbar\mathfrak{t}_+).$$

Proof of (2) and (3) These statements are a minor modification of the statement of Theorem A.7 of [13]. For the sake of completeness, let us recall the main ingredients, beginning with the proof that $\hbar G$ (and thus $\hbar\mu(\mathfrak{t}_+)$) is contained in $\dot{Y}_{\hbar}(\mathfrak{g})$, given in Lemmas A.5 and A.6 of [13].

For each $n \in \mathbb{N}$, $x \in \mathfrak{g}$, and $i \in I$, the formulas of Proposition 3.2 imply that

$$\Delta^n(x) = \sum_{a=1}^n x^{(a)} \quad \text{and} \quad \Delta^n(t_{i1}) = \sum_{a=1}^n t_{i1}^{(a)} + \hbar \sum_{a < b} r_i^{ab},$$

where $y^{(b)} = 1^{\otimes(b-1)} \otimes y \otimes 1^{\otimes(n-b)} \in Y_{\hbar}(\mathfrak{g})^{\otimes n}$ for any $y \in Y_{\hbar}(\mathfrak{g})$ and, for each $a < b$, $B \mapsto B^{ab}$ is the algebra homomorphism $Y_{\hbar}(\mathfrak{g})^{\otimes 2} \rightarrow Y_{\hbar}(\mathfrak{g})^{\otimes n}$ given on simple tensors by $(x \otimes y)^{ab} = x^{(a)}y^{(b)}$. Since $(1 - \varepsilon)$ projects $Y_{\hbar}(\mathfrak{g})$ onto $\text{Ker}(\varepsilon)$, it follows readily from these formulas that

$$\hbar\mathfrak{g} \cup \{\hbar t_{i1}\}_{i \in I} \subset \dot{Y}_{\hbar}(\mathfrak{g}).$$

We may now deduce that $\hbar G \subset \dot{Y}_{\hbar}(\mathfrak{g})$ as follows. By (5.1) and the above, $\dot{Y}_{\hbar}(\mathfrak{g})$ is a \mathfrak{g} -submodule of $Y_{\hbar}(\mathfrak{g})$ which is preserved by the operators $\{\text{ad}(t_{i1})\}_{i \in I}$. Hence, the elements

$$\hbar x_{ik}^{\pm} = (\pm 1)^k \text{ad}(T_i)^k(\hbar x_i^{\pm}) \quad \text{and} \quad \hbar h_{ik} = [x_i^+, \hbar x_{ik}^-]$$

necessarily belong to $\dot{Y}_{\hbar}(\mathfrak{g})$ for each $i \in I$ and $k \in \mathbb{N}$, where $T_i = (2d_i)^{-1}t_{i1}$. As $\hbar x_{\alpha,k}^{\pm}$ belongs to the \mathfrak{g} -submodule of $Y_{\hbar}(\mathfrak{g})$ generated by $\hbar x_{ik}^{\pm}$, we can conclude that $\hbar G \subset \dot{Y}_{\hbar}(\mathfrak{g})$.

Since $R_{\hbar}(U(\mathfrak{t}_+))$ has basis given by the set of ordered monomials in $\hbar q(G)$, to complete the proof of both Parts (2) and (3), it suffices to see that $B(\hbar G)$ spans $\dot{Y}_{\hbar}(\mathfrak{g})$.

This follows from the fact that $B(\mathfrak{G})$ is a basis of $Y_{\hbar}(\mathfrak{g})$ together with the crucial Lemma 3.3 of [19] (see also [11, Lemma 4.12]). We refer the reader to the proof of [13, Proposition A.2] for complete details. ■

Proof of (1) We have already seen that $\dot{Y}_{\hbar}(\mathfrak{g})$ is an \mathbb{N} -graded subalgebra of $Y_{\hbar}(\mathfrak{g})$. That it is a Hopf subalgebra of $Y_{\hbar}(\mathfrak{g})$ is a special case of Proposition 3.3 of [20], which passes to completions and makes use of a modification of a technical result for quantized formal series Hopf algebras established in [9, Proposition 2.1]. It is worth pointing out that, in our specialized setting, it is possible to give a concise direct proof that $\dot{Y}_{\hbar}(\mathfrak{g})$ is a subcoalgebra of $Y_{\hbar}(\mathfrak{g})$ stable under the antipode S .

Indeed, by Proposition 3.2, one has $\Delta(\hbar x) \in \dot{Y}_{\hbar}(\mathfrak{g})^{\otimes 2}$ for $x \in \hbar\mathfrak{g} \cup \{t_{i1}\}_{i \in I}$. As $\dot{Y}_{\hbar}(\mathfrak{g})^{\otimes 2}$ is a \mathfrak{g} -submodule of $Y_{\hbar}(\mathfrak{g})^{\otimes 2}$ stable under all operators $\text{ad}(t_{i1} \otimes 1 + 1 \otimes t_{i1})$, the inclusion $\Delta(\hbar\mathfrak{G}) \subset \dot{Y}_{\hbar}(\mathfrak{g})^{\otimes 2}$ will hold provided $\dot{Y}_{\hbar}(\mathfrak{g})^{\otimes 2}$ is preserved by the operators $\text{ad}(\hbar r_i)$. This is itself a consequence of the fact that $r_i \in \mathfrak{g} \otimes \mathfrak{g}$, the inclusion $\hbar\mathfrak{g} \subset \dot{Y}_{\hbar}(\mathfrak{g})$, that $\dot{Y}_{\hbar}(\mathfrak{g})^{\otimes 2}$ is a \mathfrak{g} -submodule of $Y_{\hbar}(\mathfrak{g})^{\otimes 2}$, and the relation

$$[\hbar r_i, x \otimes y] = [\hbar r_i, x^{(1)}]y^{(2)} + x^{(1)}[\hbar r_i, y^{(2)}] \quad \forall x, y \in \dot{Y}_{\hbar}(\mathfrak{g}).$$

Hence, by Part (3), we can conclude that $\Delta(\dot{Y}_{\hbar}(\mathfrak{g})) \subset \dot{Y}_{\hbar}(\mathfrak{g})^{\otimes 2}$. A similar argument, using the formulas of Proposition 3.2 and that $\dot{Y}_{\hbar}(\mathfrak{g})$ is a \mathfrak{g} -submodule of $\dot{Y}_{\hbar}(\mathfrak{g})$ stable under the operators $\text{ad}(S(t_{i1}))$, implies that $S(\dot{Y}_{\hbar}(\mathfrak{g})) \subset \dot{Y}_{\hbar}(\mathfrak{g})$. ■

Proof of (4) By Part (2), \mathfrak{q} is an \mathbb{N} -graded $\mathbb{C}[[\hbar]]$ -linear epimorphism with kernel $\hbar\dot{Y}_{\hbar}(\mathfrak{g})$, and thus gives rise to an isomorphism of graded vector spaces $\mathfrak{q} : \dot{Y}_{\hbar}(\mathfrak{g})/\hbar\dot{Y}_{\hbar}(\mathfrak{g}) \xrightarrow{\sim} S(\hbar\mathfrak{t}_+)$. To conclude, it suffices to prove that \mathfrak{q} is an algebra homomorphism. By (5.1), $\dot{Y}_{\hbar}(\mathfrak{g})/\hbar\dot{Y}_{\hbar}(\mathfrak{g})$ is commutative, and so the linear map $\hbar\mathfrak{t}_+ \rightarrow \dot{Y}_{\hbar}(\mathfrak{g})/\hbar\dot{Y}_{\hbar}(\mathfrak{g})$ sending any $\hbar x \in \hbar\mathfrak{t}_+$ to the image of $\hbar\mu(x)$ in $\dot{Y}_{\hbar}(\mathfrak{g})/\hbar\dot{Y}_{\hbar}(\mathfrak{g})$ uniquely extends to an algebra homomorphism $\mathfrak{p} : S(\hbar\mathfrak{t}_+) \rightarrow \dot{Y}_{\hbar}(\mathfrak{g})/\hbar\dot{Y}_{\hbar}(\mathfrak{g})$. Since $\mathfrak{q} \circ \mathfrak{p} = 1$, we can conclude that \mathfrak{q} is the inverse of \mathfrak{p} , and thus an algebra homomorphism. ■

Recall from (3.14) that $Y_{\hbar}\mathfrak{g}$ denotes the \hbar -adically complete Yangian associated with \mathfrak{g} , and let $\dot{Y}_{\hbar}\mathfrak{g}$ denote the \hbar -adic completion of the Hopf algebra $\dot{Y}_{\hbar}(\mathfrak{g})$:

$$\dot{Y}_{\hbar}\mathfrak{g} := \varprojlim_n (\dot{Y}_{\hbar}(\mathfrak{g})/\hbar^n \dot{Y}_{\hbar}(\mathfrak{g})).$$

As an immediate consequence of the above proposition and Corollary 2.6, we obtain the following result.

Corollary 5.3 $\dot{Y}_{\hbar}\mathfrak{g}$ is a topologically free, \mathbb{N} -graded topological Hopf algebra over $\mathbb{C}[[\hbar]]$ of finite type. Moreover:

- (1) $\dot{Y}_{\hbar}\mathfrak{g}$ is a flat deformation of the \mathbb{N} -graded algebra $S(\hbar\mathfrak{t}_+)$ over $\mathbb{C}[[\hbar]]$. In particular, $\dot{Y}_{\hbar}\mathfrak{g} \cong S(\hbar\mathfrak{t}_+)[[\hbar]]$ as an \mathbb{N} -graded topological $\mathbb{C}[[\hbar]]$ -module.
- (2) $\dot{Y}_{\hbar}\mathfrak{g}$ is a topological Hopf subalgebra of the completed Yangian $Y_{\hbar}\mathfrak{g}$.

The statement that $\dot{Y}_{\hbar}\mathfrak{g}$ is of finite type reduces to the fact that the homogeneous components $S_n(\hbar\mathfrak{t}_+)$ of $S(\hbar\mathfrak{t}_+)$ are all finite-dimensional complex vector spaces (see below Corollary 2.9).

Henceforth, we will identify $S(\hbar\mathfrak{t}_+)$ with the semiclassical limit of $\dot{Y}_{\hbar}(\mathfrak{g})$ as an \mathbb{N} -graded Hopf algebra. We emphasize that this is a non-cocommutative Hopf algebra;

in particular, it is not isomorphic to the standard symmetric Hopf algebra on $\hbar\mathfrak{t}_+$, which we denote by $\mathbb{S}(\hbar\mathfrak{t}_+)$. However, $\dot{Y}_\hbar(\mathfrak{g})$ and $S(\hbar\mathfrak{t}_+)$ are filtered deformations of $R_\hbar(U(\mathfrak{t}_+))$ and $\mathbb{S}(\hbar\mathfrak{t}_+)$, respectively, as we now explain. Consider the Hopf ideal

$$(5.2) \quad \mathbf{J} := \hbar Y_\hbar(\mathfrak{g}) \cap \dot{Y}_\hbar(\mathfrak{g}) \subset \dot{Y}_\hbar(\mathfrak{g})$$

Here, we note that it follows from Part (3) of the above proposition that, for each pair of positive integers $k, n \in \mathbb{N}_+$, one has

$$(5.3) \quad \hbar^k Y_\hbar(\mathfrak{g})^{\otimes n} \cap \dot{Y}_\hbar(\mathfrak{g})^{\otimes n} = \sum_{k_1+\dots+k_n=k} \mathbf{J}^{k_1} \otimes \dots \otimes \mathbf{J}^{k_n}.$$

In particular, $\mathbf{J}^k = \hbar^k Y_\hbar(\mathfrak{g}) \cap \dot{Y}_\hbar(\mathfrak{g})$ for each $k \in \mathbb{N}$, and so the associated graded Hopf algebra $\text{gr}_\mathbf{J}(\dot{Y}_\hbar(\mathfrak{g}))$ with respect to the \mathbf{J} -adic filtration embeds inside the associated graded Hopf algebra $\text{gr}_\hbar(Y_\hbar(\mathfrak{g}))$ of $Y_\hbar(\mathfrak{g})$ with respect to the \hbar -adic filtration. Since $Y_\hbar(\mathfrak{g})$ is a torsion-free Hopf algebra deformation of $U(\mathfrak{t}_+)$ over $\mathbb{C}[[\hbar]]$, we have isomorphisms of graded Hopf algebras

$$\begin{aligned} \text{gr}_\hbar(Y_\hbar(\mathfrak{g})) &= \bigoplus_{n \in \mathbb{N}} \hbar^n Y_\hbar(\mathfrak{g}) / \hbar^{n+1} Y_\hbar(\mathfrak{g}) \cong \bigoplus_{n \in \mathbb{N}} \hbar^n U(\mathfrak{t}_+) = U(\mathfrak{t}_+)[[\hbar]], \\ \text{gr}_\mathbf{J}(\dot{Y}_\hbar(\mathfrak{g})) &= \bigoplus_{n \in \mathbb{N}} \mathbf{J}^n / \mathbf{J}^{n+1} \cong \bigoplus_{n \in \mathbb{N}} \hbar^n \mathbf{F}_n(U(\mathfrak{t}_+)) = R_\hbar(U(\mathfrak{t}_+)), \end{aligned}$$

where the second isomorphism follows from the first and Proposition 5.2. In fact, the module isomorphisms v_G and \dot{v}_G are filtered, and the above identifications can be realized as the associated graded maps $\text{gr}(v_G)$ and $\text{gr}(\dot{v}_G)$, respectively.

The image of \mathbf{J} under the quotient map \dot{q} is the Hopf ideal

$$(5.4) \quad \mathbb{J} := \bigoplus_{n>0} S^n(\hbar\mathfrak{t}_+) \subset S(\hbar\mathfrak{t}_+)$$

where $S^n(\hbar\mathfrak{t}_+)$ denotes the n th symmetric power of $\hbar\mathfrak{t}_+$. This is also a Hopf ideal in $\mathbb{S}(\hbar\mathfrak{t}_+)$, and one has a canonical isomorphism $\text{gr}_\mathbb{J}\mathbb{S}(\hbar\mathfrak{t}_+) \cong \mathbb{S}(\hbar\mathfrak{t}_+)$ of graded Hopf algebras. Since the elements of $\mu(\mathfrak{t}_+)$ are primitive modulo $\hbar Y_\hbar(\mathfrak{g})^{\otimes 2}$, the subspace $\hbar\mathfrak{t}_+ = \mathbb{J}/\mathbb{J}^2$ of $\text{gr}_\mathbb{J}\mathbb{S}(\hbar\mathfrak{t}_+)$ consists of primitive elements, and we can conclude that

$$\text{gr}_\mathbb{J}\mathbb{S}(\hbar\mathfrak{t}_+) \cong \mathbb{S}(\hbar\mathfrak{t}_+) \cong \text{gr}_\mathbb{J}\mathbb{S}(\hbar\mathfrak{t}_+)$$

as graded Hopf algebras.

Remark 5.4 Since $\mathbf{J}^k = \hbar^k Y_\hbar(\mathfrak{g}) \cap \dot{Y}_\hbar(\mathfrak{g})$ for each $k \in \mathbb{N}$, the closure of $\dot{Y}_\hbar(\mathfrak{g})$ in the quantized enveloping algebra $Y_\hbar\mathfrak{g}$ coincides with the \mathbf{J} -adic completion

$$Y_\hbar\mathfrak{g}' := \varprojlim_n (\dot{Y}_\hbar(\mathfrak{g})/\mathbf{J}^n) \subset Y_\hbar\mathfrak{g}.$$

This is precisely the quantized formal series Hopf algebra $U_\hbar\mathfrak{b}'$ from Section 5.1 associated with $U_\hbar\mathfrak{b} = Y_\hbar\mathfrak{g}$. As \mathbf{J} surjects onto the ideal \mathbb{J} from (5.4), the quotient map \dot{q} induces an isomorphism of $\mathbb{C}[[\hbar]]$ -algebras

$$Y_\hbar\mathfrak{g}' / \hbar Y_\hbar\mathfrak{g}' \xrightarrow{\sim} \widehat{S(\hbar\mathfrak{t}_+)} := \prod_{n \in \mathbb{N}} S^n(\hbar\mathfrak{t}_+) \cong \varprojlim_n (S(\hbar\mathfrak{t}_+) / \mathbb{J}^n).$$

5.3 Triangular decomposition

Proposition 5.2 implies that the triangular decomposition of $Y_{\hbar}(\mathfrak{g})$, reviewed in Section 3.4, induces a triangular decomposition on $\dot{Y}_{\hbar}(\mathfrak{g})$, with the \mathbb{N} -graded $\mathbb{C}[\hbar]$ -algebras

$$\dot{Y}_{\hbar}^{\pm}(\mathfrak{g}) := Y_{\hbar}^{\pm}(\mathfrak{g}) \cap \dot{Y}_{\hbar}(\mathfrak{g}) \quad \text{and} \quad \dot{Y}_{\hbar}^0(\mathfrak{g}) := Y_{\hbar}^0(\mathfrak{g}) \cap \dot{Y}_{\hbar}(\mathfrak{g}),$$

playing the roles of $Y_{\hbar}^{\pm}(\mathfrak{g})$ and $Y_{\hbar}^0(\mathfrak{g})$, respectively. In this subsection, we spell this out explicitly. Let us set

$$\mathfrak{t}_{\pm}^{\pm} := \mathfrak{n}_{\pm}[t] \subset \mathfrak{t}_{\pm} \quad \text{and} \quad \mathfrak{t}_{+}^0 := \mathfrak{h}[t] \subset \mathfrak{t}_{+}$$

and recall from Section 3.4 that ν_{\pm} and ξ are the isomorphisms

$$\nu_{\pm} : Y_{\hbar}^{\pm}(\mathfrak{g}) \xrightarrow{\sim} U(\mathfrak{t}_{\pm}^{\pm})[\hbar] \quad \text{and} \quad \xi : Y_{\hbar}^0(\mathfrak{g}) \xrightarrow{\sim} S(\mathfrak{t}_{+}^0)[\hbar]$$

defined above (3.13) and in the statement of Proposition 3.5, respectively. We further recall from (3.17) that, for each $\beta \in Q_{+}$, $\nu(\beta) \in \mathbb{N}$ is defined by

$$\nu(\beta) = \min\{k \in \mathbb{N} : \exists \beta_1, \dots, \beta_k \in \Delta^{+} \text{ with } \beta = \beta_1 + \dots + \beta_k\}.$$

The following corollary provides an analogue of Proposition 5.2 for the subalgebras $\dot{Y}_{\hbar}^{\pm}(\mathfrak{g})$ and $\dot{Y}_{\hbar}^0(\mathfrak{g})$.

Corollary 5.5 *Let $\dot{\nu}_{\pm}$ and $\dot{\xi}$ denote the restrictions of ν_{\pm} and ξ to $\dot{Y}_{\hbar}^{\pm}(\mathfrak{g})$ and $\dot{Y}_{\hbar}^0(\mathfrak{g})$, respectively. Then:*

- (1) $\dot{\nu}_{\pm}$ is an isomorphism of \mathbb{N} -graded $\mathbb{C}[\hbar]$ -modules

$$\dot{\nu}_{\pm} : \dot{Y}_{\hbar}^{\pm}(\mathfrak{g}) \xrightarrow{\sim} R_{\hbar}(U(\mathfrak{t}_{\pm}^{\pm})) \subset U(\mathfrak{t}_{\pm}^{\pm})[\hbar].$$

- (2) $\dot{\xi}$ is an isomorphism of \mathbb{N} -graded $\mathbb{C}[\hbar]$ -algebras

$$\dot{\xi} : \dot{Y}_{\hbar}^0(\mathfrak{g}) \xrightarrow{\sim} S(\hbar\mathfrak{t}_{+}^0)[\hbar] \subset S(\mathfrak{t}_{+}^0)[\hbar].$$

- (3) $\dot{Y}_{\hbar}^{\pm}(\mathfrak{g})$ is a torsion-free, \mathbb{N} -graded $\mathbb{C}[\hbar]$ -algebra deformation of $S(\hbar\mathfrak{t}_{\pm}^{\pm})$. In particular, there is an isomorphism of graded $\mathbb{C}[\hbar]$ -modules

$$\dot{Y}_{\hbar}^{\pm}(\mathfrak{g}) \cong S(\hbar\mathfrak{t}_{\pm}^{\pm})[\hbar].$$

- (4) $\dot{Y}_{\hbar}^{\pm}(\mathfrak{g})$ is a Q -graded subalgebra of $\dot{Y}_{\hbar}^{\pm}(\mathfrak{g})$ with

$$\dot{Y}_{\hbar}^{\pm}(\mathfrak{g})_{\pm\beta} \subset J^{\nu(\beta)} \subset \hbar^{\nu(\beta)} Y_{\hbar}^{\pm}(\mathfrak{g})_{\pm\beta} \quad \forall \beta \in Q_{+}.$$

We note that each of these results follows readily from Propositions 3.5 and 5.2, in addition to Corollary 2.6 in the case of Part (3). We leave the details as an exercise to the interested reader. As a consequence of this corollary and Proposition 5.2, we obtain the following analogue of Part (3) from Proposition 3.5.

Corollary 5.6 *The multiplication map*

$$m : \dot{Y}_{\hbar}^{+}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}^0(\mathfrak{g}) \otimes \dot{Y}_{\hbar}^{-}(\mathfrak{g}) \rightarrow \dot{Y}_{\hbar}(\mathfrak{g})$$

is an isomorphism of \mathbb{N} -graded $\mathbb{C}[\hbar]$ -modules.

Remark 5.7 As in Remark 3.7, the above results can easily be lifted to the \hbar -adic setting using Corollary 2.6 (see also Corollary 5.3). We especially note that, for each choice of the symbol χ , the \hbar -adic completion

$$\dot{Y}_{\hbar}^{\chi} \mathfrak{g} := \varprojlim_n \dot{Y}_{\hbar}^{\chi}(\mathfrak{g}) / \hbar^n \dot{Y}_{\hbar}^{\chi}(\mathfrak{g})$$

is a topologically free \mathbb{N} -graded $\mathbb{C}[[\hbar]]$ -algebra of finite type, which provides a flat deformation of the symmetric algebra $S(\hbar t_+^{\chi})$ over $\mathbb{C}[[\hbar]]$. In addition, $\dot{Y}_{\hbar}^{\chi} \mathfrak{g}$ embeds inside the completed Yangian $Y_{\hbar} \mathfrak{g}$, and the multiplication map induces an isomorphism of \mathbb{N} -graded $\mathbb{C}[[\hbar]]$ -modules

$$m : \dot{Y}_{\hbar}^+ \mathfrak{g} \otimes \dot{Y}_{\hbar}^0 \mathfrak{g} \otimes \dot{Y}_{\hbar}^- \mathfrak{g} \xrightarrow{\sim} \dot{Y}_{\hbar} \mathfrak{g},$$

where \otimes is now the topological tensor product $\widehat{\otimes}$ over $\mathbb{C}[[\hbar]]$ (see Remark 2.11).

5.4 The adjoint and coadjoint actions

We now prove two lemmas concerning the Drinfeld–Gavarini subalgebra $\dot{Y}_{\hbar}(\mathfrak{g})$ which will play an important role in the main results and constructions of Sections 7 and 8. The first of these, Lemma 5.8, will be used to construct the quantum double of the Yangian in Section 8.2.

Let $\blacktriangledown : Y_{\hbar}(\mathfrak{g}) \otimes Y_{\hbar}(\mathfrak{g}) \rightarrow Y_{\hbar}(\mathfrak{g})$ and $\blacktriangle : Y_{\hbar}(\mathfrak{g}) \rightarrow Y_{\hbar}(\mathfrak{g}) \otimes Y_{\hbar}(\mathfrak{g})$ denote the left adjoint action of $Y_{\hbar}(\mathfrak{g})$ on itself, and the right adjoint coaction of $Y_{\hbar}(\mathfrak{g})$ on itself, respectively. That is,

$$\begin{aligned} \blacktriangledown &= m^3 \circ (\mathbf{1}^{\otimes 2} \otimes S) \circ (23) \circ (\Delta \otimes \mathbf{1}), \\ \blacktriangle &= (\mathbf{1} \otimes m) \circ (12) \circ (S \otimes \mathbf{1}^{\otimes 2}) \circ \Delta^3. \end{aligned}$$

The \hbar -adic analogue of the below result, for a quantized enveloping algebra $U_{\hbar} \mathfrak{b}$ with $\dot{Y}_{\hbar}(\mathfrak{g})$ replaced by $U_{\hbar} \mathfrak{b}'$, was established in Propositions 4.3 and 4.4 of [1] (see also Proposition A.5 therein).

Lemma 5.8 *One has*

$$\blacktriangledown(Y_{\hbar}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}(\mathfrak{g})) \subset \dot{Y}_{\hbar}(\mathfrak{g}) \quad \text{and} \quad \blacktriangle(Y_{\hbar}(\mathfrak{g})) \subset Y_{\hbar}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}(\mathfrak{g}).$$

Proof Since \blacktriangledown makes $Y_{\hbar}(\mathfrak{g})$ a left module, to prove the first inclusion, it suffices to show that $x \cdot y := \blacktriangledown(x \otimes y) \in \dot{Y}_{\hbar}(\mathfrak{g})$ for all $x \in \mathfrak{g} \cup \{t_{i1}\}_{i \in \mathbf{I}}$ and $y \in \dot{Y}_{\hbar}(\mathfrak{g})$. Since

$$x \cdot y = [x, y] \quad \text{and} \quad t_{i1} \cdot y = [t_{i1}, y] + m([y^{(1)}, \hbar r_i]) \quad \forall x \in \mathfrak{g}, i \in \mathbf{I}, y \in \dot{Y}_{\hbar}(\mathfrak{g}),$$

this follows from the observation that $\hbar r_i \in \mathfrak{g} \otimes \hbar \mathfrak{g} \subset \mathfrak{g} \otimes \dot{Y}_{\hbar}(\mathfrak{g})$ and the fact that $\dot{Y}_{\hbar}(\mathfrak{g})$ is a \mathfrak{g} -submodule of $Y_{\hbar}(\mathfrak{g})$ stable under $\text{ad}(t_{i1})$ for all $i \in \mathbf{I}$.

As for the second inclusion, note that \blacktriangle is a homomorphism of right modules:

$$(5.5) \quad \blacktriangle \circ m = \gamma \circ (\blacktriangle \otimes \mathbf{1}),$$

where $\gamma : Y_{\hbar}(\mathfrak{g})^{\otimes 2} \otimes Y_{\hbar}(\mathfrak{g}) \rightarrow Y_{\hbar}(\mathfrak{g})^{\otimes 2}$ is the right action defined by

$$\gamma = m \otimes m^2 \circ (\mathbf{1}^{\otimes 2} \otimes S \otimes \mathbf{1}^{\otimes 2}) \circ (24) \circ (\mathbf{1}^{\otimes 2} \otimes \Delta^3).$$

This right action preserves $Y_{\hbar}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}(\mathfrak{g}) \subset Y_{\hbar}(\mathfrak{g})^{\otimes 2}$. Indeed, if $y \in Y_{\hbar}(\mathfrak{g}), z \in \dot{Y}_{\hbar}(\mathfrak{g})$, and $x \in \mathfrak{g}$, we have

$$\gamma(y \otimes z \otimes x) = y \otimes [z, x] + yx \otimes z \in Y_{\hbar}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}(\mathfrak{g}),$$

and replacing x by t_{i1} , for any $i \in I$, yields instead

$$\gamma(y \otimes z \otimes t_{i1}) = y \otimes ([z, t_{i1}] + m([z^{(2)}, \hbar r_i])) + yt_{i1} \otimes z + y^{(1)}(z^{(2)} \hbar r_i - \hbar r_i^{21} z^{(2)})$$

which again belongs to $Y_{\hbar}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}(\mathfrak{g})$ as $\hbar r_i^{21}, \hbar r_i \in (\hbar \mathfrak{g} \otimes \mathfrak{g}) \cap (\mathfrak{g} \otimes \hbar \mathfrak{g})$ and $\dot{Y}_{\hbar}(\mathfrak{g})$ is an $\text{ad}(t_{i1})$ -stable \mathfrak{g} -submodule of $Y_{\hbar}(\mathfrak{g})$.

Since $Y_{\hbar}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}(\mathfrak{g})$ is a right submodule of $Y_{\hbar}(\mathfrak{g})^{\otimes 2}$ under γ , the condition (5.5) guarantees that $\blacktriangle(Y_{\hbar}(\mathfrak{g})) \subset Y_{\hbar}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}(\mathfrak{g})$ will hold provided it holds on $\mathfrak{g} \cup \{t_{i1}\}_{i \in I}$. To conclude, it suffices to note that, for each $x \in \mathfrak{g}$ and $i \in I$, one has

$$\begin{aligned} \blacktriangle(x) &= x \otimes 1 \in Y_{\hbar}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}(\mathfrak{g}), \\ \blacktriangle(t_{i1}) &= t_{i1} \otimes 1 + \hbar(r_i - r_i^{21}) \in Y_{\hbar}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}(\mathfrak{g}). \end{aligned} \quad \blacksquare$$

5.5 The R-matrix

The second lemma we will need concerns the universal R-matrix $\mathcal{R}(z)$ of the Yangian, and will play a crucial role in identifying the dual Yangian as a subalgebra of the Yangian double $DY_{\hbar} \mathfrak{g}$ in Section 7. In what follows, all notation is as in Section 3.6.

Lemma 5.9 *The factors $\mathcal{R}^{\pm}(z)$ and $\mathcal{R}^0(z)$ of the universal R-matrix $\mathcal{R}(z)$ have coefficients in $(\dot{Y}_{\hbar}(\mathfrak{g}) \otimes Y_{\hbar}(\mathfrak{g})) \cap (Y_{\hbar}(\mathfrak{g}) \cap \dot{Y}_{\hbar}(\mathfrak{g}))$. Consequently,*

$$\mathcal{R}(z) \in (\dot{Y}_{\hbar}(\mathfrak{g}) \otimes Y_{\hbar}(\mathfrak{g}))[[z^{-1}]] \cap (Y_{\hbar}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}(\mathfrak{g}))[[z^{-1}]].$$

Proof Since $\dot{Y}_{\hbar}(\mathfrak{g})$ is preserved by $\text{ad}(t_{i1})$ for any $i \in I$ and, for each $\alpha \in \Delta^+$, the simple tensor $\hbar x_{\alpha}^{-} \otimes x_{\alpha}^{+} = x_{\alpha}^{-} \otimes \hbar x_{\alpha}^{+}$ belongs to the intersection of $\dot{Y}_{\hbar}(\mathfrak{g}) \otimes Y_{\hbar}(\mathfrak{g})$ and $Y_{\hbar}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}(\mathfrak{g})$, it follows from (3.18) and induction on $\text{ht}(\beta)$ that

$$\mathcal{R}_{\beta}^{-}(z) \in (\dot{Y}_{\hbar}(\mathfrak{g}) \otimes Y_{\hbar}(\mathfrak{g}))[[z^{-1}]] \cap (Y_{\hbar}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}(\mathfrak{g}))[[z^{-1}]] \quad \forall \beta \in Q_+.$$

Consequently, both $\mathcal{R}^{-}(z)$ and $\mathcal{R}^{+}(z) = \mathcal{R}_{21}^{-}(-z)^{-1}$ belong to the intersection of $(\dot{Y}_{\hbar}(\mathfrak{g}) \otimes Y_{\hbar}(\mathfrak{g}))[[z^{-1}]]$ and $(Y_{\hbar}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}(\mathfrak{g}))[[z^{-1}]]$. It is thus enough to prove that

$$\mathcal{R}^0(z) \in (\dot{Y}_{\hbar}(\mathfrak{g}) \otimes Y_{\hbar}(\mathfrak{g}))[[z^{-1}]] \cap (Y_{\hbar}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}(\mathfrak{g}))[[z^{-1}]].$$

Recall from (5.2) that $J = \dot{Y}_{\hbar}(\mathfrak{g}) \cap \hbar Y_{\hbar}(\mathfrak{g})$. Since $\hbar h_i(u) \in J[[u^{-1}]]$, the logarithm $t_i(u) = \log(1 + \hbar h_i(u))$ and its Borel transform $B_i(u)$ both have coefficients in J . Therefore, we have

$$\hbar^{-1} B_i(u) \otimes B_j(-u) \in (\dot{Y}_{\hbar}(\mathfrak{g}) \otimes Y_{\hbar}(\mathfrak{g}))[[u]] \cap (Y_{\hbar}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}(\mathfrak{g}))[[u]] \quad \forall i, j \in I.$$

Since $g(z/2\kappa\hbar)$ is divisible by \hbar , it follows from (3.19) that the logarithm $\mathcal{S}(z)$ of $\mathcal{R}^0(z)$, and thus $\mathcal{R}^0(z)$ itself, has coefficients belonging to the intersection of $\dot{Y}_{\hbar}(\mathfrak{g}) \otimes Y_{\hbar}(\mathfrak{g})$ and $Y_{\hbar}(\mathfrak{g}) \otimes \dot{Y}_{\hbar}(\mathfrak{g})$. ■

6 The dual Yangian

With Section 5 at our disposal, we are now in a position to introduce the dual Yangian $\dot{Y}_{\hbar}\mathfrak{g}^*$. After defining $\dot{Y}_{\hbar}\mathfrak{g}^*$ and spelling out some of its basic properties in Sections 6.1 and 6.2, we prove in Sections 6.3 and 6.4 that it is a homogeneous quantization of the Lie bialgebra $(\mathfrak{t}_-, \delta_-)$ defined in Section 2.5. We conclude in Section 6.5 by identifying a family of generators for $\dot{Y}_{\hbar}\mathfrak{g}^*$ and establishing a triangular decomposition.

6.1 The dual Yangian $\dot{Y}_{\hbar}\mathfrak{g}^*$

By Corollary 5.3, the \hbar -adic completion $\dot{Y}_{\hbar}\mathfrak{g}$ of $\dot{Y}_{\hbar}(\mathfrak{g})$ is an \mathbb{N} -graded topological Hopf algebra of finite type. We may thus apply the machinery from Section 2.3 to obtain a \mathbb{Z} -graded topological Hopf structure on its restricted dual.

In more detail, by Proposition 2.10, the restricted dual $\dot{Y}_{\hbar}\mathfrak{g}^*$, as defined in Definition 2.7, is a \mathbb{Z} -graded topological Hopf algebra over $\mathbb{C}[[\hbar]]$ with product, unit, coproduct, counit, and antipode given by the transposes

$$\begin{aligned} \Delta^t : \dot{Y}_{\hbar}\mathfrak{g}^* \otimes \dot{Y}_{\hbar}\mathfrak{g}^* &\rightarrow \dot{Y}_{\hbar}\mathfrak{g}^*, & \varepsilon^t : \mathbb{C}[[\hbar]] &\rightarrow \dot{Y}_{\hbar}\mathfrak{g}^*, \\ m^t : \dot{Y}_{\hbar}\mathfrak{g}^* &\rightarrow \dot{Y}_{\hbar}\mathfrak{g}^* \otimes \dot{Y}_{\hbar}\mathfrak{g}^*, & \iota^t : \dot{Y}_{\hbar}\mathfrak{g}^* &\rightarrow \mathbb{C}[[\hbar]], & S^t : \dot{Y}_{\hbar}\mathfrak{g}^* &\rightarrow \dot{Y}_{\hbar}\mathfrak{g}^*, \end{aligned}$$

respectively, where Δ , ε , m , ι , and S are the coproduct, counit, product, unit, and antipode of $\dot{Y}_{\hbar}\mathfrak{g}$, respectively, and \otimes is the topological tensor product over $\mathbb{C}[[\hbar]]$ (see Remark 2.11).

Definition 6.1 The topological Hopf algebra $\dot{Y}_{\hbar}\mathfrak{g}^*$ introduced above is called *the dual Yangian* of \mathfrak{g} .

Explicitly, $\dot{Y}_{\hbar}\mathfrak{g}^*$ is the subspace of the $\mathbb{C}[[\hbar]]$ -linear dual $\dot{Y}_{\hbar}\mathfrak{g}^*$ consisting of those $f : \dot{Y}_{\hbar}\mathfrak{g} \rightarrow \mathbb{C}[[\hbar]]$ which are continuous with respect to the gradation topology on $\dot{Y}_{\hbar}\mathfrak{g}$. It can be recovered as the \hbar -adic completion of the \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -algebra

$$(\dot{Y}_{\hbar}\mathfrak{g}^*)_{\mathbb{Z}} := \bigoplus_{a \in \mathbb{Z}} \dot{Y}_{\hbar}\mathfrak{g}_a^* \cong \dot{Y}_{\hbar}(\mathfrak{g})^*,$$

where $\dot{Y}_{\hbar}(\mathfrak{g})^*$ is the graded dual of $\dot{Y}_{\hbar}(\mathfrak{g})$ over $\mathbb{C}[[\hbar]]$ which, as explained in Remark 2.8, coincides with $(\dot{Y}_{\hbar}\mathfrak{g}^*)_{\mathbb{Z}}$ under the natural identification of $\dot{Y}_{\hbar}\mathfrak{g}^*$ with $\text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}(\mathfrak{g}), \mathbb{C}[[\hbar]])$. Here, we have set

$$\dot{Y}_{\hbar}\mathfrak{g}_a^* := \text{Hom}_{\mathbb{C}[[\hbar]]}^a(\dot{Y}_{\hbar}\mathfrak{g}, \mathbb{C}[[\hbar]]) \cong \text{Hom}_{\mathbb{C}[[\hbar]]}^a(\dot{Y}_{\hbar}(\mathfrak{g}), \mathbb{C}[[\hbar]]) \quad \forall a \in \mathbb{Z}.$$

By Corollary 2.9, $\dot{Y}_{\hbar}\mathfrak{g}^*$ is a flat deformation of the graded Hopf algebra

$$\dot{Y}_{\hbar}\mathfrak{g}^* / \hbar \dot{Y}_{\hbar}\mathfrak{g}^* \cong S(\hbar\mathfrak{t}_+)^*.$$

In particular, the dual Yangian $\dot{Y}_{\hbar}\mathfrak{g}^*$ is isomorphic to $S(\hbar\mathfrak{t}_+)^*[[\hbar]]$ as a \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -module. Here, we recall that $S(\hbar\mathfrak{t}_+)^*$ is the graded dual of the \mathbb{N} -graded Hopf algebra $S(\hbar\mathfrak{t}_+)$ over \mathbb{C} . As a vector space, one has

$$S(\hbar\mathfrak{t}_+)^* = \bigoplus_{n \in \mathbb{N}} S(\hbar\mathfrak{t}_+)^*_{-n},$$

where $S(\hbar t_+)^*_{-n} = S_n(\hbar t_+)^* \subset \text{Hom}_{\mathbb{C}}(S(\hbar t_+), \mathbb{C})$. We shall identify this Hopf algebra with the enveloping algebra of the Lie algebra $\mathfrak{t}_- = t^{-1}\mathfrak{g}[t^{-1}]$ in Section 6.3 below.

Now let us make a few comments which concern the restricted dual $Y_{\hbar\mathfrak{g}}^*$ of the full, completed, Yangian $Y_{\hbar\mathfrak{g}}$. Since $Y_{\hbar\mathfrak{g}}$ is an \mathbb{N} -graded topological Hopf algebra over $\mathbb{C}[[\hbar]]$, the formalism of Section 2.3 implies that $\dot{Y}_{\hbar\mathfrak{g}}^*$ is a \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -algebra, which provides a flat deformation of the algebra $U(\mathfrak{t}_+)^*$ over $\mathbb{C}[[\hbar]]$. In particular, there is an isomorphism of \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -modules

$$Y_{\hbar\mathfrak{g}}^* \cong U(\mathfrak{t}_+)^*[[\hbar]].$$

However, $Y_{\hbar\mathfrak{g}}^*$ is not itself a topological Hopf algebra over $\mathbb{C}[[\hbar]]$ with respect to the \hbar -adic topology. It is, however, naturally a subalgebra of the topological Hopf algebra $\dot{Y}_{\hbar\mathfrak{g}}^*$. This is made explicit by the below result.

Proposition 6.2 *The $\mathbb{C}[[\hbar]]$ -linear map*

$$Y_{\hbar\mathfrak{g}}^* \rightarrow \dot{Y}_{\hbar\mathfrak{g}}^*, \quad f \mapsto f|_{\dot{Y}_{\hbar\mathfrak{g}}} \quad \forall f \in Y_{\hbar\mathfrak{g}}^*,$$

is an injective homomorphism of \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -algebras.

Proof Since $\dot{Y}_{\hbar\mathfrak{g}}$ is a \mathbb{Z} -graded Hopf subalgebra of $Y_{\hbar\mathfrak{g}}$, the map $f \mapsto f|_{\dot{Y}_{\hbar\mathfrak{g}}}$ respects the underlying \mathbb{Z} -graded algebra structures. Moreover, if $f \in Y_{\hbar\mathfrak{g}}^*$ vanishes on the basis $B(\hbar\mathfrak{G})$ from Proposition 5.2, then it vanishes on the basis $B(\mathfrak{G})$ of $\dot{Y}_{\hbar}(\mathfrak{g})$ as $\mathbb{C}[[\hbar]]$ is torsion-free. This yields the injectivity. ■

Remark 6.3 Let \mathfrak{J} be as in (5.2). Then, since \mathfrak{J} satisfies

$$\mathfrak{J}^n \subset \bigoplus_{k \geq n} \dot{Y}_{\hbar}(\mathfrak{g})_k \quad \forall n \in \mathbb{N},$$

every $f \in \dot{Y}_{\hbar\mathfrak{g}}^*$ is automatically continuous with respect to the \mathfrak{J} -adic topology on $\dot{Y}_{\hbar}(\mathfrak{g})$, and so uniquely extends to an element of the associated topological dual $Y_{\hbar\mathfrak{g}}^{\circ}$ to $Y_{\hbar\mathfrak{g}}$ (see Section 5.1 and Remark 5.4). In this sense, the notion of duality considered here is compatible with that for a general quantized enveloping algebra $U_{\hbar}\mathfrak{b}$ outlined in Section 5.1, despite the fact that we did not need to leave the category of topological Hopf algebras over $\mathbb{C}[[\hbar]]$ to define $\dot{Y}_{\hbar\mathfrak{g}}^*$.

Remark 6.4 An alternative description of $\dot{Y}_{\hbar\mathfrak{g}}^*$ using the formalism of Remark 5.1 can be found in [12, Section 3.1].

6.2 Chevalley involution and \mathfrak{g} -action

We now make a handful of simple observations which will play an important role in the remainder of this article. In what follows, we shall freely make use of the fact that $\dot{Y}_{\hbar\mathfrak{g}}^*$ can be naturally viewed as a subspace of $\text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}(\mathfrak{g}), \mathbb{C}[[\hbar]])$.

Since the Chevalley involution ω defined in Lemma 3.3 is an anti-automorphism of the graded Hopf algebra $Y_{\hbar}(\mathfrak{g})$, it follows from the definition of $\dot{Y}_{\hbar}(\mathfrak{g})$ (or, alternatively, from Proposition 5.2) that it restricts to an anti-automorphism of the graded Hopf algebra $\dot{Y}_{\hbar}(\mathfrak{g})$, which we again denote by ω . Consequently, the transpose ω^t of ω , uniquely determined by

$$\omega^t(f)(x) = f(\omega(x)) \quad \forall f \in \dot{Y}_{\hbar\mathfrak{g}}^* \text{ and } x \in \dot{Y}_{\hbar}(\mathfrak{g}),$$

is an involutive Hopf algebra anti-automorphism of $\dot{Y}_\hbar \mathfrak{g}^*$. We call ω^t the *Chevalley involution* of $\dot{Y}_\hbar \mathfrak{g}^*$.

Next, recall that the adjoint action of \mathfrak{g} on $Y_\hbar(\mathfrak{g})$ preserves the Drinfeld–Gavarini subalgebra $\dot{Y}_\hbar(\mathfrak{g})$. Since each graded component $\dot{Y}_\hbar(\mathfrak{g})_k$ is also a submodule, the restricted dual $\dot{Y}_\hbar \mathfrak{g}^*$ is a \mathfrak{g} -module equipped with the coadjoint action

$$(x \cdot f)(y) = f(S(x) \cdot y) \quad \forall x \in U(\mathfrak{g}), y \in \dot{Y}_\hbar(\mathfrak{g}) \text{ and } f \in \dot{Y}_\hbar \mathfrak{g}^*.$$

We now introduce a topological Q -grading on $\dot{Y}_\hbar \mathfrak{g}^*$ compatible with the above action which is analogous to that obtained for $DY_\hbar \mathfrak{g}$ in Section 4.2. For each $\beta \in Q$, define the closed $\mathbb{C}[[\hbar]]$ -submodule $\dot{Y}_\hbar \mathfrak{g}_\beta^* \subset \dot{Y}_\hbar \mathfrak{g}^*$ by

$$\dot{Y}_\hbar \mathfrak{g}_\beta^* = \{f \in \dot{Y}_\hbar \mathfrak{g}^* : f(\dot{Y}_\hbar(\mathfrak{g})_\alpha) \subset \mathbb{C}[[\hbar]]_{\alpha+\beta} \quad \forall \alpha \in Q\},$$

where $\mathbb{C}[[\hbar]]_\alpha$ is $\mathbb{C}[[\hbar]]$ if $\alpha = 0$ and is $\{0\}$ otherwise. It is easy to see that $\dot{Y}_\hbar \mathfrak{g}_\beta^*$ is just the β -weight space of $\dot{Y}_\hbar \mathfrak{g}^*$ with respect to the \mathfrak{g} -module structure introduced above. That is, one has

$$\dot{Y}_\hbar \mathfrak{g}_\beta^* = \{f \in \dot{Y}_\hbar \mathfrak{g}^* : h \cdot f = \beta(h)f \quad \forall h \in \mathfrak{h}\}.$$

As $\dot{Y}_\hbar(\mathfrak{g})$ is Q -graded as a Hopf algebra, the direct sum

$$\dot{Y}_\hbar \mathfrak{g}_Q^* = \bigoplus_{\beta \in Q} \dot{Y}_\hbar \mathfrak{g}_\beta^* \subset \dot{Y}_\hbar \mathfrak{g}^*$$

is a Q -graded $\mathbb{C}[[\hbar]]$ -subalgebra of $\dot{Y}_\hbar \mathfrak{g}^*$. Moreover, the counit, coproduct, and antipode of $\dot{Y}_\hbar \mathfrak{g}^*$ are all Q -graded, degree zero, maps. It is not difficult to prove that $\dot{Y}_\hbar \mathfrak{g}_Q^*$ is a dense subalgebra of $\dot{Y}_\hbar \mathfrak{g}^*$ whose subspace topology coincides with its \hbar -adic topology. Hence, we obtain the following result:

Corollary 6.5 $\dot{Y}_\hbar \mathfrak{g}^*$ is Q -graded as a topological Hopf algebra over $\mathbb{C}[[\hbar]]$.

6.3 Classical duality

We now wish to identify the graded dual $S(\hbar \mathfrak{t}_+)^*$ of the \mathbb{N} -graded Hopf algebra $S(\hbar \mathfrak{t}_+)$ with the enveloping algebra $U(\mathfrak{t}_-)$, where we recall that $\mathfrak{t}_- = t^{-1} \mathfrak{g}[t^{-1}]$.

To formulate this result optimally, we must first give a few preliminary remarks. To begin, we note that the semiclassical limit of the Chevalley involution ω^t of $\dot{Y}_\hbar \mathfrak{g}^*$ coincides, by definition, with the transpose $\hat{\omega}^t$ of the automorphism $\hat{\omega}$ of $S(\hbar \mathfrak{t}_+)$ given on $\hbar \mathfrak{t}_+$ by

$$\hat{\omega}(\hbar x) = \hbar \bar{\omega}(x) \quad \forall x \in \mathfrak{t}_+,$$

where $\bar{\omega}$ is as in (3.9). Similarly, the coadjoint action of \mathfrak{g} on $\dot{Y}_\hbar \mathfrak{g}^*$ introduced in Section 6.2 specializes to an action of \mathfrak{g} on $S(\hbar \mathfrak{t}_+)^*$. By definition, this action is dual to that of \mathfrak{g} on $S(\hbar \mathfrak{t}_+)$ inherited from the adjoint action of \mathfrak{g} on $\dot{Y}_\hbar(\mathfrak{g})$.

On the other hand, the Chevalley involution $\bar{\omega}$ of $U(\mathfrak{g}[t^{\pm 1}])$, defined as in (3.9) with r taking values in \mathbb{Z} , and the adjoint action of \mathfrak{g} on $U(\mathfrak{g}[t^{\pm 1}])$ both preserve $U(\mathfrak{t}_-)$. The resulting involution and \mathfrak{g} -module structure on $U(\mathfrak{t}_-)$ will be compared to those of $S(\hbar \mathfrak{t}_+)^*$ described in the previous paragraph in (3) of Proposition 6.6.

Consider now the standard symmetric algebra grading

$$S(\hbar\mathfrak{t}_+) = \bigoplus_{n \in \mathbb{N}} S^n(\hbar\mathfrak{t}_+),$$

where $S^n(\hbar\mathfrak{t}_+)$ is the n th symmetric power of $\hbar\mathfrak{t}_+$, as in (5.4). Since $S^1(\hbar\mathfrak{t}_+) = \hbar\mathfrak{t}_+$, every linear functional f in $(\hbar\mathfrak{t}_+)^*$ trivially extends to an element of $S(\hbar\mathfrak{t}_+)^*$ satisfying $f(S^n(\hbar\mathfrak{t}_+)) = 0$ for all $n \neq 1$, which is contained in $S(\hbar\mathfrak{t}_+)^*$ provided $f \in (\hbar\mathfrak{t}_+)^*$. That is, we have $(\hbar\mathfrak{t}_+)^* \subset S(\hbar\mathfrak{t}_+)^*$. In addition, we have a homogeneous, degree zero, isomorphism of graded vector spaces

$$\text{Res}_-^{\hbar} : \mathfrak{t}_- \xrightarrow{\sim} (\hbar\mathfrak{t}_+)^*, \quad \text{Res}_-^{\hbar}(x)(\hbar y) = \text{Res}_-(x)(y) = \langle x, y \rangle \quad \forall x \in \mathfrak{t}_-, y \in \mathfrak{t}_+,$$

where Res_- and \langle, \rangle are as defined in Section 2.5. With the above at our disposal, we are now prepared to identify $S(\hbar\mathfrak{t}_+)^*$ and $U(\mathfrak{t}_-)$.

Proposition 6.6 *The restricted dual $S(\hbar\mathfrak{t}_+)^*$ has the following properties:*

(1) $(\hbar\mathfrak{t}_+)^*$ is the Lie algebra of primitive elements in $S(\hbar\mathfrak{t}_+)^*$, with bracket

$$[f, g] = (f \otimes g) \circ \hbar\delta_+ \quad \forall f, g \in (\hbar\mathfrak{t}_+)^*.$$

(2) Res_-^{\hbar} uniquely extends to an isomorphism of graded Hopf algebras

$$\varphi : U(\mathfrak{t}_-) \xrightarrow{\sim} S(\hbar\mathfrak{t}_+)^*.$$

(3) φ is a \mathfrak{g} -module intertwiner commuting with Chevalley involutions.

Parts (1) and (2) of this proposition can be viewed as a variant of [25, Corollary 3.4] applied to a restricted version of the setting in [25, Section 3E]. They can be seen as a consequence of a graded generalization of (a special case of) Theorem 4.8 in [20]. It is not difficult to prove this variant directly using a fairly general argument, as we illustrate below.

Proof of (1) An element $f \in S(\hbar\mathfrak{t}_+)^*$ is primitive precisely when it satisfies

$$(6.1) \quad f(xy) = f(x)\varepsilon(y) + \varepsilon(x)f(y) \quad \forall x, y \in S(\hbar\mathfrak{t}_+).$$

Since the counit ε of $S(\hbar\mathfrak{t}_+)$ vanishes on $\mathbb{J} = \bigoplus_{n>0} S^n(\hbar\mathfrak{t}_+)$, it follows readily that f must vanish on $\mathbb{J}^2 = \bigoplus_{n>1} S^n(\hbar\mathfrak{t}_+)$. Moreover, the above condition gives $f(1) = 2f(1)$, and hence f vanishes on \mathbb{C} . It follows that $f \in S^1(\hbar\mathfrak{t}_+)^* = (\hbar\mathfrak{t}_+)^*$.

Conversely, if $f \in (\hbar\mathfrak{t}_+)^*$, then $\Delta(f)$ vanishes on $S^n(\hbar\mathfrak{t}_+) \otimes S^m(\hbar\mathfrak{t}_+)$ unless $n + m = 1$, and on $\hbar\mathfrak{t}_+ \otimes \mathbb{C}$ and $\mathbb{C} \otimes \hbar\mathfrak{t}_+$ the identity (6.1) trivially holds. This completes the proof that the Lie algebra $\text{Prim}S(\hbar\mathfrak{t}_+)^*$ coincides with $(\hbar\mathfrak{t}_+)^*$ as a vector space. Let us now prove that its bracket is given by

$$[f, g] = (f \otimes g) \circ \hbar\delta_+ \quad \forall f, g \in (\hbar\mathfrak{t}_+)^*.$$

Since $[f, g] \in (\hbar\mathfrak{t}_+)^*$, it is enough to establish this equality on $\hbar\mathfrak{t}_+$. By definition, we have

$$[f, g](\hbar x) = (fg - gf)(\hbar x) = (f \otimes g)(\Delta - \Delta^{\text{op}})(\hbar x) \quad \forall \hbar x \in \hbar\mathfrak{t}_+.$$

It thus suffices to prove that $(\Delta - \Delta^{\text{op}})|_{\hbar\mathfrak{t}_+} - \hbar\delta_+$ has image in $\text{Ker}(f \otimes g)$. As $Y_{\hbar}(\mathfrak{g})$ is a quantization of $(\mathfrak{t}_+, \delta_+)$, we have

$$(\Delta_{Y_{\hbar}(\mathfrak{g})} - \Delta_{Y_{\hbar}(\mathfrak{g})}^{\text{op}})(\hbar\mu(x)) - \hbar^2(\mu \otimes \mu)\delta_+(x) \in \hbar^3 Y_{\hbar}(\mathfrak{g})^{\otimes 2} \cap \dot{Y}_{\hbar}(\mathfrak{g})^{\otimes 2},$$

where we recall that $\mu = \nu^{-1}$, with ν as in (3.13). Applying $\dot{q} \otimes \dot{q}$ and taking note of (5.3), we obtain

$$(6.2) \quad (\Delta - \Delta^{\text{op}})(\hbar x) - \hbar\delta_+(\hbar x) \in \sum_{n+m=3} \mathbb{J}^n \otimes \mathbb{J}^m \subset \text{Ker}(f \otimes g),$$

which completes the proof of (1). ■

Proof of (2) Consider now (2). By Part (1), $\text{Res}_{\hbar}^{\hbar}$ is an isomorphism of graded Lie algebras $\mathfrak{t}_- \xrightarrow{\sim} \text{PrimS}(\hbar\mathfrak{t}_+)^* \subset \text{S}(\hbar\mathfrak{t}_+)^*$. By the universal property of $U(\mathfrak{t}_-)$, it extends uniquely to a homomorphism of graded Hopf algebras

$$\varphi : U(\mathfrak{t}_-) \rightarrow \text{S}(\hbar\mathfrak{t}_+)^*,$$

which is necessarily injective (by [37, Lemma 5.3.3], for instance). As the finite-dimensional graded components $U(\mathfrak{t}_-)_{-n}$ and $\text{S}(\hbar\mathfrak{t}_+)^*_{-n} = \text{S}_n(\hbar\mathfrak{t}_+)^*$ have the same dimension for each $n \in \mathbb{N}$, it follows that φ is an isomorphism. ■

Proof of (3) If $x \in \mathfrak{t}_-$, then $\varphi(\bar{\omega}(x))$ is the element of $(\hbar\mathfrak{t}_+)^*$ determined by

$$\varphi(\bar{\omega}(x))(\hbar y) = \langle \bar{\omega}(x), y \rangle = \langle x, \bar{\omega}(y) \rangle = \dot{\omega}^t(\varphi(x))(\hbar y) \quad \forall y \in \mathfrak{t}_+,$$

where the second equality follows from the fact that the bilinear form (\cdot, \cdot) on \mathfrak{g} is ω -invariant. As $(\hbar\mathfrak{t}_+)^*$ is stable under $\dot{\omega}^t$ and $U(\mathfrak{t}_-)$ is generated by \mathfrak{t}_- , we may conclude that $\varphi \circ \bar{\omega} = \dot{\omega}^t \circ \varphi$.

Similarly, if $x \in \mathfrak{g}$ and $y \in \mathfrak{t}_-$, then $\varphi([x, y]) \in (\hbar\mathfrak{t}_+)^*$ is determined by

$$\varphi([x, y])(\hbar z) = \langle [x, y], z \rangle = \langle y, [z, x] \rangle = \varphi(y)(\hbar[z, x]) \quad \forall z \in \mathfrak{t}_+.$$

On the other hand, since \mathbb{J}^k is a \mathfrak{g} -submodule of $\text{S}(\hbar\mathfrak{t}_+)$ for each $k \in \mathbb{N}$, $x \cdot \varphi(y)$ also belongs to $(\hbar\mathfrak{t}_+)^*$. Moreover, we have

$$(x \cdot \varphi(y))(\hbar z) = \varphi(y)(S(x) \cdot \hbar z) = \varphi(y)(\hbar[z, x]) \quad \forall z \in \mathfrak{t}_+,$$

where we have used that $S(x) \cdot \hbar z - \hbar[z, x] \in \mathbb{J}^2 \subset \text{Ker}(\varphi(y))$, which is proven analogously to (6.2). Since \mathfrak{g} acts on both $U(\mathfrak{t}_-)$ and $\text{S}(\hbar\mathfrak{t}_+)^*$ by derivations and $U(\mathfrak{t}_-)$ is generated by \mathfrak{t}_- , the above computation proves that φ is a \mathfrak{g} -module homomorphism. ■

6.4 $\dot{Y}_{\hbar}\mathfrak{g}^*$ as a quantization

Since the dual Yangian $\dot{Y}_{\hbar}\mathfrak{g}^*$ is a \mathbb{Z} -graded topologically free Hopf algebra over $\mathbb{C}[[\hbar]]$ with semiclassical limit that, by Proposition 6.6, can be identified with $U(\mathfrak{t}_-)$ as a graded Hopf algebra, it is a quantized enveloping algebra which provides a homogeneous quantization of a \mathbb{Z} -graded Lie bialgebra structure (\mathfrak{t}_-, δ) on the graded Lie algebra $\mathfrak{t}_- = t^{-1}\mathfrak{g}[t^{-1}]$.

The following theorem asserts that this Lie bialgebra structure on \mathfrak{t}_- is precisely that associated with the Manin triple $(\mathfrak{t}, \mathfrak{t}_+, \mathfrak{t}_-)$ from Section 2.5.

Theorem 6.7 $\dot{Y}_{\hbar}\mathfrak{g}^*$ is a homogeneous quantization of the Lie bialgebra $(\mathfrak{t}_-, \delta_-)$.

Proof In light of the above discussion, the definition δ_- given in Section 2.5, and the identification of Proposition 6.6, it is sufficient to prove that the Lie bialgebra structure on $\mathfrak{t}_- \cong (\hbar\mathfrak{t}_+)^*$ quantized by $\dot{Y}_{\hbar}\mathfrak{g}^*$ has cobracket δ given by

$$\delta(f)(\hbar x \otimes \hbar y) = f(\hbar[x, y]) \quad \forall x, y \in \mathfrak{t}_+, f \in (\hbar\mathfrak{t}_+)^*.$$

By definition, δ is given by the formula

$$\delta(f) := \hbar^{-1}(m^t(\hat{f}) - m_{21}^t(\hat{f})) \pmod{\hbar\dot{Y}_{\hbar}\mathfrak{g}^* \otimes \dot{Y}_{\hbar}\mathfrak{g}^*},$$

where $\hat{f} \in \dot{Y}_{\hbar}\mathfrak{g}^*$ is any lift of $f \in (\hbar\mathfrak{t}_+)^*$ and m is the product on $\dot{Y}_{\hbar}\mathfrak{g}$. For any two elements $x, y \in \mathfrak{t}_+$, we have

$$(6.3) \quad [\mu(x), \hbar\mu(y)] - \hbar\mu[x, y] \in \mathfrak{J}^2 \subset \hat{f}^{-1}(\hbar\mathbb{C}[[\hbar]]),$$

which implies the desired result:

$$\begin{aligned} (\delta(f) - f \circ \hbar^{-1}[\cdot, \cdot])(\hbar x \otimes \hbar y) \\ = \hat{f}([\mu(x), \hbar\mu(y)] - \hbar\mu[x, y]) \pmod{\hbar\mathbb{C}[[\hbar]]} = 0. \end{aligned} \quad \blacksquare$$

6.5 The dual triangular decomposition

Our main goal in this subsection is to establish a triangular decomposition for $\dot{Y}_{\hbar}\mathfrak{g}^*$ dual to that for $\dot{Y}_{\hbar}\mathfrak{g}$ established in Corollaries 5.5 and 5.6 (see also Remark 5.7). Along the way, we shall identify a family of generators for $\dot{Y}_{\hbar}\mathfrak{g}^*$ (see Lemma 6.8).

For each choice of the symbol χ , consider the restricted dual $\dot{Y}_{\hbar}^{\chi}\mathfrak{g}^*$ of the \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -algebra $\dot{Y}_{\hbar}^{\chi}\mathfrak{g}$. By Corollaries 2.9 and 5.5, this is a \mathbb{Z} -graded topologically free $\mathbb{C}[[\hbar]]$ -module with semiclassical limit equal to the graded dual

$$S(\hbar\mathfrak{t}_+^{\chi})^* = \bigoplus_{n \in \mathbb{N}} S(\hbar\mathfrak{t}_+^{\chi})_{-n}^*$$

of the symmetric algebra $S(\hbar\mathfrak{t}_+^{\chi})$. Moreover, by Remark 2.8, $\dot{Y}_{\hbar}^{\chi}\mathfrak{g}^*$ coincides with the \hbar -adic completion of the graded dual to $\dot{Y}_{\hbar}^{\chi}(\mathfrak{g})$ taken in the category of \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -modules. Now let $\pi^{\chi} : \dot{Y}_{\hbar}\mathfrak{g} \rightarrow \dot{Y}_{\hbar}^{\chi}\mathfrak{g}$ denote the $\mathbb{C}[[\hbar]]$ -linear projection associated with the identification of $\dot{Y}_{\hbar}\mathfrak{g}$ with $\dot{Y}_{\hbar}^{\chi}\mathfrak{g} \otimes \dot{Y}_{\hbar}^0\mathfrak{g} \otimes \dot{Y}_{\hbar}^{-}\mathfrak{g}$ established in Remark 5.7. That is, we have

$$\pi^+ = \mathbf{1}_{\dot{Y}_{\hbar}^+\mathfrak{g}} \otimes \varepsilon \otimes \varepsilon, \quad \pi^0 = \varepsilon \otimes \mathbf{1}_{\dot{Y}_{\hbar}^0\mathfrak{g}} \otimes \varepsilon \quad \text{and} \quad \pi^- = \varepsilon \otimes \varepsilon \otimes \mathbf{1}_{\dot{Y}_{\hbar}^-\mathfrak{g}}.$$

Taking the transpose of π^{χ} yields a \mathbb{Z} -graded embedding of $\mathbb{C}[[\hbar]]$ -modules

$$(\pi^{\chi})^t : \dot{Y}_{\hbar}^{\chi}\mathfrak{g}^* \hookrightarrow \dot{Y}_{\hbar}\mathfrak{g}^*, \quad f \mapsto f \circ \pi^{\chi} \quad \forall f \in \dot{Y}_{\hbar}^{\chi}\mathfrak{g}^*,$$

with image consisting of precisely those $g \in \dot{Y}_{\hbar}\mathfrak{g}^*$ for which $g \circ \pi^{\chi} = g$. Note that if $\hbar^n g$ is contained in this image for some $n \in \mathbb{N}$, then g itself is, and so the subspace topology on $(\pi^{\chi})^t(\dot{Y}_{\hbar}^{\chi}\mathfrak{g}^*)$ coincides with its \hbar -adic topology. Furthermore, the semiclassical limit of $(\pi^{\chi})^t$ is the embedding $S(\hbar\mathfrak{t}_+^{\chi})^* \hookrightarrow S(\hbar\mathfrak{t}_+)^*$ induced by the projection $\mathfrak{t}_+ = \mathfrak{t}_+^+ \oplus \mathfrak{t}_+^0 \oplus \mathfrak{t}_+^- \rightarrow \mathfrak{t}_+^{\chi}$.

We shall henceforth adopt the viewpoint that $\dot{Y}_{\hbar}^{\chi} \mathfrak{g}^*$ is a \mathbb{Z} -graded topological submodule of $\dot{Y}_{\hbar} \mathfrak{g}^*$, with the above identification assumed. We further note that the Chevalley involution ω^t of $\dot{Y}_{\hbar} \mathfrak{g}^*$ satisfies

$$\omega^t(\dot{Y}_{\hbar}^{\pm} \mathfrak{g}^*) = \dot{Y}_{\hbar}^{\mp} \mathfrak{g}^* \quad \text{and} \quad \omega^t|_{\dot{Y}_{\hbar}^0 \mathfrak{g}^*} = \mathbf{1}_{\dot{Y}_{\hbar}^0 \mathfrak{g}^*}.$$

Let us now identify a set of elements which generate $\dot{Y}_{\hbar} \mathfrak{g}^*$ as a topological $\mathbb{C}[[\hbar]]$ -algebra. These generators are constructed so as to naturally correspond to the coefficients of the $DY_{\hbar} \mathfrak{g}$ -valued series $\dot{h}_i(u)$ and $\dot{x}_{\beta}^{\pm}(u)$ defined in Section 4.2, and such an identification will be made precise in the proof of Lemma 7.4 (see (7.2)).

Given $\beta \in Q$, let $\pi_{\beta} : \dot{Y}_{\hbar}^{-}(\mathfrak{g}) \rightarrow \dot{Y}_{\hbar}^{-}(\mathfrak{g})_{\beta}$ denote the natural projection onto the β -component of $\dot{Y}_{\hbar}^{-}(\mathfrak{g})$. Since the $\mathbb{C}[[\hbar]]$ -module isomorphism

$$\dot{\nu} = \nu|_{\dot{Y}_{\hbar}(\mathfrak{g})} : \dot{Y}_{\hbar}(\mathfrak{g}) \xrightarrow{\sim} R_{\hbar}(U(\mathfrak{t}_+)) \subset U(\mathfrak{t}_+)[\hbar]$$

is Q -graded and respects the underlying triangular decompositions, we have

$$\dot{\nu}(\pi_{-\alpha_i}(\dot{Y}_{\hbar}^{-}(\mathfrak{g}))) \subset \bigoplus_{k \in \mathbb{N}} \mathbb{C}[[\hbar]] \cdot \hbar x_i^{-} t^k \subset \hbar \mathfrak{t}_+[\hbar] \quad \forall i \in I.$$

We may thus compose $\dot{\nu} \circ \pi_{-\alpha_i}$ with $\text{Res}_{-}^{\hbar}(x_i^{+} t^{-k-1}) \in (\hbar \mathfrak{t}_+)^*$, as defined above Proposition 6.6, for any fixed $k \in \mathbb{N}$. This outputs a degree $-k - 1$ element

$$(6.4) \quad X_{i,-k-1}^{+} := \text{Res}_{-}^{\hbar}(x_i^{+} t^{-k-1}) \circ \dot{\nu} \circ \pi_{-\alpha_i} \in \dot{Y}_{\hbar}^{-} \mathfrak{g}^* \subset \dot{Y}_{\hbar} \mathfrak{g}^*,$$

where we work through the identification of $\text{Hom}_{\mathbb{C}[[\hbar]]}^a(\dot{Y}_{\hbar}^{-}(\mathfrak{g}), \mathbb{C}[[\hbar]])$ with the a th component $\text{Hom}_{\mathbb{C}[[\hbar]]}^a(\dot{Y}_{\hbar}^{-} \mathfrak{g}, \mathbb{C}[[\hbar]])$ of $\dot{Y}_{\hbar}^{-} \mathfrak{g}^*$ (see Remark 2.8). We now enlarge this family of elements to a generating set for the $\mathbb{C}[[\hbar]]$ -algebra $\dot{Y}_{\hbar} \mathfrak{g}^*$ using the coadjoint action of \mathfrak{g} on $\dot{Y}_{\hbar} \mathfrak{g}^*$ and the Chevalley involution ω^t from Section 6.2. For each $i \in I$, $\beta \in \Delta^+$, and $k \in \mathbb{N}$, we introduce the degree $-k - 1$ elements

$$h_{i,-k-1} = -x_i^{-} \cdot X_{i,-k-1}^{+},$$

$$X_{\beta,-k-1}^{+} := X^{\beta} \cdot X_{i(\beta),-k-1}^{+} \in \dot{Y}_{\hbar} \mathfrak{g}_{\beta}^*, \quad X_{\beta,-k-1}^{-} := \omega^t(X_{\beta,-k-1}^{+}) \in \dot{Y}_{\hbar} \mathfrak{g}_{-\beta}^*,$$

where $i(\beta) \in I$ and $X^{\beta} \in U(\mathfrak{n}_+)_{\beta - \alpha_{i(\beta)}}$ are as in (3.11). In the same spirit as in Section 4.2, we organize these elements into generating series in $\dot{Y}_{\hbar} \mathfrak{g}^*[[u]]$ by setting

$$h_i(u) := - \sum_{r < 0} h_{i,r} u^{-r-1} \quad \text{and} \quad X_{\beta}^{\pm}(u) := - \sum_{r < 0} X_{\beta,r}^{\pm} u^{-r-1}$$

for each $i \in I$ and $\beta \in \Delta^+$. We then have the following lemma.

Lemma 6.8 *The dual Yangian $\dot{Y}_{\hbar} \mathfrak{g}^*$ is topologically generated as a $\mathbb{C}[[\hbar]]$ -algebra by the coefficients of $\{X_{\beta}^{\pm}(u)\}_{\beta \in \Delta^+}$ and $\{h_i(u)\}_{i \in I}$. Moreover, their images under the quotient map $\dot{Y}_{\hbar} \mathfrak{g}^* \twoheadrightarrow \dot{Y}_{\hbar} \mathfrak{g}^* / \hbar \dot{Y}_{\hbar} \mathfrak{g}^* \cong U(\mathfrak{t}_-)$ are given by*

$$h_{i,r} \mapsto h_i t^r \quad \text{and} \quad X_{\beta,r}^{\pm} \mapsto x_{\beta}^{\pm} t^r \quad \forall i \in I, \beta \in \Delta^+ \text{ and } r < 0.$$

Proof As $\dot{Y}_{\hbar} \mathfrak{g}^*$ is a flat deformation of the algebra $U(\mathfrak{t}_-)$ over $\mathbb{C}[[\hbar]]$ and the elements $h_i t^{-k-1}$ and $x_{\beta}^{\pm} t^{-k-1}$ generate $\mathfrak{t}_- = t^{-1} \mathfrak{g}[t^{-1}]$ as a Lie algebra, it is sufficient to prove the second assertion of the proposition.

That the element X_{ir}^+ coincides with $x_i^+ t^r$ modulo $\hbar \dot{Y}_{\hbar \mathfrak{g}}^*$ is an immediate consequence of its definition and the identification of $U(\mathfrak{t}_-)$ with $\dot{Y}_{\hbar \mathfrak{g}}^* / \hbar \dot{Y}_{\hbar \mathfrak{g}}^* \cong S(\hbar \mathfrak{t}_-)^*$ provided by Part (2) of Proposition 6.6. The remaining equivalences now follow from the definitions of h_i and x_β^\pm (see (3.11)) and Part (3) of Proposition 6.6, which implies that the quotient map $\dot{Y}_{\hbar \mathfrak{g}}^* \twoheadrightarrow U(\mathfrak{t}_-)$ is a \mathfrak{g} -module homomorphism intertwining Chevalley involutions. ■

Let us now turn toward establishing a triangular decomposition for the dual Yangian $\dot{Y}_{\hbar \mathfrak{g}}^*$. Following the notation from Section 5.3, let us introduce the Lie subalgebras \mathfrak{t}^χ of $\mathfrak{t} = t^{-1} \mathfrak{g}[t^{-1}]$ by setting

$$\mathfrak{t}^\pm := t^{-1} \mathfrak{n}_\pm[t^{-1}] \subset \mathfrak{t}_- \quad \text{and} \quad \mathfrak{t}^0 := t^{-1} \mathfrak{h}[t^{-1}] \subset \mathfrak{t}_-.$$

The below proposition provides a strengthening of the \hbar -adic analogues of Proposition 3.2 and Theorem 4.2 (i) from [32].

Proposition 6.9 $\dot{Y}_{\hbar \mathfrak{g}}^\pm$ and $\dot{Y}_{\hbar \mathfrak{g}}^0$ are $\mathbb{C}[[\hbar]]$ -subalgebras of $\dot{Y}_{\hbar \mathfrak{g}}^*$. Moreover:

- (1) $\dot{Y}_{\hbar \mathfrak{g}}^\pm$ is a flat deformation of the \mathbb{Z} -graded algebra $U(\mathfrak{t}^\mp)$ over $\mathbb{C}[[\hbar]]$. In particular, there is an isomorphism of \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -modules

$$\dot{Y}_{\hbar \mathfrak{g}}^\pm \cong U(\mathfrak{t}^\mp)[[\hbar]].$$

- (2) $\dot{Y}_{\hbar \mathfrak{g}}^0$ is commutative and isomorphic to $U(\mathfrak{t}^0)[[\hbar]] \cong S(\mathfrak{t}^0)[[\hbar]]$ as a \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -algebra.
- (3) $\dot{Y}_{\hbar \mathfrak{g}}^\pm$ and $\dot{Y}_{\hbar \mathfrak{g}}^0$ are topologically generated as $\mathbb{C}[[\hbar]]$ -algebras by the coefficients of $\{X_\beta^\mp(u)\}_{\beta \in \Delta^+}$ and $\{h_i(u)\}_{i \in I}$, respectively.

Proof One can deduce that $\dot{Y}_{\hbar \mathfrak{g}}^\chi$ is a subalgebra of $\dot{Y}_{\hbar \mathfrak{g}}^*$ using properties of the Yangian coproduct, as in [32, Proposition 3.2]. We shall give an alternate simple proof of this fact in Section 7.3 which illustrates that it follows naturally from properties of $\mathcal{R}(z)$ (see Corollary 7.7 and Remark 7.8).

Let us complete the proof of the proposition assuming Corollary 7.7, which, as explained in Remark 7.8, also implies that the coefficients of $\{X_\beta^\mp(u)\}_{\beta \in \Delta^+}$ and $\{h_i(u)\}_{i \in I}$ belong to $\dot{Y}_{\hbar \mathfrak{g}}^\mp$ and $\dot{Y}_{\hbar \mathfrak{g}}^0$, respectively, and that $\dot{Y}_{\hbar \mathfrak{g}}^0$ is commutative. ■

Proof of (1) and (2) The graded Lie bialgebra isomorphism $\text{Res}_-^h : \mathfrak{t}_- \xrightarrow{\sim} (\hbar \mathfrak{t}_-)^*$ of Proposition 6.6 restricts to an isomorphism $\mathfrak{t}_-^\chi \xrightarrow{\sim} (\hbar \mathfrak{t}_+^\chi)^* \subset S(\hbar \mathfrak{t}_+^\chi)^*$. Since $\dot{Y}_{\hbar \mathfrak{g}}^\chi$ is a subalgebra of $\dot{Y}_{\hbar \mathfrak{g}}^*$, its semiclassical limit $S(\hbar \mathfrak{t}_+^\chi)^*$ is a \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -subalgebra of $S(\hbar \mathfrak{t}_-)^*$ and not just a submodule. It follows from these observations that the graded Hopf algebra isomorphism φ of Proposition 6.6 restricts to an injective \mathbb{Z} -graded algebra homomorphism

$$(6.5) \quad \varphi_\chi : U(\mathfrak{t}_-^\chi) \rightarrow S(\hbar \mathfrak{t}_+^\chi)^* \subset S(\hbar \mathfrak{t}_-)^*$$

which is surjective by the same argument as used in the proof of Part (2) of Proposition 6.6. Since $\dot{Y}_{\hbar \mathfrak{g}}^\chi$ is a \mathbb{Z} -graded topologically free $\mathbb{C}[[\hbar]]$ -algebra with semiclassical limit $S(\hbar \mathfrak{t}_+^\chi)^* \cong U(\mathfrak{t}_-^\chi)$, taking $\chi = \pm$ recovers Part (1) of the proposition.

As for Part (2), since $\dot{Y}_\hbar^0 \mathfrak{g}^*$ is a commutative topological $\mathbb{C}[[\hbar]]$ -algebra containing $\{h_{ik}\}_{i \in I, k < 0}$ (by Corollary 7.7), there is a $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$\eta : S(\mathfrak{t}_-^0)[[\hbar]] \rightarrow \dot{Y}_\hbar^0 \mathfrak{g}^*$$

uniquely determined by $\eta(h_i t^k) = h_{ik}$ for all $i \in I$ and $k < 0$. By Lemma 6.8, the semiclassical limit $\bar{\eta} : S(\mathfrak{t}_-^0) \rightarrow U(\mathfrak{t}_-^0)$ of η satisfies $\bar{\eta}(h_i t^k) = h_{ik}$ for all $i \in I$ and $k < 0$, and thus coincides with the canonical isomorphism $S(\mathfrak{t}_-^0) \xrightarrow{\sim} U(\mathfrak{t}_-^0)$. Here, we have assumed the identification of $\dot{Y}_\hbar^0 \mathfrak{g}^* / \hbar \dot{Y}_\hbar^0 \mathfrak{g}^*$ with $U(\mathfrak{t}_-^0)$ provided by φ_0 from (6.5) above. As $S(\mathfrak{t}_-^0)$ and $\dot{Y}_\hbar^0 \mathfrak{g}^*$ are both topologically free, we can conclude from Lemma 2.1 that η is an isomorphism of topological $\mathbb{C}[[\hbar]]$ -algebras. ■

Proof of (3) For $\dot{Y}_\hbar^0 \mathfrak{g}^*$, this follows from the definition of the isomorphism η given in the proof of (2) above.

Similarly, by Lemma 6.8 and Corollary 7.7, $X_{\beta,k}^\mp$ belongs to $\dot{Y}_\hbar^\pm \mathfrak{g}^*$ and specializes to $x_\beta^\mp t^k$ in $\dot{Y}_\hbar^\pm \mathfrak{g}^* / \hbar \dot{Y}_\hbar^\pm \mathfrak{g}^* \cong U(\mathfrak{t}_\mp) \subset U(\mathfrak{t}_-)$, for each $\beta \in \Delta^+$ and $k < 0$. Since the elements $x_\beta^\mp t^k$ generate the Lie algebra \mathfrak{t}_\mp and, by Part (1), $\dot{Y}_\hbar^\pm \mathfrak{g}^*$ is a flat deformation of the algebra $U(\mathfrak{t}_\mp)$ over $\mathbb{C}[[\hbar]]$, this completes the proof of the proposition. ■

Remark 6.10 We caution that $\dot{Y}_\hbar^\pm \mathfrak{g}^*$ is *not* generated by the elements $X_{i,-k-1}^\mp$, for $i \in I$ and $k \in \mathbb{N}$, unless $\mathfrak{g} = \mathfrak{sl}_2$, just as $t^{-1}n_\mp[t^{-1}]$ is not generated as a Lie algebra by the elements $x_i^\mp t^{-k-1}$ outside of the rank one case. In particular, the statement of Part (i) in [32, Theorem 4.2], which is the analogue of Part (2) above, should be adjusted.

As an application of Lemma 2.1, Proposition 6.9, the decomposition $\mathfrak{t}_- = \mathfrak{t}_-^- \oplus \mathfrak{t}_-^0 \oplus \mathfrak{t}_-^+$, and the Poincaré–Birkhoff–Witt theorem for enveloping algebras, we obtain the following variant of Theorem 3.1(ii) in [32].

Corollary 6.11 *The multiplication map*

$$m : \dot{Y}_\hbar^+ \mathfrak{g}^* \otimes \dot{Y}_\hbar^0 \mathfrak{g}^* \otimes \dot{Y}_\hbar^- \mathfrak{g}^* \rightarrow \dot{Y}_\hbar \mathfrak{g}^*$$

is an isomorphism of \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -modules.

Remark 6.12 We note that for the statement of [32, Theorem 3.1] to hold, the tensor product \otimes must be taken to be a completion of the algebraic tensor product $\otimes_{\mathbb{C}}$ compatible with the underlying \mathbb{Z} -filtrations.

7 DY_ħg as a quantization

In this section, we construct a \mathbb{Z} -graded topological Hopf algebra structure on $DY_\hbar \mathfrak{g}$ which quantizes the graded Lie bialgebra structure on the loop algebra $\mathfrak{t} = \mathfrak{g}[t^{\pm 1}]$ defined in Section 2.5. This will be achieved in Theorem 7.5 using Proposition 7.1 and Corollary 7.3. As a consequence of these results, we obtain in Section 7.3 a characterization of the restricted duals $\dot{Y}_\hbar^\pm \mathfrak{g}^*$ and $\dot{Y}_\hbar^0 \mathfrak{g}^*$ in terms of the universal R -matrix $\mathcal{R}(z)$ which completes the proof of Proposition 6.9.

7.1 The morphism Φ_z^\vee

Henceforth, the notation $Y_\hbar \mathfrak{g}^\vee$ will be used to denote the topological Hopf algebra $(\dot{Y}_\hbar \mathfrak{g}^*)^{\text{cop}}$ over $\mathbb{C}[[\hbar]]$. That is, $Y_\hbar \mathfrak{g}^\vee$ coincides with the dual Yangian $\dot{Y}_\hbar \mathfrak{g}^*$ as an algebra,

and has coproduct Δ^\vee , counit ε^\vee , and antipode S^\vee given by

$$\begin{aligned} \Delta^\vee &:= (12) \circ m^t : Y_{\hbar} \mathfrak{g}^\vee \rightarrow Y_{\hbar} \mathfrak{g}^\vee \otimes Y_{\hbar} \mathfrak{g}^\vee, \\ \varepsilon^\vee &:= \iota^t : Y_{\hbar} \mathfrak{g}^\vee \rightarrow \mathbb{C}[[\hbar]], \quad S^\vee := (S^{-1})^t : Y_{\hbar} \mathfrak{g}^\vee \rightarrow Y_{\hbar} \mathfrak{g}^\vee, \end{aligned}$$

where m , ι , and S are the product, unit, and antipode of the Drinfeld–Gavarini Yangian $\check{Y}_{\hbar} \mathfrak{g} \subset Y_{\hbar} \mathfrak{g}$ (see Section 6.1). By Theorem 6.7, $Y_{\hbar} \mathfrak{g}^\vee$ is a flat deformation of the enveloping algebra $U(\mathfrak{t}_-) = U(t^{-1} \mathfrak{g}[t^{-1}])$ over $\mathbb{C}[[\hbar]]$ which provides a homogeneous quantization of the Lie bialgebra $(\mathfrak{t}_-, -\delta_-)$.

Our present goal is to construct a homomorphism of $\mathbb{C}[[\hbar]]$ -algebras

$$\Phi_z^\vee : Y_{\hbar} \mathfrak{g}^\vee \rightarrow Y_{\hbar} \mathfrak{g}^\vee[[z^{-1}]]$$

which is compatible with both the formal shift operator Φ_z of Theorem 4.6 and the Hopf structures on $Y_{\hbar} \mathfrak{g}^\vee$ and $Y_{\hbar} \mathfrak{g}$. This is achieved using the universal R -matrix $\mathcal{R}(z)$ of the Yangian as follows. By Lemma 5.9, $\mathcal{R}(z)$ is an element of $(\check{Y}_{\hbar}(\mathfrak{g}) \otimes Y_{\hbar}(\mathfrak{g}))[[z^{-1}]]$, and thus gives rise to a $\mathbb{C}[[\hbar]]$ -module homomorphism

$$\Phi_z^\vee : Y_{\hbar} \mathfrak{g}^\vee \rightarrow Y_{\hbar} \mathfrak{g}^\vee[[z^{-1}]], \quad f \mapsto (f \otimes 1)\mathcal{R}(-z) \quad \forall f \in Y_{\hbar} \mathfrak{g}^\vee.$$

Now recall that $\dot{h}_i(u), \dot{x}_\beta^\pm(u) \in DY_{\hbar} \mathfrak{g}[[u]]$ and $h_i(u), X_\beta^\pm(u) \in Y_{\hbar} \mathfrak{g}^\vee[[u]]$ are the generating series defined in Sections 4.2 and 6.5, respectively. In addition, we let \mathcal{E}_z denote the canonical $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$\mathcal{E}_z : Y_{\hbar} \mathfrak{g}[[w^{-1}]] \otimes Y_{\hbar} \mathfrak{g}[[z^{-1}]] \rightarrow Y_{\hbar} \mathfrak{g}^{\otimes 2}[[z^{-1}]]$$

given by evaluating w to z . The following proposition asserts that Φ_z^\vee indeed has the desired properties.

Proposition 7.1 Φ_z^\vee has the following properties:

(1) It is a homomorphism of $\mathbb{C}[[\hbar]]$ -algebras satisfying

$$\Delta \circ \Phi_z^\vee = \mathcal{E}_z \circ (\Phi_w^\vee \otimes \Phi_z^\vee) \circ \Delta^\vee, \quad \varepsilon \circ \Phi_z^\vee = \varepsilon^\vee \quad \text{and} \quad S \circ \Phi_z^\vee = \Phi_z^\vee \circ S^\vee.$$

(2) It is a $U(\mathfrak{g})$ -module homomorphism compatible with Chevalley involutions:

$$\Phi_z^\vee \circ \omega^t = \omega \circ \Phi_z^\vee.$$

(3) Its restriction to $\check{Y}_{\hbar}(\mathfrak{g})^*$ is a \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$\Phi_z^\vee|_{\check{Y}_{\hbar}(\mathfrak{g})^*} : \check{Y}_{\hbar}(\mathfrak{g})^* \rightarrow \mathbb{L}\widehat{Y_{\hbar} \mathfrak{g}}_z.$$

(4) For each $\beta \in \Delta^+$ and $i \in I$, one has

$$\begin{aligned} \Phi_z^\vee(X_\beta^\pm(u)) &= \sum_{n \in \mathbb{N}} (-1)^n u^n \partial_z^{(n)} x_\beta^\pm(-z) = \Phi_z(\dot{x}_\beta^\pm(u)), \\ \Phi_z^\vee(h_i(u)) &= \sum_{n \in \mathbb{N}} (-1)^n u^n \partial_z^{(n)} h_i(-z) = \Phi_z(\dot{h}_i(u)). \end{aligned}$$

Proof of (1) This is a modification of the standard result that if H is a finite-dimensional quasitriangular Hopf algebra over a field \mathbb{K} with R -matrix $R \in H \otimes_{\mathbb{K}} H$, then the map $f \mapsto (f \otimes 1_H)R$ defines a homomorphism of Hopf algebras $(H^*)^{\text{cop}} \rightarrow H$ (see, for instance, [4, Section 4.2.B] or [39, Section 2]). The only novelty in the present

setting is the appearance of the formal parameter z . Nonetheless, for the sake of completeness, we shall give a full proof.

As the product on $Y_{\hbar}\mathfrak{g}^\vee$ is the transpose of the coproduct Δ on $\dot{Y}_{\hbar}\mathfrak{g}$ and $\Delta \otimes \mathbf{1}(\mathcal{R}(z)) = \mathcal{R}_{13}(z)\mathcal{R}_{23}(z)$, we have

$$\Phi_z^\vee(fg) = f \otimes g \otimes \mathbf{1}(\mathcal{R}_{13}(-z)\mathcal{R}_{23}(-z)) = \Phi_z^\vee(f)\Phi_z^\vee(g) \quad \forall f, g \in Y_{\hbar}\mathfrak{g}^\vee.$$

Since, in addition, $(\varepsilon \otimes \mathbf{1})\mathcal{R}(z) = 1$, we can conclude that Φ_z^\vee is a homomorphism of unital, associative $\mathbb{C}[[\hbar]]$ -algebras.

Let us now verify the coproduct identity $\Delta \circ \Phi_z^\vee = \mathcal{E}_z \circ (\Phi_w^\vee \otimes \Phi_z^\vee) \circ \Delta^\vee$ on $f \in Y_{\hbar}\mathfrak{g}^\vee$. Using the cabling identity $\mathbf{1} \otimes \Delta(\mathcal{R}(z)) = \mathcal{R}_{13}(z)\mathcal{R}_{12}(z)$, we obtain

$$\begin{aligned} \Delta(\Phi_z^\vee(f)) &= (f \otimes \Delta)(\mathcal{R}(-z)) \\ &= (f \otimes \mathbf{1} \otimes \mathbf{1})(\mathcal{R}_{13}(-z)\mathcal{R}_{12}(-z)) \\ &= (\Delta^\vee(f) \otimes \mathbf{1} \otimes \mathbf{1})(\mathcal{R}_{13}(-z)\mathcal{R}_{24}(-z)) \\ &= (\mathcal{E}_z \circ (\Delta^\vee(f) \otimes \mathbf{1} \otimes \mathbf{1}))(\mathcal{R}_{13}(-w)\mathcal{R}_{24}(-z)) \\ &= (\mathcal{E}_z \circ (\Phi_w^\vee \otimes \Phi_z^\vee) \circ \Delta^\vee)(f). \end{aligned}$$

Finally, the remaining two identities follow from the relations $(\mathbf{1} \otimes \varepsilon)\mathcal{R}(z) = 1$ and $(S^{-1} \otimes \mathbf{1})\mathcal{R}(z) = (\mathbf{1} \otimes S)\mathcal{R}(z)$. Indeed, for each $f \in Y_{\hbar}\mathfrak{g}^\vee$, we have

$$\varepsilon \circ \Phi_z^\vee(f) = (f \otimes \varepsilon)\mathcal{R}(-z) = f(1) = \varepsilon^\vee(f),$$

$$S \circ \Phi_z^\vee(f) = (f \otimes S)\mathcal{R}(-z) = (S^\vee(f) \otimes \mathbf{1})\mathcal{R}(-z) = \Phi_z^\vee \circ S^\vee(f). \quad \blacksquare$$

Proof of (2) Since τ_z restricts to the identity on $U(\mathfrak{g})$, the intertwiner equation (3.20) implies that $\mathcal{R}(z)$ is a \mathfrak{g} -invariant element of $Y_{\hbar}(\mathfrak{g})^{\otimes 2}[[z^{-1}]]$:

$$[x \otimes \mathbf{1} + \mathbf{1} \otimes x, \mathcal{R}(z)] = 0 \quad \forall x \in \mathfrak{g}.$$

It follows readily from this fact, and the definition of the \mathfrak{g} -module structure on $Y_{\hbar}\mathfrak{g}^\vee = \dot{Y}_{\hbar}\mathfrak{g}^*$ introduced in Section 6.2, that Φ_z^\vee is a $U(\mathfrak{g})$ -module homomorphism.

Similarly, by Corollary 3.10, the Chevalley involution ω satisfies $(\omega \otimes \omega)\mathcal{R}(z) = \mathcal{R}(z)$, and we thus have

$$\Phi_z^\vee(\omega^t(f)) = ((f \circ \omega) \otimes \omega^2)\mathcal{R}(-z) = (f \otimes \omega)\mathcal{R}(-z) = \omega(\Phi_z^\vee(f)) \quad \forall f \in Y_{\hbar}\mathfrak{g}^\vee. \quad \blacksquare$$

Proof of (3) Suppose that f is a degree k element of $\dot{Y}_{\hbar}(\mathfrak{g})^*$ for some $k \in \mathbb{Z}$. Then $f \otimes \mathbf{1} : \dot{Y}_{\hbar}(\mathfrak{g}) \otimes Y_{\hbar}(\mathfrak{g}) \rightarrow Y_{\hbar}(\mathfrak{g})$ is homogeneous of degree k , and hence

$$f \otimes \mathbf{1}(\widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)} \cap (\dot{Y}_{\hbar}(\mathfrak{g}) \otimes Y_{\hbar}(\mathfrak{g})))[[z^{-1}]] \subset z^k \prod_{n \in \mathbb{N}} Y_{\hbar}(\mathfrak{g})_{n+k} z^{-n-k} \subset z^k \widehat{Y_{\hbar}\mathfrak{g}}_z,$$

where $\widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)} = \prod_{n \in \mathbb{N}} (Y_{\hbar}(\mathfrak{g})^{\otimes 2})_n z^{-n}$, as in Section 3.6. The assertion now follows from Part (3) of Theorem 3.8 and Lemma 5.9, which yield

$$\mathcal{R}(z) \in \widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)} \cap (\dot{Y}_{\hbar}(\mathfrak{g}) \otimes Y_{\hbar}(\mathfrak{g})))[[z^{-1}]]. \quad \blacksquare$$

Proof of (4) Since Φ_z^\vee is a $U(\mathfrak{g})$ -module homomorphism intertwining Chevalley involutions, it is sufficient to establish that

$$(71) \quad \Phi_z^\vee(X_{i,-k-1}^+) = (-1)^{k+1} \partial_z^{(k)} x_i^+(-z) = \Phi_z^\vee(x_{i,-k-1}^+) \quad \forall i \in \mathbf{I} \text{ and } k \in \mathbb{N},$$

where we recall that $X_{i,-k-1}^+$ was defined explicitly in (6.4). Since $\varepsilon \otimes 1$ sends both $\mathcal{R}^0(z)$ and $\mathcal{R}^+(z)$ to 1 and $X_{i,-k-1}^+$ vanishes on $\check{Y}_{\hbar}^-(\mathfrak{g})_{\beta}$ for $\beta \neq -\alpha_i$, we have

$$\Phi_z^{\vee}(X_{i,-k-1}^+) = (X_{i,-k-1}^+ \otimes 1)\mathcal{R}_{\alpha_i}^-(z).$$

Using (3.18), we deduce that the element $\mathcal{R}_{\alpha_i}^-(z)$ is given by

$$\mathcal{R}_{\alpha_i}^-(z) = \sum_{p \geq 0} \frac{\alpha_i(\zeta)\mathbf{T}(\zeta)^p}{(z\alpha_i(\zeta))^{p+1}} (\hbar x_{i,0}^- \otimes x_{i,0}^+) = \sum_{p \geq 0} \sum_{n=0}^p \binom{p}{n} (-1)^n \hbar x_{i,n}^- \otimes x_{i,p-n}^+ z^{-p-1},$$

where we have used $\text{ad}(\mathbf{T}(\zeta))^p(x_{i,0}^{\pm}) = (\pm 1)^p \alpha_i(\zeta)^p x_{i,p}^{\pm}$. This can be rewritten as

$$\mathcal{R}_{\alpha_i}^-(z) = \sum_{n \in \mathbb{N}} \hbar x_{i,n}^- \otimes \partial_z^{(n)} x_i^+(z) = - \sum_{n \in \mathbb{N}} \hbar x_{i,n}^- \otimes \Phi_{-z}(x_{i,-n-1}^+),$$

where the second equality is due to Part (1) of Theorem 4.6. As $X_{i,-k-1}^+(\hbar x_{i,n}^-) = \text{Res}_-^{\hbar}(x_i^+ t^{-k-1})(\hbar x_i^- t^n) = -\delta_{kn}$ for all $k, n \in \mathbb{N}$, this implies the identity (7.1). ■

Remark 7.2 As $Y_{\hbar} \mathfrak{g}^{\vee}$ and $\widehat{LY_{\hbar} \mathfrak{g}_z}$ are \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -algebras with

$$(Y_{\hbar} \mathfrak{g}^{\vee})_{\mathbb{Z}} = (\check{Y}_{\hbar} \mathfrak{g}^*)_{\mathbb{Z}} \cong \check{Y}_{\hbar}(\mathfrak{g})^* \quad \text{and} \quad (\widehat{LY_{\hbar} \mathfrak{g}_z})_{\mathbb{Z}} = \widehat{LY_{\hbar} \mathfrak{g}_z},$$

Part (3) of the proposition is equivalent to the assertion that Φ_z^{\vee} is a \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -algebra homomorphism $\Phi_z^{\vee} : Y_{\hbar} \mathfrak{g}^{\vee} \rightarrow \widehat{LY_{\hbar} \mathfrak{g}_z}$.

Since Φ_z^{\vee} has image in $\widehat{LY_{\hbar} \mathfrak{g}_z}$, we may compose it with $\mathcal{E}\mathcal{U}$ from (4.2) to obtain a $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$\Phi^{\vee} := \mathcal{E}\mathcal{U} \circ \Phi_z^{\vee} : Y_{\hbar} \mathfrak{g}^{\vee} \rightarrow \widehat{Y_{\hbar} \mathfrak{g}}.$$

Our present goal is to apply Part (1) of Proposition 7.1 to interpret Φ^{\vee} as a homomorphism of topological Hopf algebras, where the topological structure on the completed Yangian is that induced by the gradation topology on $Y_{\hbar} \mathfrak{g}$. To make this precise, let us define $\widehat{Y_{\hbar} \mathfrak{g}}^{(n)}$, for any $n \in \mathbb{N}$, to be the formal completion of $Y_{\hbar}(\mathfrak{g})^{\otimes n}$ with respect to its \mathbb{N} -grading:

$$\widehat{Y_{\hbar} \mathfrak{g}}^{(n)} := \prod_{k \in \mathbb{N}} (Y_{\hbar}(\mathfrak{g})^{\otimes n})_k.$$

Equivalently, it is the completion of the \mathbb{N} -graded topological $\mathbb{C}[[\hbar]]$ -algebra $Y_{\hbar} \mathfrak{g}^{\otimes n}$ with respect to the filtration defining its gradation topology, as defined in Section 2.3. By [41, Proposition A.1], this is a topologically free $\mathbb{C}[[\hbar]]$ -algebra containing $Y_{\hbar} \mathfrak{g}^{\otimes n}$ as a subalgebra (see also [41, Lemma 4.1] and Section 4.3 above). Furthermore, we can (and shall) view

$$(\widehat{Y_{\hbar} \mathfrak{g}})^{\otimes n} \subset \widehat{Y_{\hbar} \mathfrak{g}}^{(n)} \quad \forall n \in \mathbb{N},$$

where \otimes denotes the topological tensor product over $\mathbb{C}[[\hbar]]$.

Next, observe that the \mathbb{N} -graded Hopf algebra structure on $Y_{\hbar}(\mathfrak{g})$ induces a topological Hopf structure on $\widehat{Y_{\hbar} \mathfrak{g}}$, equipped with the grading-completed tensor product.

More precisely, as the multiplication m , coproduct Δ , counit ε , and antipode S on $Y_{\hbar}(\mathfrak{g})$ are \mathbb{N} -graded, they uniquely extend to $\mathbb{C}[[\hbar]]$ -module homomorphisms

$$m : \widehat{Y_{\hbar}\mathfrak{g}}^{(2)} \rightarrow \widehat{Y_{\hbar}\mathfrak{g}}, \quad \Delta : \widehat{Y_{\hbar}\mathfrak{g}} \rightarrow \widehat{Y_{\hbar}\mathfrak{g}}^{(2)}, \quad \varepsilon : \widehat{Y_{\hbar}\mathfrak{g}} \rightarrow \mathbb{C}[[\hbar]], \quad S : \widehat{Y_{\hbar}\mathfrak{g}} \rightarrow \widehat{Y_{\hbar}\mathfrak{g}},$$

which collectively satisfy the axioms of a Hopf algebra. Proposition 7.1 then admits the following corollary.

Corollary 7.3 *The $\mathbb{C}[[\hbar]]$ -algebra homomorphism Φ^\vee is a morphism of topological Hopf algebras. That is, it satisfies*

$$\Delta \circ \Phi^\vee = (\Phi^\vee \otimes \Phi^\vee) \circ \Delta^\vee, \quad \varepsilon \circ \Phi^\vee = \varepsilon^\vee \quad \text{and} \quad S \circ \Phi^\vee = \Phi^\vee \circ S^\vee.$$

In particular, one has $\text{Im}(\Delta \circ \Phi^\vee) \subset \widehat{Y_{\hbar}\mathfrak{g}} \otimes \widehat{Y_{\hbar}\mathfrak{g}}$.

Proof The counit and antipode relations are obtained by applying \mathcal{E}^ν to the corresponding relations of Part (1) of Proposition 7.1 and appealing to the identity $\Phi^\vee = \mathcal{E}^\nu \circ \Phi_z^\vee$. The idea now is that the relation $\Delta \circ \Phi^\vee = (\Phi^\vee \otimes \Phi^\vee) \circ \Delta^\vee$ should follow by applying $\mathcal{E}^\nu \otimes \mathcal{E}^\nu$ to both sides of the identity $\Delta \circ \Phi_z^\vee = \mathcal{E}_z \circ (\Phi_w^\vee \otimes \Phi_z^\vee) \circ \Delta^\vee$.

However, to make this precise, we must first make a few technical observations. Recall from (3.16) that $L(\widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)})$ is the \mathbb{Z} -graded subalgebra of $Y_{\hbar}(\mathfrak{g})^{\otimes 2}[[z, z^{-1}]]$ with k th homogeneous component $z^k \widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)}$, where

$$\widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)} = \prod_{n \in \mathbb{N}} (Y_{\hbar}(\mathfrak{g})^{\otimes 2})_n z^{-n} \subset Y_{\hbar}(\mathfrak{g})^{\otimes 2}[[z^{-1}]].$$

Following Section 4.3, we shall write $L(\widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)})$ for the \hbar -adic completion of $L(\widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)})$. This is a \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -algebra contained in the formal series space $(Y_{\hbar}\mathfrak{g} \otimes Y_{\hbar}\mathfrak{g})[[z^{\pm 1}]]$ (see Lemma 4.5). As in (4.2), evaluation at $z = 1$ yields a $\mathbb{C}[[\hbar]]$ -algebra epimorphism

$$\mathcal{E}^{\nu(2)} : L(\widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)}) \twoheadrightarrow \widehat{Y_{\hbar}\mathfrak{g}}^{(2)}, \quad f(z) \mapsto f(1).$$

Next, let \mathcal{E}_z^L denote the natural $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$\mathcal{E}_z^L : L\widehat{Y_{\hbar}\mathfrak{g}}_w \otimes L\widehat{Y_{\hbar}\mathfrak{g}}_z \rightarrow L(\widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)})$$

given by evaluating $w \mapsto z$. Then, since Φ_z^\vee is a \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -algebra homomorphism $Y_{\hbar}\mathfrak{g}^\vee \rightarrow L\widehat{Y_{\hbar}\mathfrak{g}}_z$ and Δ is homogeneous of degree zero, the first relation of Part (1) in Proposition 7.1 is equivalent to the identity

$$\Delta \circ \Phi_z^\vee = \mathcal{E}_z^L \circ (\Phi_w^\vee \otimes \Phi_z^\vee) \circ \Delta^\vee$$

in $\text{Hom}_{\mathbb{C}[[\hbar]]}(Y_{\hbar}\mathfrak{g}^\vee, L(\widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)}))$. As \mathcal{E}_z^L satisfies the relation $\mathcal{E}^{\nu(2)} \circ \mathcal{E}_z^L = \mathcal{E}^\nu \otimes \mathcal{E}^\nu$, applying $\mathcal{E}^{\nu(2)}$ to the above identity recovers $\Delta \circ \Phi^\vee = (\Phi^\vee \otimes \Phi^\vee) \circ \Delta^\vee$. ■

7.2 $DY_{\hbar}\mathfrak{g}$ as a quantization

With Proposition 7.1 and Corollary 7.3 in hand, we now turn to proving that $DY_{\hbar}\mathfrak{g}$ provides a homogeneous quantization of the \mathbb{Z} -graded Lie bialgebra $\mathfrak{t} = \mathfrak{g}[t^{\pm 1}]$, equipped with the Lie cobracket δ defined in Section 2.5.

To begin, observe that the homomorphisms Φ_z^\vee and Φ^\vee satisfy the relation

$$\Gamma \circ \Phi^\vee = \Gamma_z \circ \Phi_z^\vee,$$

where Γ and Γ_z are as in Theorem 4.8 and (4.3), respectively. The $\mathbb{C}[[\hbar]]$ -algebra homomorphism defined by either side of this relation shall be denoted \check{i} :

$$\check{i} := \Gamma_z \circ \Phi_z^\vee : Y_{\hbar}\mathfrak{g}^\vee \rightarrow \widehat{DY_{\hbar}\mathfrak{g}}.$$

The following lemma shows that \check{i} is injective, \mathbb{Z} -graded, and has image contained in the Yangian double $DY_{\hbar}\mathfrak{g}$.

Lemma 7.4 *The morphism \check{i} is an embedding of \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -algebras*

$$\check{i} : Y_{\hbar}\mathfrak{g}^\vee \hookrightarrow DY_{\hbar}\mathfrak{g}$$

satisfying $\Phi \circ \check{i} = \Phi^\vee$ and $\Phi_z \circ \check{i} = \Phi_z^\vee$. In particular, Φ^\vee and Φ_z^\vee are both injective.

Proof Since $\Gamma_z \circ \Phi_z = \mathbf{1}_{DY_{\hbar}\mathfrak{g}}$, Part (4) of Proposition 7.1 yields

$$(7.2) \quad \begin{aligned} \check{i}(X_\beta^\pm(u)) &= \Gamma_z(\Phi_z(\dot{x}_\beta^\pm(u))) = \dot{x}_\beta^\pm(u) \quad \forall \beta \in \Delta^+, \\ \check{i}(h_i(u)) &= \Gamma_z(\Phi_z(\dot{h}_i(u))) = \dot{h}_i(u) \quad \forall i \in \mathbf{I}. \end{aligned}$$

By Remark 4.9, $DY_{\hbar}\mathfrak{g}$ is a closed subspace of its \mathcal{J} -adic completion, viewed as a topological $\mathbb{C}[[\hbar]]$ -module. As $Y_{\hbar}\mathfrak{g}^\vee$ is topologically generated as a $\mathbb{C}[[\hbar]]$ -algebra by the coefficients of all the series $X_\beta^\pm(u)$ and $h_i(u)$ (by Lemma 6.8), the equalities (7.2) imply that \check{i} has image in $DY_{\hbar}\mathfrak{g}$ and thus can be viewed as a $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$\check{i} : Y_{\hbar}\mathfrak{g}^\vee \rightarrow DY_{\hbar}\mathfrak{g}$$

which necessarily satisfies $\Phi \circ \check{i} = \Phi^\vee$. Similarly, since $\Phi_z(DY_{\hbar}\mathfrak{g})$ is closed in $\widehat{LY_{\hbar}\mathfrak{g}_z}$ (see Remark 4.9), Part (4) of Proposition 7.1 and (7.2) give $\Phi_z \circ \check{i} = \Phi_z^\vee$.

As $Y_{\hbar}\mathfrak{g}^\vee$ and $DY_{\hbar}\mathfrak{g}$ are both topologically free, it follows from Lemma 2.1 that \check{i} will be injective provided its semiclassical limit $\check{i} : U(\mathfrak{t}_-) \rightarrow U(\mathfrak{t})$ is. That this is indeed the case is a consequence of Lemma 6.8, the relations (7.2), and the definitions of $\dot{x}_\beta^\pm(u)$ and $\dot{h}_i(u)$ given in Section 4.2, which imply that \check{i} coincides with the natural inclusion of $U(\mathfrak{t}_-)$ into $U(\mathfrak{t})$ induced by the polarization $\mathfrak{t} = \mathfrak{t}_+ \oplus \mathfrak{t}_- = \mathfrak{g}[t] \oplus t^{-1}\mathfrak{g}[t^{-1}]$.

Finally, by Part (2) of Theorem 4.6 and Part (3) of Proposition 7.1, Φ_z and Φ_z^\vee are both \mathbb{Z} -graded (see Remarks 4.7 and 7.2). As Φ_z is injective, it follows automatically that \check{i} is \mathbb{Z} -graded. ■

The embedding \check{i} is at the heart of the following theorem, which provides the first main result of this article.

Theorem 7.5 *There is a unique \mathbb{Z} -graded topological Hopf algebra structure on $DY_{\hbar}\mathfrak{g}$ such that the inclusions*

$$Y_{\hbar}\mathfrak{g} \xrightarrow{i} DY_{\hbar}\mathfrak{g} \xleftarrow{\check{i}} Y_{\hbar}\mathfrak{g}^\vee$$

are morphisms of \mathbb{Z} -graded topological Hopf algebras over $\mathbb{C}[[\hbar]]$. Equipped with this Hopf structure, $DY_{\hbar}\mathfrak{g}$ is a homogeneous quantization of the Lie bialgebra (\mathfrak{t}, δ) .

Proof If $DY_{\hbar}\mathfrak{g}$ can be equipped with a coproduct $\dot{\Delta}$, counit $\dot{\varepsilon}$, and antipode \dot{S} which give it the structure of a topological Hopf algebra over $\mathbb{C}[[\hbar]]$ and, in addition, make ι and \check{i} morphisms of \mathbb{Z} -graded topological Hopf algebras, then this structure is necessarily unique and \mathbb{Z} -graded. Indeed, $\dot{\Delta}$, $\dot{\varepsilon}$, and \dot{S} are determined by their values on any set of generators of the topological algebra $DY_{\hbar}\mathfrak{g}$, and thus by their values on $\iota(Y_{\hbar}\mathfrak{g}) \cup \check{i}(Y_{\hbar}\mathfrak{g}^{\vee})$.

Let us now establish the existence of a topological Hopf structure on $DY_{\hbar}\mathfrak{g}$ over $\mathbb{C}[[\hbar]]$ with the claimed properties. We begin by observing that ι and \check{i} satisfy the relations

$$(7.3) \quad \begin{aligned} \Delta \circ \Phi \circ \iota &= (\Phi \otimes \Phi) \circ (\iota \otimes \iota) \circ \Delta, \\ \Delta \circ \Phi \circ \check{i} &= (\Phi \otimes \Phi) \circ (\check{i} \otimes \check{i}) \circ \Delta^{\vee}. \end{aligned}$$

The first relation follows from the identity $\Phi \circ \iota = \tau_1$ and that, for each $c \in \mathbb{C}$, τ_c is a Hopf algebra automorphism (see Section 3.3). As for the second relation, since $\Phi \circ \check{i} = \Phi^{\vee}$, Corollary 7.3 yields

$$\Delta \circ \Phi \circ \check{i} = \Delta \circ \Phi^{\vee} = (\Phi^{\vee} \otimes \Phi^{\vee}) \circ \Delta^{\vee} = (\Phi \otimes \Phi) \circ (\check{i} \otimes \check{i}) \circ \Delta^{\vee}.$$

Since $DY_{\hbar}\mathfrak{g}$ is generated as a topological $\mathbb{C}[[\hbar]]$ -algebra by $\iota(Y_{\hbar}\mathfrak{g}) \cup \check{i}(Y_{\hbar}\mathfrak{g}^{\vee})$, these relations imply that $\Delta \circ \Phi$ has image satisfying

$$\text{Im}(\Delta \circ \Phi) \subset \text{Im}(\Phi \otimes \Phi) \subset \widehat{Y_{\hbar}\mathfrak{g}} \otimes \widehat{Y_{\hbar}\mathfrak{g}},$$

where we view the right-hand side as subspace of $\widehat{Y_{\hbar}\mathfrak{g}}^{(2)}$, as in Corollary 7.3. We may therefore introduce a $\mathbb{C}[[\hbar]]$ -algebra homomorphism $\dot{\Delta}$ by

$$\dot{\Delta} := (\Gamma \otimes \Gamma) \circ \Delta \circ \Phi : DY_{\hbar}\mathfrak{g} \rightarrow DY_{\hbar}\mathfrak{g} \otimes DY_{\hbar}\mathfrak{g},$$

where Γ is the inverse of $\widehat{\Phi}$, as in Theorem 4.8. Since $\Gamma \circ \Phi = 1$, it follows from this definition and the relations (7.3) that

$$(7.4) \quad \dot{\Delta} \circ \iota = (\iota \otimes \iota) \circ \Delta \quad \text{and} \quad \dot{\Delta} \circ \check{i} = (\check{i} \otimes \check{i}) \circ \Delta^{\vee}.$$

Similarly, from Corollary 7.3 and the relations $\tau_1 \circ S = S \circ \tau_1$ and $\varepsilon \circ \tau_1 = \varepsilon$, we find that ι and \check{i} satisfy

$$\begin{aligned} S \circ \Phi \circ \iota &= \Phi \circ \iota \circ S, & S \circ \Phi \circ \check{i} &= \Phi \circ \check{i} \circ S^{\vee}, \\ \varepsilon \circ \Phi \circ \iota &= \varepsilon, & \varepsilon \circ \Phi \circ \check{i} &= \varepsilon^{\vee}. \end{aligned}$$

In particular, these relations imply that $\text{Im}(S \circ \Phi) \subset \text{Im}(\Phi)$. We may therefore define morphisms \dot{S} and $\dot{\varepsilon}$ by

$$\dot{S} := \Gamma \circ S \circ \Phi : DY_{\hbar}\mathfrak{g} \rightarrow DY_{\hbar}\mathfrak{g} \quad \text{and} \quad \dot{\varepsilon} := \varepsilon \circ \Phi : DY_{\hbar}\mathfrak{g} \rightarrow \mathbb{C}[[\hbar]]$$

which by construction satisfy the compatibility relations

$$(7.5) \quad \dot{S} \circ \iota = \iota \circ S, \quad \dot{S} \circ \check{i} = \check{i} \circ S^{\vee}, \quad \dot{\varepsilon} \circ \iota = \varepsilon \quad \text{and} \quad \dot{\varepsilon} \circ \check{i} = \varepsilon^{\vee}.$$

Since $\Gamma|_{\text{Im}(\Phi)} = \Phi^{-1}$ and $Y_{\hbar}\mathfrak{g}$ is a topological Hopf algebra with coproduct Δ , antipode S , and counit ε , the above definitions imply that $DY_{\hbar}\mathfrak{g}$ is a topological Hopf algebra over $\mathbb{C}[[\hbar]]$ with coproduct $\dot{\Delta}$, antipode \dot{S} , and counit $\dot{\varepsilon}$. Moreover, the relations

(7.4) and (7.5) prove that, when $DY_{\hbar}\mathfrak{g}$ is given this Hopf structure, the \mathbb{Z} -graded embeddings ι and \tilde{i} are homomorphisms of topological Hopf algebras.

We are left to establish the second assertion of the theorem. By what we have shown so far and Theorem 4.3, $DY_{\hbar}\mathfrak{g}$ is a flat, \mathbb{Z} -graded Hopf algebra deformation of the universal enveloping $U(\mathfrak{t})$ of $\mathfrak{t} = \mathfrak{g}[t^{\pm 1}]$ over $\mathbb{C}[[\hbar]]$. It thus provides a homogeneous quantization of a \mathbb{Z} -graded Lie bialgebra structure on the Lie algebra \mathfrak{t} with cobracket $\delta_{\mathfrak{t}}$ given by the formula (2.2). From Theorems 6.7 and 6.7 and the first part of the theorem, $\delta_{\mathfrak{t}}$ satisfies $\delta_{\mathfrak{t}}|_{\mathfrak{t}_+} = \delta_+$ and $\delta_{\mathfrak{t}}|_{\mathfrak{t}_-} = -\delta_-$. As $\mathfrak{t} = \mathfrak{t}_+ \oplus \mathfrak{t}_-$, we can conclude that $\delta_{\mathfrak{t}}$ coincides with the Lie cobracket δ defined in Section 2.5. ■

Remark 7.6 The proof of Theorem 7.5 shows that the coproduct $\hat{\Delta}$, counit $\hat{\epsilon}$, and antipode \hat{S} on $DY_{\hbar}\mathfrak{g}$ are uniquely determined by the requirement that Φ is a homomorphism of topological Hopf algebras

$$\Phi : DY_{\hbar}\mathfrak{g} \rightarrow \widehat{Y_{\hbar}\mathfrak{g}},$$

where $\widehat{Y_{\hbar}\mathfrak{g}}$ is given the topological Hopf structure defined above Corollary 7.3. Here, we emphasize that although $DY_{\hbar}\mathfrak{g}$ is a genuine topological Hopf algebra over $\mathbb{C}[[\hbar]]$ in the sense of Section 2.1, the completed Yangian $\widehat{Y_{\hbar}\mathfrak{g}}$ is not. In the same breath, the Hopf algebra structure on $DY_{\hbar}\mathfrak{g}$ is uniquely characterized by the requirement that Φ_z satisfies the relations

$$\Delta \circ \Phi_z = \mathcal{E}_z^L \circ (\Phi_w \otimes \Phi_z) \circ \hat{\Delta}, \quad \epsilon \circ \Phi_z = \hat{\epsilon} \quad \text{and} \quad S \circ \Phi_z = \Phi_z \circ \hat{S},$$

where \mathcal{E}_z^L is as in the proof of Corollary 7.3. In particular, this makes precise the uniqueness statement in the first assertion of Theorem I.

7.3 The dual triangular decomposition revisited

We conclude this section by giving an equivalent characterization of the restricted dual $\check{Y}_{\hbar}^{\chi}\mathfrak{g}^*$ to $\check{Y}_{\hbar}^{\chi}\mathfrak{g}$ considered in Section 6.5, where we recall that χ takes value $+$, $-$ or 0 .

Corollary 7.7 For each choice of χ , $\check{Y}_{\hbar}^{\chi}\mathfrak{g}^*$ satisfies

$$\check{Y}_{\hbar}^{\chi}\mathfrak{g}^* = \{f \in \check{Y}_{\hbar}\mathfrak{g}^* : \Phi_z^{\vee}(f) = (f \otimes 1)\mathcal{R}^{\chi}(-z)\} = (\Phi_z^{\vee})^{-1}(Y_{\hbar}^{-\chi}\mathfrak{g}[[z^{-1}]]).$$

Proof From the definitions of $\check{Y}_{\hbar}^{\chi}\mathfrak{g}$ and Φ_z^{\vee} , and the Gauss decomposition of the universal R -matrix, we obtain the sequence of inclusions

$$\check{Y}_{\hbar}^{\chi}\mathfrak{g}^* \subset \{f \in \check{Y}_{\hbar}\mathfrak{g}^* : \Phi_z^{\vee}(f) = (f \otimes 1)\mathcal{R}^{\chi}(-z)\} \subset (\Phi_z^{\vee})^{-1}(Y_{\hbar}^{-\chi}\mathfrak{g}[[z^{-1}]]).$$

It therefore suffices to show that $(\Phi_z^{\vee})^{-1}(Y_{\hbar}^{-\chi}\mathfrak{g}[[z^{-1}]]) \subset \check{Y}_{\hbar}^{\chi}\mathfrak{g}^*$. We shall establish this for $\chi = -$. The proof in the remaining cases is identical, and hence omitted.

Let $\pi_+ : Y_{\hbar}\mathfrak{g} \rightarrow Y_{\hbar}^+\mathfrak{g}$ be the $\mathbb{C}[[\hbar]]$ -linear projection associated with the opposite triangular decomposition $Y_{\hbar}\mathfrak{g} \cong Y_{\hbar}^-\mathfrak{g} \otimes Y_{\hbar}^0\mathfrak{g} \otimes Y_{\hbar}^+\mathfrak{g}$ from Remark 3.7, and suppose $f \in \check{Y}_{\hbar}\mathfrak{g}^*$ satisfies $\Phi_z(f) \in Y_{\hbar}^+\mathfrak{g}[[z^{-1}]]$. We wish to show that $f \in \check{Y}_{\hbar}^-\mathfrak{g}^*$.

To begin, note that since $(1 \otimes \epsilon)\mathcal{R}^{\chi}(z) = 1$, we have

$$\Phi_z^{\vee}(f) = \pi_+(\Phi_z^{\vee}(f)) = (f \otimes 1)\mathcal{R}^-(z).$$

Consider now the element $f \circ \pi^- \in \dot{Y}_{\hbar}^- \mathfrak{g}^* \subset \dot{Y}_{\hbar} \mathfrak{g}^*$, where $\pi^- : \dot{Y}_{\hbar} \mathfrak{g} \rightarrow \dot{Y}_{\hbar}^- \mathfrak{g}$ is as in Section 6.5. Since $(\pi^- \otimes 1)\mathcal{R}(z) = \mathcal{R}^-(z)$, it satisfies

$$\Phi_z^\vee(f \circ \pi^-) = (f \otimes 1)\mathcal{R}^-(z) = \Phi_z^\vee(f).$$

As Φ_z^\vee is injective (by Lemma 7.4), we can conclude that $f = f \circ \pi^- \in \dot{Y}_{\hbar}^- \mathfrak{g}^*$. ■

Remark 7.8 Since $Y_{\hbar}^\pm \mathfrak{g}$ and $Y_{\hbar}^0 \mathfrak{g}$ are $\mathbb{C}[[\hbar]]$ -subalgebras of $Y_{\hbar} \mathfrak{g}$, it follows from this corollary that $\dot{Y}_{\hbar}^\chi \mathfrak{g}^*$ is a $\mathbb{C}[[\hbar]]$ -subalgebra of $\dot{Y}_{\hbar} \mathfrak{g}^*$ for each choice of χ . In addition, it is immediate from this characterization and Part (4) of Proposition 7.1 that the coefficients of the series $X_{\beta}^\pm(u)$ and $h_i(u)$ belong to $\dot{Y}_{\hbar}^\mp \mathfrak{g}^*$ and $\dot{Y}_{\hbar}^0 \mathfrak{g}^*$, respectively, for each $\beta \in \Delta^+$ and $i \in I$. Similarly, since Φ_z^\vee is injective and $Y_{\hbar}^0 \mathfrak{g}$ is commutative, we deduce that $\dot{Y}_{\hbar}^0 \mathfrak{g}^*$ is commutative.

In particular, these observations complete the proof of Proposition 6.9. We emphasize that we have not applied that proposition in establishing any of the results in Section 7.

8 DY_ħg as a quantum double

We now turn to reframing Theorem 7.5 in the context of the quantum double. Our central objective is to prove the second main result of this article, Theorem 8.4, which establishes that the Yangian double $DY_{\hbar} \mathfrak{g}$ is isomorphic, as a \mathbb{Z} -graded topological Hopf algebra over $\mathbb{C}[[\hbar]]$, to the restricted quantum double of the Yangian, which is defined explicitly in Section 8.2.

8.1 The quantum double of $U_{\hbar} \mathfrak{b}$

Let us begin by recalling, in broad strokes, the general construction of the quantum double of a quantized enveloping algebra, as was first outlined by Drinfeld in [7, Section 13].

Suppose that $U_{\hbar} \mathfrak{b}$ is a quantization of a finite-dimensional Lie bialgebra \mathfrak{b} , and let $U_{\hbar} \mathfrak{b}^\vee$ denote the quantized enveloping algebra $(U_{\hbar} \mathfrak{b}^\circ)^{\text{cop}}$, where $U_{\hbar} \mathfrak{b}^\circ$ is the topological dual to $U_{\hbar} \mathfrak{b}'$ introduced in Section 5.1. Then there exists a unique topological Hopf algebra $D(U_{\hbar} \mathfrak{b})$ over $\mathbb{C}[[\hbar]]$, the *quantum double* of $U_{\hbar} \mathfrak{b}$, satisfying the following three properties:

- (1) There are embeddings of topological Hopf algebras

$$U_{\hbar} \mathfrak{b} \xrightarrow{i} D(U_{\hbar} \mathfrak{b}) \xleftarrow{j} U_{\hbar} \mathfrak{b}^\vee.$$

- (2) The composite $m \circ (j \otimes i) : U_{\hbar} \mathfrak{b}^\vee \otimes U_{\hbar} \mathfrak{b} \rightarrow D(U_{\hbar} \mathfrak{b})$ is an isomorphism of $\mathbb{C}[[\hbar]]$ -modules.
- (3) The canonical element $R \in U_{\hbar} \mathfrak{b}' \otimes U_{\hbar} \mathfrak{b}^\circ \subset D(U_{\hbar} \mathfrak{b}) \otimes D(U_{\hbar} \mathfrak{b})$ associated with the pairing between $U_{\hbar} \mathfrak{b}'$ and $U_{\hbar} \mathfrak{b}^\circ$, which coincides with the canonical tensor in $U_{\hbar} \mathfrak{b} \otimes U_{\hbar} \mathfrak{b}^*$, defines a quasitriangular structure on $D(U_{\hbar} \mathfrak{b})$. That is, one has

$$\begin{aligned} \Delta^{\text{op}}(x) &= R\Delta(x)R^{-1} \quad \forall x \in D(U_{\hbar} \mathfrak{b}), \\ \Delta \otimes \mathbf{1}(R) &= R_{13}R_{23}, \quad \mathbf{1} \otimes \Delta(R) = R_{13}R_{12}. \end{aligned}$$

In addition, $D(U_{\hbar}\mathfrak{b})$ provides a quantization of the Drinfeld double of the finite-dimensional Lie bialgebra \mathfrak{b} .

The quantum double $D(U_{\hbar}\mathfrak{b})$ can be realized explicitly as the tensor product of topological coalgebras $U_{\hbar}\mathfrak{b}^{\vee} \otimes U_{\hbar}\mathfrak{b}$, with multiplication determined from the cross relations

$$(8.1) \quad \iota(x)\check{i}(f) = f_1(x_1)f_3(S^{-1}(x_3))f_2 \otimes x_2 \quad \forall x \in U_{\hbar}\mathfrak{b}', f \in U_{\hbar}\mathfrak{b}^{\vee},$$

where we have used the sumless Sweedler notation for iterated coproducts on $U_{\hbar}\mathfrak{b}$ and $U_{\hbar}\mathfrak{b}^{\vee}$, and ι and \check{i} are now given by $\iota(x) = 1 \otimes x$ and $\check{i}(f) = f \otimes 1$ for all $x \in U_{\hbar}\mathfrak{b}$ and $f \in U_{\hbar}\mathfrak{b}^{\vee}$. As spelled out in detail in [1, Section A.5], this can be realized as a special instance of the *double cross-product* construction. Namely, one has

$$D(U_{\hbar}\mathfrak{b}) = U_{\hbar}\mathfrak{b}^{\vee} \bowtie U_{\hbar}\mathfrak{b},$$

with respect to the left coadjoint action \triangleright of $U_{\hbar}\mathfrak{b}$ on $U_{\hbar}\mathfrak{b}^{\vee}$ and the right coadjoint action \triangleleft on $U_{\hbar}\mathfrak{b}^{\vee}$ on $U_{\hbar}\mathfrak{b}$. Given the $U_{\hbar}\mathfrak{b}$ analogue of Lemma 5.8, established in [1, Proposition A.5], this construction of $D(U_{\hbar}\mathfrak{b})$ proceeds identically to the analogous construction for the quantum double of a finite-dimensional Hopf algebra; we refer the reader to the texts [36, Section 7], [26, Section IX.4], [33, Section 8.2] and [37, Section 10.3], for instance, as well as the articles [34, 35, 39].

8.2 The restricted quantum double of the Yangian

In our case, $U_{\hbar}\mathfrak{b} = Y_{\hbar}\mathfrak{g}$ is not a quantization of a finite-dimensional Lie bialgebra, but rather a homogeneous quantization of an \mathbb{N} -graded Lie bialgebra \mathfrak{b} with finite-dimensional graded components \mathfrak{b}_k . In this setting, the double cross-product construction alluded to above remains valid provided all duals are taken in the category of \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -modules; that is, we replace $U_{\hbar}\mathfrak{b}^{\vee}$ with $Y_{\hbar}\mathfrak{g}^{\vee}$. This produces the *restricted quantum double* $D(U_{\hbar}\mathfrak{b})$ of $U_{\hbar}\mathfrak{b}$.

Let us now give the detailed construction of this topological Hopf algebra. Following Section 5.4, let \blacktriangledown and \blacktriangle denote the left adjoint action of $Y_{\hbar}\mathfrak{g}$ on itself, and the right adjoint coaction of $Y_{\hbar}\mathfrak{g}$ on itself, respectively:

$$\begin{aligned} \blacktriangledown &= m^3 \circ (\mathbf{1}^{\otimes 2} \otimes S) \circ (23) \circ (\Delta \otimes \mathbf{1}), \\ \blacktriangle &= (\mathbf{1} \otimes m) \circ (12) \circ (S \otimes \mathbf{1}^{\otimes 2}) \circ \Delta^3, \end{aligned}$$

where all tensor products are now taken to be the topological tensor product over $\mathbb{C}[[\hbar]]$. By Lemma 5.8, we have

$$\blacktriangledown(Y_{\hbar}\mathfrak{g} \otimes \check{Y}_{\hbar}\mathfrak{g}) \subset \check{Y}_{\hbar}\mathfrak{g} \quad \text{and} \quad \blacktriangle(Y_{\hbar}\mathfrak{g}) \subset Y_{\hbar}\mathfrak{g} \otimes \check{Y}_{\hbar}\mathfrak{g}.$$

We may thus dualize \blacktriangledown and \blacktriangle to obtain the so-called left and right coadjoint actions

$$\triangleright: Y_{\hbar}\mathfrak{g} \otimes Y_{\hbar}\mathfrak{g}^{\vee} \rightarrow Y_{\hbar}\mathfrak{g}^{\vee} \quad \text{and} \quad \triangleleft: Y_{\hbar}\mathfrak{g} \otimes Y_{\hbar}\mathfrak{g}^{\vee} \rightarrow Y_{\hbar}\mathfrak{g},$$

respectively, of $Y_{\hbar}\mathfrak{g}$ on $Y_{\hbar}\mathfrak{g}^{\vee}$ and of $Y_{\hbar}\mathfrak{g}^{\vee}$ on $Y_{\hbar}\mathfrak{g}$. These are defined on simple tensors by the formulas

$$x \triangleleft f = \mathbf{1} \otimes f \circ (\mathbf{1} \otimes S^{-1}) \circ \blacktriangle(x) \quad \text{and} \quad (x \triangleright f)(y) = f(S^{-1}(x) \cdot y)$$

for all $x \in Y_{\hbar}\mathfrak{g}$, $f \in Y_{\hbar}\mathfrak{g}^{\vee}$, and $y \in \check{Y}_{\hbar}\mathfrak{g}$, where the action of $S^{-1}(x)$ on y is given by \blacktriangledown and we have written $x \triangleright f$ for $\triangleright(x \otimes f)$ and $x \triangleleft f$ for $\triangleleft(x \otimes f)$.

The tuple $(Y_{\hbar}\mathfrak{g}, Y_{\hbar}\mathfrak{g}^{\vee}, \triangleright, \triangleleft)$ forms a *matched pair* of \mathbb{Z} -graded topological Hopf algebras over $\mathbb{C}[[\hbar]]$. Explicitly, \triangleright and \triangleleft are \mathbb{Z} -graded homomorphisms of topological $\mathbb{C}[[\hbar]]$ -modules which satisfy the following set of conditions:

(M1) $(Y_{\hbar}\mathfrak{g}^{\vee}, \triangleright)$ is a left $Y_{\hbar}\mathfrak{g}$ -module coalgebra and $(Y_{\hbar}\mathfrak{g}, \triangleleft)$ is a right $Y_{\hbar}\mathfrak{g}^{\vee}$ -module coalgebra. Equivalently, \triangleright and \triangleleft are morphisms of topological coalgebras:

$$\Delta^{\vee} \circ \triangleright = \triangleright \otimes \triangleright \circ (23) \circ \Delta \otimes \Delta^{\vee}, \quad \Delta \circ \triangleleft = \triangleleft \otimes \triangleleft \circ (23) \circ \Delta \otimes \Delta^{\vee},$$

$$\varepsilon \otimes \varepsilon^{\vee} = \varepsilon^{\vee} \circ \triangleright, \quad \varepsilon \otimes \varepsilon^{\vee} = \varepsilon \circ \triangleleft,$$

which make $Y_{\hbar}\mathfrak{g}^{\vee}$ a left $Y_{\hbar}\mathfrak{g}$ -module and $Y_{\hbar}\mathfrak{g}$ a right $Y_{\hbar}\mathfrak{g}^{\vee}$ -module.

(M2) \triangleleft and \triangleright are compatible with the products m and m^{\vee} on $Y_{\hbar}\mathfrak{g}$ and $Y_{\hbar}\mathfrak{g}^{\vee}$:

$$\triangleleft \circ (m \otimes 1) = m \circ (\triangleleft \otimes 1) \circ (1 \otimes \triangleright \otimes \triangleleft) \circ (34) \circ 1 \otimes \Delta \otimes \Delta^{\vee},$$

$$\triangleright \circ (1 \otimes m^{\vee}) = m^{\vee} \circ (1 \otimes \triangleright) \circ (\triangleright \otimes \triangleleft \otimes 1) \circ (23) \circ \Delta \otimes \Delta^{\vee} \otimes 1.$$

(M3) The unit maps ι and ι^{\vee} are module homomorphisms:

$$\triangleleft \circ (\iota \otimes 1) = \iota \otimes \varepsilon^{\vee} \quad \text{and} \quad \triangleright \circ (1 \otimes \iota^{\vee}) = \varepsilon \otimes \iota^{\vee}.$$

(M4) \triangleleft and \triangleright satisfy the compatibility relation

$$(\triangleleft \otimes \triangleright) \circ (23) \circ \Delta \otimes \Delta^{\vee} = (12) \circ (\triangleright \otimes \triangleleft) \circ (23) \circ \Delta \otimes \Delta^{\vee}.$$

Here, we note that the above conditions coincide with those from [26, Definition IX.2.2], [39, Section 2], and [1, Section A.1], which agree with those from [36, Definition 7.2.1] up to conventions on the order in which the tensor factors appear.

Since $Y_{\hbar}\mathfrak{g}^{\vee}$ and $Y_{\hbar}\mathfrak{g}$ are matched, we may form the double cross-product Hopf algebra $Y_{\hbar}\mathfrak{g}^{\vee} \bowtie Y_{\hbar}\mathfrak{g}$ in the category of \mathbb{Z} -graded topological Hopf algebras over $\mathbb{C}[[\hbar]]$ by following the standard procedure (see [26, Theorem IX.2.3] or [36, Theorem 7.2.2], for instance). As a \mathbb{Z} -graded topological coalgebra, $Y_{\hbar}\mathfrak{g}^{\vee} \bowtie Y_{\hbar}\mathfrak{g}$ coincides with the tensor product of $Y_{\hbar}\mathfrak{g}^{\vee}$ and $Y_{\hbar}\mathfrak{g}$:

$$Y_{\hbar}\mathfrak{g}^{\vee} \bowtie Y_{\hbar}\mathfrak{g} = Y_{\hbar}\mathfrak{g}^{\vee} \otimes Y_{\hbar}\mathfrak{g},$$

$$\Delta_D = (23) \circ \Delta^{\vee} \otimes \Delta \quad \text{and} \quad \varepsilon_D = \varepsilon^{\vee} \otimes \varepsilon,$$

where Δ_D denotes the coproduct and ε_D denotes the counit. The algebra structure on $Y_{\hbar}\mathfrak{g}^{\vee} \bowtie Y_{\hbar}\mathfrak{g}$ is uniquely determined by the requirement that the inclusions

$$\check{i}_D : Y_{\hbar}\mathfrak{g}^{\vee} \rightarrow Y_{\hbar}\mathfrak{g}^{\vee} \otimes Y_{\hbar}\mathfrak{g} \quad \text{and} \quad \iota_D : Y_{\hbar}\mathfrak{g} \rightarrow Y_{\hbar}\mathfrak{g}^{\vee} \otimes Y_{\hbar}\mathfrak{g}$$

are $\mathbb{C}[[\hbar]]$ -algebra homomorphisms satisfying the relations

$$(8.2) \quad m_D \circ (\check{i}_D \otimes \iota_D) = 1_{Y_{\hbar}\mathfrak{g}^{\vee}} \otimes 1_{Y_{\hbar}\mathfrak{g}},$$

$$m_D \circ (\iota_D \otimes \check{i}_D) = (\triangleright \otimes \triangleleft) \circ (23) \circ \Delta \otimes \Delta^{\vee},$$

where m_D denotes the multiplication on $D(Y_{\hbar}\mathfrak{g})$. In particular, the unit map is $\iota^{\vee} \otimes \iota$. Finally, the antipode S_D on $Y_{\hbar}\mathfrak{g}^{\vee} \bowtie Y_{\hbar}\mathfrak{g}$ is given by

$$S_D = m_D \circ (\iota_D \otimes \check{i}_D) \circ (S \otimes S^{\vee}) \circ (12).$$

This double cross-product Hopf algebra is the restricted quantum double of the Yangian, as we formally record in the below definition.

Definition 8.1 The \mathbb{Z} -graded topological Hopf algebra $Y_{\hbar}\mathfrak{g}^\vee \bowtie Y_{\hbar}\mathfrak{g}$ is called the *restricted quantum double* of the Yangian $Y_{\hbar}\mathfrak{g}$, and is denoted

$$D(Y_{\hbar}\mathfrak{g}) := Y_{\hbar}\mathfrak{g}^\vee \bowtie Y_{\hbar}\mathfrak{g}.$$

This definition, together with the general theory, implies that $D(Y_{\hbar}\mathfrak{g})$ provides a homogeneous quantization of the restricted Drinfeld double $\mathfrak{t} \cong D(\mathfrak{t}_+)$ of the graded Lie bialgebra $(\mathfrak{t}_+, \delta_+)$, realized on the space $\mathfrak{t}_+^* \oplus \mathfrak{t}_+ \cong \mathfrak{t}_- \oplus \mathfrak{t}_+.$ ²

One stipulation to carrying out the quantum double construction in the restricted homogeneous setting is that the canonical element R associated with the pairing between $\dot{Y}_{\hbar}\mathfrak{g}$ and $\dot{Y}_{\hbar}\mathfrak{g}^*$, although formally satisfying the relations of (3) in Section 8.1, does not converge in the \hbar -adically complete tensor product $\dot{Y}_{\hbar}\mathfrak{g} \otimes \dot{Y}_{\hbar}\mathfrak{g}^* \subset D(Y_{\hbar}\mathfrak{g}) \otimes D(Y_{\hbar}\mathfrak{g})$, and so $DY_{\hbar}\mathfrak{g}$ is only *topologically* quasitriangular. Here, the prefix “topologically” is a bit subtle: it does not refer to the \hbar -adic topology and must be handled with care. In Section 9, we shall identify R with the universal R -matrix $\mathcal{R}(w - z) \in Y_{\hbar}(\mathfrak{g})^{\otimes 2}[[w][[z^{-1}]]$. This viewpoint allows for a precise interpretation of the relations of (3) in terms of those from Theorem 3.8.

Remark 8.2 There are many competing, though equivalent, variants of the definition of the quantum double. For instance, it may be realized on the tensor product $Y_{\hbar}\mathfrak{g} \otimes Y_{\hbar}\mathfrak{g}^\vee$, as in [7, Section 13]. Here, we follow the conventions from [26, 39].

8.3 $DY_{\hbar}\mathfrak{g}$ as a quantum double

We now turn to proving the \hbar -adic variant of the main conjecture from [32, Section 2], which postulates that $DY_{\hbar}\mathfrak{g}$ and $D(Y_{\hbar}\mathfrak{g})$ are one and the same (see Theorem 8.4 below).

Recall from the proof of Corollary 7.3 that $L(\widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)}) \subset (Y_{\hbar}\mathfrak{g} \otimes Y_{\hbar}\mathfrak{g})[[z^{\pm 1}]]$ denotes the \mathbb{Z} -graded topological $\mathbb{C}[[\hbar]]$ -algebra obtained by completing the graded subalgebra $L(\widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)})$ of $Y_{\hbar}(\mathfrak{g})^{\otimes 2}[[z, z^{-1}]]$ defined in (3.16). Given this notation, we have $\mathbb{C}[[\hbar]]$ -linear maps

$$\begin{aligned} \pi_z &: L\widehat{Y_{\hbar}\mathfrak{g}}_z \otimes Y_{\hbar}\mathfrak{g} \rightarrow L\widehat{Y_{\hbar}\mathfrak{g}}_z, & \pi_z &:= m_z \circ (1 \otimes \tau_z), \\ \pi_z^* &: L\widehat{Y_{\hbar}\mathfrak{g}}_z \rightarrow L(\widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)}), & \pi_z^* &(y) = (1 \otimes y)\mathcal{R}(-z), \end{aligned}$$

where m_z denotes the multiplication in $L\widehat{Y_{\hbar}\mathfrak{g}}_z$. These obey a Drinfeld–Yetter right action/left coaction compatibility condition, as the next lemma makes explicit.

Lemma 8.3 *The pair (π_z, π_z^*) satisfy the relation*

$$\pi_z^* \circ \pi_z \circ (12) = m \otimes \pi_z \circ (432) \circ \blacktriangle \otimes \blacktriangledown \otimes I \circ \Delta^{\text{op}} \otimes \pi_z^*$$

in $\text{Hom}_{\mathbb{C}[[\hbar]]}(Y_{\hbar}\mathfrak{g} \otimes L\widehat{Y_{\hbar}\mathfrak{g}}_z, L(\widehat{Y_{\hbar}\mathfrak{g}}_z^{(2)}))$, where we have set

$$\blacktriangle := (S^{-1} \otimes I) \circ (12) \circ \blacktriangle \quad \text{and} \quad \blacktriangledown := \blacktriangledown \circ (S^{-1} \otimes I).$$

²This also follows immediately from Theorem 8.4.

Proof This follows from a straightforward modification of the proof of (9) in [39, Lemma 1], using the antipode relations

$$m \circ (1 \otimes S^{-1}) \circ \Delta^{\text{op}} = \iota \circ \varepsilon = m \circ (S^{-1} \otimes 1) \circ \Delta^{\text{op}},$$

the counit relation $m \circ (1 \otimes \varepsilon) \circ \Delta = 1$, and the intertwining relation

$$(8.3) \quad 1 \otimes \tau_z \circ \Delta(x) = \mathcal{R}(-z)^{-1} \cdot 1 \otimes \tau_z \circ \Delta^{\text{op}}(x) \cdot \mathcal{R}(-z) \quad \forall x \in Y_{\hbar}\mathfrak{g},$$

which itself follows from the identities (1) and (3.20) of Theorem 3.8.

To illustrate this, we apply both sides of the claimed identity to a simple tensor $x \otimes y \in Y_{\hbar}\mathfrak{g} \otimes L\widehat{Y_{\hbar}\mathfrak{g}_z}$, while exploiting the sumless Sweedler notation $\Delta^n(x) = x_1 \otimes \dots \otimes x_n$ for iterated coproducts. Expanding the right-hand side of the resulting expression, while using that $\blacktriangledown(x \otimes y) = S^{-1}(x_2)yx_1$ and $\blacktriangle(x) = S^{-1}(x_3)x_1 \otimes x_2$, we obtain

$$\begin{aligned} S^{-1}(x_5)x_3S^{-1}(x_2) \otimes y \cdot \mathcal{R}(-z) \cdot x_1 \otimes \tau_z(x_4) &= S^{-1}(x_4) \otimes y \cdot \mathcal{R}(-z) \cdot x_1\varepsilon(x_2) \otimes \tau_z(x_3) \\ &= S^{-1}(x_3) \otimes y \cdot \mathcal{R}(-z) \cdot x_1 \otimes \tau_z(x_2) \\ &= S^{-1}(x_3)x_2 \otimes y\tau_z(x_1) \cdot \mathcal{R}(-z) \\ &= 1 \otimes y\tau_z(x_1\varepsilon(x_2)) \cdot \mathcal{R}(-z) \\ &= 1 \otimes y\tau_z(x) \cdot \mathcal{R}(-z), \end{aligned}$$

where we have applied the intertwining relation (8.3) in the third equality, and the appropriate Hopf relations listed above in each of the remaining equalities. Since

$$\pi_z^* \circ \pi_z(y \otimes x) = \pi_z^*(y\tau_z(x)) = (1 \otimes y\tau_z(x))\mathcal{R}(-z),$$

the above computation implies the lemma. ■

We now come to the main theorem of this section. Recall from Definition 8.1 that we may identify the restricted quantum double $D(Y_{\hbar}\mathfrak{g})$ with $Y_{\hbar}\mathfrak{g}^{\vee} \otimes Y_{\hbar}\mathfrak{g}$ as a topological coalgebra over $\mathbb{C}[[\hbar]]$. We further recall that

$$\iota : Y_{\hbar}\mathfrak{g} \hookrightarrow DY_{\hbar}\mathfrak{g} \quad \text{and} \quad \check{\iota} : Y_{\hbar}\mathfrak{g}^{\vee} \hookrightarrow DY_{\hbar}\mathfrak{g}$$

are the \mathbb{Z} -graded embeddings of topological Hopf algebras over $\mathbb{C}[[\hbar]]$ which featured prominently in Theorem 7.5.

Theorem 8.4 *The $\mathbb{C}[[\hbar]]$ -module homomorphism*

$$Y := m \circ (\check{\iota} \otimes \iota) : Y_{\hbar}\mathfrak{g}^{\vee} \otimes Y_{\hbar}\mathfrak{g} \rightarrow DY_{\hbar}\mathfrak{g}$$

is an isomorphism of \mathbb{Z} -graded topological Hopf algebras $D(Y_{\hbar}\mathfrak{g}) \xrightarrow{\sim} DY_{\hbar}\mathfrak{g}$.

Proof We shall first prove that Y is an isomorphism of \mathbb{Z} -graded topological coalgebras over $\mathbb{C}[[\hbar]]$, which follows from Theorem 7.5. Afterward, we will use Lemma 8.3 to complete the proof of the theorem by proving that Y is an algebra homomorphism $D(Y_{\hbar}\mathfrak{g}) \rightarrow DY_{\hbar}\mathfrak{g}$. Here, we note that it follows automatically that Y intertwines the underlying antipodes, though this is straightforward to verify. ■

Proof that Y is an isomorphism of \mathbb{Z} -graded coalgebras Since m , ι , and $\check{\iota}$ are all \mathbb{Z} -graded $\mathbb{C}[[\hbar]]$ -module homomorphisms, the same is true of Y . Moreover, as the

coproduct $\dot{\Delta}$ of $DY_{\hbar}\mathfrak{g}$ is an algebra homomorphism, it satisfies

$$\begin{aligned} \dot{\Delta} \circ Y &= \dot{\Delta} \circ m \circ (\check{i} \otimes \iota) \\ &= m \otimes m \circ (23) \circ (\dot{\Delta} \otimes \dot{\Delta}) \circ (\check{i} \otimes \iota) \\ &= m \otimes m \circ (23) \circ (\check{i} \otimes \check{i} \otimes \iota \otimes \iota) \circ (\Delta^\vee \otimes \Delta) \\ &= Y \otimes Y \circ (23) \circ \Delta^\vee \otimes \Delta. \end{aligned}$$

Similarly, as the counit $\dot{\varepsilon}$ of $DY_{\hbar}\mathfrak{g}$ is an algebra homomorphism, one has

$$\dot{\varepsilon} \circ Y = \dot{\varepsilon} \circ m \circ (\check{i} \otimes \iota) = m \circ (\dot{\varepsilon} \otimes \dot{\varepsilon}) \circ (\check{i} \otimes \iota) = m \circ \varepsilon^\vee \otimes \varepsilon.$$

This proves that Y is a \mathbb{Z} -graded homomorphism of topological coalgebras. Now let us turn to establishing the bijectivity of Y . As $Y_{\hbar}\mathfrak{g}^\vee \otimes Y_{\hbar}\mathfrak{g}$ is a topologically free $\mathbb{C}[[\hbar]]$ -module with semiclassical limit $U(\mathfrak{t}_-) \otimes_{\mathbb{C}} U(\mathfrak{t}_+)$ and $DY_{\hbar}\mathfrak{g}$ is a flat deformation of $U(\mathfrak{t})$, Lemma 2.1 asserts that it is sufficient to establish that the semiclassical limit

$$\check{Y} = \check{m} \circ (\check{i} \otimes \check{\iota}) : U(\mathfrak{t}_-) \otimes_{\mathbb{C}} U(\mathfrak{t}_+) \rightarrow U(\mathfrak{t})$$

of Y is an isomorphism, where \check{m} , \check{i} , and $\check{\iota}$ are the semiclassical limits of m , i , and ι , respectively. As \check{i} and $\check{\iota}$ quantize the natural inclusions of \mathfrak{t}_- and \mathfrak{t}_+ into \mathfrak{t} (as was shown in the proof of Lemma 7.4 for \check{y}) and \check{m} is the multiplication map on $U(\mathfrak{t})$, this follows from the Poincaré–Birkhoff–Witt theorem for enveloping algebras and the decomposition $\mathfrak{t} = \mathfrak{t}_- \oplus \mathfrak{t}_+$. ■

Proof that Y is an algebra homomorphism By (8.2), we must show that $Y \circ \iota_D$ and $Y \circ \check{i}_D$ are $\mathbb{C}[[\hbar]]$ -algebra homomorphisms and that, in addition, Y satisfies the relations

$$(8.4) \quad \begin{aligned} Y \circ m_D \circ (\check{i}_D \otimes \iota_D) &= m \circ (Y \otimes Y) \circ (\check{i}_D \otimes \iota_D), \\ m \circ (Y \otimes Y) \circ (\iota_D \otimes \check{i}_D) &= m \circ (Y \otimes Y) \circ (\triangleright \otimes \triangleleft) \circ (23) \circ \Delta \otimes \Delta^\vee. \end{aligned}$$

By definition of Y , we have $Y \circ \iota_D = \iota$ and $Y \circ \check{i}_D = \check{i}$. Moreover, both sides of the identity

$$Y \circ m_D \circ (\check{i}_D \otimes \iota_D) = m \circ (Y \otimes Y) \circ (\check{i}_D \otimes \iota_D)$$

coincide with Y , viewed as a map $Y_{\hbar}\mathfrak{g}^\vee \otimes Y_{\hbar}\mathfrak{g} \rightarrow DY_{\hbar}\mathfrak{g}$. Hence, we are left to verify the second relation of (8.4), which we shall establish by appealing to the injective $\mathbb{C}[[\hbar]]$ -algebra homomorphism Φ_z from Theorem 4.6. Namely, it is enough to show that

$$\Phi_z \circ m \circ (Y \otimes Y) \circ (\iota_D \otimes \check{i}_D) = \Phi_z \circ m \circ (Y \otimes Y) \circ (\triangleright \otimes \triangleleft) \circ (23) \circ \Delta \otimes \Delta^\vee.$$

Since Φ_z is an algebra homomorphism satisfying $\Phi_z \circ \iota = \tau_z$ and $\Phi_z \circ \check{i} = \Phi_z^\vee$ (see Lemma 7.4), this is equivalent to

$$(8.5) \quad m_z \circ (\tau_z \otimes \Phi_z^\vee) = m_z \circ (\Phi_z^\vee \otimes \tau_z) \circ (\triangleright \otimes \triangleleft) \circ (23) \circ \Delta \otimes \Delta^\vee,$$

where m_z is the product in $L\widehat{Y_{\hbar}\mathfrak{g}_z}$. The proof that (8.5) is satisfied follows an argument parallel to that employed to establish Part (a) of Theorem 2 in [39], using Lemma 8.3 in place of [39, (9)]. For the sake of completeness, we give a complete argument below.

Applying the left-hand side of (8.5) to an arbitrary simple tensor $x \otimes f$, we obtain

$$\tau_z(x)\Phi_z^\vee(f) = (f \otimes \tau_z(x))\mathcal{R}(-z) = (f \otimes \mathbf{1}) \circ \pi_z^* \circ \pi_z(1 \otimes x),$$

which, by Lemma 8.3, may be rewritten as

$$(8.6) \quad \tau_z(x)\Phi_z^\vee(f) = f \otimes 1 \circ m \otimes \pi_z \circ (432) \circ \blacktriangle \otimes \blacktriangledown \otimes 1 \circ \Delta^{\text{op}}(x) \otimes \mathcal{R}(-z).$$

On the other hand, by definition of Φ_z^\vee , \triangleright , and \triangleleft , the right-hand side of (8.5) evaluated on $x \otimes f$ is

$$\Phi_z^\vee(x_1 \triangleright f_1) \cdot \tau_z(x_2 \triangleleft f_2) = (f_1 \otimes 1 \circ \blacktriangledown \otimes 1(x_1 \otimes \mathcal{R}(-z))) \cdot (f_2 \otimes \tau_z \circ \blacktriangle(x_2)),$$

where we have employed the sumless Sweedler notation $\Delta(x) = x_1 \otimes x_2$ and $\Delta^\vee(f) = f_1 \otimes f_2$. That this coincides with the expression (8.6) for $\tau_z(x)\Phi_z^\vee(f)$ is a consequence of the following general computation. For each $a \in \dot{Y}_{\hbar\mathfrak{g}}$ and $b \in Y_{\hbar\mathfrak{g}}$, we have

$$\begin{aligned} (f_1(x_1 \bullet a) \otimes b) \cdot (f_2 \otimes \tau_z \circ \blacktriangle(x_2)) &= f(x_2^{(1)}(x_1 \bullet a))b\tau_z(x_2^{(2)}) \\ &= f \otimes 1 \circ m \otimes \pi_z(x_2^{(1)} \otimes (x_1 \bullet a) \otimes b \otimes x_2^{(2)}) \\ &= (f \otimes 1 \circ m \otimes \pi_z \circ (432) \circ \blacktriangle \otimes \blacktriangledown \otimes 1 \circ \Delta^{\text{op}} \otimes 1 \otimes 1)(x \otimes a \otimes b), \end{aligned}$$

where we have set $x \bullet a = \blacktriangledown(x \otimes a)$ and in the first and second equalities we have used the (sumless) Sweedler-type notation $\blacktriangle(x) = x^{(1)} \otimes x^{(2)}$. This completes the proof of (8.5), and thus the proof that Υ is an algebra homomorphism. ■

9 The universal R-matrix

In this final section, we establish the last assertion of Theorem I, which identifies the universal R-matrix \mathcal{R} of the Yangian double $DY_{\hbar\mathfrak{g}} \cong D(Y_{\hbar\mathfrak{g}})$ with Drinfeld’s universal R-matrix $\mathcal{R}(z)$ (see Theorem 9.6). Though this in fact follows without too much effort from the results of Sections 7 and 8, constructing \mathcal{R} precisely does require some care. After laying the groundwork in Sections 9.1 and 9.2, we define \mathcal{R} and prove our last main result in Section 9.3. The final two subsections – Sections 9.4 and 9.5 – are devoted to providing additional context pertinent to this theorem.

9.1 The isomorphism Θ

Let θ denote the canonical $\mathbb{C}[[\hbar]]$ -module injection

$$\theta : Y_{\hbar\mathfrak{g}} \otimes \dot{Y}_{\hbar\mathfrak{g}}^* \hookrightarrow \text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar\mathfrak{g}}, Y_{\hbar\mathfrak{g}})$$

determined on simple tensors by $\theta(x \otimes f)(y) = f(y)x$. This injection is a homomorphism of topological $\mathbb{C}[[\hbar]]$ -algebras provided we equip $\text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar\mathfrak{g}}, Y_{\hbar\mathfrak{g}})$ with the convolution product

$$\varphi_1 \star \varphi_2 := m \circ (\varphi_1 \otimes \varphi_2) \circ \Delta|_{\dot{Y}_{\hbar\mathfrak{g}}} \quad \forall \varphi_1, \varphi_2 \in \text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar\mathfrak{g}}, Y_{\hbar\mathfrak{g}})$$

and identity element $\iota \circ \varepsilon$, where m , Δ , ι , and ε are the multiplication, coproduct, unit, and counit of $Y_{\hbar\mathfrak{g}}$, respectively. Our main goal in this subsection is to identify a natural extension of θ which is an isomorphism.

Let us begin by noting some topological properties of the homomorphism space $\text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar\mathfrak{g}}, Y_{\hbar\mathfrak{g}})$. As $Y_{\hbar\mathfrak{g}}$ and $\dot{Y}_{\hbar\mathfrak{g}}$ are topologically free with semiclassical

limits $U(t_+)$ and $S(\hbar t_+)$, respectively, $\text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar\mathfrak{g}}, Y_{\hbar\mathfrak{g}})$ is a topologically free $\mathbb{C}[[\hbar]]$ -modules with

$$(9.1) \quad \text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar\mathfrak{g}}, Y_{\hbar\mathfrak{g}}) \cong (\text{Hom}_{\mathbb{C}}(S(\hbar t_+), U(t_+)))[[\hbar]].$$

In addition, $\text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar\mathfrak{g}}, Y_{\hbar\mathfrak{g}})$ is a Hausdorff and complete topological space with respect to the topology associated with the descending filtration \mathbb{E}_\bullet of closed $\mathbb{C}[[\hbar]]$ -submodules defined by

$$(9.2) \quad \mathbb{E}_n := \{f \in \text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar\mathfrak{g}}, Y_{\hbar\mathfrak{g}}) : f(J_n) = 0\},$$

where we have set $J_n = \bigoplus_{k < n} \dot{Y}_{\hbar\mathfrak{g}}(g)_k$ with $J_0 = \{0\}$. Said in more algebraic terms, the natural $\mathbb{C}[[\hbar]]$ -module homomorphism

$$\text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar\mathfrak{g}}, Y_{\hbar\mathfrak{g}}) \rightarrow \varprojlim_n (\text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar\mathfrak{g}}, Y_{\hbar\mathfrak{g}})/\mathbb{E}_n)$$

is an isomorphism, as is readily verified. Moreover, as the coproduct Δ on $Y_{\hbar\mathfrak{g}}$ is graded, \mathbb{E}_\bullet is a descending filtration of ideals and the above is an isomorphism of $\mathbb{C}[[\hbar]]$ -algebras.

We now turn toward enlarging the domain of θ . For each $n \in \mathbb{Z}$, let $\dot{Y}_{\hbar\mathfrak{g}}^{*(n)}$ denote the closure of the $\mathbb{C}[[\hbar]]$ -submodule of $\dot{Y}_{\hbar\mathfrak{g}}^*$ generated by $\bigoplus_{k \geq n} \dot{Y}_{\hbar\mathfrak{g}}^*_{-k}$. Since $\dot{Y}_{\hbar\mathfrak{g}}^*_k \subset \hbar \dot{Y}_{\hbar\mathfrak{g}}^*_{k-1}$ for $k > 0$, $\dot{Y}_{\hbar\mathfrak{g}}^*_{(0)} = \dot{Y}_{\hbar\mathfrak{g}}^*$ and we have a descending \mathbb{N} -filtration

$$\dot{Y}_{\hbar\mathfrak{g}}^* = \dot{Y}_{\hbar\mathfrak{g}}^*_{(0)} \supset \dot{Y}_{\hbar\mathfrak{g}}^*_{(1)} \supset \dots \supset \dot{Y}_{\hbar\mathfrak{g}}^*_{(n)} \supset \dots$$

We may thus introduce the topological tensor product

$$Y_{\hbar\mathfrak{g}} \tilde{\otimes} \dot{Y}_{\hbar\mathfrak{g}}^* := \varprojlim_n (Y_{\hbar\mathfrak{g}} \otimes \dot{Y}_{\hbar\mathfrak{g}}^* / Y_{\hbar\mathfrak{g}} \otimes \dot{Y}_{\hbar\mathfrak{g}}^*_{(n)}).$$

Since $\dot{Y}_{\hbar\mathfrak{g}}^*$ is topologically generated as a $\mathbb{C}[[\hbar]]$ -algebra by the space $\bigoplus_{k > 0} \dot{Y}_{\hbar\mathfrak{g}}^*_{-k}$ and, for each $n \in \mathbb{N}$, $\bigoplus_{k \geq n} \dot{Y}_{\hbar\mathfrak{g}}^*_{-k}$ is stable under multiplication by any x in this space, we see that each $\dot{Y}_{\hbar\mathfrak{g}}^*_{(n)}$ is an ideal in $\dot{Y}_{\hbar\mathfrak{g}}^*$. It follows that $Y_{\hbar\mathfrak{g}} \tilde{\otimes} \dot{Y}_{\hbar\mathfrak{g}}^*$ is a $\mathbb{C}[[\hbar]]$ -algebra.

To see that $Y_{\hbar\mathfrak{g}} \tilde{\otimes} \dot{Y}_{\hbar\mathfrak{g}}^*$ is a topologically free $\mathbb{C}[[\hbar]]$ -module, let us introduce the classical spaces $S(\hbar t_+)^*_{(n)}$ and $U(t_+) \tilde{\otimes} S(\hbar t_+)^*$ by setting

$$S(\hbar t_+)^*_{(n)} := \bigoplus_{k \geq n} S(\hbar t_+)^*_{-k},$$

$$U(t_+) \tilde{\otimes} S(\hbar t_+)^* := \prod_{n \in \mathbb{N}} U(t_+) \otimes_{\mathbb{C}} S(\hbar t_+)^*_{-n} \cong \varprojlim_n U(t_+) \otimes_{\mathbb{C}} (S(\hbar t_+)^* / S(\hbar t_+)^*_{(n)}).$$

Lemma 9.1 *The $\mathbb{C}[[\hbar]]$ -algebra $Y_{\hbar\mathfrak{g}} \tilde{\otimes} \dot{Y}_{\hbar\mathfrak{g}}^*$ is topologically free with*

$$Y_{\hbar\mathfrak{g}} \tilde{\otimes} \dot{Y}_{\hbar\mathfrak{g}}^* / \hbar(Y_{\hbar\mathfrak{g}} \tilde{\otimes} \dot{Y}_{\hbar\mathfrak{g}}^*) \cong U(t_+) \tilde{\otimes} S(\hbar t_+)^*.$$

Proof Let $v_* : \dot{Y}_{\hbar\mathfrak{g}}^* \xrightarrow{\sim} S(\hbar t_+)^*[[\hbar]]$ be any fixed \mathbb{Z} -graded isomorphism of topological $\mathbb{C}[[\hbar]]$ -modules. We claim that

$$(9.3) \quad v_*(\dot{Y}_{\hbar\mathfrak{g}}^*_{(n)}) = S(\hbar t_+)^*_{(n)}[[\hbar]].$$

To see this, fix $n \in \mathbb{N}$. Since v_* is graded, we have

$$v_*(\dot{Y}_{\hbar\mathfrak{g}}^*_{-k}) = \prod_{a \in \mathbb{N}} S(\hbar t_+)^*_{-k-a} \hbar^a \subset S(\hbar t_+)^*_{(n)}[[\hbar]] \quad \forall k \geq n.$$

As $S(\hbar t_+)^*_{(n)}[[\hbar]]$ is a closed $\mathbb{C}[[\hbar]]$ -submodule of $S(\hbar t_+)^*[[\hbar]]$, we get $v_*(\dot{Y}_{\hbar\mathfrak{g}}^*_{(n)}) \subset S(\hbar t_+)^*_{(n)}[[\hbar]]$. Conversely, if $\sum_{a \geq 0} x_a \hbar^a \in S(\hbar t_+)^*_{(n)}[[\hbar]]$, then $y_a := v_*^{-1}(x_a) \in \bigoplus_{k \geq n} \dot{Y}_{\hbar\mathfrak{g}}^*_{-k}$ for each $a \in \mathbb{N}$. Therefore, we have

$$v_* \left(\sum_{a \geq 0} x_a \hbar^a \right) = \sum_{a \geq 0} y_a \hbar^a \in \dot{Y}_{\hbar\mathfrak{g}}^*_{(n)},$$

which completes the proof of (9.3). The statement of the lemma now follows from the sequence of isomorphisms

$$\begin{aligned} Y_{\hbar\mathfrak{g}} \otimes \dot{Y}_{\hbar\mathfrak{g}}^* / Y_{\hbar\mathfrak{g}} \otimes \dot{Y}_{\hbar\mathfrak{g}}^*_{(n)} &\cong (U(t_+) \otimes_{\mathbb{C}} S(\hbar t_+)^*)[[\hbar]] / (U(t_+) \otimes_{\mathbb{C}} S(\hbar t_+)^*_{(n)})[[\hbar]] \\ &\cong (U(t_+) \otimes_{\mathbb{C}} (S(\hbar t_+)^* / S(\hbar t_+)^*_{(n)}))[[\hbar]]. \quad \blacksquare \end{aligned}$$

The next result outputs the desired extensions of θ .

Proposition 9.2 *The injection θ extends to an isomorphism of $\mathbb{C}[[\hbar]]$ -algebras*

$$\Theta : Y_{\hbar\mathfrak{g}} \widetilde{\otimes} \dot{Y}_{\hbar\mathfrak{g}}^* \xrightarrow{\sim} \text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar\mathfrak{g}}, Y_{\hbar\mathfrak{g}}).$$

Proof Let \mathbb{E}_\bullet be the filtration of $\mathbb{E} := \text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar\mathfrak{g}}, Y_{\hbar\mathfrak{g}})$ defined in (9.2). Then

$$\theta(Y_{\hbar\mathfrak{g}} \otimes \dot{Y}_{\hbar\mathfrak{g}}^*_{(n)}) \subset \mathbb{E}_n \quad \forall n \in \mathbb{N}.$$

Indeed, this follows from the definition of θ and the fact that if $f \in \bigoplus_{k \geq n} \dot{Y}_{\hbar\mathfrak{g}}^*_{-k}$ and $x \in J_n$, then $f(x) = 0$. Hence, we obtain

$$\Theta : Y_{\hbar\mathfrak{g}} \widetilde{\otimes} \dot{Y}_{\hbar\mathfrak{g}}^* = \varprojlim (Y_{\hbar\mathfrak{g}} \otimes \dot{Y}_{\hbar\mathfrak{g}}^* / Y_{\hbar\mathfrak{g}} \otimes \dot{Y}_{\hbar\mathfrak{g}}^*_{(n)}) \rightarrow \varprojlim \mathbb{E} / \mathbb{E}_n = \mathbb{E}.$$

As $Y_{\hbar\mathfrak{g}}$ and \mathbb{E} are topologically free, to show Θ is an isomorphism, it is sufficient to show its semiclassical limit $\tilde{\Theta}$ is (see Lemma 2.1). Employing the identifications of (9.1) and Lemma 9.1, we see that $\tilde{\Theta}$ coincides with the isomorphism

$$U(t_+) \widetilde{\otimes} S(\hbar t_+)^* \xrightarrow{\sim} \text{Hom}_{\mathbb{C}}(S(\hbar t_+), U(t_+))$$

uniquely extending the canonical map $U(t_+) \otimes_{\mathbb{C}} S(\hbar t_+)^* \hookrightarrow \text{Hom}_{\mathbb{C}}(S(\hbar t_+), U(t_+))$, which is the semiclassical limit $\tilde{\theta}$ of θ , by continuity. \blacksquare

We now give two remarks pertinent to the above discussion.

Remark 9.3 By Proposition 9.2, we may introduce the $\mathbb{C}[[\hbar]]$ -subalgebra

$$\dot{Y}_{\hbar\mathfrak{g}} \widetilde{\otimes} \dot{Y}_{\hbar\mathfrak{g}}^* := \Theta^{-1}(\text{End}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar\mathfrak{g}})) \subset Y_{\hbar\mathfrak{g}} \widetilde{\otimes} \dot{Y}_{\hbar\mathfrak{g}}^*.$$

By definition, it comes equipped with an isomorphism

$$\Theta|_{\dot{Y}_{\hbar\mathfrak{g}} \widetilde{\otimes} \dot{Y}_{\hbar\mathfrak{g}}^*} : \dot{Y}_{\hbar\mathfrak{g}} \widetilde{\otimes} \dot{Y}_{\hbar\mathfrak{g}}^* \xrightarrow{\sim} \text{End}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar\mathfrak{g}})$$

extending the injection $\theta|_{\dot{Y}_{\hbar}\mathfrak{g} \otimes \dot{Y}_{\hbar}\mathfrak{g}^*}$. It is worth noting that, by following the above arguments, it is easy to see that $\dot{Y}_{\hbar}\mathfrak{g} \widetilde{\otimes} \dot{Y}_{\hbar}\mathfrak{g}^*$ admits the equivalent description

$$\dot{Y}_{\hbar}\mathfrak{g} \widetilde{\otimes} \dot{Y}_{\hbar}\mathfrak{g}^* \cong \varprojlim_n (\dot{Y}_{\hbar}\mathfrak{g} \otimes \dot{Y}_{\hbar}\mathfrak{g}^* / \dot{Y}_{\hbar}\mathfrak{g} \otimes \dot{Y}_{\hbar}\mathfrak{g}_{(n)}^*).$$

Suppose now that $\varphi, \gamma \in \text{End}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}\mathfrak{g})$ with γ homogeneous of degree $a \in \mathbb{Z}$. Then $\varphi \otimes \gamma^t$ uniquely extends to an element of $\text{End}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}\mathfrak{g} \widetilde{\otimes} \dot{Y}_{\hbar}\mathfrak{g}^*)$ and Θ has the property that

$$(9.4) \quad \varphi \circ \Theta(\gamma) \circ \gamma = (\Theta \circ (\varphi \otimes \gamma^t))(\gamma) \quad \forall \gamma \in \dot{Y}_{\hbar}\mathfrak{g} \widetilde{\otimes} \dot{Y}_{\hbar}\mathfrak{g}^*.$$

Remark 9.4 The argument used to establish Proposition 6.2 implies that the natural $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$\text{End}_{\mathbb{C}[[\hbar]]}(Y_{\hbar}\mathfrak{g}) \rightarrow \text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}\mathfrak{g}, Y_{\hbar}\mathfrak{g}), \quad \varphi \mapsto \varphi|_{\dot{Y}_{\hbar}\mathfrak{g}},$$

is an embedding. Hence, we can (and shall) view $\text{End}_{\mathbb{C}[[\hbar]]}(Y_{\hbar}\mathfrak{g})$ as a subalgebra of $\text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}\mathfrak{g}, Y_{\hbar}\mathfrak{g})$. We may thus introduce $Y_{\hbar}\mathfrak{g} \widetilde{\otimes} Y_{\hbar}\mathfrak{g}^* \subset Y_{\hbar}\mathfrak{g} \widetilde{\otimes} \dot{Y}_{\hbar}\mathfrak{g}^*$ by setting

$$Y_{\hbar}\mathfrak{g} \widetilde{\otimes} Y_{\hbar}\mathfrak{g}^* := \Theta^{-1}(\text{End}_{\mathbb{C}[[\hbar]]}(Y_{\hbar}\mathfrak{g})).$$

We then have

$$(Y_{\hbar}\mathfrak{g} \widetilde{\otimes} Y_{\hbar}\mathfrak{g}^*) \cap (\dot{Y}_{\hbar}\mathfrak{g} \widetilde{\otimes} \dot{Y}_{\hbar}\mathfrak{g}^*) \cong \text{End}_{\mathbb{C}[[\hbar]]}^{\dot{Y}_{\hbar}\mathfrak{g}}(Y_{\hbar}\mathfrak{g}),$$

where the right-hand side consists of all $f \in \text{End}_{\mathbb{C}[[\hbar]]}(Y_{\hbar}\mathfrak{g})$ for which $f(\dot{Y}_{\hbar}\mathfrak{g}) \subset \dot{Y}_{\hbar}\mathfrak{g}$.

9.2 The restriction Θ_{χ}

We now give a few comments which concern the triangular decompositions of $\dot{Y}_{\hbar}\mathfrak{g}$ and $\dot{Y}_{\hbar}\mathfrak{g}^*$. For each choice of the symbol χ , let $\pi^{\chi} : \dot{Y}_{\hbar}\mathfrak{g} \rightarrow \dot{Y}_{\hbar}^{\chi}\mathfrak{g}$ denote the $\mathbb{C}[[\hbar]]$ -linear projection associated with the triangular decomposition of $\dot{Y}_{\hbar}\mathfrak{g}$, as defined in Section 6.5. These projections give rise to $\mathbb{C}[[\hbar]]$ -module embeddings

$$\text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}^{\chi}\mathfrak{g}, Y_{\hbar}^{\chi}\mathfrak{g}) \hookrightarrow \text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}\mathfrak{g}, Y_{\hbar}\mathfrak{g}), \quad \varphi^{\chi} \mapsto \varphi^{\chi} \circ \pi^{\chi}.$$

We shall henceforth adopt the viewpoint that $\text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}^{\chi}\mathfrak{g}, Y_{\hbar}^{\chi}\mathfrak{g})$ is a submodule of $\text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}\mathfrak{g}, Y_{\hbar}\mathfrak{g})$, with the above identification assumed. The restriction θ_{χ} of θ to $Y_{\hbar}^{\chi}\mathfrak{g} \otimes \dot{Y}_{\hbar}^{\chi}\mathfrak{g}^*$ is then a $\mathbb{C}[[\hbar]]$ -module injection

$$\theta_{\chi} : Y_{\hbar}^{\chi}\mathfrak{g} \otimes \dot{Y}_{\hbar}^{\chi}\mathfrak{g}^* \hookrightarrow \text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}^{\chi}\mathfrak{g}, Y_{\hbar}^{\chi}\mathfrak{g}).$$

Following the above procedure, we may introduce the $\mathbb{C}[[\hbar]]$ -algebra

$$Y_{\hbar}^{\chi}\mathfrak{g} \widetilde{\otimes} \dot{Y}_{\hbar}^{\chi}\mathfrak{g}^* := \varprojlim_n (Y_{\hbar}^{\chi}\mathfrak{g} \otimes \dot{Y}_{\hbar}^{\chi}\mathfrak{g}^* / Y_{\hbar}^{\chi}\mathfrak{g} \otimes \dot{Y}_{\hbar}^{\chi}\mathfrak{g}_{(n)}^*),$$

where $\dot{Y}_{\hbar}^{\chi}\mathfrak{g}_{(n)}^*$ is the closure of the $\mathbb{C}[[\hbar]]$ -submodule of $\dot{Y}_{\hbar}^{\chi}\mathfrak{g}^*$ generated by the direct sum $\bigoplus_{k \geq n} \dot{Y}_{\hbar}^{\chi}\mathfrak{g}_{-k}^*$. The above arguments show that this completed tensor product is a

flat $\mathbb{C}[[\hbar]]$ -algebra deformation of

$$U(t^{\chi}) \otimes S(\hbar t^{\chi})^* = \prod_{n \in \mathbb{N}} U(t^{\chi}) \otimes_{\mathbb{C}} S(\hbar t^{\chi})_{-n}^*.$$

It follows that the natural algebra homomorphism $Y_{\hbar}^{\chi} \mathfrak{g} \otimes \dot{Y}_{\hbar}^{\chi} \mathfrak{g}^* \rightarrow Y_{\hbar} \mathfrak{g} \otimes \dot{Y}_{\hbar} \mathfrak{g}^*$ is injective and, by the proof of Proposition 9.2, that θ_{χ} uniquely extends to an isomorphism of $\mathbb{C}[[\hbar]]$ -modules

$$(9.5) \quad \Theta_{\chi} : Y_{\hbar}^{\chi} \mathfrak{g} \otimes \dot{Y}_{\hbar}^{\chi} \mathfrak{g}^* \xrightarrow{\sim} \text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}^{\chi} \mathfrak{g}, Y_{\hbar}^{\chi} \mathfrak{g})$$

which coincides with the restriction of Θ to $Y_{\hbar}^{\chi} \mathfrak{g} \otimes \dot{Y}_{\hbar}^{\chi} \mathfrak{g}^*$. In particular, since $Y_{\hbar}^{\chi} \mathfrak{g} \otimes \dot{Y}_{\hbar}^{\chi} \mathfrak{g}^*$ is a subalgebra of $Y_{\hbar} \mathfrak{g} \otimes \dot{Y}_{\hbar} \mathfrak{g}^*$, the space $\text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}^{\chi} \mathfrak{g}, Y_{\hbar}^{\chi} \mathfrak{g})$ is a subalgebra of $\text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar} \mathfrak{g}, Y_{\hbar} \mathfrak{g})$ and Θ_{χ} is an isomorphism of $\mathbb{C}[[\hbar]]$ -algebras.

Finally, as in Remarks 9.3 and 9.4, we set

$$\begin{aligned} \dot{Y}_{\hbar}^{\chi} \mathfrak{g} \otimes \dot{Y}_{\hbar}^{\chi} \mathfrak{g}^* &:= \Theta_{\chi}^{-1}(\text{End}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}^{\chi} \mathfrak{g})) \cong \varprojlim_n (\dot{Y}_{\hbar}^{\chi} \mathfrak{g} \otimes \dot{Y}_{\hbar}^{\chi} \mathfrak{g}^* / \dot{Y}_{\hbar}^{\chi} \mathfrak{g} \otimes \dot{Y}_{\hbar}^{\chi} \mathfrak{g}_{(n)}^*), \\ Y_{\hbar}^{\chi} \mathfrak{g} \otimes Y_{\hbar}^{\chi} \mathfrak{g}^* &:= \Theta_{\chi}^{-1}(\text{End}_{\mathbb{C}[[\hbar]]}(Y_{\hbar}^{\chi} \mathfrak{g})). \end{aligned}$$

9.3 Canonical tensors and universal R-matrices

By Proposition 9.2, we may introduce R and R^{χ} in $Y_{\hbar} \mathfrak{g} \otimes \dot{Y}_{\hbar} \mathfrak{g}^*$, for each choice of χ , as the elements

$$R := \Theta^{-1}(\mathbf{1}_{\dot{Y}_{\hbar} \mathfrak{g}}) \in Y_{\hbar} \mathfrak{g} \otimes \dot{Y}_{\hbar} \mathfrak{g}^* \quad \text{and} \quad R^{\chi} := \Theta^{-1}(\mathbf{1}_{\dot{Y}_{\hbar}^{\chi} \mathfrak{g}}) \in Y_{\hbar}^{\chi} \mathfrak{g} \otimes \dot{Y}_{\hbar}^{\chi} \mathfrak{g}^*.$$

That is, R and R^{χ} are the canonical elements associated with the Hopf pairing $\dot{Y}_{\hbar} \mathfrak{g} \times \dot{Y}_{\hbar} \mathfrak{g}^* \rightarrow \mathbb{C}[[\hbar]]$ and its restriction to $\dot{Y}_{\hbar}^{\chi} \mathfrak{g} \times \dot{Y}_{\hbar}^{\chi} \mathfrak{g}^*$, respectively.

As $\mathbf{1}_{\dot{Y}_{\hbar} \mathfrak{g}}$ and $\mathbf{1}_{\dot{Y}_{\hbar}^{\chi} \mathfrak{g}}$ coincide with $\mathbf{1}_{Y_{\hbar} \mathfrak{g}}$ and $\mathbf{1}_{Y_{\hbar}^{\chi} \mathfrak{g}}$ under the identifications of Remark 9.4 and Section 9.2, respectively, R (resp. R^{χ}) is also the canonical element defined by the pairing between $Y_{\hbar} \mathfrak{g}$ and its restricted dual $Y_{\hbar} \mathfrak{g}^*$ (resp. $Y_{\hbar}^{\chi} \mathfrak{g}$ and $Y_{\hbar}^{\chi} \mathfrak{g}^*$). In particular, we have

$$R \in (Y_{\hbar} \mathfrak{g} \otimes Y_{\hbar} \mathfrak{g}^*) \cap (\dot{Y}_{\hbar} \mathfrak{g} \otimes \dot{Y}_{\hbar} \mathfrak{g}^*) \quad \text{and} \quad R^{\chi} \in (Y_{\hbar}^{\chi} \mathfrak{g} \otimes Y_{\hbar}^{\chi} \mathfrak{g}^*) \cap (\dot{Y}_{\hbar}^{\chi} \mathfrak{g} \otimes \dot{Y}_{\hbar}^{\chi} \mathfrak{g}^*).$$

It is worth pointing out that one can immediately deduce a number of properties that R and R^{χ} satisfy using only elementary properties of Θ . For instance:

- (R1) Applying (9.4) with $\gamma = 1$ while using that $\Theta(1 \otimes f) = \iota \circ f$ recovers the characteristic identities

$$(\iota \circ f) \otimes \mathbf{1} \cdot R = 1 \otimes f \quad \text{and} \quad (\iota \circ f^{\chi}) \otimes \mathbf{1} \cdot R^{\chi} = 1 \otimes f^{\chi}$$

for all $f \in \dot{Y}_{\hbar} \mathfrak{g}^*$ and $f^{\chi} \in \dot{Y}_{\hbar}^{\chi} \mathfrak{g}^*$.

- (R2) Taking $\varphi = \gamma = \omega$ in (9.4) with $\gamma = \mathbf{1}_{\dot{Y}_{\hbar} \mathfrak{g}}$ or $\mathbf{1}_{\dot{Y}_{\hbar}^{\chi} \mathfrak{g}}$ while using that $\mathbf{1}_{\dot{Y}_{\hbar}^{\pm} \mathfrak{g}} = \omega \circ \mathbf{1}_{\dot{Y}_{\hbar}^{\mp} \mathfrak{g}} \circ \omega$ gives rise to the identities

$$R = (\omega \otimes \omega^t)R \quad \text{and} \quad R^{\pm} = (\omega \otimes \omega^t)(R^{\mp}).$$

(R3) Since $\mathbf{1}_{\dot{Y}_{\hbar}\mathfrak{g}}$ is a \mathfrak{g} -invariant element of $\text{End}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}\mathfrak{g})$ and Θ is a \mathfrak{g} -module intertwiner, R is a \mathfrak{g} -invariant element $\dot{Y}_{\hbar}\mathfrak{g} \widetilde{\otimes} \dot{Y}_{\hbar}\mathfrak{g}^*$. Using the multiplication in $D(Y_{\hbar}\mathfrak{g})$, this can be written as

$$[x \otimes 1 + 1 \otimes x, R] = 0 \quad \forall x \in \mathfrak{g}.$$

Similarly, R^\pm and R^0 are both \mathfrak{h} -invariant.

In addition, the standard quantum double arguments show that R satisfies the quasitriangularity relations of (3) in Section 8.1 in suitable completions of tensor powers of $D(Y_{\hbar}\mathfrak{g})$. For instance, the cabling identities $\Delta \otimes \mathbf{1}(R) = R_{13}R_{23}$ and $\mathbf{1} \otimes \Delta^\vee(R) = R_{13}R_{12}$ are equivalent to the simple identities

$$\mathbf{1}_{\dot{Y}_{\hbar}\mathfrak{g}}^{(1)} * \mathbf{1}_{\dot{Y}_{\hbar}\mathfrak{g}}^{(2)} = \Delta \quad \text{and} \quad (\varepsilon \otimes \mathbf{1}_{\dot{Y}_{\hbar}\mathfrak{g}}) * (\mathbf{1}_{\dot{Y}_{\hbar}\mathfrak{g}} \otimes \varepsilon) = (12) \circ m$$

in the convolution algebras

$$\begin{aligned} (\dot{Y}_{\hbar}\mathfrak{g}^{\otimes 2}) \widetilde{\otimes} \dot{Y}_{\hbar}\mathfrak{g}^* &:= \text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}\mathfrak{g}, \dot{Y}_{\hbar}\mathfrak{g}^{\otimes 2}), \\ \dot{Y}_{\hbar}\mathfrak{g} \widetilde{\otimes} (\dot{Y}_{\hbar}\mathfrak{g}^*)^{\otimes 2} &:= \text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_{\hbar}\mathfrak{g}^{\otimes 2}, \dot{Y}_{\hbar}\mathfrak{g}), \end{aligned}$$

respectively, where $\mathbf{1}_{\dot{Y}_{\hbar}\mathfrak{g}}^{(a)} : \dot{Y}_{\hbar}\mathfrak{g} \rightarrow \dot{Y}_{\hbar}\mathfrak{g}^{\otimes 2}$ is given by $x \mapsto x^{(a)}$.

Our main goal in this section is to establish the remaining assertion of Theorem 1 from Section 1.1, which claims that R can be identified with the universal R -matrix of the Yangian. This interpretation allows for a precise framework for understanding the topological quasitriangular structure on $D(Y_{\hbar}\mathfrak{g})$ alluded to above. To achieve this, we will need the next lemma, where Φ_z^\vee is as in Proposition 7.1.

Lemma 9.5 $\tau_w \otimes \Phi_z^\vee$ extends to an injective $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$\tau_w \widetilde{\otimes} \Phi_z^\vee : Y_{\hbar}\mathfrak{g} \widetilde{\otimes} Y_{\hbar}\mathfrak{g}^\vee \hookrightarrow (Y_{\hbar}\mathfrak{g}^{\otimes 2}[[w]])[[z^{-1}]].$$

Proof It is sufficient to show $\mathbf{1} \otimes \Phi_z^\vee$ extends to an injective $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$\mathbf{1} \widetilde{\otimes} \Phi_z^\vee : Y_{\hbar}\mathfrak{g} \widetilde{\otimes} Y_{\hbar}\mathfrak{g}^\vee \rightarrow Y_{\hbar}\mathfrak{g}^{\otimes 2}[[z^{-1}]].$$

Since $\Phi_z^\vee(\dot{Y}_{\hbar}\mathfrak{g}_{-k}^*) \subset z^{-k} \widehat{Y_{\hbar}\mathfrak{g}}_z \subset z^{-k} Y_{\hbar}\mathfrak{g}[[z^{-1}]]$, $\mathbf{1} \otimes \Phi_z^\vee$ induces a family of compatible algebra homomorphisms

$$(\mathbf{1} \otimes \Phi_z^\vee)_n : Y_{\hbar}\mathfrak{g} \otimes \dot{Y}_{\hbar}\mathfrak{g}^* / \dot{Y}_{\hbar}\mathfrak{g} \otimes \dot{Y}_{\hbar}\mathfrak{g}_{(n)}^* \rightarrow Y_{\hbar}\mathfrak{g}^{\otimes 2}[[z^{-1}]] / z^{-n} Y_{\hbar}\mathfrak{g}^{\otimes 2}[[z^{-1}]].$$

Taking the projective limit gives the desired extension $\mathbf{1} \widetilde{\otimes} \Phi_z^\vee$. We are left to verify that it is indeed injective, which is perhaps a bit subtle (given that Φ_z^\vee is injective), but not difficult. As $Y_{\hbar}\mathfrak{g} \widetilde{\otimes} \dot{Y}_{\hbar}\mathfrak{g}^*$ is separated with respect to the \hbar -adic topology and $Y_{\hbar}\mathfrak{g}^{\otimes 2}[[z^{-1}]]$ is torsion free, this can be done by verifying that the semiclassical limit of $\mathbf{1} \widetilde{\otimes} \Phi_z^\vee$ is injective. This is the linear map

$$U(\mathfrak{t}_+) \widetilde{\otimes} S(\hbar \mathfrak{t}_+)^* \rightarrow \prod_{n \in \mathbb{N}} z^{-n} (U(\mathfrak{t}_+) \otimes_{\mathbb{C}} \widehat{U(\mathfrak{t}_+)}_z) \subset U(\mathfrak{t}_+)^{\otimes 2}[[z^{-1}]]$$

which is $\mathbf{1} \otimes \bar{\Phi}_z^\vee : U(\mathfrak{t}_+) \widetilde{\otimes} S(\hbar \mathfrak{t}_+)^*_{-n} \rightarrow z^{-n} (U(\mathfrak{t}_+) \otimes_{\mathbb{C}} \widehat{U(\mathfrak{t}_+)}_z)$ on the n -th component of the direct product, where $\bar{\Phi}_z^\vee$ is the semiclassical limit of Φ_z^\vee and $\widehat{U(\mathfrak{t}_+)}_z =$

$\prod_{n \in \mathbb{N}} U(\mathfrak{t}_+) n z^{-n}$. The desired result now follows from the fact that Φ_z^\vee is injective, which can be seen as a consequence of the equality $\mathcal{E} \circ \Phi_z^\vee = \Phi \circ i$ and that, by [41, Theorem 6.2] and the proof of Lemma 7.4, both Φ and i have injective semiclassical limits. ■

To recover the desired result as stated in (2) of Theorem I, we now translate some of the above constructions and results from $D(Y_{\hbar}\mathfrak{g})$ to $DY_{\hbar}\mathfrak{g}$ using the isomorphism Y from Theorem 8.4 or, equivalently, the Hopf algebra embeddings ι and i from Theorem 7.5. Let us set

$$Y_{\hbar}\mathfrak{g} \dot{\otimes} Y_{\hbar}\mathfrak{g}^\vee := \varprojlim_n (\iota(Y_{\hbar}\mathfrak{g}) \otimes i(Y_{\hbar}\mathfrak{g}^\vee) / \iota(Y_{\hbar}\mathfrak{g}) \otimes i(Y_{\hbar}\mathfrak{g}_{(n)}^\vee)).$$

This definition is such that isomorphism

$$\iota \otimes i : Y_{\hbar}\mathfrak{g} \otimes Y_{\hbar}\mathfrak{g}^\vee \xrightarrow{\sim} \iota(Y_{\hbar}\mathfrak{g}) \otimes i(Y_{\hbar}\mathfrak{g}^\vee) \subset DY_{\hbar}\mathfrak{g} \otimes DY_{\hbar}\mathfrak{g}$$

extends to an isomorphism of $\mathbb{C}[[\hbar]]$ -algebras

$$\iota \dot{\otimes} i : Y_{\hbar}\mathfrak{g} \widetilde{\otimes} Y_{\hbar}\mathfrak{g}^\vee \xrightarrow{\sim} Y_{\hbar}\mathfrak{g} \dot{\otimes} Y_{\hbar}\mathfrak{g}^\vee.$$

Armed with these preliminaries, we may now introduce the *universal R-matrix* of the Yangian double $DY_{\hbar}\mathfrak{g}$ as the element

$$\mathcal{R} := (\iota \dot{\otimes} i)(R) \in (Y_{\hbar}\mathfrak{g} \dot{\otimes} Y_{\hbar}\mathfrak{g}^\vee) \cap (\dot{Y}_{\hbar}\mathfrak{g} \dot{\otimes} Y_{\hbar}\mathfrak{g}^\vee) \subset Y_{\hbar}\mathfrak{g} \dot{\otimes} Y_{\hbar}\mathfrak{g}^\vee,$$

where, for each choice of A and B , $A \dot{\otimes} B$ denotes the image of $A \widetilde{\otimes} B$ under $\iota \dot{\otimes} i$. Similarly, we define the $DY_{\hbar}\mathfrak{g}$ analogues of R^\pm and R^0 by setting

$$\mathcal{R}^\pm := (\iota \dot{\otimes} i)(R^\pm) \in Y_{\hbar}^\pm \mathfrak{g} \dot{\otimes} \dot{Y}_{\hbar}^\pm \mathfrak{g}^* \quad \text{and} \quad \mathcal{R}^0 := (\iota \dot{\otimes} i)(R^0) \in Y_{\hbar}^0 \mathfrak{g} \dot{\otimes} \dot{Y}_{\hbar}^0 \mathfrak{g}^*.$$

By Theorem 4.6 and Lemma 7.4, Φ_z satisfies $\Phi_z \circ \iota = \tau_z$ and $\Phi_z \circ i = \Phi_z^\vee$. Lemma 9.5 therefore implies that the restriction of $\Phi_w \otimes \Phi_z$ to $\iota(Y_{\hbar}\mathfrak{g}) \otimes i(Y_{\hbar}\mathfrak{g}^\vee) \subset DY_{\hbar}\mathfrak{g}^{\otimes 2}$ extends to an injective $\mathbb{C}[[\hbar]]$ -algebra homomorphism

$$\Phi_w \dot{\otimes} \Phi_z : Y_{\hbar}\mathfrak{g} \dot{\otimes} Y_{\hbar}\mathfrak{g}^\vee \hookrightarrow Y_{\hbar}\mathfrak{g}^{\otimes 2}[[w]][[z^{-1}]].$$

Indeed, we may set $\Phi_w \dot{\otimes} \Phi_z := (\tau_w \widetilde{\otimes} \Phi_z^\vee) \circ (\iota \dot{\otimes} i)^{-1}$. With this homomorphism at our disposal, we are now in a position to state and prove the last main result outlined in Section 1.1.

Theorem 9.6 *The following identities hold in $Y_{\hbar}(\mathfrak{g})^{\otimes 2}[[w]][[z^{-1}]]$:*

$$\begin{aligned} (\Phi_w \dot{\otimes} \Phi_z)(\mathcal{R}) &= \mathcal{R}(w - z), \\ (\Phi_w \dot{\otimes} \Phi_z)(\mathcal{R}^\pm) &= \mathcal{R}^\pm(w - z), \quad (\Phi_w \dot{\otimes} \Phi_z)(\mathcal{R}^0) = \mathcal{R}^0(w - z). \end{aligned}$$

Consequently, \mathcal{R} admits the factorization $\mathcal{R} = \mathcal{R}^+ \cdot \mathcal{R}^- \cdot \mathcal{R}^0$ in $Y_{\hbar}\mathfrak{g} \dot{\otimes} Y_{\hbar}\mathfrak{g}^\vee$.

Proof By Theorems 4.1 and 6.7 of [17] and Theorem 3.8 above, $\mathcal{R}(z)$ and $\mathcal{R}^\chi(z)$ satisfy $(\tau_w \otimes 1)\mathcal{R}(z) = \mathcal{R}(w + z)$ and $(\tau_w \otimes 1)\mathcal{R}^\chi(z) = \mathcal{R}^\chi(w + z)$ in $Y_{\hbar}(\mathfrak{g})^{\otimes 2}[[w]][[z^{-1}]]$, for each choice of the symbol χ . Therefore, to prove the first assertion of the theorem, it is sufficient to establish the equalities

$$(1 \widetilde{\otimes} \Phi_z^\vee)(R) = \mathcal{R}(-z) \quad \text{and} \quad (1 \widetilde{\otimes} \Phi_z^\vee)(R^\chi) = \mathcal{R}^\chi(-z)$$

in $(\dot{Y}_\hbar \mathfrak{g} \otimes Y_\hbar \mathfrak{g})[[z^{-1}]]$ and $(\dot{Y}_\hbar^\chi \mathfrak{g} \otimes Y_\hbar^{-\chi} \mathfrak{g})[[z^{-1}]]$, respectively. These relations will hold provided their images under the evaluations

$$\begin{aligned} f \otimes 1 &: (\dot{Y}_\hbar \mathfrak{g} \otimes Y_\hbar \mathfrak{g})[[z^{-1}]] \rightarrow Y_\hbar \mathfrak{g}[[z^{-1}]], \\ f^\chi \otimes 1 &: (\dot{Y}_\hbar^\chi \mathfrak{g} \otimes Y_\hbar^{-\chi} \mathfrak{g})[[z^{-1}]] \rightarrow Y_\hbar^{-\chi} \mathfrak{g}[[z^{-1}]] \end{aligned}$$

are satisfied in $Y_\hbar \mathfrak{g}[[z^{-1}]]$ for each $f \in \dot{Y}_\hbar \mathfrak{g}^*$ and $f^\chi \in \dot{Y}_\hbar^\chi \mathfrak{g}^*$, respectively. This follows from the definition of Φ_z^\vee , (R1), and Corollary 7.7, which collectively output the relations

$$\begin{aligned} (f \otimes \Phi_z^\vee)(R) &= \Phi_z^\vee(f) = (f \otimes 1)\mathcal{R}(-z), \\ (f^\chi \otimes \Phi_z^\vee)(R) &= \Phi_z^\vee(f) = (f^\chi \otimes 1)\mathcal{R}^\chi(-z). \end{aligned}$$

This completes the proof of the first statement of the theorem. The second assertion now follows immediately from the decomposition $\mathcal{R}(z) = \mathcal{R}^+(z)\mathcal{R}^0(z)\mathcal{R}^-(z)$ (see Theorem 3.8) and the injectivity of $\Phi_w \otimes \Phi_z^\vee$. ■

Remark 9.7 In the closely related setting of [32], a formal version (*i.e.*, suppressing convergence issues) of the factorization $\mathcal{R} = \mathcal{R}^+ \cdot \mathcal{R}^0 \cdot \mathcal{R}^-$ from Theorem 9.6 was established in [32, Proposition 5.1] by proving directly that the Hopf pairing between the Yangian and its dual splits with respect to the underlying triangular decompositions (see Theorem 3.1 therein).

Remark 9.8 Since Φ_z intertwines the Chevalley involutions of $DY_\hbar \mathfrak{g}$ and $Y_\hbar \mathfrak{g}$, it follows from Part (2) of Proposition 7.1 and the relation $\Phi_z \circ \check{i} = \Phi_z^\vee$ of Lemma 7.4 that \check{i} satisfies $\omega \circ \check{i} = \check{i} \circ \omega^t$. Consequently, the property (R2) translates to

$$\mathcal{R} = (\omega \otimes \omega)(\mathcal{R}) \quad \text{and} \quad \mathcal{R}^\pm = (\omega \otimes \omega)(\mathcal{R}^\mp).$$

Hence, by Theorem 9.6, $\mathcal{R}(z)$ and its components $\mathcal{R}^\pm(z)$ satisfy the relations

$$\mathcal{R}(z) = (\omega \otimes \omega)(\mathcal{R}(z)) \quad \text{and} \quad \mathcal{R}^\pm(z) = (\omega \otimes \omega)(\mathcal{R}^\mp(z)).$$

In particular, this observation recovers the first identity of Corollary 3.10.

One can now interpret the quasitriangularity relations for $DY_\hbar \mathfrak{g}$ in terms of the relations of Theorem 3.8. Namely, setting $v = w - z$, we see that the cabling identities of \mathcal{R} correspond to the relations

$$\begin{aligned} \Delta \otimes 1(\mathcal{R}(v)) &= \mathcal{R}_{13}(v)\mathcal{R}_{23}(v), \\ 1 \otimes \Delta(\mathcal{R}(v)) &= \mathcal{R}_{13}(v)\mathcal{R}_{12}(v) \end{aligned}$$

in $Y_\hbar(\mathfrak{g})^{\otimes 3}[[w]][[z^{-1}]]$, which are satisfied by Theorem 3.8. In more detail, these are obtained heuristically by applying $\Phi_u \otimes \Phi_w \otimes \Phi_z$ and $\Phi_w \otimes \Phi_y \otimes \Phi_z$ to

$$(\dot{\Delta} \otimes 1)(\mathcal{R}) = \mathcal{R}_{13}\mathcal{R}_{23} \quad \text{and} \quad (1 \otimes \dot{\Delta})(\mathcal{R}) = \mathcal{R}_{13}\mathcal{R}_{12},$$

respectively, where $\dot{\Delta}$ is the coproduct on $DY_\hbar \mathfrak{g}$, and then evaluating $u \mapsto w$ and $y \mapsto z$ while using $(\Phi_u \otimes \Phi_z) \circ \dot{\Delta}|_{u=z} = \dot{\Delta} \circ \Phi_z$. Similarly, applying $\Phi_w \otimes \Phi_z$ to the intertwiner equation $\Delta^{\text{op}}(x) = \mathcal{R} \Delta(x) \mathcal{R}^{-1}$ for $x \in Y_\hbar(\mathfrak{g})$ leads to the identity

$$\tau_v \otimes 1 \circ \Delta^{\text{op}}(\tau_z(x)) = \mathcal{R}(v) \cdot \tau_v \otimes 1 \circ \Delta(\tau_z(x)) \cdot \mathcal{R}(v)^{-1}$$

in $Y_{\hbar}(\mathfrak{g})^{\otimes 2}[w][z; z^{-1}]$, which is satisfied by Theorem 3.8. We caution, however, that the situation here is more subtle for general $x \in DY_{\hbar}\mathfrak{g}$.

9.4 Remarks

The identifications established in Theorem 9.6 provide a rigorous framework for understanding the motivating remarks given in [17, Section 1.6]. We now give a few comments related to this point.

First, we note that the diagonal factor $\mathcal{R}^0(z)$ of $\mathcal{R}(z)$ was explicitly obtained in [17, Section 6.6] by computing the common asymptotic expansion of the two $GL(V_1 \otimes V_2)$ -valued meromorphic abelian R -matrices constructed in [16, Section 5], where V_1, V_2 are an arbitrary pair of finite-dimensional representations of the Yangian. As explained in [16, Section 5.2], their construction was motivated by the heuristic formula for \mathcal{R}^0 given in [32, Theorem 5.2]. Theorem 9.6 makes this relation precise, and shows that the explicit formula for $\mathcal{R}^0(z)$ obtained in [17], and recalled in Section 3.6, does indeed compute the canonical element defined by the pairing on $\dot{Y}_{\hbar}^0\mathfrak{g} \times \dot{Y}_{\hbar}^0\mathfrak{g}^*$.

Next, we emphasize that the factor \mathcal{R}^{\pm} (equivalently, $\mathcal{R}^{\pm}(z)$) now admits two distinct characterizations. On the one hand, it is uniquely determined by the recurrence relations (3.18) which were at the heart of [17, Theorem 4.1]. On the other hand, it arises as the canonical element defined by the pairing on $\dot{Y}_{\hbar}^{\pm}\mathfrak{g} \times \dot{Y}_{\hbar}^{\pm}\mathfrak{g}^*$, and can thus be realized explicitly by computing the dual set to any fixed homogeneous basis of the \mathbb{N} -graded torsion free $\mathbb{C}[[\hbar]]$ -module $\dot{Y}_{\hbar}^{\pm}(\mathfrak{g})$. In more detail, if $B_k^{\pm} \subset \dot{Y}_{\hbar}^{\pm}(\mathfrak{g})_k$ is a lift of any basis of the finite-dimensional k th component $S_k(\hbar t_{\pm}^{\pm})$ of $S(\hbar t_{\pm}^{\pm}) \cong \dot{Y}_{\hbar}^{\pm}(\mathfrak{g})/\hbar\dot{Y}_{\hbar}^{\pm}(\mathfrak{g})$ (see Section 5.3), then $B^{\pm} = \cup_k B_k^{\pm}$ is a basis of $\dot{Y}_{\hbar}^{\pm}(\mathfrak{g})$ and

$$\mathcal{R}^{\pm} = \sum_{x \in B^{\pm}} \iota(x) \otimes \check{\iota}(f_x) \in Y_{\hbar}\mathfrak{g} \otimes Y_{\hbar}\mathfrak{g}^{\vee},$$

where $\{f_x\}_{x \in B^{\pm}} \subset \dot{Y}_{\hbar}^{\pm}\mathfrak{g}^*$ is the dual set to B^{\pm} , uniquely determined by $f_x(y) = \delta_{x,y}$ for all $x, y \in B^{\pm}$. Here, we note if x is of degree k , then it follows automatically that $f_x \in \dot{Y}_{\hbar}\mathfrak{g}_{-k}^* \cong \text{Hom}_{\mathbb{C}[[\hbar]]}^{-k}(\dot{Y}_{\hbar}(\mathfrak{g}), \mathbb{C}[[\hbar]])$. This implies that the right-hand side of the above expression defines a unique element in $Y_{\hbar}\mathfrak{g} \otimes Y_{\hbar}\mathfrak{g}^{\vee}$, which coincides with \mathcal{R}^{\pm} as its image under $\Theta_{\pm} \circ (\iota \otimes \check{\iota})^{-1}$ is $1_{Y_{\hbar}^{\pm}\mathfrak{g}}$ (see (9.5) and Section 9.3).

In the special case where $\mathfrak{g} = \mathfrak{sl}_2$, there are two closed-form expressions for \mathcal{R}^{\pm} (equivalently, $\mathcal{R}^{\pm}(z)$) which have arisen from these two separate viewpoints (see [32, Theorem 5.1] and [17, Theorem 5.5]). For \mathfrak{g} of arbitrary rank, no such expressions are known, though an infinite-product formula for \mathcal{R}^{\pm} was conjectured in [32, (5.43)], motivated by the earlier works [28–31].

9.5 On the blocks of \mathcal{R}^{\pm}

To conclude, we wish to highlight that the dual bases approach discussed in the previous subsection provides a natural interpretation of some of the basic properties of $\mathcal{R}^{\pm}(z)$ discovered in [17].

Given $\beta \in Q$, let $\pi_{\beta} : DY_{\hbar}\mathfrak{g} \rightarrow DY_{\hbar}\mathfrak{g}_{\beta}$ denote the $\mathbb{C}[[\hbar]]$ -linear projection associated with the topological Q -grading on $DY_{\hbar}\mathfrak{g}$ defined in Section 4.2, and let

$\dot{\pi}_\beta : \dot{Y}_\hbar \mathfrak{g} \rightarrow \dot{Y}_\hbar \mathfrak{g}_\beta$ denote its restriction to $\dot{Y}_\hbar \mathfrak{g}$. Consider the element

$$\dot{\pi}_\beta \circ \mathbf{1}_{\dot{Y}_\hbar \mathfrak{g}} = \mathbf{1}_{\dot{Y}_\hbar \mathfrak{g}_\beta} \circ \dot{\pi}_\beta \in \text{End}_{\mathbb{C}[[\hbar]]}(\dot{Y}_\hbar \mathfrak{g}),$$

where we view $\text{End}_{\mathbb{C}[[\hbar]]}(\dot{Y}_\hbar \mathfrak{g}) \subset \text{End}_{\mathbb{C}[[\hbar]]}(\dot{Y}_\hbar \mathfrak{g})$, as in Section 9.2. Note that under the identification provided by the natural inclusion

$$\text{End}_{\mathbb{C}[[\hbar]]}(\dot{Y}_\hbar \mathfrak{g}_\beta) \hookrightarrow \text{Hom}_{\mathbb{C}[[\hbar]]}(\dot{Y}_\hbar \mathfrak{g}, Y_\hbar^\pm \mathfrak{g}), \quad \varphi \mapsto \varphi \circ \dot{\pi}_\beta,$$

it coincides with the identity transformation of $\text{End}_{\mathbb{C}[[\hbar]]}(\dot{Y}_\hbar \mathfrak{g}_\beta)$. For each $\beta \in Q_+$, we may therefore define $R_\beta^\pm := \Theta^{-1}(\mathbf{1}_\beta^\pm)$, where $\mathbf{1}_\beta^\pm = \mathbf{1}_{\dot{Y}_\hbar \mathfrak{g}_{\pm\beta}}$. By (9.4) and the reasoning used in (R2) of Section 9.3, we have

$$(9.6) \quad R_\beta^\pm = (\dot{\pi}_{\pm\beta} \otimes \mathbf{1})(R^\pm) = (\mathbf{1} \otimes \dot{\pi}_{\pm\beta}^t)(R^\pm) \quad \text{and} \quad R_\beta^\pm = (\omega \otimes \omega^t)(R_\beta^\mp),$$

where we note that, for each $\alpha \in Q$, $\dot{\pi}_\alpha^t$ is just the projection $\dot{Y}_\hbar \mathfrak{g}^* \rightarrow \dot{Y}_\hbar \mathfrak{g}_{-\alpha}^*$ associated with the topological Q -grading on $\dot{Y}_\hbar \mathfrak{g}^*$ (see Corollary 6.5). In the topological tensor product $\dot{Y}_\hbar \mathfrak{g} \widetilde{\otimes} \dot{Y}_\hbar \mathfrak{g}^*$, we have the equality $R^\pm = \sum_{\beta \in Q_+} R_\beta^\pm$.

For the sake of the below discussion, it is worth pointing out that the convergence of the infinite sum $\sum_{\beta \in Q_+} R_\beta^\pm$ is also clear from the point of view of dual bases. Indeed, by Corollary 5.5, we have $\dot{Y}_\hbar^\pm(\mathfrak{g})_{\pm\beta} \subset \hbar^{v(\beta)} Y_\hbar^\pm(\mathfrak{g})_{\pm\beta}$, where we recall that $v(\beta) \in \mathbb{N}$ is defined by (3.17). Hence, any homogeneous element in $\dot{Y}_\hbar^\pm(\mathfrak{g})_{\pm\beta}$ with respect to the underlying \mathbb{N} -grading belongs to $\dot{Y}_\hbar(\mathfrak{g})_{v(\beta)+\ell}$ for some $\ell \in \mathbb{N}$. It follows that the dual set to any homogeneous basis of the \mathbb{N} -graded $\mathbb{C}[[\hbar]]$ -module $\dot{Y}_\hbar^\pm(\mathfrak{g})_{\pm\beta}$ belongs to

$$\bigoplus_{k \geq v(\beta)} \dot{Y}_\hbar^\pm \mathfrak{g}_{-k}^* \subset \dot{Y}_\hbar^\pm \mathfrak{g}_{(v(\beta))}^*,$$

where the space on the right-hand side is the closure of the $\mathbb{C}[[\hbar]]$ -module generated by the left-hand side (see Section 9.2). This implies the convergence of $\sum_{\beta \in Q_+} R_\beta^\pm$ in $\dot{Y}_\hbar^\pm \mathfrak{g} \widetilde{\otimes} \dot{Y}_\hbar \mathfrak{g}^*$ while establishing that $R_\beta^\pm \in \hbar^{v(\beta)} Y_\hbar^\pm \mathfrak{g} \widetilde{\otimes} \dot{Y}_\hbar \mathfrak{g}^*$. Next, following the procedure from Section 9.3, let us set

$$\mathcal{R}_\beta^\pm := (\iota \dot{\otimes} i)(R_\beta^\pm) \quad \forall \beta \in Q_+.$$

The relations of (9.6) then translate to

$$\mathcal{R}_\beta^\pm = (\pi_{\pm\beta} \otimes \mathbf{1})(\mathcal{R}^\pm) = (\mathbf{1} \otimes \pi_{\mp\beta})(\mathcal{R}^\pm) \quad \text{and} \quad \mathcal{R}_\beta^\pm = (\omega \otimes \omega)(\mathcal{R}_\beta^\mp),$$

where ω is the Chevalley involution on $DY_\hbar \mathfrak{g}$ (see Section 4.2). Now recall that $\mathcal{R}_\beta^-(z)$ is the $Y_\hbar(\mathfrak{g})_{-\beta} \otimes Y_\hbar(\mathfrak{g})_\beta$ component of $\mathcal{R}^-(z)$, characterized by the recurrence relation (3.18). The $Y_\hbar(\mathfrak{g})_\beta \otimes Y_\hbar(\mathfrak{g})_{-\beta}$ block of $\mathcal{R}^+(z)$ is then

$$\mathcal{R}_\beta^+(z) = (\omega \otimes \omega)\mathcal{R}_\beta^-(z) \quad \forall \beta \in Q_+,$$

and we have the following corollary of Theorem 9.6.

Corollary 9.9 For each $\beta \in Q_+$, the element \mathcal{R}_β^\pm satisfies

$$(\Phi_w \dot{\otimes} \Phi_z)(\mathcal{R}_\beta^\pm) \in (\hbar/z)^{v(\beta)} Y_\hbar(\mathfrak{g})^{\otimes 2} [w][[z^{-1}]]$$

in addition to the relation $(\Phi_w \dot{\otimes} \Phi_z)(\mathcal{R}_\beta^\pm) = \mathcal{R}_\beta^\pm(w - z)$.

Of course, the second assertion is immediate from Theorem 9.6 as Φ_w is the identity on \mathfrak{g} and thus satisfies $\pi_\alpha \circ \Phi_w = \Phi_w \circ \pi_\alpha|_{Y_{\hbar}\mathfrak{g}}$ for all $\alpha \in Q$. The first statement then follows from the properties of $\mathcal{R}^-(z)$ established in [17] and recalled in Section 3.6. However, we wish to point out that this assertion is a natural consequence of the above discussion on dual bases. Indeed, we have seen that any homogeneous basis of $Y_{\hbar}^{\pm}(\mathfrak{g})_{\beta}$ lies in $\hbar^{v(\beta)} Y_{\hbar}(\mathfrak{g})$, and that the image of its dual set under \check{i} is contained in $\bigoplus_{k \geq v(\beta)} DY_{\hbar}\mathfrak{g}_{-k}$. As Φ_z is graded, we have

$$\Phi_z(DY_{\hbar}\mathfrak{g}_{-k}) \subset z^{-k} \widehat{Y_{\hbar}\mathfrak{g}_z} \subset z^{-v(\beta)} Y_{\hbar}(\mathfrak{g})[[z^{-1}]] \quad \forall k \geq v(\beta),$$

which yields the first statement of the corollary.

A Homogenization of the R-matrix

In this appendix, we show that the results of [17, Section 7.4] imply Theorem 3.8, as promised in Remark 3.9. Let $Y(\mathfrak{g})$ denote the Yangian defined over \mathbb{C} with \hbar specialized to 1:

$$Y(\mathfrak{g}) := Y_{\hbar}(\mathfrak{g})/(\hbar - 1)Y_{\hbar}(\mathfrak{g}).$$

Slightly abusing notation, we shall denote the images of x_{ir}^{\pm} and h_{ir} again by x_{ir}^{\pm} and h_{ir} , respectively. The graded Hopf algebra structure on $Y_{\hbar}(\mathfrak{g})$ induces on $Y(\mathfrak{g})$ the structure of an \mathbb{N} -filtered Hopf algebra over the complex numbers with filtration \mathbb{F}_{\bullet} defined by letting \mathbb{F}_k denote the image of $\bigoplus_{n \leq k} Y_{\hbar}(\mathfrak{g})_k$. The Yangian $Y(\mathfrak{g})$ is then a filtered deformation of the graded Hopf algebra $U(\mathfrak{t}_+)$:

$$\text{gr}_{\mathbb{F}} Y(\mathfrak{g}) \cong U(\mathfrak{t}_+).$$

One can recover $Y_{\hbar}(\mathfrak{g})$ from $Y(\mathfrak{g})$ using the Rees algebra formalism (see [22, Proposition 2.2] and [23, Theorem 6.10], for example). In more detail, there is an isomorphism of \mathbb{N} -graded Hopf algebras

$$\begin{aligned} \varphi_{\hbar} : Y_{\hbar}(\mathfrak{g}) &\xrightarrow{\sim} R_{\hbar}(Y(\mathfrak{g})) = \bigoplus_{k \in \mathbb{N}} \hbar^k \mathbb{F}_k(Y(\mathfrak{g})) \subset Y(\mathfrak{g})[[\hbar]] \\ x_{ir}^{\pm} &\mapsto \hbar^r x_{ir}^{\pm}, \quad h_{ir} \mapsto \hbar^r h_{ir} \quad \forall i \in \mathbf{I}, r \in \mathbb{N}. \end{aligned}$$

Here, the Hopf algebra structure on $R_{\hbar}(Y(\mathfrak{g}))$ is obtained by extending that of $Y(\mathfrak{g})$ by $\mathbb{C}[[\hbar]]$ -linearity, and we note that $R_{\hbar}(Y(\mathfrak{g})) \otimes_{\mathbb{C}} Y(\mathfrak{g}) \cong R_{\hbar}(Y(\mathfrak{g})) \otimes_{\mathbb{C}[[\hbar]]} R_{\hbar}(Y(\mathfrak{g}))$.

In [17], the universal R -matrix $\mathbb{R}(z)$ of the Yangian $Y(\mathfrak{g})$ is constructed as a product

$$\mathbb{R}(z) = \mathbb{R}^+(z) \mathbb{R}^0(z) \mathbb{R}^-(z) \in 1 + z^{-1} Y(\mathfrak{g})^{\otimes 2}[[z^{-1}]],$$

where $\mathbb{R}^+(z) = \mathbb{R}_{21}^-(z)^{-1}$ and the factors $\mathbb{R}^-(z)$ and $\mathbb{R}^0(z)$ are as in Sections 4.1 and 6.6 of [17], respectively. In particular, $\mathbb{R}^{\pm}(z) - 1$, $\mathbb{R}^0(z) - 1$, and $\mathbb{R}(z) - 1$ lay in the subspace

$$z^{-1} \prod_{n \in \mathbb{N}} \mathbb{F}_n(Y(\mathfrak{g})^{\otimes 2}) z^{-n} \subset Y(\mathfrak{g})^{\otimes 2}[[z^{-1}]]$$

and hence $\mathbb{R}^\pm(z/\hbar)$, $\mathbb{R}^0(z/\hbar)$, and $\mathbb{R}(z/\hbar)$ are elements of $\mathbb{R}_\hbar(Y(\mathfrak{g}))^{\otimes 2}[[z^{-1}]]$. The definitions of $\mathcal{R}^\pm(z)$, $\mathcal{R}^0(z)$ and $\mathcal{R}(z)$ given in Section 3.6 are such that one has the equalities

$$(A.1) \quad \begin{aligned} \varphi_\hbar^{\otimes 2}(\mathcal{R}^\pm(z)) &= \mathbb{R}^\pm(z/\hbar), & \varphi_\hbar^{\otimes 2}(\mathcal{R}^0(z)) &= \mathbb{R}^0(z/\hbar), \\ \varphi_\hbar^{\otimes 2}(\mathcal{R}(z)) &= \mathbb{R}(z/\hbar). \end{aligned}$$

Using this fact and the results of [17], we can recover the below proposition, which is a restatement of Theorem 3.8.

Proposition A.1 $\mathcal{R}(z)$ is the unique formal series in $1 + z^{-1}Y_\hbar(\mathfrak{g})^{\otimes 2}[[z^{-1}]]$ satisfying the intertwiner equation

$$\tau_z \otimes \mathbf{1} \circ \Delta^{\text{op}}(x) = \mathcal{R}(z) \cdot \tau_z \otimes \mathbf{1} \circ \Delta(x) \cdot \mathcal{R}(z)^{-1} \quad \forall x \in Y_\hbar(\mathfrak{g})$$

in $Y_\hbar(\mathfrak{g})^{\otimes 2}[[z; z^{-1}]]$, in addition to the cabling identities

$$\begin{aligned} \Delta \otimes \mathbf{1}(\mathcal{R}(z)) &= \mathcal{R}_{13}(z)\mathcal{R}_{23}(z), \\ \mathbf{1} \otimes \Delta(\mathcal{R}(z)) &= \mathcal{R}_{13}(z)\mathcal{R}_{12}(z) \end{aligned}$$

in $Y_\hbar(\mathfrak{g})^{\otimes 3}[[z^{-1}]]$. Moreover, $\mathcal{R}(z)$ satisfies the properties (1)–(3) of Theorem 3.8.

Proof That $\mathcal{R}(z)$ satisfies the properties (1)–(3) of Theorem 3.8 follows from (A.1) and the corresponding properties of $\mathbb{R}(z)$ established in (3)–(5) of [17, Theorem 7.4]. Similarly, that $\mathcal{R}(z)$ satisfies the cabling identities in $Y_\hbar(\mathfrak{g})^{\otimes 3}[[z^{-1}]]$ follows from the equality $\varphi_\hbar^{\otimes 2}(\mathcal{R}(z)) = \mathbb{R}(z/\hbar)$ and Theorem 7.4(2) of [17], which asserts that $\mathbb{R}(z)$ satisfies the cabling identities in $Y(\mathfrak{g})^{\otimes 3}[[z^{-1}]]$. As for the intertwiner equation, upon applying the isomorphism $\varphi_\hbar^{\otimes 2}$, we deduce that it will hold provided $\mathbb{R}(z)$ satisfies

$$\tau_z^\varphi \otimes \mathbf{1} \circ \Delta_{Y(\mathfrak{g})}^{\text{op}}(x) = \mathbb{R}(z/\hbar) \cdot \tau_z^\varphi \otimes \mathbf{1} \circ \Delta_{Y(\mathfrak{g})}(x) \cdot \mathbb{R}(z/\hbar)^{-1} \quad \forall x \in \mathbb{R}_\hbar(Y(\mathfrak{g}))$$

in $\mathbb{R}_\hbar(Y(\mathfrak{g}))^{\otimes 2}[[z; z^{-1}]] \subset (Y(\mathfrak{g})^{\otimes 2})[[\hbar]][[z; z^{-1}]]$, where $\tau_z^\varphi = \varphi_\hbar \circ \tau_z \circ \varphi_\hbar^{-1}$. Since τ_z^φ is determined by

$$\tau_z^\varphi(x_i^\pm(u/\hbar)) = x_i^\pm((u - z)/\hbar), \quad \tau_z^\varphi(h_i(u/\hbar)) = h_i((u - z)/\hbar) \quad \forall i \in \mathbf{I},$$

the above equality will hold provided that, for each $x \in Y(\mathfrak{g})$, and $\zeta \in \mathbb{C}^\times$, one has

$$(A.2) \quad \mathring{\tau}_{z/\zeta} \otimes \mathbf{1} \circ \Delta_{Y(\mathfrak{g})}^{\text{op}}(x) = \mathbb{R}(z/\zeta) \cdot \mathring{\tau}_{z/\zeta} \otimes \mathbf{1} \circ \Delta_{Y(\mathfrak{g})}(x) \cdot \mathbb{R}(z/\zeta)^{-1},$$

where $\mathring{\tau}_z : Y(\mathfrak{g}) \rightarrow Y(\mathfrak{g})[[z]]$ is obtained by specializing the algebra homomorphism τ_z defined in (3.8). This equality is immediate from Part (1) of [17, Theorem 7.4].

As for the uniqueness assertion; the argument given in Appendix B of [17] translates naturally to the formal setting. Alternatively, one can see this as a consequence of the uniqueness of $\mathbb{R}(z)$ itself, as proven therein. Indeed, if $\mathcal{R}(z) \in 1 + z^{-1}Y_\hbar(\mathfrak{g})^{\otimes 2}[[z^{-1}]]$ is another solution of the intertwiner equation satisfying the cabling identities, then to see that $\mathcal{R}(z) = \mathbb{R}(z)$, it suffices to show that $\mathbb{X}(z)|_{\hbar=\zeta} = \mathbb{R}(z/\zeta)$ for each $\zeta \in \mathbb{C}^\times$, where

$$\mathbb{X}(z) := \varphi_\hbar^{\otimes 2}(\mathcal{R}(z)) \in Y(\mathfrak{g})[[\hbar]][[z^{-1}]].$$

This follows from the fact that $\mathbb{X}(z)|_{\hbar=\zeta}$ and $\mathbb{R}(z/\zeta)$ both satisfy the cabling identities in $Y(\mathfrak{g})^{\otimes 3}[[z^{-1}]]$ and the intertwiner equation (A.2), and so coincide by the uniqueness of $\mathbb{R}(z)$, as established in Appendix B of [17]. ■

Acknowledgments I would like to thank Alex Weekes, Andrea Appel, Sachin Gautam, and Valerio Toledano Laredo for the many insightful discussions and helpful comments they have been the source of over the last few years. These have played a significant role in shaping this article.

References

- [1] A. Appel and V. Toledano Laredo, *A 2-categorical extension of Etingof–Kazhdan quantisation*. *Selecta Math. (N.S.)* 24(2018), no. 4, 3529–3617.
- [2] A. Appel and V. Toledano Laredo, *Coxeter categories and quantum groups*. *Selecta Math. (N.S.)* 25(2019), no. 3, Article no. 44, 97 pp.
- [3] V. Chari and A. Pressley, *Fundamental representations of Yangians and singularities of R-matrices*. *J. Reine Angew. Math.* 417(1991), 87–128.
- [4] V. Chari and A. Pressley, *A guide to quantum groups*, Cambridge University Press, Cambridge, 1994.
- [5] I. Damiani, *La R-matrice pour les algèbres quantiques de type affine non tordu*. *Ann. Sci. École Norm. Sup.* 31(1998), no. 4, 493–523.
- [6] V. Drinfeld, *Hopf algebras and the quantum Yang–Baxter equation*. *Soviet Math. Dokl.* 32(1985), no. 1, 254–258.
- [7] V. Drinfeld, *Quantum groups*. In: *Proceedings of the international congress of mathematicians*, vol. 1, 2 (Berkeley, Calif., 1986), American Mathematical Society, Providence, RI, 1987, pp. 798–820.
- [8] V. Drinfeld, *A new realization of Yangians and quantum affine algebras*. *Soviet Math. Dokl.* 36(1988), no. 2, 212–216.
- [9] B. Enriquez and G. Halbout, *An \hbar -adic valuation property of universal R-matrices*. *J. Algebra* 261(2003), no. 2, 434–447.
- [10] P. Etingof, I. Frenkel, and A. Kirillov, *Lectures on representation theory and Knizhnik–Zamolodchikov equations*, *Mathematical Surveys and Monographs*, 58, American Mathematical Society, Providence, RI, 1998.
- [11] P. Etingof and D. Kazhdan, *Quantization of Lie bialgebras. I*. *Selecta Math. (N.S.)* 2(1996), no. 1, 1–41.
- [12] P. Etingof and D. Kazhdan, *Quantization of Lie bialgebras. III*. *Selecta Math. (N.S.)* 4(1998), no. 2, 233–269.
- [13] M. Finkelberg and A. Tsymbaliuk, *Shifted quantum affine algebras: integral forms in type A*. *Arnold Math. J.* 5(2019), nos. 2–3, 197–283.
- [14] I. Frenkel and N. Reshetikhin, *Quantum affine algebras and holonomic difference equations*. *Commun. Math. Phys.* 146(1992), no. 1, 1–60.
- [15] S. Gautam and V. Toledano Laredo, *Yangians and quantum loop algebras*. *Selecta Math. (N.S.)* 19(2013), no. 2, 271–336.
- [16] S. Gautam and V. Toledano Laredo, *Meromorphic tensor equivalence for Yangians and quantum loop algebras*. *Publ. Math. Inst. Hautes Études Sci.* 125(2017), 267–337.
- [17] S. Gautam, V. Toledano Laredo, and C. Wendlandt, *The meromorphic R-matrix of the Yangian*. In: A. Alekseev, E. Frenkel, M. Rosso, B. Webster, and M. Yakimov (eds.), *Representation theory, mathematical physics, and integrable systems*, Honor of Nicolai Reshetikhin, *Progress in Mathematics*, 340, Birkhäuser/Springer, Cham, 2021, pp. 201–269.
- [18] S. Gautam and C. Wendlandt, *Poles of finite-dimensional representations of Yangians*. *Selecta Math. (N.S.)* 29(2023), no. 1, Article no. 13, 68 pp.
- [19] F. Gavarini, *The quantum duality principle*. *Ann. Inst. Fourier (Grenoble)* 52(2002), no. 3, 809–834.
- [20] F. Gavarini, *The global quantum duality principle*. *J. Reine Angew. Math.* 612(2007), 17–33.
- [21] N. Guay, H. Nakajima, and C. Wendlandt, *Coproduct for Yangians of affine Kac–Moody algebras*. *Adv. Math.* 338(2018), 865–911.

- [22] N. Guay, V. Regelskis, and C. Wendlandt, *Equivalences between three presentations of orthogonal and symplectic Yangians*. Lett. Math. Phys. **109**(2019), no. 2, 327–379.
- [23] N. Guay, V. Regelskis, and C. Wendlandt, *Vertex representations for Yangians of Kac–Moody algebras*. J. Éc. Polytech. Math. **6**(2019), 665–706.
- [24] V. Kac, *Infinite-dimensional Lie algebras*, 3rd ed., Cambridge University Press, Cambridge, 1990.
- [25] J. Kamnitzer, B. Webster, A. Weekes, and O. Yacobi, *Yangians and quantizations of slices in the affine Grassmannian*. Algebra Number Theory **8**(2014), no. 4, 857–893.
- [26] C. Kassel, *Quantum groups*, Graduate Texts in Mathematics, 155, Springer, New York, 1995.
- [27] C. Kassel and V. Turaev, *Biquantization of Lie bialgebras*. Pacific J. Math. **195**(2000), no. 2, 297–369.
- [28] S. Khoroshkin, A. Stolin, and V. Tolstoy, *Generalized gauss decomposition of trigonometric R-matrices*. Modern Phys. Lett. A **10**(1995), no. 19, 1375–1392.
- [29] S. Khoroshkin and V. Tolstoy, *Universal R-matrix for quantized (super)algebras*. Commun. Math. Phys. **141**(1991), no. 3, 599–617.
- [30] S. Khoroshkin and V. Tolstoy, *The Cartan–Weyl basis and the universal R-matrix for quantum Kac–Moody algebras and superalgebras*. In: H.-D. Doebner and V. Dobrev (eds.), Proceedings of the international workshop on mathematical physics (Clausthal, 1991), Quantum Symmetries, World Scientific, River Edge, NJ, 1993, pp. 336–351.
- [31] S. Khoroshkin and V. Tolstoy, *Twisting of quantum (super-) algebras*. In: H.-D. Doebner, V. Dobrev, and A. Ushveridze (eds.), Proceedings of the international symposium on mathematical physics (Arnold Sommerfeld Institute, Clausthal, 1993), Generalized Symmetries in Physics, World Scientific, River Edge, NJ, 1994, pp. 42–54.
- [32] S. Khoroshkin and V. Tolstoy, *Yangian double*. Lett. Math. Phys. **36**(1996), no. 4, 373–402.
- [33] A. Klimyk and K. Schmüdgen, *Quantum groups and their representations*, Texts and Monographs in Physics, Springer, Berlin, 1997.
- [34] S. Majid, *Physics for algebraists: noncommutative and noncocommutative Hopf algebras by a bicrossproduct construction*. J. Algebra **130**(1990), no. 1, 17–64.
- [35] S. Majid, *Some remarks on the quantum double*. Czechoslov. J. Phys. **44**(1994), 1059–1071.
- [36] S. Majid, *Foundations of quantum group theory*, Cambridge University Press, Cambridge, 1995.
- [37] S. Montgomery, *Hopf algebras and their actions on rings*, CBMS Regional Conference Series in Mathematics, 82, published for the Conference Board of the Mathematical Sciences, Washington, DC, American Mathematical Society, Providence, RI, 1993.
- [38] M. Nazarov, *Double Yangian and the universal R-matrix*. Japan J. Math. **15**(2020), no. 1, 169–221.
- [39] D. Radford, *Minimal quasitriangular Hopf algebras*. J. Algebra **157**(1993), no. 2, 285–315.
- [40] V. Stukopin, *The quantum double of the Yangian of a Lie superalgebra of type $a(m,n)$ and the computation of the universal R-matrix*. Fundam. Prikl. Mat. **11**(2005), no. 2, 185–208.
- [41] C. Wendlandt, *The formal shift operator on the Yangian double*. Int. Math. Res. Not. IMRN **2022**(2021), no. 14, 10952–11010.
- [42] Y. Yang and G. Zhao, *Cohomological Hall algebras and affine quantum groups*. Selecta Math. (N.S.) **24**(2018), no. 2, 1093–1119.

Department of Mathematics and Statistics, University of Saskatchewan, Saskatoon, SK, Canada
e-mail: wendlandt@math.usask.ca