

ON FINE SELMER GROUPS AND SIGNED SELMER GROUPS OF ELLIPTIC MODULAR FORMS

ANTONIO LEI  and MENG FAI LIM 

(Received 17 August 2021; accepted 19 October 2021; first published online 19 April 2022)

Abstract

Let f be an elliptic modular form and p an odd prime that is coprime to the level of f . We study the link between divisors of the characteristic ideal of the p -primary fine Selmer group of f over the cyclotomic \mathbb{Z}_p extension of \mathbb{Q} and the greatest common divisor of signed Selmer groups attached to f defined using the theory of Wach modules. One of the key ingredients of our proof is a generalisation of a result of Wingberg on the structure of fine Selmer groups of abelian varieties with supersingular reduction at p to the context of modular forms.

2020 *Mathematics subject classification*: primary 11R23; secondary 11F11, 11R18.

Keywords and phrases: elliptic modular forms, fine Selmer groups, signed Selmer groups.

1. Introduction

Let p be a fixed odd prime number and f a fixed normalised eigen-cusppform of level N and weight $k \geq 2$ with $p \nmid N$. Let K denote the completion of the Hecke field of f at a prime above p and write \mathcal{O} for its ring of integers. Let V_f denote the K -adic $G_{\mathbb{Q}}$ -representation of f defined by Deligne [8]. We fix a Galois-stable \mathcal{O} -lattice T_f inside V_f and write $A_f = V_f/T_f$. We write \bar{f} for the conjugate modular form of f and we write $V_{\bar{f}}$, $T_{\bar{f}}$ and $A_{\bar{f}}$ for the corresponding $G_{\mathbb{Q}}$ -modules attached to \bar{f} with $T_{\bar{f}}$ chosen to be $\text{Hom}_{\mathcal{O}}(T_f, \mathcal{O})(1-k)$, where $M(j)$ denotes the j th Tate twist of a $G_{\mathbb{Q}}$ -module M .

Let \mathbb{Q}_{cyc} be the cyclotomic \mathbb{Z}_p -extension of \mathbb{Q} and let Γ denote the Galois group $\text{Gal}(\mathbb{Q}_{\text{cyc}}/\mathbb{Q})$. The Iwasawa algebra $\Lambda = \mathcal{O}[[\Gamma]]$ is defined to be $\varprojlim \mathcal{O}[\Gamma/\Gamma^{p^n}]$, where the connecting maps are projections. After fixing a topological generator γ of Γ , there is an isomorphism of rings $\Lambda \cong \mathcal{O}[[X]]$ sending γ to $X+1$. Given a Λ -module M , denote its Pontryagin dual by $M^{\vee} := \text{Hom}_{\mathcal{O}}(M, K/\mathcal{O})$. We write M^t for the Λ -module that is M as an \mathcal{O} -module equipped with a Γ -action given by $\gamma \cdot m = \gamma^{-1}m$. Finally, if $F \in \Lambda = \mathcal{O}[[X]]$, we write F^t for the power series $F(1/(1+X) - 1)$.

The authors' research is partially supported by the NSERC Discovery Grants Program RGPIN-2020-04259 and RGPAS-2020-00096 (Lei), and the National Natural Science Foundation of China under Grant No. 11771164 and the Fundamental Research Funds for the Central Universities of CCNU under Grant CCNU20TD002 (Lim).

© The Author(s), 2022. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

For $g = f$ or \bar{f} and $j \in \mathbb{Z}$, let $\text{Sel}_0(A_g(j)/\mathbb{Q}_{\text{cyc}})$ denote the fine Selmer group of $A_g(j)$ over \mathbb{Q}_{cyc} (whose definition will be reviewed in Section 2.1). The fine Selmer groups of abelian varieties were first systematically studied by Coates and Sujatha in [7] and a little later by Wuthrich [29]. Various conjectures on the structure of these groups have been formulated and they are still wide open to this date.

It has been proved by Kato [12] that $\text{Sel}_0(A_g(j)/\mathbb{Q}_{\text{cyc}})^\vee$ is a finitely generated torsion Λ -module. Let $\mathfrak{F}_{g,j}$ denote a choice of characteristic element (that is, a generator of the characteristic ideal $\text{Char}_\Lambda \text{Sel}_0(A_g(j)/\mathbb{Q}_{\text{cyc}})^\vee$). The main goal of the present article is to study the divisors of $\mathfrak{F}_{f,i}$ and $\mathfrak{F}_{\bar{f},k-i}$ for a fixed integer i . These specific twists are considered due to the perfect pairing

$$T_f(i) \times A_{\bar{f}}(k-i) \rightarrow \mu_{p^\infty}$$

of $G_{\mathbb{Q}}$ -modules.

We are able to relate the divisors of $\mathfrak{F}_{f,i}$ and $\mathfrak{F}_{\bar{f},k-i}$ to the greatest common divisors of the signed Selmer groups attached to $\bar{f}(k-i)$ defined in [16, 17] by the theory of Wach modules (see Section 2.3, where the definitions of these groups are reviewed). Let us write $\mathfrak{F}_{\bar{f},k-i}^\sharp$ and $\mathfrak{F}_{\bar{f},k-i}^\flat$ for a choice of characteristic elements of the Pontryagin duals of these signed Selmer groups. Under a mild hypothesis on the local representations at bad primes and a hypothesis on the validity of Kato's Iwasawa main conjecture for $\bar{f}(k-i)$ (respectively labelled **(H0)** and **(H-IMC)** below), we show that if F is an irreducible element of Λ that is outside a certain explicit set of linear factors, $F \nmid \mathfrak{F}_{f,i}$ and $F \nmid \mathfrak{F}_{\bar{f},k-i}$ if and only if $F \nmid \gcd(\mathfrak{F}_{\bar{f},k-i}^\sharp, \mathfrak{F}_{\bar{f},k-i}^\flat)$. (See Theorem 4.4 for the precise statement of this result.)

In the case where T_f is the Tate module of an elliptic curve E/\mathbb{Q} with good supersingular reduction at p , a similar result has been proved in [18]. One of the key ingredients of the proof given in [18] is a link between the fine Selmer group of E over \mathbb{Q}_{cyc} and the maximal Λ -torsion submodule of the Pontryagin dual of the p -primary Selmer group of E over \mathbb{Q}_{cyc} that was proved by Wingberg [28] (see also Matar [19], where an alternative proof is given). In the present paper, we prove Theorem 4.4 by first establishing analogues of Wingberg's result in the context of modular forms (see Theorems 3.1 and 3.4). Wingberg worked with Selmer groups defined using flat cohomology, whereas Matar worked with Selmer groups defined using Galois cohomology. In the present paper, the latter definition is used. Our proof is very different from the ones employed in both [19] and [28]. We make use of Nekovář's spectral sequence that seems to give a somewhat simpler and more general proof than the previous proofs available in the literature. We hope that these results may be of independent interest.

2. Definitions of Selmer groups and related objects

2.1. Fine Selmer groups. Let L_n be the unique subextension of \mathbb{Q}_{cyc} such that $[L_n : \mathbb{Q}] = p^n$. Given an algebraic extension L of \mathbb{Q} , we write $S(L)$ for the primes of

L lying above pN as well as the archimedean primes. We write $G_S(L)$ for the Galois group of the maximal algebraic extension of L that is unramified outside $S(L)$.

For $g \in \{f, \bar{f}\}$ and $j \in \mathbb{Z}$, we define the fine Selmer group of $A_g(j)$ over L to be

$$\text{Sel}_0(A_g(j)/L) = \ker \left(H^1(G_S(L), A_g(j)) \rightarrow \prod_{v \in S(L)} H^1(L_v, A_g(j)) \right).$$

Recall that $\text{Sel}_0(A_g(j)/\mathbb{Q}_{\text{cyc}})^\vee$ is torsion over Λ ; we write $\mathfrak{F}_{g,j}$ for a choice of a generator of the characteristic ideal $\text{Char}_\Lambda \text{Sel}_0(A_g(j)/\mathbb{Q}_{\text{cyc}})^\vee$.

The classical Selmer group of $A_g(j)$ over L is defined to be

$$\text{Sel}(A_g(j)/L) = \ker \left(H^1(G_S(L), A_g(j)) \rightarrow \prod_{v \in S(L)} \frac{H^1(L_v, A_g(j))}{H^1_f(L_v, A_g(j))} \right),$$

where $H^1_f(L_v, A_g(j))$ is defined as in [4, Section 3].

2.2. Wach modules and signed Coleman maps. We recall the definition of signed Coleman maps from [16, 17] that generalise those studied in [13, 27] in the context of elliptic curves with supersingular reduction at p . Here, we do not require p to be a nonordinary prime for f . The only requirement is that p is coprime to N , so that the representation V_f is crystalline at p .

For $g \in \{f, \bar{f}\}$, $j \in \mathbb{Z}$ and $m \in \{1, 2\}$, we shall write

$$\mathfrak{H}_{g,j}^m = \varprojlim_n H^m(L_n, T_g(j)), \quad \mathbb{H}_{g,j}^m = \varprojlim H^m(L_{n,p}, T_g(j)),$$

where the connecting maps are corestrictions and we have abused notation by writing p for the unique prime of L_n above p . We write $\text{loc}_p : \mathfrak{H}_{g,j}^1 \rightarrow \mathbb{H}_{g,j}^1$ for the localisation map.

Let $T = T_g(k-1)$, where $g \in \{f, \bar{f}\}$. We write $\mathbb{N}(T)$ for the Wach module of T . (See [3, Section II.1] for the precise definition of $\mathbb{N}(T)$.) Roughly speaking, $\mathbb{N}(T)$ is a filtered φ -module over the ring $\mathcal{O}[[\pi]]$, where π is an element in the ring of Witt vectors of $\varprojlim_{\chi \mapsto \chi^p} \mathbb{C}_p$, given by $[(1, \zeta_p, \zeta_{p^2}, \dots)] - 1$, and ζ_{p^n} is a primitive p^n th root of unity in \mathbb{C}_p such that $\zeta_{p^{n+1}}^p = \zeta_{p^n}$. One may regard π as a formal variable equipped with an action of φ and $\Gamma_0 = \text{Gal}(\mathbb{Q}_p(\mu_{p^\infty})/\mathbb{Q}_p)$ given by $\varphi(\pi) = (1 + \pi)^p - 1$ and $\sigma(\pi) = (1 + \pi)^{\chi_{\text{cyc}}(\sigma)} - 1$ for $\sigma \in \Gamma_0$, where χ_{cyc} is the p -adic cyclotomic character. Let $H_{\text{Iw}}^1(\mathbb{Q}_p(\mu_{p^\infty}), T) = \varprojlim_n H^1(\mathbb{Q}_p(\mu_{p^n}), T)$, where the connecting maps are corestrictions. Berger proved in [2, Appendix A] that there is an isomorphism of $\mathcal{O}[[\Gamma_0]]$ -modules

$$H_{\text{Iw}}^1(\mathbb{Q}_p(\mu_{p^\infty}), T) \cong \mathbb{N}(T)^{\psi=1},$$

where ψ is a canonical left inverse of φ and the superscript $\psi = 1$ denotes the kernel of $\psi - 1$.

We recall from [16, 17] that for $\bullet \in \{\sharp, \flat\}$, on choosing an appropriate basis of $\mathbb{N}(T_g(k-1))$, one can define signed Coleman maps

$$\mathrm{Col}_g^\bullet : H_{\mathrm{Iw}}^1(\mathbb{Q}_p(\mu_{p^\infty}), T_g(k-1)) \rightarrow \mathcal{O}[\Gamma_0],$$

decomposing Perrin-Riou's big logarithm map defined in [22].

We define the twisted Coleman maps

$$\mathrm{Col}_{g,j}^\bullet : \mathbb{H}_{g,j}^1 \rightarrow \Lambda$$

as follows. We have the \mathcal{O} -isomorphism

$$H_{\mathrm{Iw}}^1(\mathbb{Q}_p(\mu_{p^\infty}), T_g(j)) \xrightarrow{e^{k-1-j}} H_{\mathrm{Iw}}^1(\mathbb{Q}_p(\mu_{p^\infty}), T_g(k-1)),$$

where e is a basis of the $G_{\mathbb{Q}_p}$ -representation $\mathbb{Z}_p(1)$ as given in [23, Section A.4]. Let Tw be the \mathcal{O} -linear automorphism on $\mathcal{O}[\Gamma_0]$ sending $\sigma \in \Gamma_0$ to $\chi_{\mathrm{cyc}}(\sigma)\sigma$. We can define a Λ -morphism

$$\mathrm{Tw}^{i-k+1} \circ \mathrm{Col}_g^\bullet \circ e^{k-1-j} : H_{\mathrm{Iw}}^1(\mathbb{Q}_p(\mu_{p^\infty}), T_g(j)) \rightarrow \mathcal{O}[\Gamma_0]$$

and obtain $\mathrm{Col}_{g,j}^\bullet$ on taking the trivial isotypic component of $\Delta := \mathrm{Gal}(\mathbb{Q}_p(\mu_p)/\mathbb{Q}_p)$.

2.3. Signed Selmer groups and main conjectures. Let $H_\bullet^1(\mathbb{Q}_{\mathrm{cyc},p}, A_{\bar{f}}(k-i))$ denote the orthogonal complement of $\ker \mathrm{Col}_{f,i}^\bullet$ under the local Tate pairing

$$H_{\mathrm{Iw}}^1(\mathbb{Q}_p(\mu_{p^\infty}), T_f(i)) \times H^1(\mathbb{Q}_{\mathrm{cyc},p}, A_{\bar{f}}(k-i)) \rightarrow \mathbb{Q}_p/\mathbb{Z}_p.$$

We define the signed Selmer groups $\mathrm{Sel}^\bullet(A_{\bar{f}}(k-i)/\mathbb{Q}_{\mathrm{cyc}})$ as

$$\ker \left(H^1(G_S(\mathbb{Q}_{\mathrm{cyc}}), A_{\bar{f}}(k-i)) \rightarrow \frac{H^1(\mathbb{Q}_{\mathrm{cyc},p}, A_{\bar{f}}(k-i))}{H_\bullet^1(\mathbb{Q}_{\mathrm{cyc},p}, A_{\bar{f}}(k-i))} \times \prod_{v \nmid p} \frac{H^1(\mathbb{Q}_{\mathrm{cyc},v}, A_{\bar{f}}(k-i))}{H_{\mathrm{f}}^1(\mathbb{Q}_{\mathrm{cyc},v}, A_{\bar{f}}(k-i))} \right).$$

Note that when $A_{\bar{f}}(k-i)$ is given by $E[p^\infty]$ for some elliptic curve E/\mathbb{Q} with good supersingular reduction at p or when $a_p(\bar{f}) = 0$, one may choose an appropriate basis of the Wach module so that these Selmer groups coincide with the ones studied in [13, 14, 27] (see [16, Sections 5.2–5.4]).

Let $\mathbf{z}_{f,i}$ be Kato's zeta element inside $H_{\mathrm{Iw}}^1(\mathbb{Q}_{\mathrm{cyc}}, T_f(i)) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$ as defined in [12]. It is conjectured that $\mathbf{z}_{f,i} \in H_{\mathrm{Iw}}^1(\mathbb{Q}_{\mathrm{cyc}}, T_f(i))$ (see [12, Conjecture 12.10]). Assuming this (which we shall do for the rest of the article), recall from [16, Equation (61)] the Poitou–Tate exact sequence

$$0 \rightarrow \frac{\mathfrak{H}_{f,i}^1}{\Lambda \mathbf{z}_{f,i}} \rightarrow \frac{\mathrm{Image}(\mathrm{Col}_{f,i}^\bullet)}{(L_p^\bullet)} \rightarrow \mathrm{Sel}^\bullet(A_{\bar{f}}(k-i)/\mathbb{Q}_{\mathrm{cyc}})^\vee \rightarrow \mathrm{Sel}_0(A_{\bar{f}}(k-i)/\mathbb{Q}_{\mathrm{cyc}})^\vee \rightarrow 0, \quad (2.1)$$

where $L_p^\bullet = \mathrm{Col}_{f,i}^\bullet \circ \mathrm{loc}_p(\mathbf{z}_{f,i})$ is the signed p -adic L -function associated to $\bar{f}(k-i)$. If $L_p^\bullet \neq 0$, then $\mathrm{Sel}^\bullet(A_{\bar{f}}(k-i)/\mathbb{Q}_{\mathrm{cyc}})^\vee$ is a finitely generated torsion Λ -module. We write $\mathfrak{F}_{\bar{f},k-i}^\bullet$ for a characteristic element of this module.

Kato [12] proved that the first and fourth terms of (2.1) are Λ -torsion. He further formulated the Iwasawa main conjecture:

$$\mathrm{Char}_{\Lambda} \mathfrak{S}_{f,i}^1 / \Lambda \mathbf{z}_{f,i} = \mathrm{Char}_{\Lambda} \mathrm{Sel}_0(A_{\overline{f}}(k-i)/\mathbb{Q}_{\mathrm{cyc}})^{\vee} = (\mathfrak{F}_{\overline{f},k-i}). \quad (2.2)$$

It follows from (2.1) that this is equivalent to

$$\mathrm{Char}_{\Lambda} \mathrm{Sel}^{\bullet}(A_{\overline{f}}(k-i)/\mathbb{Q}_{\mathrm{cyc}})^{\vee} = \mathrm{Char}_{\Lambda} \mathrm{Image}(\mathrm{Col}_{f,i}^{\bullet}) / (L_p^{\bullet}), \quad (2.3)$$

provided that $L_p^{\bullet} \neq 0$.

For the rest of the article, we shall assume that the following hypothesis holds.

(H-IMC) Kato's Iwasawa main conjecture (2.2) holds and both L_p^{\sharp} and L_p^{\flat} are nonzero.

REMARK 2.1. We say a few words on the hypothesis **(H-IMC)**. In his seminal work [12], Kato established that the inclusion ' \subseteq ' in (2.2) holds after tensoring by \mathbb{Q}_p under the assumption that the image of the representation $T_f|_{G_{\mathbb{Q}(\mu_{p^{\infty}})}}$ contains a copy of $\mathrm{SL}_2(\mathbb{Z}_p)$, provided that certain local terms vanish (see, in particular, [12, Theorem 12.5(4)]). Note that in [12, Conjecture 12.10], Kato's main conjecture is formulated in terms of étale cohomology groups. Kobayashi and Kurihara showed that it can be recast in terms of Galois cohomology groups and fine Selmer groups (see, for example, [13, Proposition 7.1]). The reverse inclusion of (2.2) has been established in the monumental work of Skinner and Urban [26] for a p -ordinary modular form (under certain hypotheses; see [26, Theorem 1]). In the nonordinary form, there have been several recent breakthroughs in this reverse direction (see [5, 6, 9]). Finally, the work [16] has supplied many sufficient conditions for the nonvanishing of L_p^{\sharp} and L_p^{\flat} (see [16, Corollary 3.29 and Proposition 3.39]). In view of these developments, it seems reasonable to assume hypothesis **(H-IMC)**; our main results rely on it.

2.4. Images of signed Coleman maps. We review an explicit description of the images of these Coleman maps. Recall that γ is a fixed topological generator of Γ giving the identification $\Lambda = \mathcal{O}[[X]]$ via $X = \gamma - 1$. Let $u = \chi_{\mathrm{cyc}}(\gamma)$. From [17, Section 5A], if η is a character on Δ , then there exist constants $c_{\eta,j} \in K$ such that

$$e_{\eta} \mathrm{Image}(\mathrm{Col}_f^{\sharp} \oplus \mathrm{Col}_f^{\flat}) \otimes_{\mathcal{O}} K = \{(F, G) \in \Lambda^2 : F(u^j - 1) = c_{\eta,j} G(u^j - 1), 0 \leq j \leq k-2\}.$$

(In [17], it is assumed that f is nonordinary at p , meaning that $a_p(f)$ is a nonunit in \mathcal{O} . But the same calculations still apply to the ordinary case. See [17, Remark 1.10].) In particular, this tells us that there is an isomorphism of Λ -modules

$$\frac{\Lambda^2 \otimes_{\mathcal{O}} K}{e_{\eta} \mathrm{Image}(\mathrm{Col}_f^{\sharp} \oplus \mathrm{Col}_f^{\flat}) \otimes_{\mathcal{O}} K} \cong \frac{\Lambda \otimes_{\mathcal{O}} K}{\prod_{j=0}^{k-2} (X - u^j + 1)},$$

where e_{η} denotes the idempotent attached to η .

LEMMA 2.2. *There is a pseudo-isomorphism of Λ -modules*

$$\frac{\Lambda^2}{e_\eta \text{Image}(\text{Col}_f^\# \oplus \text{Col}_f^b)} \sim \frac{\Lambda}{\prod_{j=0}^{k-2} (X - u^j + 1)}.$$

PROOF. It is enough to show that the μ -invariant of the quotient on the left-hand side vanishes. For simplicity, let us write

$$C = e_\eta \text{Image}(\text{Col}_f^\# \oplus \text{Col}_f^b) \quad \text{and} \quad C_\bullet = e_\eta \text{Image} \text{Col}_f^\bullet \quad \text{for } \bullet \in \{\#, b\}.$$

Consider the tautological short exact sequence $0 \rightarrow C \rightarrow \Lambda^2 \rightarrow \Lambda^2/C \rightarrow 0$. This gives the exact sequence

$$0 \rightarrow (\Lambda^2/C)[\varpi] \rightarrow C/\varpi \xrightarrow{\Phi} \Lambda^2/\varpi,$$

where ϖ is a fixed uniformiser of \mathcal{O} . Similarly, we have the exact sequences

$$0 \rightarrow (\Lambda/C_\bullet)[\varpi] \rightarrow C_\bullet/\varpi \xrightarrow{\Phi_\bullet} \Lambda/\varpi \quad \text{for } \bullet \in \{\#, b\}.$$

Recall from [17, Theorem 5.10] that the μ -invariants of Λ/C_\bullet are zero. Therefore, $\ker \Phi_\bullet = (\Lambda/C_\bullet)[\varpi]$ is finite for both choices of \bullet . Note that $C \subset C_\# \oplus C_b$ and $\Phi = (\Phi_\# \oplus \Phi_b)|_{C/\varpi}$ by definition, which implies that

$$\ker \Phi \subset \ker \Phi_\# \oplus \ker \Phi_b.$$

Hence, $\ker \Phi = (\Lambda^2/C)[\varpi]$ is finite. In particular, the μ -invariant of Λ^2/C is zero, which finishes the proof of the lemma. \square

REMARK 2.3. In the case where T_f is the p -adic Tate module of an elliptic curve with good supersingular reduction at p , we can in fact describe the set C explicitly. See [15, Proposition 2.2].

It follows from Lemma 2.2 that there is an exact sequence

$$0 \rightarrow \mathbb{H}_{f,i}^1 \rightarrow \Lambda^2 \rightarrow C_{k,i} \rightarrow 0, \quad (2.4)$$

where the first map is given by $\text{Col}_{f,i}^\# \oplus \text{Col}_{f,i}^b$ and $C_{k,i}$ is a Λ -module, which is pseudo-isomorphic to $\Lambda/\eta_{k,i}$, with $\eta_{k,i}$ being the image of $\prod_{j=0}^{k-2} (X - u^j + 1)$ under the twisting map Tw^{i-k+1} .

3. A generalisation of a result of Wingberg

Let \mathcal{A} be an abelian variety over \mathbb{Q} with supersingular reduction at p . Wingberg proved that there is a pseudo-isomorphism of Λ -modules

$$(\text{Sel}(\mathcal{A}[p^\infty]/\mathbb{Q}_{\text{cyc}})^\vee)_{\Lambda\text{-tor}} \sim (\text{Sel}_0(\mathcal{A}^t[p^\infty]/\mathbb{Q}_{\text{cyc}})^\vee)^\iota,$$

where \mathcal{A}^t is the dual abelian variety of \mathcal{A} (see [28, Corollary 2.5]). Here, $M_{\Lambda\text{-tor}}$ denotes the maximal torsion submodule of a Λ -module M , $\text{Sel}(\mathcal{A}[p^\infty]/\mathbb{Q}_{\text{cyc}})$ is the p -primary Selmer group of \mathcal{A} over \mathbb{Q}_{cyc} and $\text{Sel}_0(\mathcal{A}^t[p^\infty]/\mathbb{Q}_{\text{cyc}})$ is the p -primary fine

Selmer group of \mathcal{A}' over \mathbb{Q}_{cyc} defined in a similar manner to the fine Selmer groups for $A_g(j)$ in Section 2.1 above. We prove the following analogue of Wingberg's result in the context of modular forms.

THEOREM 3.1. *We have a pseudo-isomorphism of Λ -modules*

$$(H^1(\mathbb{Q}_{\text{cyc}}, A_g(j))^\vee)_{\Lambda\text{-tor}} \sim (\mathfrak{S}_{g,j}^2)^\iota,$$

where $g \in \{f, \bar{f}\}$ and $j \in \mathbb{Z}$.

PROOF. From the low-degree terms of Nekovář's spectral sequence (see [20, Lemma 9.1.5]),

$$\text{Ext}'_\Lambda(H_{\text{Iw}}^{3-s}(\mathbb{Q}_{\text{cyc}}, T_g(j)), \Lambda) \Rightarrow H^{3-r-s}(\mathbb{Q}_{\text{cyc}}, A_g(j))^\vee,$$

we obtain the exact sequence of Λ -modules

$$0 \longrightarrow \text{Ext}'_\Lambda(H_{\text{Iw}}^2(\mathbb{Q}_{\text{cyc}}, T_g(j)), \Lambda) \longrightarrow H^1(\mathbb{Q}_{\text{cyc}}, A_g(j))^\vee \longrightarrow \text{Hom}_\Lambda(H_{\text{Iw}}^1(\mathbb{Q}_{\text{cyc}}, T_g(j)), \Lambda).$$

Since $\text{Hom}_\Lambda(H_{\text{Iw}}^1(\mathbb{Q}_{\text{cyc}}, T_g(j)), \Lambda)$ is a reflexive Λ -module by [21, Corollary 5.1.3], it is Λ -torsion-free. Hence, it follows from the exact sequence that there is an isomorphism

$$(H^1(\mathbb{Q}_{\text{cyc}}, A_g(j))^\vee)_{\Lambda\text{-tor}} \cong \text{Ext}'_\Lambda(H_{\text{Iw}}^2(\mathbb{Q}_{\text{cyc}}, T_g(j)), \Lambda)$$

of Λ -modules. By [21, Proposition 5.5.13], the latter is pseudo-isomorphic to $(\mathfrak{S}_{g,j}^2)^\iota$, which concludes the proof of the theorem. \square

When f is nonordinary at p , we shall establish a direct analogue of Wingberg's theorem on the level of Selmer groups under the following hypothesis on the local representation $A_{\bar{f}}$ (see Theorem 3.4 below).

(H0) For all $v \in S(\mathbb{Q}_{\text{cyc}})$, the group $H^0(\mathbb{Q}_{\text{cyc},v}, A_{\bar{f}}(k-i))$ is finite.

REMARK 3.2. Note that when $v \mid p$, the group $H^0(\mathbb{Q}_{\text{cyc},v}, A_{\bar{f}}(k-i))$ is always finite by [10, Lemma 3.3]. See also [10, Section 5], where sufficient conditions and explicit examples of the finiteness of $H^0(\mathbb{Q}_{\text{cyc},v}, A_{\bar{f}}(k-i))$ for $v \mid N$ are studied.

LEMMA 3.3. *Under (H0), we have a pseudo-isomorphism of Λ -modules*

$$\text{Sel}_0(A_f(i)/\mathbb{Q}_{\text{cyc}})^\vee \sim \mathfrak{S}_{\bar{f},k-i}^2.$$

PROOF. Let $n \geq 0$ be an integer. By the Poitou–Tate exact sequence (see, for example, [23, Section A.3.1]), we have the exact sequence

$$\bigoplus_{v \in S(L_n)} H^0(L_{n,v}, A_{\bar{f}}(k-i)) \longrightarrow H^2(G_S(L_n), T_f(i))^\vee \xrightarrow{\delta} H^1(G_S(L_n), A_{\bar{f}}(k-i)) \xrightarrow{\tau} \bigoplus_{v \in S(L_n)} H^1(L_{n,v}, A_{\bar{f}}(k-i)).$$

By definition,

$$\text{Image}(\delta) = \ker(\tau) = \text{Sel}_0(A_{\bar{f}}(k-i)/L_n).$$

This gives the exact sequence

$$\bigoplus_{v \in S(L_n)} H^0(L_{n,v}, A_{\bar{f}}(k-i)) \longrightarrow H^2(G_S(L_n), T_f(i))^\vee \longrightarrow \text{Sel}_0(A_{\bar{f}}(k-i)/L_n) \longrightarrow 0.$$

By the local Tate duality,

$$H^0(L_{n,v}, A_{\bar{f}}(k-i))^\vee \cong H^2(L_{n,v}, T_f(i))$$

for all $v \in S$. Thus, after taking Pontryagin duals and inverse limits,

$$0 \longrightarrow \text{Sel}_0(A_{\bar{f}}(k-i)/\mathbb{Q}_{\text{cyc}})^\vee \longrightarrow H_{\text{Iw}}^2(G_S(\mathbb{Q}_{\text{cyc}}), T_f(i)) \longrightarrow \bigoplus_{v \in S(\mathbb{Q}_{\text{cyc}})} H^0(\mathbb{Q}_{\text{cyc},v}, A_{\bar{f}}(k-i))^\vee.$$

Therefore, the lemma follows from **(H0)**. \square

THEOREM 3.4. *Suppose that f is nonordinary at p and **(H0)** holds. We have a pseudo-isomorphism of Λ -modules*

$$(\text{Sel}(A_f(i)/\mathbb{Q}_{\text{cyc}})^\vee)_{\Lambda\text{-tor}} \sim (\text{Sel}_0(A_{\bar{f}}(k-i)/\mathbb{Q}_{\text{cyc}})^\vee)^t.$$

PROOF. Consider the defining sequence of the Selmer group

$$0 \longrightarrow \text{Sel}(A_f(i)/\mathbb{Q}_{\text{cyc}}) \longrightarrow H^1(\mathbb{Q}_{\text{cyc}}, A_f(i)) \longrightarrow \bigoplus_{w \in S(\mathbb{Q}_{\text{cyc}})} \frac{H^1(\mathbb{Q}_{\text{cyc},w}, A_f(i))}{H_f^1(\mathbb{Q}_{\text{cyc},w}, A_f(i))}.$$

By [4, Proposition 3.8], there is an isomorphism

$$\varprojlim H^1(L_{n,w_n}, T_{\bar{f}}(k-i)) \cong \left(\frac{H^1(\mathbb{Q}_{\text{cyc},w}, A_f(i))}{H_f^1(\mathbb{Q}_{\text{cyc},w}, A_f(i))} \right)^\vee,$$

where $w_n \in S(L_n)$ is such that w_n lies below w and w_{n+1} . When $w \nmid p$, this is zero by [12, Section 17.10]. When $w \mid p$, it is also zero by [24, Theorem 0.6] under the hypothesis that f is nonordinary at p . Hence, we have an isomorphism

$$\text{Sel}(A_f(i)/\mathbb{Q}_{\text{cyc}}) \cong H^1(\mathbb{Q}_{\text{cyc}}, A_f(i))$$

of Λ -modules. Combining this with Theorem 3.1, we obtain a pseudo-isomorphism

$$(\text{Sel}(A_f(i)/\mathbb{Q}_{\text{cyc}})^\vee)_{\Lambda\text{-tor}} \sim (\mathfrak{S}_{f,i}^2)^t$$

of Λ -modules. But the latter is pseudo-isomorphic to $\text{Sel}_0(A_f(i)/\mathbb{Q}_{\text{cyc}})^\vee$ by Lemma 3.3 under **(H0)**. This concludes the proof of the theorem. \square

4. Comparison of characteristic elements

4.1. Preliminary lemmas. We prove several preliminary lemmas that will be used in the proofs of Theorems 4.3 and 4.4 below.

LEMMA 4.1. Assume that **(H0)** holds. If F is an irreducible element of Λ such that $F \nmid \mathfrak{F}_{f,k-i}$, then we have a pseudo-isomorphism

$$(\mathbb{H}_{f,i}^1/\mathfrak{S}_{f,i}^1)[F^\infty] \sim \text{Sel}_0(A_f(i)/\mathbb{Q}_{\text{cyc}})^{\vee,\iota}[F^\infty]$$

of Λ -modules. In particular, $(\mathbb{H}_{f,i}^1/\mathfrak{S}_{f,i}^1)[F]$ is finite if and only if $F \nmid \mathfrak{F}_{f,i}$.

PROOF. We have the Poitou–Tate exact sequence

$$0 \rightarrow \mathfrak{S}_{f,i}^1 \rightarrow \mathbb{H}_{f,i}^1 \rightarrow H^1(\mathbb{Q}_{\text{cyc}}, A_{\bar{f}}(k-i))^\vee \rightarrow \text{Sel}_0(A_{\bar{f}}(k-i)/\mathbb{Q}_{\text{cyc}})^\vee \rightarrow 0.$$

This gives the short exact sequence

$$0 \rightarrow (\mathbb{H}_{f,i}^1/\mathfrak{S}_{f,i}^1) \rightarrow H^1(\mathbb{Q}_{\text{cyc}}, A_{\bar{f}}(k-i))^\vee \rightarrow \text{Sel}_0(A_{\bar{f}}(k-i)/\mathbb{Q}_{\text{cyc}})^\vee \rightarrow 0,$$

which in turn induces the exact sequence

$$0 \rightarrow (\mathbb{H}_{f,i}^1/\mathfrak{S}_{f,i}^1)[F^\infty] \rightarrow H^1(\mathbb{Q}_{\text{cyc}}, A_{\bar{f}}(k-i))^\vee[F^\infty] \rightarrow \text{Sel}_0(A_{\bar{f}}(k-i)/\mathbb{Q}_{\text{cyc}})^\vee[F^\infty].$$

By assumption, $\text{Sel}_0(A_{\bar{f}}(k-i)/\mathbb{Q}_{\text{cyc}})^\vee[F^\infty]$ is finite. Therefore, we obtain a pseudo-isomorphism

$$(\mathbb{H}_{f,i}^1/\mathfrak{S}_{f,i}^1)[F^\infty] \sim H^1(\mathbb{Q}_{\text{cyc}}, A_{\bar{f}}(k-i))^\vee[F^\infty].$$

By Theorem 3.1 and Lemma 3.3, we have the pseudo-isomorphism of Λ -modules

$$H^1(\mathbb{Q}_{\text{cyc}}, A_{\bar{f}}(k-i))^\vee[F^\infty] \sim \text{Sel}_0(A_f(i)/\mathbb{Q}_{\text{cyc}})^{\vee,\iota}[F^\infty].$$

Combining these two pseudo-isomorphisms finishes the proof of the lemma. \square

LEMMA 4.2. Assume that **(H-IMC)** holds. Let F be an irreducible element of Λ such that $F \nmid \eta_{k,i}$. Recall that loc_p denotes the localisation map from $\mathfrak{S}_{f,i}^1$ to $\mathbb{H}_{f,i}^1$. Then one has a pseudo-isomorphism

$$(\Lambda^2/\Lambda(\mathfrak{F}_{f,k-i}^\#, \mathfrak{F}_{f,k-i}^b))[F^\infty] \sim (\mathbb{H}_{f,i}^1/(\text{loc}_p(\mathbf{z}_{f,i}))) [F^\infty]$$

of Λ -modules. Hence, $F \nmid \gcd(\mathfrak{F}_{f,k-i}^\#, \mathfrak{F}_{f,k-i}^b)$ if and only if $(\mathbb{H}_{f,i}^1/(\text{loc}_p(\mathbf{z}_{f,i}))) [F]$ is finite.

PROOF. By (2.4) and (2.3), we have the exact sequence

$$0 \rightarrow \mathbb{H}_{f,i}^1/(\text{loc}_p(\mathbf{z}_{f,i})) \rightarrow \Lambda^2/\Lambda(\mathfrak{F}_{f,k-i}^\#, \mathfrak{F}_{f,k-i}^b) \rightarrow \Lambda/\eta_{k,i} \rightarrow 0.$$

This gives the exact sequence

$$0 \rightarrow (\mathbb{H}_{f,i}^1/(\text{loc}_p(\mathbf{z}_{f,i}))) [F^\infty] \rightarrow (\Lambda^2/\Lambda(\mathfrak{F}_{f,k-i}^\#, \mathfrak{F}_{f,k-i}^b)) [F^\infty] \rightarrow \Lambda/\eta_{k,i} [F^\infty].$$

The last term is finite since $F \nmid \eta_{k,i}$. Hence, the result follows. \square

4.2. Comparison between characteristic ideals of fine Selmer groups and signed Selmer groups. The goal of this section is to prove a generalisation of [18, Theorem 1.2] (see Theorem 4.4 below). We shall do so via the following intermediate result.

THEOREM 4.3. Assume that **(H-IMC)** and **(H0)** hold. Let F be an irreducible element of Λ such that $F \nmid \eta_{k,i}$ and $F \nmid \mathfrak{F}_{\bar{f},k-i}$. There is a pseudo-isomorphism of Λ -modules

$$(\Lambda^2/\Lambda(\mathfrak{F}_{\bar{f},k-i}^\sharp, \mathfrak{F}_{\bar{f},k-i}^\flat))[F^\infty] \sim \text{Sel}_0(A_f(i)/\mathbb{Q}_{\text{cyc}})^{\vee, \iota}[F^\infty].$$

In particular, $F^\iota \nmid \mathfrak{F}_{f,i}$ if and only if $F \nmid \gcd(\mathfrak{F}_{\bar{f},k-i}^\sharp, \mathfrak{F}_{\bar{f},k-i}^\flat)$.

PROOF. By Lemma 4.2, we have a pseudo-isomorphism

$$(\Lambda^2/\Lambda(\mathfrak{F}_{\bar{f},k-i}^\sharp, \mathfrak{F}_{\bar{f},k-i}^\flat))[F^\infty] \sim (\mathbb{H}_{f,i}^1/(\text{loc}_p(\mathbf{z}_{f,i}))) [F^\infty]$$

of Λ -modules. Consider the short exact sequence

$$0 \rightarrow \mathfrak{H}_{f,i}^1/\Lambda \mathbf{z}_{f,i} \rightarrow \mathbb{H}_{f,i}^1/(\text{loc}_p(\mathbf{z}_{f,i})) \rightarrow \mathbb{H}_{f,i}^1/\mathfrak{H}_{f,i}^1 \rightarrow 0.$$

Since $\mathfrak{H}_{f,i}^1/\Lambda \mathbf{z}_{f,i}$ is a torsion Λ -module with $\text{Char}_\Lambda(\mathfrak{H}_{f,i}^1/\Lambda \mathbf{z}_{f,i}) = \mathfrak{F}_{\bar{f},k-i}$ not divisible by F , we have the following pseudo-isomorphism of Λ -modules:

$$(\mathbb{H}_{f,i}^1/(\text{loc}_p(\mathbf{z}_{f,i}))) [F^\infty] \sim (\mathbb{H}_{f,i}^1/\mathfrak{H}_{f,i}^1) [F^\infty].$$

By Lemma 4.1, the latter is pseudo-isomorphic to $\text{Sel}_0(A_f(i)/\mathbb{Q}_{\text{cyc}})^{\vee, \iota}[F^\infty]$. On combining these pseudo-isomorphisms, the theorem follows. \square

We can now state and prove the main result of the article.

THEOREM 4.4. Assume that **(H-IMC)** and **(H0)** hold. Let F be an irreducible element of Λ such that $F \nmid \eta_{k,i}$. Then $F^\iota \nmid \mathfrak{F}_{f,i}$ and $F \nmid \mathfrak{F}_{\bar{f},k-i}$ if and only if $F \nmid \gcd(\mathfrak{F}_{\bar{f},k-i}^\sharp, \mathfrak{F}_{\bar{f},k-i}^\flat)$.

PROOF. Theorem 4.3 tells us that if $F^\iota \nmid \mathfrak{F}_{f,i}$ and $F \nmid \mathfrak{F}_{\bar{f},k-i}$, then $F \nmid \gcd(\mathfrak{F}_{\bar{f},k-i}^\sharp, \mathfrak{F}_{\bar{f},k-i}^\flat)$. This proves the ‘only if’ implication.

Conversely, suppose that $F \nmid \gcd(\mathfrak{F}_{\bar{f},k-i}^\sharp, \mathfrak{F}_{\bar{f},k-i}^\flat)$. From the inclusion

$$\text{Sel}_0(A_{\bar{f}}(k-i)) \subset \text{Sel}^\bullet(A_{\bar{f}}(k-i)),$$

we have $F \nmid \mathfrak{F}_{\bar{f},k-i}$. The remaining assertion that $F^\iota \nmid \mathfrak{F}_{f,i}$ now follows from Theorem 4.3. \square

REMARK 4.5. Let E/\mathbb{Q} be an elliptic curve with good supersingular reduction at p . In [18], it is stated in the proof of Proposition 3.1 that by [28, Corollary 2.5], we have the equality

$$\text{Char}_\Lambda(\text{Sel}_{p^\infty}(E/\mathbb{Q}_{\text{cyc}})^\vee)_{\Lambda\text{-tor}} = \text{Char}_\Lambda \text{Sel}_0(E/\mathbb{Q}_{\text{cyc}})^\vee.$$

However, one of the two Selmer groups should be twisted by ι in order for the equality to hold. As such, for the rest of the proof to go through, the additional hypothesis that the irreducible element f (not to be confused with the notation for a modular form in the present article) satisfies $(f) = (f^\iota)$ is required. Consequently, the statement of [18, Theorem 1.2] should also be modified.

Specialising our Theorem 4.4 to the case $F = \varpi$, where ϖ is a uniformiser of \mathcal{O} , we may deduce the following result.

COROLLARY 4.6. *Assume that (H-IMC) and (H0) hold. Then the following statements are equivalent.*

- (a) *The μ -invariants of $\text{Sel}_0(A_f(i)/\mathbb{Q}_{\text{cyc}})^\vee$ and $\text{Sel}_0(A_{\bar{f}}(k-i)/\mathbb{Q}_{\text{cyc}})^\vee$ are zero.*
- (b) *At least one of $\text{Sel}^\sharp(A_{\bar{f}}(k-i)/\mathbb{Q}_{\text{cyc}})^\vee$ and $\text{Sel}^\flat(A_{\bar{f}}(k-i)/\mathbb{Q}_{\text{cyc}})^\vee$ has trivial μ -invariant.*

REMARK 4.7. From [7, Conjecture A], [11, Conjecture A] and [1, Conjecture 1.2], the μ -invariants of $\text{Sel}_0(A_f(i)/\mathbb{Q}_{\text{cyc}})^\vee$ and $\text{Sel}_0(A_{\bar{f}}(k-i)/\mathbb{Q}_{\text{cyc}})^\vee$ are predicted always to vanish. Corollary 4.6 gives an alternative formulation of these conjectures in terms of signed Selmer groups. In fact, numerical calculations carried out by Pollack (see [25, Section 7] and <http://math.bu.edu/people/rpollack/Data/data.html>) suggest that the μ -invariants of both $\text{Sel}^\sharp(A_{\bar{f}}(k-i)/\mathbb{Q}_{\text{cyc}})^\vee$ and $\text{Sel}^\flat(A_{\bar{f}}(k-i)/\mathbb{Q}_{\text{cyc}})^\vee$ are zero when $A_{\bar{f}}(k-i)$ is given by $E[p^\infty]$ for some elliptic curve E/\mathbb{Q} with good supersingular reduction at p .

Acknowledgement

We would like to thank the anonymous referee for providing various helpful comments that have helped us improve the exposition of this article.

References

- [1] C. S. Aribam, ‘On the μ -invariant of fine Selmer groups’, *J. Number Theory* **135** (2014), 284–300.
- [2] L. Berger, ‘Bloch and Kato’s exponential map: three explicit formulas’, *Doc. Math.* **Extra Vol.** (2003), 99–129.
- [3] L. Berger, ‘Limites de représentations cristallines’, *Compos. Math.* **140**(6) (2004), 1473–1498.
- [4] S. Bloch and K. Kato, ‘ L -functions and Tamagawa numbers of motives’, in: *The Grothendieck Festschrift*, P. Cartier, L. Illusie, N. M. Katz, G. Laumon, Y. Manin and K. A. Ribet (eds.), Vol. I, Progress in Mathematics, 86 (Birkhäuser, Boston, MA, 1990), 333–400.
- [5] F. Castella, M. Çiperiani, C. Skinner and F. E. I. Sprung, ‘On the Iwasawa main conjectures for modular forms at non-ordinary primes’, Preprint, 2018, [arXiv:1804.10993v2](https://arxiv.org/abs/1804.10993v2).
- [6] F. Castella, Z. Liu and X. Wan, ‘Iwasawa–Greenberg main conjecture for non-ordinary modular forms and Eisenstein congruences on $\text{GU}(3, 1)$ ’, Preprint, 2021, [arXiv:2109.08375](https://arxiv.org/abs/2109.08375).
- [7] J. Coates and R. Sujatha, ‘Fine Selmer groups of elliptic curves over p -adic Lie extensions’, *Math. Ann.* **331**(4) (2005), 809–839.
- [8] P. Deligne, ‘Formes modulaires et représentations ℓ -adiques’, *Sémin. Bourbaki* **21** (1968–1969), Exp. no. 355, 139–172.
- [9] O. Fouquet and X. Wan, ‘The Iwasawa main conjecture for modular motives’, Preprint, 2021, [arXiv:2107.13726](https://arxiv.org/abs/2107.13726).
- [10] J. Hatley, D. Kundu, A. Lei and J. Ray, ‘On the fine Selmer groups of modular forms and duality’, Preprint, 2021.
- [11] S. Jha and R. Sujatha, ‘On the Hida deformations of fine Selmer groups’, *J. Algebra* **338** (2011), 180–196.
- [12] K. Kato, ‘ p -adic Hodge theory and values of zeta functions of modular forms’, in: *Cohomologies p -adiques et applications arithmétiques. III*, Astérisque **295**(ix) (2004), 117–290.

- [13] S.-I. Kobayashi, 'Iwasawa theory for elliptic curves at supersingular primes', *Invent. Math.* **152**(1) (2003), 1–36.
- [14] A. Lei, 'Iwasawa theory for modular forms at supersingular primes', *Compos. Math.* **147**(3) (2011), 803–838.
- [15] A. Lei and M. F. Lim, 'Mordell–Weil ranks and Tate–Shafarevich groups of elliptic curves with mixed-reduction type over cyclotomic extensions', *Int. J. Number Theory*, to appear. <https://doi.org/10.1142/S1793042122500208>.
- [16] A. Lei, D. Loeffler and S. L. Zerbes, 'Wach modules and Iwasawa theory for modular forms', *Asian J. Math.* **14**(4) (2010), 475–528.
- [17] A. Lei, D. Loeffler and S. L. Zerbes, 'Coleman maps and the p -adic regulator', *Algebra Number Theory* **5**(8) (2011), 1095–1131.
- [18] A. Lei and R. Sujatha, 'On fine Selmer groups and the greatest common divisor of signed and chromatic p -adic L -functions', *Proc. Amer. Math. Soc.* **149**(8) (2021), 3235–3243.
- [19] A. Matar, 'On the Λ -cotorsion subgroup of the Selmer group', *Asian J. Math.* **24**(3) (2020), 437–456.
- [20] J. Nekovář, 'Selmer complexes', *Astérisque* **310** (2006), 559 pages.
- [21] J. Neukirch, A. Schmidt and K. Wingberg, *Cohomology of Number Fields*, 2nd edn, Grundlehren der mathematischen Wissenschaften, 323 (Springer, Berlin, 2008).
- [22] B. Perrin-Riou, 'Théorie d'Iwasawa des représentations p -adiques sur un corps local', *Invent. Math.* **115**(1) (1994), 81–161.
- [23] B. Perrin-Riou, 'Fonctions L_p -adiques des représentations p -adiques', *Astérisque* **198**(229) (1995), 212 pages.
- [24] B. Perrin-Riou, 'Représentations p -adiques et normes universelles. I. Le cas cristallin', *J. Amer. Math. Soc.* **13**(3) (2000), 533–551.
- [25] R. Pollack, 'On the p -adic L -function of a modular form at a supersingular prime', *Duke Math. J.* **118**(3) (2003), 523–558.
- [26] C. Skinner and E. Urban, 'The Iwasawa main conjectures for GL_2 ', *Invent. Math.* **195**(1) (2014), 1–277.
- [27] F. E. I. Sprung, 'Iwasawa theory for elliptic curves at supersingular primes: a pair of main conjectures', *J. Number Theory* **132**(7) (2012), 1483–1506.
- [28] K. Wingberg, 'Duality theorems for Abelian varieties over Z_p -extensions', in: *Algebraic Number Theory*, R. Coates, B. Greenberg, B. Mazur and I. Satake (eds.), Advanced Studies in Pure Mathematics, 17 (Academic Press, Boston, MA, 1989), 471–492.
- [29] C. Wuthrich, 'Iwasawa theory of the fine Selmer group', *J. Algebraic Geom.* **16**(1) (2007), 83–108.

ANTONIO LEI, Département de Mathématiques et de Statistique,
 Université Laval, Pavillion Alexandre-Vachon,
 1045 Avenue de la Médecine, Québec, QC G1V 0A6, Canada
 e-mail: antonio.lei@mat.ulaval.ca

MENG FAI LIM, School of Mathematics and Statistics
 and Hubei Key Laboratory of Mathematical Sciences,
 Central China Normal University, Wuhan 430079, PR China
 e-mail: limmf@mail.ccnu.edu.cn