

Crops and Soils Research Paper

Cite this article: Nurdianti RR, Nuryana RS, Handoko A, Hernaman I, Ramdani D, Jayanegara A, Dickhoefer U, Böttger C, Südekum K-H (2023). Nutritional compositions of Katuk leaves and their supplementation to hays of different quality: an *in vitro* study. *The Journal of Agricultural Science* **161**, 428–437. <https://doi.org/10.1017/S002185962300031X>

Received: 30 September 2022

Revised: 1 May 2023

Accepted: 4 May 2023

First published online: 6 June 2023

Keywords:

fibre digestibility; fibre fractions; Hohenheim gas test; proximate nutrients; *Sauropus androgynus* (L.) Merr

Corresponding author:

R.R. Nurdianti; Email: risma_rizkia.nurdianti@uni-hohenheim.de

Nutritional compositions of Katuk leaves and their supplementation to hays of different quality: an *in vitro* study

R.R. Nurdianti^{1,2,3} , R. S. Nuryana² , A. Handoko² , I. Hernaman² ,
D. Ramdani^{2,3} , A. Jayanegara^{3,4} , U. Dickhoefer^{1,5} , C. Böttger⁶
and K.-H. Südekum⁶ 

¹Institute of Agricultural Sciences in the Tropics (Hans-Ruthenberg-Institute), University of Hohenheim, 70599

Stuttgart, Germany; ²Faculty of Animal Husbandry, University of Padjadjaran, 45363 Sumedang, Indonesia;

³Faculty of Animal Science, Animal Feed and Nutrition Modelling (AFENUE) Research Group, IPB University, 16680

Bogor, Indonesia; ⁴Department of Nutrition and Feed Technology, IPB University, 16680 Bogor, Indonesia;

⁵Institute of Animal Nutrition and Physiology, Kiel University, 24098 Kiel, Germany and ⁶Institute of Animal

Science, University of Bonn, 53115 Bonn, Germany.

Abstract

Katuk leaves (*Sauropus androgynus* (L.) Merr.; KL) are widely consumed by breast-feeding Indonesian mothers as it has been reported to increase breast milk production. It is hypothesized that supplementing KL in diets might increase crude protein (CP) concentration and fibre digestibility in the diet. The KL had high CP and non-fibre carbohydrate concentrations (333 and 332 g/kg dry matter; DM, respectively), but low neutral detergent fibre assayed with heat, a stable amylase and expressed exclusive of residual ash (aNDFom; 200 g/kg DM). Fibre digestibility linearly increased with increasing of KL supplementation in low-quality hay (LQH) diets. The KL did not contain a considerable amount of tannins. In LQH diets, gas production after 24 h incubation (GP₂₄) linearly increased with increasing of KL supplementation ($P < 0.001$). Meanwhile, GP₂₄ linearly decreased with increasing of KL supplementation in medium- and high-quality hays (MQH and HQH; $P < 0.001$). Metabolizable energy tended to linearly increase in LQH diets, but tended to linearly decrease with increasing of KL supplementation in MQH and HQH diets ($P = 0.078$). Therefore, this study suggested that KL can be a potential supplement in the ruminant diet due to its abundant dietary proteins but low fibre concentration in its leaves. However, further studies (e.g. *in vitro* or *in vivo*) investigating other rumen parameters after incubation should be performed to validate how KL can be supplemented in the diet of ruminant livestock.

Introduction

Dairy cattle in developing countries often produce less milk and have shorter lactation periods than cattle in other regions. Poor animal performance in small-scale dairy systems in developing countries is the result of factors such as climate (high ambient temperature and humidity), low-quality feed, low levels of concentrate supplementation, low genetic potential for milk production of multi-purpose animals (in addition to milk and meat these cattle also often provide draught power) and high incidence of diseases (Food and Agriculture Organization, 2014). In tropical and sub-tropical developing countries, there is a gap between available and required animal feeds. Typically, ruminants in those countries are fed lignified forages and crop residues that are low in available energy and nitrogen (N; Nasser *et al.*, 2009).

Sauropus androgynus (L.) Merr. (Katuk) is a perennial shrub found growing wild in South East Asia and widely cultivated in Indonesia and Malaysia (Padmavathi and Rao, 1990). Katuk, a plant species of the Euphorbiaceae family, is rich in fatty acids, flavonoids and polyphenols as the main bioactive constituents (Zhang *et al.*, 2020) and is commonly utilized as a medicinal herb in the treatment of diabetes, cancer, inflammation, microbial infection, ulcers, obesity and allergies (Paul and Beena Anto, 2011; Zhang *et al.*, 2020). However, it is not recommended to consume excessive amount of freshly uncooked Katuk leaves (KL) over a period of time, because it could be associated with the occurrence of bronchiolitis obliterans disease (Zhang *et al.*, 2020). Consumption of extract of KL is considered to increase milk production in human mothers during lactation up to 50.7 g/100 g (Sa'roni *et al.*, 2012). Supplementing KL in the diet might also increase the milk yield of ruminants. In addition, Noach *et al.* (2020) reported that KL inclusion (with or without Zn bio complex) in a concentrate diet of late pregnant Ettawah could improve the milk production and birth weight of kid during kidding period.

© The Author(s), 2023. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Evaluating the nutritive value of the available local feed resources is important to assess their potential role in the nutrition of livestock (Taghizadeh *et al.*, 2008) which can influence the livestock productivity (Geremew *et al.*, 2017). For this, *in vitro* techniques are commonly used to screen a variety of feedstuffs that are not available in sufficient quantity for *in vivo* experiments (Getachew *et al.*, 2002). Menke and Steingass (1988) reported a strong correlation between metabolizable energy (ME) values measured *in vivo* and predicted from 24 h *in vitro* gas production (GP) and chemical composition of feeds. *In vitro* GP is an alternative technique used to determine the nutritive value of feedstuffs, since rate and extent of degradation and rumen fermentation can be simply determined by measurement of cumulative GP (Sommart *et al.*, 2000). *In vitro* GP technique is less expensive, easy to determine and appropriate for use in developing countries (Salamatazar *et al.*, 2012). Fibre digestibility can also be determined using a modified *in vitro* technique (Tilley and Terry, 1963; Robertson and Van Soest, 1981). With respect to KL as animal feed supplement, limited information is available on their nutritive value and ruminal degradation. Therefore, the present study aimed at testing a hypothesis that the nutritive value of KL and their supplementation to hays of different quality could improve the crude protein (CP) concentration, fibre digestibility and ME, especially in low-quality hay diet. The preliminary findings of this study were published in abstract form (Nurdianti *et al.*, 2017).

Materials and methods

Sample collection and experimental design

Approximately 0.5 kg of fresh leaves of KL without stem were harvested for the sample collection. Samples of KL with old maturity were collected from 80–90 cm herbaceous-shrub plant in Bandung, Indonesia. In Bandung, annual precipitation ranges from 2500 to 3000 mm and temperature ranges from 24 to 28 °C (BMKG, 2013).

After collection, KL samples were stored at 4 °C overnight, oven dried at 60 °C for 24 h (F115, Binder GmbH, Tuttlingen, Germany), and ground to pass a 1 mm sieve using a laboratory mill (model SM100, Retsch GmbH, Haan, Germany). As a substrate for KL to be added to, three grass hays from temperate regions (grown in Germany and Switzerland) were selected based on their CP concentration to represent a wide range of qualities: hay with low quality (LQH), hay with medium quality (MQH) and hay with high quality (HQH). Hays were ground to pass a 1 mm sieve using the same mill as for KL. Samples of KL and each grass hay were incubated alone or in KL + grass hay combinations with four different inclusion levels of KL (i.e. 0, 50, 100 and 200 g/kg dry matter (DM)). The cows used as donor animals were housed in accordance with the German Animal Welfare legislation. All procedures for animal handling within the present study were performed according to the National Committee for the Protection of Animals Used for Scientific Purposes for the Federal Republic of Germany. The experimental diets were arranged in a completely randomized design in six replicates. The feedstuff combinations, i.e. diets, are presented in Table 1.

Chemical analyses

Chemical composition of the individual sample was determined according to the Association of German Agricultural Analytic and Research Institutes (Verband Deutscher Landwirtschaftlicher

Untersuchungs- und Forschungsanstalten, 2007). The DM concentration was determined by oven-drying at 105 °C for 24 h (method 3.1) and ash concentration was determined by incinerating samples at 550 °C for 5 h (method 8.1). The N concentration was determined by Kjeldahl procedure (method 4.1.2) and CP was calculated as N×6.25. Ether extract (EE) concentration was analysed using method 5.1.1. Neutral detergent fibre, assayed with a heat-stable amylase and expressed exclusive of residual ash (aNDFom), acid detergent fibre, expressed exclusive of residual ash (ADFom) and lignin determined by solubilization of cellulose with sulphuric acid (Lignin (sa)) concentrations were measured using an ANKOM200 Fibre Analyzer (Ankom Technology Corp., Macedon, NY, USA) according to methods 6.5.1, 6.5.2 and 6.5.3, respectively. Sodium sulphite was used for the aNDFom analysis. Concentration of hemicelluloses was determined by subtracting ADFom concentration from aNDFom concentration and concentration of cellulose by subtracting Lignin (sa) concentration from ADFom concentration (Rinne *et al.*, 1997). All samples were analysed in duplicate for each chemical constituent and analysis repeated, if the coefficient of variation exceeded 5%.

Non-fibre carbohydrates (NFC) were calculated according to National Research Council (2001), as:

$$\text{NFC} = 1000 - (\text{aNDFom} + \text{CP} + \text{EE} + \text{ash}) \quad (1)$$

where all nutrient concentrations are in g/kg DM.

Tannin bioassay

To investigate whether the KL sample contained tannins or not, an indirect approach was selected. Katuk leaves were evaluated using polyethylene glycol (PEG) supplementation in an *in vitro* GP experiment as mentioned by Jayanegara *et al.* (2009). The PEG has a high affinity for tannins and makes tannins inactive by binding to them. Hence, KL samples (220–250 mg) were incubated without and with 750 mg PEG supplementation with 10 ml of rumen fluid and 20 ml of buffer solution (Menke *et al.*, 1979) using Hohenheim gas test (HGT; Makkar *et al.*, 1995).

The syringes were placed in a rotor inside the incubator (39 °C) with about one rotation per minute. The GP was recorded after 48 h of incubation (GP₄₈). Blanks were prepared by incubating syringes containing rumen fluid and buffer solution, but without any sample. Katuk leave samples were incubated in triplicate in two different *in vitro* incubation runs. The difference between the *in vitro* GP with and without PEG supplementation is an indicator of tannin effect. The increase in gas on addition of PEG is a measure of tannin activity, where the protocol used was similar to that described in Makkar *et al.* (1995).

Fibre fractionation and *in vitro* fibre digestibility

The fibre fractionation and *in vitro* fibre digestibility experiment were evaluated using the modified Tilley and Terry technique (Tilley and Terry, 1963; Robertson and Van Soest, 1981). For this, rumen fluid was collected from two rumen-fistulated Jersey cows, fed on a total mixed ration containing grass silage (180 g/kg DM), corn silage (167 g/kg DM), grass hay (184 g/kg DM), barley straw (21 g/kg DM), and concentrate mixture (426 g/kg DM), urea (2 g/kg DM) and mineral mixture (20 g/kg DM). The total mixed ration had a forage to concentrate ratio of 55:45 (on DM basis) and contained 148 g CP/kg DM. Rumen fluid was collected immediately before morning feeding and

Table 1. Ingredient composition and calculated chemical composition ($n = 2$) of the experimental diets

	Forage												
	Low-quality hay				Medium-quality hay				High-quality hay				
					Katuk leaves (g/kg DM)								
Level	0	50	100	200	0	50	100	200	0	50	100	200	1000
Ingredient composition of diets (g/kg DM)													
Katuk leaves	0	50	100	200	0	50	100	200	0	50	100	200	1000
Low-quality grass hay	1000	950	900	800	0	0	0	0	0	0	0	0	0
Medium-quality grass hay	0	0	0	0	1000	950	900	800	0	0	0	0	0
High-quality grass hay	0	0	0	0	0	0	0	0	1000	950	900	800	0
Chemical composition of the diets (g/kg DM)													
Dry matter (g/kg FM)	928	927	926	925	922	921	921	920	907	907	907	908	911
Ash	56	59	62	67	86	87	88	91	102	102	103	104	111
Crude protein	45	60	74	103	89	101	113	137	147	156	166	184	333
Ether extract	3.6	4.7	5.7	7.8	13	13.6	14.2	15.3	30	29.7	29.5	28.9	24.5
aNDFom	700	675	650	600	550	533	515	480	512	496	481	450	200
ADFom	450	437	424	398	301	296	290	279	280	276	271	262	192
Lignin (sa)	33.9	32.8	31.7	29.5	31.9	30.9	29.9	27.9	31.8	30.8	29.8	27.8	11.9
Hemicelluloses ^a	250	238	226	202	249	237	225	201	232	221	210	187	8.0
Cellulose ^a	416	404	393	369	269	265	260	251	248	245	241	235	180
NFC ^a	195	202	208	222	263	266	270	276	209	215	221	234	332

ADFom, acid detergent fibre expressed exclusive of residual ash; aNDFom, neutral detergent fibre assayed with a heat-stable amylase and expressed exclusive of residual ash; DM, dry matter; FM, fresh matter; Lignin (sa), lignin determined by solubilization of cellulose with sulphuric acid; NFC, non-fibrous carbohydrate.

^aCellulose = ADFom – Lignin (sa); hemicellulose = aNDFom – ADFom; NFC = 1000 – (aNDFom + crude protein + ether extract + ash).

strained through two layers of cheesecloth into pre-warmed and isolated bottle. All laboratory handling of rumen fluid was carried out under a continuous flow of CO₂. The rumen fluid of both cows was mixed, homogenized and filtered using a nylon net (pore size 100 µm).

The ground initial sample (500 mg) was weighed into a 250 ml flask (Goering and Van Soest, 1970). The rumen fluid (10 ml) and the buffer solution (40 ml) were added to each flask. The flasks were then placed in a preheated (39 °C) water bath under CO₂ positive pressure to ensure anaerobiosis, and incubated for 240 h, which corresponds to the maximum extent of fibre digestion in an anaerobic environment *in vitro* (Fox *et al.*, 2004; Raffrenato and Van Amburgh, 2010; Raffrenato *et al.*, 2018). Re-inoculation of the flasks with 10 ml rumen fluid and 40 ml buffer solution was conducted after 120 h of incubation to preserve the microbial activity during the whole incubation process (Palmonari *et al.*, 2014). Blanks were prepared by incubating flasks containing buffer and rumen fluid but without any sample to correct for any feed particles introduced into the *in vitro* system with the rumen fluid (Raffrenato *et al.*, 2018). Each sample was incubated in triplicate in two runs resulting in six observations per experimental diet.

At the end of the incubation, the whole content of each flask was moved to a 600 ml beaker that was covered by a round cold-water condenser to minimize evaporation (Mertens, 2002) and determine the aNDFom concentration of the residue (Goering and Van Soest, 1970). About 0.5 g of sodium sulphite and 50 ml of neutral detergent solution were added to each

refluxing beaker and refluxed for 60 min at boiling temperature to create vigorous particle movement. After refluxing, the content of each beaker was filtered through crucibles (40 µm porosity, Duran™ Borosilicate Glass Filter Crucibles number 2, DWK Life Sciences, Wertheim, Germany) and the water removed with a vacuum pump. Filtered residues were dried in a forced-air oven (105 °C) for 3 h, and the weights of the crucibles were recorded. Ash correction was done by incineration of the residue at 550 °C for 4 h.

The *in vitro* neutral detergent fibre digestibility (IVNDFD of the incubated samples after 240 h; IVNDFD₂₄₀) was calculated as

$$\text{IVNDFD}_{240} = \frac{\text{aNDFom}_r - \text{aNDFom}_b}{\text{aNDFom}_i} \quad (2)$$

where aNDFom_r is the residual aNDFom after 240 h *in vitro* fermentation (g/kg DM), aNDFom_b is the blank correction after 240 h *in vitro* fermentation (g/kg DM) and aNDFom_i represents the initial aNDFom concentration from samples (g/kg DM).

The uNDF concentration (g/kg DM) after 240 h *in vitro* fermentation (uNDF₂₄₀) was calculated as

$$\text{uNDF}_{240} \text{ (g/kg DM)} = \frac{(1000 - \text{IVNDFD}_{240}) \times \text{aNDFom}_i}{1000} \quad (3)$$

where aNDFom_i (g/kg DM) is the aNDFom concentration of the sample (g/kg DM) and IVNDFD₂₄₀ is *in vitro* fibre digestibility (IVNDFD) of the incubated samples after 240 h.

The pdNDF concentration (g/kg DM) was calculated as

$$\text{pdNDF (g/kg DM)} = \text{aNDFom}_i - \text{uNDF}_{240} \quad (4)$$

where aNDFom_i (g/kg DM) is the aNDFom concentration of the sample (g/kg DM) and uNDF_{240} is the uNDF in the residue after 240 h *in vitro* fermentation (g/kg DM).

In vitro gas production, metabolizable energy and net energy for lactation

The HGT was performed according to Menke and Steingass (1988) but modified regarding the incubation duration. Rumen fluid was collected from two castrated male adult Blackface sheep, fed on a standard diet of grass hay, a commercial compound feed and barley (650, 200 and 150 g/kg DM, respectively) that covered maintenance energy requirements. The animals never received tanniferous and/or tropical feeds before. Rumen fluid was collected immediately before feeding and strained through two layers of cheesecloth into pre-warmed and isolated bottle. All laboratory handling of rumen fluid was carried out under a continuous flow of CO_2 .

The ground samples (220–250 mg) of the air-dried feedstuffs and the respective mixtures were accurately weighed into 100 ml glass syringes and the syringe pistons were lubricated with Vaseline and inserted into the syringes. Triplicates of syringes without substrate (blanks) and of standard hay and concentrate were included as laboratory controls. According to Menke and Steingass (1988), GP from the blank was subtracted from all samples incubated to obtain the net GP. Subsequently, GP from the hay standard was divided by the measured net value of the hay standard to provide the correction factor. Similarly, GP from the concentrate standard was divided by the measured net GP of the concentrate standard. The average value of correction factor and concentrate standard was used for the adjustment. Each sample was incubated in triplicate in two different *in vitro* incubation runs. Incubations were repeated when gas volumes of the standards deviated by more than 10% and when coefficient of variation between repetitions exceeded 5% from the reference values.

Syringes were filled with 30 ml of medium consisting of rumen fluid (10 ml) and 20 buffer solution (20 ml) as described by Menke and Steingass (1988), except that the concentration of NaHCO_3 was reduced to 33 g/l and that of $(\text{NH}_4)\text{HCO}_3$ increased to 6 g/l to prevent a shortage in N during prolonged incubation times. The syringes were placed in a rotor inside the incubator (39 °C) with about one rotation per minute. The GP was recorded after 2, 4, 6, 8, 12, 16, 24, 36, 48, 60, 72 and 96 h of incubation. The ME and net energy for lactation (NEL) values were calculated according to Menke and Steingass (1988) as:

$$\text{ME} = 2.20 + 0.1357 \text{ GP} + 0.0057 \text{ CP} + 0.0002859 \text{ EE}^2 \quad (5)$$

$$\text{NEL} = 0.54 + 0.0959 \text{ GP} + 0.0038 \text{ CP} + 0.0001733 \text{ EE}^2 \quad (6)$$

where GP is the net GP from 200 mg dry sample after 24 h of incubation (GP_{24}) and after being corrected from its correction factor for the day-to-day variation in the activity of rumen fluid

(ml) is expressed in ml/200 mg DM, and ash, CP and EE concentrations are expressed as g/kg DM.

Data analyses

To describe the dynamics of GP over time, the following Gompertz function (Schofield *et al.*, 1994) was chosen:

$$\text{GP} = A \exp \left\{ -\exp \left[1 + \frac{b}{A} (\text{LAG} - t) \right] \right\} \quad (7)$$

where A is the theoretical maximum of GP, b is the maximum rate of GP (ml/h) that occurs at the point of inflection of the curve, LAG is the lag time (h) which is defined as the time-axis intercept of a tangent line at the point of inflection, and t is time (h). The parameters A , b and LAG were estimated by non-linear regression analysis (PROC NLIN; SAS 9.4, SAS Institute Inc., Cary, North Carolina, USA).

The fibre fractions, *in vitro* fibre digestibility, GP (i.e. 12, 24, 48 and 96 h of incubation), ME and NEL values were analysed using a mixed procedure (PROC MIXED) by SAS 9.4 (SAS Institute Inc.) according to:

$$Y_{ijk} = \mu + \alpha_i + \beta_j + (\alpha\beta)_{ij} + R_k + e_{ijk} \quad (8)$$

where Y_{ijk} = the dependent variable; μ = the overall mean; α_i = the effect of hays of different quality (i.e. LQH, MQH and HQH); β_j = the inclusion levels of KL (i.e. 0, 50, 100 and 200 g/kg DM); $(\alpha\beta)_{ij}$ = the interaction of grass hay with different quality and inclusion levels of KL; R_k = the random effect of run; and e_{ijk} = the residual error. Linear effects of level were tested by orthogonal polynomial contrasts using the CONTRAST statement. Differences between means with $P < 0.050$ were accepted as statistically significant, and differences with $0.050 < P < 0.100$ were considered to represent tendencies to significance.

Results

Chemical composition

The calculated chemical composition varied among experimental diets (Table 1). Initial sample of KL had higher CP and NFC concentrations than experimental diets. Meanwhile, LQH diet without KL supplementation had lower CP and NFC concentrations among experimental diets. Initial sample of KL had lower aNDFom, ADFom, Lignin (sa), hemicelluloses and cellulose concentrations compared to experimental diets. Meanwhile, LQH diet without KL supplementation had higher aNDFom, ADFom Lignin (sa), hemicelluloses and cellulose concentrations compared to other experimental diets.

Fibre fractions and in vitro fibre digestibility

The concentrations of undigested aNDFom after 240 h incubation (uNDF_{240}) and potentially digestible aNDFom (pdNDF; both in g/kg DM) as well as the fibre digestibility differed between the grass hays (i.e. LQH, MQH and HQH; $P < 0.001$ for all variables; Table 2). Moreover, the uNDF_{240} and pdNDF concentrations differed between levels of KL supplementation ($P = 0.001$ and $P < 0.001$, respectively). The HQH supplemented with 200 g/kg DM basis of KL had a much lower uNDF_{240} concentration than other experimental diets. Meanwhile, MQH diet without

Table 2. Calculated fibre fractions, i.e. uNDF₂₄₀ and pdNDF (g/kg DM), and fibre digestibility, i.e. iVNDFD₂₄₀, of the experimental diets ($n = 3$)

Variable	Hay												P value ^a				
	Low-quality hay				Medium-quality hay				High-quality hay								
	0	50	100	200	0	50	100	200	0	50	100	200	SEM	H	L	H×L	CL
KL level (g/kg DM)	267	256	245	223	85	83	81	78	82	80	78	75	6.666	<0.001	0.001	0.014	<0.001
uNDF ₂₄₀ (g/kg DM)	433	419	405	377	464	449	433	402	429	416	402	376	6.666	<0.001	<0.001	0.984	<0.001
pdNDF (g/kg DM)	0.618	0.621	0.623	0.628	0.844	0.835	0.826	0.809	0.839	0.830	0.822	0.805	0.0123	<0.001	0.026	0.083	0.004
iVNDFD ₂₄₀																	

aNDFom, neutral detergent fibre assayed with a heat-stable amylase and expressed exclusive of residual ash; DM, dry matter; iVNDFD₂₄₀, *in vitro* neutral detergent fibre digestibility of the incubated samples after 240 h; KL, Katuk leaves; pdNDF, potentially digestible neutral detergent fibre; SEM, standard error mean; uNDF₂₄₀, undigested neutral detergent fibre estimated after 240 h of *in vitro* incubation.

^aProbability of the effects of H, hays of different quality; L, inclusion levels of Katuk leaves; H×L, interaction of hays of different quality and inclusion levels of Katuk leaves; CL, linear effect of inclusion levels of Katuk leaves.

supplementation of KL had highest pdNDF concentration and fibre digestibility among all experimental diets, i.e. MQH diet with KL supplementation, all LQH diet and all HQH diet. The interaction between grass hay with different quality and inclusion level of KL was significant for uNDF₂₄₀ concentration ($P = 0.014$), but tended to be significant for fibre digestibility ($P = 0.083$). In all diets, uNDF₂₄₀ and pdNDF concentrations linearly decreased with increasing level of KL in the diet ($P < 0.01$). Meanwhile, fibre digestibility linearly increased with increasing KL level in LQH diet, but not in MQH and HQH diets ($P = 0.04$).

Tannin bioassay

Addition of PEG to KL did not affect the cumulative GP₂₄ (Fig. 1), averaging 36 ml/200 mg DM. Similarly, the GP₄₈ of KL did not differ irrespective of whether it was incubated with PEG or without PEG.

In vitro gas production, metabolizable energy and net energy for lactation

The cumulative GP differed between experimental diets at all incubation times, i.e. 12 h, 24 h, 48 h and 96 h ($P < 0.001$, for all variables; Table 3). Mean total GP during 12 h of incubation (GP₁₂) ranged from 21.9 to 35.7 ml/200 mg DM. The interaction between grass hay with different quality and inclusion level of KL was significant for GP₁₂ ($P = 0.007$). In all experimental diets, GP₁₂ did not linearly increase with increasing of KL supplementation in the diet ($P = 0.410$).

Total volume of GP₂₄ ranged from 33.3 to 49.6 ml/200 mg DM. The interaction between grass hay with different quality and inclusion level of KL was significant for GP₂₄ ($P = 0.004$). In LQH diets, GP₂₄ linearly increased with increasing of KL supplementation in the diet ($P < 0.001$). Meanwhile, GP₂₄ linearly decreased with increasing of KL supplementation in MQH and HQH diets ($P < 0.001$).

Total volume of GP₄₈ ranged from 42.6 to 57.6 ml/200 mg DM. There was no interaction between grass hay with different quality and inclusion level of KL for GP₄₈ ($P = 0.101$). In all experimental diets, GP₄₈ linearly decreased with increasing of KL supplementation in the diet ($P < 0.001$).

Total volume of gas produced during 96 h (GP₉₆) ranged from 48.4 to 62.2 ml/200 mg DM. There was no interaction between grass hay with different quality and inclusion level of KL for GP₉₆ ($P = 0.410$). In all experimental diets, GP₉₆ linearly decreased with increasing level of KL in the diet ($P < 0.001$).

The ME and NEL values of experimental diets varied widely (Table 4). The calculated ME and NEL values differed between experimental diets ($P < 0.001$, for all variables). The interaction between grass hay with different quality and inclusion level of KL affected ME and NEL values ($P < 0.01$, $P = 0.001$ and $P = 0.001$, respectively). In LQH diets, ME tended to linearly increase with increasing of KL supplementation in the diet ($P = 0.078$). Meanwhile, ME tended to linearly decrease with increasing of KL supplementation in MQH and HQH diets ($P < 0.01$). In all experimental diets, NEL was not observed to linearly increase or decrease with increasing of KL supplementation in the diet ($P = 0.161$).




Figure 1. Effect of polyethylene glycol (PEG) treatment on the cumulative gas production (ml/200 mg dry matter; DM) during 48 h (h) of incubation of Katuk leaves.

Discussion

Chemical composition

According to the results of this study, KL had higher CP concentration (333 g/kg DM) than the three hays (i.e. LQH, MQH and HQH), which were grasses from temperate region. The CP concentration of KL was also greater than mean CP concentrations of tropical grasses and tropical forage legumes as determined in a previous study (79 and 198 g/kg DM, respectively; Nurdianti *et al.*, 2019). Piliang and Djojosoebagio (1991) reported a CP concentration of KL of about 257 g/kg DM. Meanwhile, Yang and Guo (2002) found that CP concentration of KL grown in South China reached up to 485 g/kg DM. Moreover, a study from Naveena *et al.* (2020) has been conducted to evaluate the nutritional composition of basal whorl leaves and terminal whorl leaves of Katuk which reported that basal whorl leaves are more enriched with nutrients than terminal whorl leaves. The authors further mentioned that the CP concentration of KL increased with the age of the plant. The present research and some previous studies showed that CP concentrations of KL exceed the requirements of ruminants, for instance, of small ruminants for maintenance and growth (110–130 g/kg DM of CP concentration; National Research Council, 2007) and even of dairy cattle for maintenance and lactation (119 g/kg DM of CP concentration; National Research Council, 2001), which makes it valuable as high-protein supplement particularly to low-protein forage diets.

The aNDFom concentration of KL was lower than of the three hays (i.e. LQH, MQH and HQH) in the present study and those reported for tropical forage legumes or tropical grasses in the literature (374 and 592 g/kg DM, respectively; Nurdianti *et al.*, 2019). Hence, supplementing KL would decrease the aNDFom concentration in the high-fibre diets. Villalba *et al.* (2021) mentioned that NFC is also important in the ruminant nutrition as it

can offer energy adequately for an efficient microbial protein synthesis (Shabi *et al.*, 1998). The present study reported that supplementation of KL in the diet can increase NFC concentration in all experimental diets. Batajoo and Shaver (1994) summarized that for cows producing over 40 kg of milk, the diet should contain NFC concentration about more than 300 g/kg DM, yet little benefit was reported if the diets contain 420 g/kg DM over 360 g/kg DM of NFC concentration. In previous study, Villalba *et al.* (2021) mentioned that the decreased dietary fibre and increased NFC concentration might lead to increased intake, increased meat and milk in ruminant production and decreased methane enteric emissions and carbon dioxide. The present study showed that KL inclusion can increase CP and NFC concentrations, while reducing aNDFom concentrations in ruminant diets based on forage grasses, and that it may thus be a valuable supplement.

Tannin bioassay

According to Makkar *et al.* (1995), PEG might bind to tannins and thereby reduce their anti-nutritive activity. In response to the deactivation of secondary compounds, nutrient fermentation might improve and *in vitro* GP might increase (Batajoo and Shaver, 1994; Nocek, 1997). The greater the increase in the GP, the higher the suppressive activity of tannin in the feeds (Jayanegara *et al.*, 2009).

A study of Selvi and Bhaskar (2012) reported that the KL contain substances such as sterols, resins, tannins, saponins, alkaloids, flavonoids, terpenoids, cardiac glycosides, phenols and catechols. Previous studies reported that KL contain per 88.7 mg tannins (Singh *et al.*, 2011) and 580 mg alkaloid papaverine (Bender and Ismail, 1975; Padmavathi and Rao, 1990) per 100 g DM of KL, as well as 11.5 mg gallic acid equivalents of total phenolics

Table 3. Kinetics cumulative gas production of the experimental diets ($n = 3$)

Variable	Hay												SEM	<i>P</i> value ^a			
	Low-quality hay				Medium-quality hay				High-quality hay					<i>H</i>	<i>L</i>	<i>H</i> × <i>L</i>	CL
	0	50	100	200	0	50	100	200	0	50	100	200					
Cumulative GP (ml/200 mg DM)																	
GP ₁₂	21.9	22.7	23.2	24.1	35.7	34.8	34.8	34.7	34.1	33.6	33.2	33.3	0.664	<0.001	0.506	0.007	0.410
GP ₂₄	33.4	33.3	33.8	34.0	49.6	48.2	47.7	46.4	46.0	45.3	44.6	43.7	0.875	<0.001	0.001	0.004	<0.001
GP ₄₈	43.1	42.6	42.6	42.6	57.6	55.9	55.5	53.9	52.5	51.3	50.8	49.5	0.833	<0.001	<0.001	0.101	<0.001
GP ₉₆	50.1	49.3	48.8	48.4	62.2	60.8	60.1	58.3	56.2	55.0	54.5	52.9	0.800	<0.001	<0.001	0.410	<0.001
Parameters of Gompertz function																	
<i>A</i> ^a (ml/200 mg DM)	48.3 (1.2)	47.5 (1.4)	46.9 (2.5)	46.4 (1.4)	59.8 (1.6)	58.2 (0.3)	57.6 (2.1)	56.0 (1.3)	54.0 (0.2)	52.7 (0.4)	52.3 (1.2)	50.7 (0.6)					
<i>b</i> ^a (ml/h)	1.15 (0.0)	1.14 (0.1)	1.19 (0.0)	1.22 (0.0)	2.20 (0.1)	2.16 (0.1)	2.14 (0.1)	2.12 (0.1)	2.24 (0.1)	2.26 (0.1)	2.17 (0.2)	2.17 (0.1)					
LAG ^a (h)	−6.10 (0.1)	−6.65 (0.6)	−6.38 (0.3)	−6.61 (0.2)	−3.77 (0.1)	−3.69 (0.2)	−3.87 (0.1)	−3.69 (0.2)	−2.89 (0.1)	−2.64 (0.2)	−2.98 (0.3)	−3.18 (0.0)					

DM, dry matter; GP, gas production; GP₁₂, gas production during 12 h of incubation; GP₂₄, gas production during 24 h of incubation; GP₄₈, gas production during 48 h of incubation; GP₉₆, gas production during 96 h of incubation; KL, Katuk leaves; SEM, standard error mean.

^a*A*, theoretical maximum of gas production (ml/200 mg dry matter); *b*, maximum rate of gas production (ml/h) that occurs at the point of inflection of the curve; LAG, lag time (h), which is defined as the time-axis intercept of a tangent line at the point of inflection; *t*, time (h). Probability of the effects of *H*, hays of different quality; *L*, inclusion levels of Katuk leaves; *H*×*L*, interaction of hays of different quality and inclusion levels of Katuk leaves; CL, linear effect of inclusion levels of Katuk leaves.

Table 4. Calculated metabolizable energy (MJ/kg DM) and net energy for lactation concentrations of the experimental diets (MJ/kg DM)

Variable	Hay												P value ^a				
	Low-quality hay				Medium-quality hay				High-quality hay								
	0	50	100	200	0	50	100	200	0	50	100	200	SEM	H	L	H×L	CL
KL level (g/kg DM)	7.0	7.1	7.2	7.4	9.5	9.4	9.4	9.4	9.5	9.5	9.5	9.4	0.119	<0.001	0.207	0.001	0.078
ME (MJ/kg DM)	3.9	4.0	4.1	4.2	5.7	5.6	5.6	5.6	5.7	5.6	5.6	5.6	0.084	<0.001	0.324	0.001	0.161
NEL (MJ/kg DM)																	

DM, dry matter; KL, Katuk leaves; ME, metabolizable energy; NEL, net energy for lactation; SEM, standard error mean.

^aProbability of the effects of H, hays of different quality; L, inclusion levels of Katuk leaves; H×L, interaction of hays of different quality and inclusion levels of Katuk leaves; CL, linear effect of inclusion levels of Katuk leaves.

and 10.4 mg rutin equivalents of total flavonoids (Maisuthisakul *et al.*, 2008) per gram db. However, in contrast to these results, the present study showed that PEG addition did not increase GP of KL indicating that KL did not contain a considerable amount of tannins.

Fibre fractions and in vitro fibre digestibility

Fibre digestibility is important in assessing forage quality (Ward, 2001). Fibre digestibility and pdNDF concentration can be calculated by subtracting the uNDF₂₄₀ concentration from total aNDFom concentration (Cotanch *et al.*, 2014; Nurdianti *et al.*, 2019). The uNDF₂₄₀ concentration is the functional fibre fraction that influences physical effectiveness, gut fill and digestion/passage dynamics of forages (Cotanch *et al.*, 2014; Harper and McNeill, 2015). Fustini *et al.* (2017) reported that voluntary feed intake is influenced and improved by forage fibre digestibility and its pdNDF concentration (when it represents up to 500 g/kg DM basis of the ration).

The uNDF₂₄₀ concentration of KL in the present study was much lower than reported for tropical forage legumes and grasses (294 and 231 g/kg DM; Nurdianti *et al.*, 2019). On the other hand, the pdNDF concentration of KL interestingly was much higher compared to tropical forage legumes and grasses (134 and 360 g/kg DM, respectively; Nurdianti *et al.*, 2019), which shows that fibre of KL is more digestible than that of other tropical forages. Therefore, the inclusion of KL in diets with hays of different quality decreased the concentration of uNDF₂₄₀, but might increase the fibre digestibility of the diet.

In vitro gas production, metabolizable energy and net energy for lactation

In the exponential model, the GP rate depends on substrate availability for fermentation after a lag time has been reached (Ørskov and McDonald, 1979; McDonald, 1981). According to López *et al.* (2007), the lag time might be affected by some factors such as the nature of the feedstuff incubated, the microbial species inoculated and the quantity of inoculum added. In the present study, the Gompertz model was the most suitable curve shapes until the end of incubation. According to Lavrenčič *et al.* (1997), the Gompertz model assumes that the specific GP rate is proportional to microbial mass, which in turn depends on the concentration of digestible substrate. Moreover, the present study showed negative lag time which is similar to the finding of Jijai *et al.* (2016) which indicates that the accelerated growth of the initial anaerobic process in most batches was facilitated by favourable substrate conditions, resulting in a significant reduction in the time required to reach the exponential phase.

Related to gas volume and *in vitro* GP characteristics, Menke *et al.* (1979) suggested that GP₂₄ has indirect relationship with organic matter digestibility and ME in feedstuffs due to the stoichiometric relationships between organic matter degradation, short-chain fatty acid (SCFA) yield and GP. Therefore, ME and SCFA yield may be predicted from *in vitro* GP (Batajoo and Shaver, 1994). Yet, GP derived from protein fermentation is relatively small as compared to that from carbohydrate fermentation (Makkar, 2004). Hence, the high CP concentration of KL in the present study rather than a poor substrate degradability might have contributed to their low GP as well as the linear decline with increasing KL inclusion level in the grass hay diet.

The estimation of the ME values can be used for the purpose of ration formulation and for other purposes in setting economic value of feeds (Getachew *et al.*, 2002). The ME value of KL was 9.52 MJ/kg DM which is good compared with typical values of feedstuffs commonly fed to cattle, such as alfalfa, barley silage, corn silage, cotton seed and sorghum silage (8.20, 8.49, 9.74 and 7.49 MJ/kg DM, respectively; National Research Council, 2001) which can give benefit economically when supplementing KL in the diet. Meanwhile, the NEL value of KL was 5.62 MJ/kg DM which is sufficient as compared to some feedstuffs commonly fed to cattle, such as alfalfa and barley grain (6.36 and 7.78 MJ/kg DM, respectively; National Research Council, 2001).

Conclusion

Katuk leaves have relatively high CP concentrations with low concentration of well digestible fibre. There was no indication that KL contained a considerable amount of tannins, hampering carbohydrate fermentation. In LQH diets, increasing KL supplementation linearly increases GP₂₄ and calculated ME, with no or even negative effects in MQH and HQH diets. Hence, KL can be a potential supplement feed for ruminant livestock, in particular when fed in addition to LQH. However, further studies (e.g. *in vitro* or *in vivo*) investigating other rumen parameters after incubation are needed to validate the current observations.

Acknowledgements. The study of the first author was funded by Lembaga Pengelola Dana Pendidikan (Indonesia Endowment Fund for Education) Scholarship, Indonesia. The proximate analyses and *in vitro* HGT were performed at the Institute of Animal Science of the University of Bonn, Germany. The fibre digestibility was determined at the Department of Animal Nutrition and Rangeland Management in the Tropics and Subtropics (490i) of Institute of Agricultural Sciences in the Tropics (Hans-Ruthenberg Institute) of the University of Hohenheim, Germany and we appreciate the supply of rumen fluid by the Institute of Animal Science of the University of Hohenheim, Germany. We appreciate the technical assistance and resources of all working groups during laboratory works.

Author contributions. R. R. N., K.-H. S. and C. B. conceived and designed the study. R. R. N. conducted the experiments and data gathering. R. R. N. owned the study and research funds. R. R. N. and C. B. performed statistical analyses. K.-H. S. and U. D. provided the research facilities. R. R. N., K.-H. S., C. B., A. H. and R. S. N. provided the samples. K.-H. S., C. B., U. D., A. J., D. R., I. H., A. H. and R. S. N. reviewed the paper. K.-H. S. and C. B. supervised the research. R. R. N. wrote the original article and revised the manuscript. All authors have read and agreed to the final version of the manuscript.

Financial support. The study of the first author was funded by Lembaga Pengelola Dana Pendidikan (Indonesia Endowment Fund for Education) Scholarship, Indonesia.

Competing interests. None.

Ethical standards. The study was conducted according to the guidelines of Institute of Animal Science of the University of Bonn and the Institute of Agricultural Sciences in the Tropics (Hans-Ruthenberg Institute) of the University of Hohenheim. All procedures for animal handling within the present study were performed according to the National Committee for the Protection of Animals Used for Scientific Purposes for the Federal Republic of Germany.

References

Batajoo KK and Shaver RD (1994) Impact of nonfiber carbohydrate on intake, digestion, and milk production by dairy cows. *Journal of Dairy Science* 77, 1580–1588.

- Bender AE and Ismail KS (1975) Nutritive value & toxicity of a Malaysian food, *Sauropus albicans*. *Plants Foods for Man* 1, 139–143.
- BMKG (2013) *Pusat Database Badan Meteorologi, Klimatologi, Dan Geofisika*. Bandung, Indonesia.
- Cotanch KW, Grant RJ, Van Amburgh ME, Zontini A, Fustini M, Palmonari A and Formigoni A (2014) Applications of uNDF in ration modeling and formulation. In Cornell Nutrition Conference for Feed Manufacturers, New York, USA, 21st October 2014, p. Ithaca, New York, USA: Cornell University.
- Food and Agriculture Organization (2014) Dairy production and products: cattle. <http://www.fao.org/dairy-production-products/production/dairy-animals/cattle/en/>
- Fox DG, Tedeschi LO, Tylutki TP, Russell JB, Van Amburgh ME, Chase LE, Pell AN and Overton TR (2004) The Cornell Net Carbohydrate and Protein System model for evaluating herd nutrition and nutrient excretion. *Animal Feed Science and Technology* 112, 29–78.
- Fustini M, Palmonari A, Canestrari G, Bonfante E, Mammi L, Pacchioli MT, Sniffen GCJ, Grant RJ, Cotanch KW and Formigoni A (2017) Effect of undigested neutral detergent fiber content of alfalfa hay on lactating dairy cows: feeding behavior, fiber digestibility, and lactation performance. *Journal of Dairy Science* 100, 4475–4483.
- Geremew G, Negesse T and Abebe A (2017) Assessment of availability and nutritive values of feed resources and their contribution to livestock nutrient requirements in Chire district, Southern Ethiopia. *Agricultural Research & Technology: Open Access* 7, 00114–00118.
- Getachew G, Crovetto GM, Fondevila M, Krishnamoorthy U, Singh B, Spanghero M, Steingass H, Robinson PH and Kailas MM (2002) Laboratory variation of 24 h *in vitro* gas production and estimated metabolizable energy values of ruminant feeds. *Animal Feed Science and Technology* 102, 169–180.
- Goering HK and Van Soest PJ (1970) *Forage Fiber Analyses: (Apparatus, Reagents, Procedures, and Some Applications)*. Washington, DC, USA: Agricultural Research Service, U.S. Dept. of Agriculture.
- Harper KJ and McNeill D (2015) The role iNDF in the regulation of feed intake and the importance of its assessment in subtropical ruminant systems (the role of iNDF in the regulation of forage intake). *Agriculture* 5, 778–790.
- Jayanegara A, Togtokhbayer N, Makkar HPS and Becker K (2009) Tannins determined by various methods as predictors of methane production reduction potential of plants by an *in vitro* rumen fermentation system. *Animal Feed Science and Technology* 150, 230–237.
- Jijai S, Srisuwan G, O-Thong S, Norli I and Siripatana C (2016) Effect of substrate and granules/inocula sizes on biochemical methane potential and methane kinetics. *Iranica Journal of Energy & Environment* 7, 94–101.
- Lavrenčić A, Stefanon B and Susmel P (1997) An evaluation of the Gompertz model in degradability studies of forage chemical components. *Animal Science* 64, 423–431.
- López S, Dhanoa MS, Dijkstra J, Bannink A, Kebreab E and France J (2007) Some methodological and analytical considerations regarding application of the gas production technique. *Animal Feed Science and Technology* 135, 139–156.
- Maisuthisakul P, Pasuk S and Ritthiruangdej P (2008) Relationship between antioxidant properties and chemical composition of some Thai plants. *Journal of Food Composition and Analysis* 21, 229–240.
- Makkar HPS (2004) *Assessing Quality and Safety of Animal Feeds* (Ed S. Jutzi). Rome, Italy: FAO.
- Makkar HPS, Blümmel M and Becker K (1995) *In vitro* effects of and interactions between tannins and saponins and fate of tannins in the rumen. *Journal of the Science of Food and Agriculture* 69, 481–493.
- McDonald I (1981) A revised model for the estimation of protein degradability in the rumen. *The Journal of Agricultural Science* 96, 251–252.
- Menke KH and Steingass H (1988) Estimation of the energetic feed value obtained from chemical analysis and *in vitro* gas production using rumen fluid. *Animal Resources Development* 28, 7–55.
- Menke KH, Raab L, Salewski A, Steingass H, Fritz D and Schneider W (1979) The estimation of the digestibility and metabolizable energy content of ruminant feeding stuffs from the gas production when they are incubated with rumen liquor *in vitro*. *The Journal of Agricultural Science* 93, 217.

- Mertens DR** (2002) Measuring fiber and its effectiveness in ruminant diets. In Plains Nutrition Council: Spring Conference, San Antonio, USA, 25th–26th April 2002, pp. 40–66. San Antonio, USA: Texas A&M Research and Extension Center.
- Nasser MEA, El-Waziry AM and Sallam SMA** (2009) *In vitro* gas production measurements and estimated energy value and microbial protein to investigate associative effects of untreated or biological treated linen straw and berseem hay. In Nutritional and Foraging Ecology of Sheep and Goats, Thessaloniki, Greece, 11st–13rd October 2007 (Ed P. Papachristou, T.G., Parissi, Z.M., Ben Salem, H., Morand-Fehr), pp. 261–266. Montpellier, France: Centre International de Hautes Etudes Agronomiques Méditerranéennes.
- National Research Council** (2001) Nutrient requirements of dairy cattle: seventh revised edition, 2001, 7th ed (Eds Board on Agriculture and Natural Resources, Subcommittee on Dairy Cattle Nutrition, & Committee on Animal Nutrition). Washington DC, USA: The National Academies Press.
- National Research Council** (2007) Nutrient requirements of small ruminant: sheep, goats, cervids, and new world camelids, 4th ed (Eds Committee on Nutrient Requirement of Small Ruminants, Board on Agriculture and Natural Resources, & Division on Earth and Life Studies). Washington DC, USA: The National Academies Press.
- Naveena E, Janavi GJ, Arumugam T and Anitha T** (2020) Estimation of nutritive composition of *Sauropus androgynus* (multivitamin plant) at different growth stages and position of leaves. *International Journal of Chemical Studies* **8**, 443–447.
- Noach YR, Handayani HT and Henuk YL** (2020) Birth weight, milk production, and milk quality of Ettawah grade goat at first kidding period supplemented with concentrate contained katuk (*Sauropus androgynus*) leaf flour and Zn bio complex. In IOP Conference Series: Earth and Environmental Science, p. 012060. IOP Publishing.
- Nocek JE** (1997) Bovine acidosis: implications on laminitis. *Journal of Dairy Science* **80**, 1005–1028.
- Nurdianti RR, Südekum KH and Böttger C** (2017) *In vitro* rumen fermentation of *Sauropus androgynus* (L.) Merr. compared to hays of different quality. In Tropentag 2017 – Future Agriculture: socio-ecological transitions and bio-cultural shifts, Bonn, Germany, 20th–22nd September 2017 (Ed E. Tielkes), p. 363. Göttingen, Germany: Cuvillier Verlag.
- Nurdianti RR, Dickhöfer U and Castro-Montoya JM** (2019) Evaluation of undigested and potential digestible fiber in tropical grasses and tropical legumes. In Tropentag 2019 – Filling gaps and removing traps for sustainable resource management, Kassel, Germany, 18th–20th September 2019 (Ed E. Tielkes), p. 217. Göttingen, Germany: Cuvillier Verlag.
- Ørskov ER and McDonald I** (1979) The estimation of protein degradability in the rumen from incubation measurements weighted according to rate of passage. *The Journal of Agricultural Science* **92**, 499–503.
- Padmavathi P and Rao MP** (1990) Nutritive value of *Sauropus androgynus* leaves. *Plant Foods for Human Nutrition* **40**, 107–113.
- Palmonari A, Fustini M, Canestrari G, Grilli E and Formigoni A** (2014) Influence of maturity on alfalfa hay nutritional fractions and indigestible fiber content. *Journal of Dairy Science* **97**, 7729–7734.
- Paul M and Beena Anto K** (2011) Antibacterial activity of *Sauropus androgynus* (L.) Merr. *International Journal of Plant Sciences* **6**, 189–192.
- Piliang WG and Djojosebagio S** (1991) *Fisiologi Nutrisi* (Ed IPB). Bogor, Indonesia: Depdikbud.
- Raffrenato E and Van Amburgh M** (2010) Development of a mathematical model to predict sizes and rates of digestion of a fast and slow degrading pool and the indigestible NDF fraction. In Cornell Nutrition Conference for Feed Manufacturers, New York, USA, 19th–21st October 2010, pp. 52–65. NY, USA, 2010.
- Raffrenato E, Ross DA and Van Amburgh ME** (2018) Development of an *in vitro* method to determine rumen undigested aNDFom for use in feed evaluation. *Journal of Dairy Science* **101**, 9888–9900.
- Rinne M, Jaakkola S and Huhtanen P** (1997) Grass maturity effects on cattle fed silage-based diets. 1. Organic matter digestion, rumen fermentation and nitrogen utilization. *Animal Feed Science and Technology* **67**, 1–17.
- Robertson JB and Van Soest PJ** (1981) The detergent system of analysis and its applications to human foods. In James WPT and Theander O (eds), *The Analysis of Dietary Fiber in Food*. New York, USA: Marcel Dekker, pp. 123–158.
- Salamatazar M, Salamatdoust-nobar R and Sis NM** (2012) Evaluation of the effects of *Thymus vulgar* on degradability kinetics of canola meal for ruminant using *in vitro* gas production technique. *Journal of Cell and Animal Biology* **6**, 164–168.
- Sa'roni S, Sadjiman T, Sja'bani M and Zulaela Z** (2012) Effectiveness of the *Sauropus androgynus* (L.) Merr leaf extract in increasing mother's breast milk production. *Media Penelitian dan Pengembangan Kesehatan* **14**, 20–24.
- Schofield P, Pitt RE and Pell AN** (1994) Kinetics of fiber digestion from *in vitro* gas production. *Journal of Animal Science* **72**, 2980–2991.
- Selvi VS and Bhaskar A** (2012) Characterization of anti-inflammatory activities and antinociceptive effects of papaverine from *Sauropus androgynus* (L.) Merr. *Global Journal of Pharmacology* **6**, 186–192.
- Shabi Z, Arieli A, Bruckental I, Aharoni Y, Zamwel S, Bor A and Tagari H** (1998) Effect of the synchronization of the degradation of dietary crude protein and organic matter and feeding frequency on ruminal fermentation and flow of digesta in the abomasum of dairy cows. *Journal of Dairy Science* **81**, 1991–2000.
- Singh S, Singh DR, Salim KM, Srivastava A, Singh LB and Srivastava RC** (2011) Estimation of proximate composition, micronutrients and phytochemical compounds in traditional vegetables from Andaman and Nicobar Islands. *International Journal of Food Sciences and Nutrition* **62**, 765–773.
- Sommart K, Parker DS, Rowlinson P and Wanapat M** (2000) Fermentation characteristics and microbial protein synthesis in an *in vitro* system using cassava, rice straw and dried Ruzi grass as substrates. *Asian-Australasian Journal of Animal Sciences* **13**, 1084–1093.
- Taghizadeh A, Palangi V and Safamehr A** (2008) Determining nutritive values of alfalfa cuts using *in situ* and gas production techniques. *American Journal of Animal and Veterinary Sciences* **3**, 85–90.
- Tilley JMA and Terry RA** (1963) A two-stage technique for the *in vitro* digestion of forage crops. *Grass and Forage Science* **18**, 104–111.
- Verband Deutscher Landwirtschaftlicher Untersuchungs- und Forschungsanstalten** (2007) *VDLUFa-Methodenbuch: Bd. III Die Chemische Untersuchung von Futtermitteln*. Darmstadt, Germany: VDLUFa-Verlag.
- Villalba JJ, Ates S and MacAdam JW** (2021) Non-fiber carbohydrates in forages and their influence on beef production systems. *Frontiers in Sustainable Food Systems* **5**, 1–12.
- Ward RT** (2001) NDF digestibility and its use in predicting forage quality.
- Yang C and Guo JX** (2002) South China wild vegetable nutritive composition. *Food Science* **11**, 121–125.
- Zhang Bd, Cheng Jx, Zhang Cf, Bai Yd, Liu Wy, Li W, Koike K, Akihisa T, Feng F and Zhang J** (2020) *Sauropus androgynus* L. Merr. – a phytochemical, pharmacological and toxicological review. *Journal of Ethnopharmacology* **257**, 112778.