

ON INGHAM'S SUMMATION METHOD

S. L. SEGAL

Ingham (2) has defined the following summation method. A series $\sum a_n$ will be called summable (I) to s if

$$\lim_{x \rightarrow \infty} \sum_{n \leq x} \frac{n}{x} \left[\frac{x}{n} \right] a_n = s,$$

where as usual $[x]$ is the greatest integer $\leq x$. (An equivalent method was described somewhat earlier by Wintner (7), who called it "an Eratosthenian method"; however, the notation (I) and the name "Ingham summability" introduced by Hardy (1) seem to have become usual.)

Ingham has proved that although the method (I) is not comparable with convergence (and so, in particular, not regular), for every δ , $0 < \delta < 1$, summability $(C, -\delta)$ implies summability (I), and for every $\delta > 0$, summability (I) implies summability (C, δ) , where (C, k) denotes the Cesàro mean of order k . Pennington (4) and Rajagopal (5) have in fact given explicit constructions of convergent series whose Ingham sum is unbounded as $x \rightarrow \infty$.

In the present paper we are mainly concerned with Tauberian converses of the above results and the possibility or impossibility of inferring (I)-summability from Abel or (C, k) -summability for some $k \geq 0$ under certain auxiliary restrictions on $\{a_n\}$. Other Tauberian theorems stated by Rajagopal (5) with brief indications of proof connect generalized Lambert summability and Pennington's generalization of Ingham summability (4). A special case of one of these results is that Lambert summability (and so *a fortiori* (C, k) summability) together with a Schmidt condition on $I(t)$ (see below) implies the convergence of $I(t)$ as $t \rightarrow \infty$.

The related question of "limitation" theorems for (I)-summability is also discussed.

Throughout this paper all sequences are of real numbers. $\mu(n)$ will denote the Möbius function

$$M(x) = \sum_{n \leq x} \mu(n), \quad N(x) = \sum_{n \leq x} \frac{\mu(n)}{n},$$

$\phi(n)$ is Euler's ϕ -function, $[x]$ is the integral part of x , and $\{x\} = x - [x]$. $\sum_{d|n}$ will denote a sum over the positive divisors of n . Given a series $\sum a_n$, it will be convenient to denote the sum

$$\sum_{n \leq t} \frac{n}{t} \left[\frac{t}{n} \right] a_n$$

Received September 12, 1964. This work has been partially supported by NSF Grant GP 2499. The author is currently a Fulbright Research Fellow at the University of Vienna.

by $I(t)$, so that the (I)-sum of $\sum a_n$ is equal to $\lim_{t \rightarrow \infty} I(t)$ (if it exists). The series $\sum a_n$ will be called (I)-bounded if

$$-\infty < \liminf_{t \rightarrow \infty} I(t) \leq \limsup_{t \rightarrow \infty} I(t) < \infty.$$

Since no more than one series will be under consideration at any one time, the notation $I(t)$ (which depends implicitly on $\sum a_n$) should not lead to any confusion. All other unexplained notation or terminology is as in (1).

THEOREM 1. *Let $\sum a_n$ be a series of real numbers which is Abel-summable and such that*

$$\sum_{d|n} da_d = O(1) \quad \text{as } n \rightarrow \infty.$$

Then $\sum a_n$ is summable (I).

Proof. We show first that it is sufficient to prove the theorem if Abel-summability is replaced by summability $(C, 1)$. Let $\sum_{d|n} da_d = b_n$. Then

$$(1) \quad a_n = \frac{1}{n} \sum_{d|n} \mu\left(\frac{n}{d}\right) b_d$$

by Möbius inversion and

$$\sum_{n \leq r} a_n = \sum_{n \leq r} \sum_{d|n} \frac{\mu(r/d)}{r/d} \frac{b_d}{d} = \sum_{d \leq r} \frac{b_d}{d} N(r/d) = \sum_{d=1}^{\infty} \frac{b_d}{d} N(r/d)$$

(for $d > r$, the sum defining $N(r/d)$ is void and $N(r/d) = 0$). And so

$$\left| \sum_{n \leq r} a_n \right| \leq \sum_{d=1}^{\infty} \frac{|b_d|}{d} |N(r/d)| \leq \text{Sup}_d |b_d| \text{Sup}_r \sum_{d=1}^{\infty} \frac{1}{d} |N(r/d)|.$$

But the second factor is known to be bounded (6, correction) and b_d is bounded by hypothesis. Hence $\sum a_n$ is bounded, and so by a well-known Tauberian theorem (1, Theorem 92), $\sum a_n$ is $(C, 1)$ summable.

The condition $\sum_{d|n} da_d = O(1)$ also implies that

$$(2) \quad I(t) = \sum_{n \leq t} a_n \frac{n}{t} \left[\frac{t}{n} \right] = \frac{1}{t} \sum_{n \leq t} \sum_{d|n} da_d = O(1),$$

and that for $y > x$,

$$\begin{aligned} I(y) - I(x) &= \frac{1}{y} \sum_{n \leq y} \sum_{d|n} da_d - \frac{1}{x} \sum_{n \leq x} \sum_{d|n} da_d \\ &= \left(\frac{1}{y} - \frac{1}{x} \right) \sum_{n \leq x} \sum_{d|n} da_d + \frac{1}{y} \sum_{x < n \leq y} \sum_{d|n} da_d \\ &= -\left(\frac{y-x}{y} \right) I(x) + O\left(\frac{y-x}{y} \right) = O\left(\frac{y-x}{y} \right), \end{aligned}$$

since $I(x) = O(1)$ by (2). Hence $I(y) - I(x) \rightarrow 0$ whenever x and y are such that $y > x$ and as $x \rightarrow \infty$, $y/x \rightarrow 1$. It follows that $I(t)$ is a slowly oscillating

function defined on $(0, \infty)$ in the sense of Schmidt. (There are two definitions of slowly oscillating functions according as the function in question is defined over $(0, \infty)$ or $(-\infty, \infty)$. Here we use the definition appropriate to $(0, \infty)$; see **(1)**, pp. 124, 288.)

We could now appeal to Rajagopal's result quoted above. However, a direct proof based on the known result

$$(3) \quad \int_0^\infty \frac{I(t)}{t} N(x/t) dt = \frac{1}{x} \sum_{n \leq x} (x - n)a_n$$

(e.g. **1**, p. 377) and using the Wiener-Pitt Tauberian theory is also possible and is given below. For completeness, we give first a proof of (3) slightly different from that in **(1)**.

Since for $t < 1$, $N(t) = 0$ and $I(t) = 0$, we have

$$\begin{aligned} \int_0^\infty \frac{I(t)N(x/t)}{t} dt &= \int_1^x \frac{I(t)}{t} N(x/t) dt = \int_1^x \frac{N(x/t)}{t^2} \sum_{a \leq t} da \left[\frac{t}{d} \right] dt \\ &= \sum_{a \leq x} da \int_a^x \frac{N(x/t)}{t^2} \left[\frac{t}{d} \right] dt = \frac{1}{x} \sum_{a \leq x} da \int_1^{x/a} N(u) \left[\frac{x}{ud} \right] du \\ &= \frac{1}{x} \sum_{a \leq x} da \sum_{k \leq x/a} \int_1^{x/ka} N(u) du \\ &= \frac{1}{x} \sum_{a \leq x} da \sum_{k \leq x/a} \left\{ \frac{x}{kd} N(x/kd) - M(x/kd) \right\} \\ &= \frac{1}{x} \sum_{a \leq x} (x - d)a_d, \end{aligned}$$

on letting $x/t = u$ and using the fact that by partial summation

$$xN(x) - M(x) = x \sum_{d \leq x} \frac{\mu(d)}{d} - \sum_{d \leq x} \mu(d) = \int_1^x N(t)dt,$$

and that, as is easily verified,

$$\sum_{k \leq x} \frac{1}{k} N(x/k) = \sum_{k \leq x} M(x/k) = 1.$$

But as noted above, $\sum a_n$ is $(C, 1)$ -summable to A , say, under the conditions of the theorem. Hence writing $G(t) = (1/t)N(1/t)$, we have, by (3),

$$(4) \quad \lim_{x \rightarrow \infty} \frac{1}{x} \int_0^\infty I(t)G(t/x) dt = \lim_{x \rightarrow \infty} \int_0^\infty \frac{I(t)}{t} N(x/t) dt = \lim_{x \rightarrow \infty} \frac{1}{x} \sum_{d \leq x} (x - d)a_d = A.$$

Also, by a classical result of Landau, $|N(u)| = O(e^{-\alpha \sqrt{\log u}})$ for some $\alpha > 0$, and $N(u) = 0$ for $0 \leq u < 1$; hence $|N(u)|/u$ is integrable in $(0, \infty)$ and

$$K = \int_0^\infty \frac{|N(u)|}{u} du = \int_0^\infty \frac{1}{t} |N(1/t)| dt = \int_0^\infty |G(t)| dt,$$

on setting $t = 1/u$. Furthermore, by partial summation, we have, by classical results of Landau,

$$(5) \quad \int_0^\infty G(t)dt = \int_0^1 \frac{1}{t} N(1/t) dt = \int_1^\infty \frac{N(u)}{u} du = \lim_{k \rightarrow \infty} \left(N(k) \log k - \sum_{d \leq k} \frac{\mu(d) \log d}{d} \right) = 1.$$

Hence by (5), (4) may be written as

$$(6) \quad \lim_{x \rightarrow \infty} \frac{1}{x} \int_0^\infty I(t)G(t/x) dt = A \int_0^\infty G(t) dt.$$

If

$$\int_0^\infty G(t)t^{-iy} dt \neq 0$$

for any real y , it will then follow by a theorem of Pitt (**1**, Theorem 233) that

$$\lim_{t \rightarrow \infty} I(t) = A,$$

which will prove the theorem. But this last condition is readily verified, since for $y = 0$ it is contained in (5), while for real $y \neq 0$ we have

$$\int_0^\infty G(t)t^{-iy} dt = \int_0^1 N(1/t)t^{-1-iy} dt = \int_1^\infty N(u)u^{-1+iy} du = \lim_{k \rightarrow \infty} \left(\frac{k^{iy}N(k)}{iy} - \frac{1}{iy} \sum_{d \leq k} \frac{\mu(d)}{d^{1-iy}} \right) = -\frac{1}{iy\zeta(1-iy)} \neq 0$$

(where $\zeta(s)$ is the Riemann zeta-function). Hence the theorem follows.

Remarks. If the assumption that $\sum a_n$ is Abel-summable is replaced by the stronger hypothesis of $(C, 1)$ -summability, then the hypothesis $\sum_{d|n} da_d = O(1)$ can be altered to the weaker pair of conditions

- (i) $\sum_{d|n} da_d > -K$ for some constant K ,
- (ii) $\sum a_n$ is (I) -bounded,

and again we may deduce that $\sum a_n$ is (I) -summable since $I(x)$ can then be shown exactly as above to be slowly decreasing in the sense of Schmidt, the remainder of the proof going through as before. In fact, using (**1**, Theorem 94) instead of (**1**, Theorem 92) we may assume Abel summability. However, $(C, 1)$ -summability and condition (ii) alone are not sufficient to imply (I) -summability as the explicit example of Theorem 2 below shows. (It will be noted that the examples of Pennington and Rajagopal of convergent series that are not (I) -summable are not (I) -bounded. On the other hand, the example below, although it is (I) -bounded and $(C, 1)$ -summable, is not convergent.)

Also, under the assumption of $(C, 1)$ -summability of $\sum a_n$ and condition (i), if we replace (ii) by the weaker condition that

$$\frac{1}{x} \int_1^x I(t) dt = O(1),$$

we may deduce the weaker conclusion:

$$\lim_{x \rightarrow \infty} \frac{1}{x} \int_1^x I(t) dt$$

exists. For it is easily verified that under these conditions,

$$S(x) = \frac{1}{x} \int_1^{x^2} I(t) dt$$

is slowly decreasing in the sense of Schmidt, and it then follows by an argument of Hardy (1, p. 304) that

$$\int_1^x \frac{S(t)}{t} N(x/t) dt = A + o(1) \quad \text{as } x \rightarrow \infty,$$

the remainder of the proof being the same as in Theorem 1.

THEOREM 2. *There exists a series $\sum a_n$ that is (I)-bounded and (C, 1)-summable but not (I)-summable.*

Proof. Let

$$a_n = \sum_{d|n} \frac{\mu(d)}{d} (-1)^{n/d}.$$

That $\sum a_n$ is (I)-bounded is almost trivial since by Möbius inversion

$$\sum_{d|n} da_d = n(-1)^n$$

and hence

$$\begin{aligned} I(t) &= \sum_{d \leq t} \frac{d}{t} \left[\frac{t}{d} \right] a_d = \frac{1}{t} \sum_{n \leq t} \sum_{d|n} da_d = \frac{1}{t} \sum_{n \leq t} n(-1)^n \\ &= \frac{1}{t} \left[\frac{t+1}{2} \right] (-1)^{[t]} = O(1); \end{aligned}$$

but does not converge as $t \rightarrow \infty$. Hence $\sum a_n$ is (I)-bounded but not (I)-summable.

To show that $\sum a_n$ is (C, 1) summable, consider first the function

$$F(n) = 4 \sum_{\substack{d|n \\ d \text{ even}}} \phi\left(\frac{1}{2}d\right).$$

Then

$$F(n) = \begin{cases} 0, & \text{if } n \text{ is odd,} \\ 4 \sum_{k|n} \phi(k) = 2n, & \text{if } n \text{ is even.} \end{cases}$$

Hence

$$n(-1)^n = F(n) - n = 4 \sum_{\substack{d|n \\ d \text{ even}}} \phi\left(\frac{1}{2}d\right) - \sum_{d|n} \phi(d).$$

Hence by Möbius inversion,

$$na_n = \sum_{d|n} \mu(d) \frac{n}{d} (-1)^{n/d} = \begin{cases} -\phi(n), & \text{if } n \text{ is odd,} \\ 4\phi(\frac{1}{2}n) - \phi(n), & \text{if } n \text{ is even.} \end{cases}$$

Now by a well-known elementary theorem of Mertens,

$$\sum_{n \leq x} \phi(n) = \frac{3}{\pi^2} x^2 + R(x),$$

where $R(x) = O(x \log x)$. Hence,

$$(7) \quad \sum_{n \leq x} na_n = -\sum_{n \leq x} \phi(n) + 4 \sum_{\substack{n \leq x \\ n \text{ even}}} \phi(\frac{1}{2}n) = -\frac{3}{\pi^2} x^2 - R(x) + 4 \sum_{n \leq \frac{1}{2}x} \phi(n) \\ = 4R(\frac{1}{2}x) - R(x),$$

and also by partial summation,

$$(8) \quad \sum_{n \leq x} a_n = \frac{4R(\frac{1}{2}x) - R(x)}{x} + \int_1^x \frac{4R(\frac{1}{2}t) - R(t)}{t^2} dt$$

(where $R(t) = -3t^2/\pi^2$ for $0 \leq t < 1$). Equations (7) and (8) give for the $(C, 1)$ -mean of $\sum a_n$

$$(9) \quad \frac{1}{x} \sum_{n \leq x} (x - n)a_n = \int_1^x \frac{4R(\frac{1}{2}t)}{t^2} dt - \int_1^x \frac{R(t)}{t^2} dt \\ = \int_1^{\frac{1}{2}x} \frac{R(t)}{t^2} dt - \int_{\frac{1}{2}x}^x \frac{R(t)}{t^2} dt - 3/\pi^2.$$

It remains to show that

$$\int_1^x \frac{R(t)}{t^2} dt$$

converges as $x \rightarrow \infty$. We have on the one hand, by partial summation,

$$(10) \quad \sum_{n \leq x} \frac{\phi(n)}{n} = \frac{6}{\pi^2} x + \frac{R(x)}{x} - \frac{3}{\pi^2} + \int_1^x \frac{R(t)}{t^2} dt$$

and, on the other,

$$(11) \quad \sum_{n \leq x} \frac{\phi(n)}{n} = \sum_{n \leq x} \sum_{d|n} \frac{\mu(d)}{d} = \sum_{d \leq x} \frac{\mu(d)}{d} \left[\frac{x}{d} \right] = \frac{1}{x} \sum_{d \leq x} \mu(d) \frac{x}{d} \left[\frac{x}{d} \right] \\ = \frac{x}{2} \sum_{d \leq x} \frac{\mu(d)}{d^2} + \frac{1}{2x} \sum_{d \leq x} \mu(d) \left[\frac{x}{d} \right]^2 - \frac{1}{2x} \sum_{d \leq x} \mu(d) \left\{ \frac{x}{d} \right\}^2 \\ = \frac{3}{\pi^2} x - \frac{x}{2} \sum_{d > x} \frac{\mu(d)}{d^2} + \frac{1}{x} \sum_{d \leq x} \mu(d) \sum_{\substack{m \leq x/d \\ m \leq x/d}} m - \frac{1}{2x} \sum_{d \leq x} \mu(d) \left[\frac{x}{d} \right] \\ - \frac{1}{2x} \sum_{d \leq x} \mu(d) \left\{ \frac{x}{d} \right\}^2 \\ = \frac{3}{\pi^2} x + \frac{1}{x} \sum_{n \leq x} \phi(n) - \frac{1}{2x} \sum_{d \leq x} \mu(d) \left\{ \frac{x}{d} \right\}^2 + o(1) \\ = \frac{6}{\pi^2} x + \frac{R(x)}{x} + o(1),$$

since

$$\sum_{d \leq x} \mu(d) \left[\frac{x}{d} \right] = 1, \quad \sum_{d > x} \frac{\mu(d)}{d^2} = o(1/x), \quad \sum_{d \leq x} \mu(d) \left\{ \frac{x}{d} \right\}^2 = o(x).$$

(This last can be deduced from "Axer's Theorem" (1, pp. 378, 386).)

Comparing (10) and (11), we have that

$$\lim_{x \rightarrow \infty} \int_1^x \frac{R(t)}{t^3} dt = \frac{3}{\pi^2}$$

and hence by (9) that $\sum a_n$ is $(C, 1)$ -summable to 0, which proves the result.

Remarks. It is obvious that, as mentioned above, $\sum a_n$ is not convergent, since if p is any odd prime, $a_p = (1/p) - 1$ and hence $a_n \neq o(1)$ as $n \rightarrow \infty$.

Pennington (4, p. 79) and Hardy (1, Theorem 266) independently noted the following "limitation" theorem for (I)-summability:

If $\sum a_n$ is (I)-summable, then $a_n = o(\log \log n)$.

A limitation theorem of a somewhat different sort is:

If $\sum a_n$ is (I)-summable, then

$$\sum_{n \leq x} a_n = o(x^\delta)$$

for every $\delta > 0$.

This latter is a corollary of Ingham's result that (I)-summability implies (C, δ) -summability for every $\delta > 0$ and a well-known theorem on Cesàro means (1, Theorem 46). This latter result can be improved to:

THEOREM 3. *If $\sum a_n$ is (I)-summable, then*

$$\sum_{n \leq x} a_n = o(\log x).$$

Proof. By altering a_1 , if necessary, we need only consider the case where $\sum a_n$ is summable (I) to 0. For m a non-negative integer, let

$$K(m) = \begin{cases} mI(m) = \sum_{d \leq m} da_d \left[\frac{m}{d} \right] = \sum_{n \leq m} \sum_{d|n} da_d & \text{if } m \geq 1, \\ 0 & \text{if } m = 0. \end{cases}$$

Then $K(m) = o(m)$ as $m \rightarrow \infty$.

Subtracting and using Möbius inversion gives

$$a_m = \frac{1}{m} \sum_{d|m} \mu(m/d) (K(d) - K(d - 1)) = \sum_{d \tau = m} \frac{\mu(r)}{r} \frac{K(d) - K(d - 1)}{d}$$

and hence

$$(12) \quad \sum_{m \leq x} a_m = \sum_{d \tau \leq x} \frac{\mu(\tau) K(d) - K(d-1)}{\tau d} = \sum_{d \leq x} \frac{K(d) - K(d-1)}{d} N(x/d) \\ = \sum_{d \leq x} \frac{K(d)}{d} d \left(\frac{N(x/d)}{d} - \frac{N(x/(d+1))}{d+1} \right)$$

since $K(0) = 0$ and $N(x/[x] + 1) = 0$.

Since $K(m) = o(m)$ as $m \rightarrow \infty$, the required result will follow by a simple Abelian argument, provided we can prove that

$$(13) \quad \sum_{d \leq x} d \left| \frac{N(x/d)}{d} - \frac{N(x/(d+1))}{d+1} \right| = O(\log x) \quad \text{as } x \rightarrow \infty$$

and, for any fixed $D \geq 1$,

$$(14) \quad \sum_{d \leq D} d \left| \frac{N(x/d)}{d} - \frac{N(x/(d+1))}{d+1} \right| = O(1) \quad \text{as } x \rightarrow \infty.$$

But the left-hand sides of (13) and (14) are maximized by

$$(15) \quad \sum_{d \leq v} |N(x/d) - N(x/(d+1))| + \sum_{d \leq v} \frac{|N(x/(d+1))|}{d+1},$$

with $v = x$ and $v = D$ respectively.

But since $|\mu(m)| \leq 1$,

$$\sum_{d \leq v} |N(x/d) - N(x/(d+1))| = \left| \sum_{d \leq v} \sum_{x/(d+1) < m \leq x/d} \frac{\mu(m)}{m} \right| \\ \leq \sum_{d \leq v} \sum_{x/(d+1) < m \leq x/d} \frac{1}{m} = \sum_{x/([v]+1) < m \leq x} \frac{1}{m} = O(1 + \log v);$$

and since $|N(x)| \leq 1$,

$$\sum_{d \leq v} \frac{|N(x/(d+1))|}{d+1} \leq \sum_{d \leq v} \frac{1}{d+1} = O(1 + \log v).$$

Taking $v = x$ and $v = D \geq 1$ (D fixed) we obtain (13) and (14) respectively, and so the theorem.

If $\sum a_n$ is (I)-bounded, then $K(m) = O(m)$ and the same proof gives

$$\sum_{n \leq x} a_n = O(\log x),$$

immediately.

Remark. By Rubel's result quoted above, the second sum in (15) is in fact $O(1)$ for $v = x$ also.

REFERENCES

1. G. H. Hardy, *Divergent series* (Oxford, 1949).
2. A. E. Ingham, *Some Tauberian theorems connected with the prime number theorem*, J. London Math. Soc., 20 (1945), 171-180.

3. E. Landau, *Handbuch der Lehre von der Verteilung der Primzahlen* (New York, 1953).
4. W. B. Pennington, *On Ingham summability and summability by Lambert series*, Proc. Camb. Philos. Soc., *51*, (1955), 65–80.
5. C. T. Rajagopal, *A note on Ingham summability and summability by Lambert series*, Proc. Indian Acad. Sci., *A42*, (1955), 41–50.
6. L. A. Rubel, *An Abelian theorem for number-theoretic sums*, Acta Arith., *6* (1960), 175–177, Correction Acta Arith., *6* (1961), 523.
7. A. Wintner, *Eratosthenian averages* (Baltimore, 1943).

University of Rochester