



Weeds

VOLUME 7
OCTOBER 1959
NUMBER 4

Journal of the Weed Society of America

Weeds

Issued Quarterly by the Weed Society of America

C. J. WILLARD, *Editor*, Department of Agronomy, H. & F. Bldg.,
1827 Neil Ave., The Ohio State University, Columbus 10, Ohio.

W. C. JACOB, *Business Manager*, Department of Agronomy, Univer-
sity of Illinois, Urbana, Illinois.

EDITORIAL COMMITTEE

O. C. LEE, Department of Botany and Plant Pathology, Purdue
University, Lafayette, Indiana.

R. A. PETERS, Department of Plant Science, University of Connecti-
cut, Storrs, Connecticut.

W. C. ROBOCKER, Crops Research Division, ARS, USDA, State
College of Washington, Pullman, Washington.

E. G. RODGERS, Department of Agronomy, University of Florida,
Gainesville, Florida.

WEEDS is a quarterly journal published by the Weed Society of America. Editorial office is located at Department of Agronomy, H. and F. Bldg., The Ohio State University, 1827 Neil Ave., Columbus 10, Ohio. Printing is by the W. F. Humphrey Press Inc., Geneva, New York. Subscription price is \$6.00 yearly for four issues; single copies \$1.50. Address all communications regarding subscriptions, advertising, and reprints to W. C. Jacob, Department of Agronomy, University of Illinois, Urbana, Illinois. Inquiries concerning information on manuscripts and other material for publication should be addressed to the Editorial Office. All checks, money orders, and other remittances should be made payable to the Weed Society of America.

Entered as second-class matter at the post office at Urbana,
Illinois with additional entry at Geneva, New York.

Table of Contents

	<i>Page</i>
The Pantothenate-Synthesizing Enzyme, A Metabolic Site in the Herbicidal Action of Chlorinated Aliphatic Acids. J. L. Hilton, J. S. Ard, L. L. Jansen and W. A. Gentner.....	381
Biological Assay of Herbicides for Fish Toxicity. James T. Davis and W. S. Hardcastle.....	397
Elimination of Rodent Cover Adjacent to Apple Trees. LeRoy Holm, Franklin A. Gilbert, and Ernest Haltvick.....	405
Evaluation of Herbicides for Several Species of Ornamentals. S. K. Ries, B. H. Grigsby and H. Davidson.....	409
Mode and Rate of Release of Isopropyl <i>N</i> -(3-chlorophenyl) carbamate from Several Granular Carriers. L. L. Danielson.....	418
Susceptibility of Velvet Mesquite to an Amine and an Ester of 2,4,5-T as Related to Various Biological and Meteorological Factors. Fred H. Tschirley and Herbert M. Hull.....	427
Effect of Gibberellic Acid on Absorption, Translocation, and Degradation of 2,4-D in Red Kidney Bean. Floyd M. Ashton.....	436
Effects of Soil Type and Time on the Herbicidal Activity of CDAA, CDEC, and EPTC. T. J. Sheets.....	442
Rotary Hoeing in Combination with Herbicides and Other Cultivations for Weed Control in Soybeans. Elroy J. Peters, Dayton L. Klingman, and Russell E. Larson.....	449
Combined Use of Pre-emergence Herbicides and Cross-Cultivation in Cotton. Chester L. Foy.....	459
Effectiveness of Aerial Application of Herbicides for Hardwood Control in Louisiana. Fred A. Peevy and Paul Y. Burns.....	463
Herbicidal Activities of Halogenoalkylcarboxylic Acid Esters. I. Germination Inhibition Activities. Katsura Munakata, Katsuyuki Yokoyama, Takeshi Shibata, Akio Harada and Fumio Hara.....	470
Wild Garlic Bulblet Viability Determined by 2,3,5-triphenyltetrazolium Chloride. Glenn C. Klingman.....	474
The Metabolism of Nut Grass (<i>Cyperus rotundus</i> L.) I. The Influence of Various Oxygen and Carbon Dioxide Levels Upon Germination and Respiration. R. D. Palmer and W. K. Porter, Jr.....	481
The Metabolism of Nut Grass (<i>Cyperus rotundus</i> L.) II. The Respiratory Quotient and Its Relation to Storage Materials and Some Terminal Enzymes. R. D. Palmer and W. K. Porter, Jr.....	490
The Metabolism of Nut Grass (<i>Cyperus rotundus</i> L.) III. Polyphenoloxidase in the Dormant Tuber. R. D. Palmer and W. K. Porter, Jr.....	504
The Metabolism of Nut Grass (<i>Cyperus rotundus</i> L.) IV. The Activities of Certain Enzymes from Tubers Treated with Amitrol. R. D. Palmer and W. K. Porter.....	511
Communication:	
Donomen: A Pesticides Nomenclature System. E. C. S. Little.....	518
News and Notes.....	522
Membership List, Weed Society of America.....	525
Sustaining Members.....	539
Index, Volume 7.....	540
Bibliography of Weed Investigations, January to March 1959.....	545


Advertisers Index

Spraying Systems Co.	ii
duPont de Nemours & Co.....	iii
U. S. Borax & Chemical Co.	iv
Stauffer Chemical Co.	v
Chipman Chemical Co.	vi




the precision nozzle for effective spraying

Supplied in a full range of interchangeable orifice tip and strainer sizes to meet every capacity requirement. TeeJet Spray Nozzles for Weed Control by spraying make it possible to take maximum advantage of the chemical and sprayer unit. TeeJet nozzles are precision built and provide a flat spray with uniform distribution. Atomization is properly controlled to give coverage with an absolute minimum of driftage. Patented tip design, with set-back orifice opening protects precision orifice from accidental damage. TeeJet spray nozzles are built for use on spray booms and portable sprayers.



TEEJET SPRAY NOZZLE male pipe connection



TEEJET SPRAY NOZZLE female pipe connection

OFF-CENTER SPRAY NOZZLES

Spraying Systems Spray Nozzles with TeeJet tips are supplied in a variety of special body types to meet any unusual spraying requirement. For example, one type of off-center spray nozzle with swivel body provides a flat spray up to 35 feet wide for spraying areas with a single nozzle, that are not accessible with a boom.



INTERCHANGEABLE ORIFICE TIPS flat and cone spray types

SUPPLEMENTARY EQUIPMENT

Complete accessories relating to nozzle use are supplied. These include strainers, special nozzle fittings, and hand valve equipment.

TeeJet Spray Nozzles are supplied for Weed Control... as well as all other types of agricultural spraying. For complete information and reference data write for Catalog 30.

SPRAYING SYSTEMS CO.
Engineers and Manufacturers

3296 RANDOLPH STREET

BELLWOOD, ILLINOIS

For Agriculture and Industry . . .

Du Pont WEED KILLERS

*offer new economies
and efficiency in killing
weeds, grass and brush*

KARMEX® diuron for weed control in many crops; irrigation and drainage ditches; and industrial areas.

KARMEX® DL for pre-emergence weed control in cotton.

TELVAR® monuron for weed control in many crops; irrigation and drainage ditches; and industrial areas.

KLOBEN neburon for weed and grass control in nursery plantings and tomatoes.

DYBAR fenuron—a pelleted product for dry application to the soil for control of brush in fence rows, drainage ditches, utility and railroad right-of-ways, and other non-cultivated areas.

TRYBEN 200, a new weed killer based on trichlorobenzoic acid, for control of bindweed and other noxious weeds; mixed broad-leaf weeds; and certain woody vines and brush.

AMMATE® X for long-term, low-cost brush control.



Better Things for Better Living *Through Chemistry*



**Four
easy
ways to
Destroy
Weeds**

1. UREABOR[®]

A nonselective, granular complex of sodium borate and substituted urea. Low application rates are a feature. Apply with the special new PCB Spreader for best results.

2. DB[®] Granular


A combination of 2,4-D and sodium borates. Kills deep-rooted, noxious weeds. Low application rates for maximum control with the utmost economy; use the PCB Spreader.
(Not intended for control of grass.)

3. POLYBOR-CHLORATE[®]

Highly soluble; for spray or dry application. It gives a quick knock-down; destroys top growth and roots. A general nonselective herbicide.

4. Concentrated BORASCU[®]

A nonselective, granular material. Apply by hand or with a mechanical spreader. Long residual action.



When you want Nonselective Herbicides

for Dependable Action



*look to United States Borax
& Chemical Corporation*

PACIFIC COAST BORAX COMPANY DIVISION



630 Shatto Place, Los Angeles 5, Calif.

= NEWS ABOUT

VAPAM[®] =

New Application Technique Suggested for Some Crops

VAPAM dosage may be reduced in many cases up to 50% with equal or better results if the treated area is covered with a plastic, paper or fabric tarp. The cover should be loosely spread over the treated area immediately after application, and left on for 24 to 48 hours for best results. The amount of water used to drench VAPAM into the soil may be reduced. When a tarp is used, it may be possible to prepare the seedbed just before applying VAPAM if soil moisture is adequate. Do not apply VAPAM to dry soils.

Careful Soil Preparation a Must

Wide-scale use of VAPAM continues to demonstrate the importance of careful soil preparation. Careful cultivation of the area to be treated should be done a *week* before application when no cover is used. All clods should be broken up, and soil should be loosened deeply. The soil should then be kept moist until time of treatment. If the soil becomes crushed, it should be cultivated lightly just before application of VAPAM. Experience has shown that proper soil preparation is the most important step in applying VAPAM. Soil temperature must be between 60°-90°F. at a 3-inch depth.

Dosage Recommendations Now More Detailed

Dosage recommendations have been made more explicit on the label. Potting and top-dressing soil treatments have been added.

Copy of New Label Available

Write to Stauffer for a copy of the new label, or for further information on VAPAM. Address your inquiry to Stauffer Chemical Company, Agricultural Chemicals Division, 380 Madison Avenue, New York 17, N. Y., or 636 California Street, San Francisco 8, Calif.

® VAPAM is Stauffer Chemical Company's trade-mark (registered in principal countries) for *sodium methyl dithiocarbamate*, a soil fumigant.





ATLACIDE: Safer chlorate weed killer...widely used for non-selective eradication of bindweed, Canada thistle, quack grass, Johnson grass and other tough perennials. Kills roots...discourages regrowth. Applied dry or dissolved in water.

ATLACIDE-2,4-D: Offers dual killing action of sodium chlorate and 2,4-D.

CHLOREA: A combination of sodium chlorate, borate and monuron in powder form. Kills weeds and grasses. Combines the proven effectiveness of chlorate on deep-rooted weeds with the soil-surface action of monuron on shallow-rooted grasses and annual seedling growth. Lasting residual effect inhibits new growth. Does not create a fire hazard when used as directed. Applied dry or as a water-mixed spray.

CHLOREA GRANULAR Similar to Chlorea, but a "pelletized" material. No mixing...easy to apply by hand or with mechanical spreader.

CHLORAX "40": A composition of sodium chlorate and borate...for weed and grass control. Has lasting residual effect. Does not create a fire hazard. Applied dry or as a spray.

ATLAS "A": A 40% sodium arsenite solution (4 lbs. arsenic trioxide per gal.). Destroys vegetation and algae in ponds and lakes. Controls certain turf weeds. Used as general weed killer and to kill trees and stumps. Also used to kill potato vines prior to harvesting.

SODIUM ARSENITE: A powder containing 75% arsenic trioxide. Used for the same purposes as Atlas "A". Applied dry or as a spray.

CHIP-CAL: New low-lime calcium arsenate formulation in granular and spray powder forms. Used as crabgrass preventatives for established turf. Applied any time before crabgrass seed starts to germinate.

2,4-D & 2,4,5-T: 2,4-D Amine and 2,4-D Ester liquids; 2,4-D Ester dusts; Low Volatile 2,4,5-T and Brush Killer.

METHOXONE: Contains MCP sodium salt. Used for control in small grains, flax, rice and grass. Controls same weeds as 2,4-D; considered safer for selective spraying.

BUTOXONE: A liquid containing 4-(2,4-DB). For selective control of many broadleaf weeds in established alfalfa and certain other legumes.

Write for Weed Control Booklets

CHIPMAN CHEMICAL COMPANY, INC.

Chicago, Ill.

Bound Brook, N. J.

Portland, Ore.

Palo Alto, Calif.

Pasadena, Tex.

Bessemer, Ala.

Manufacturers of Weed Killers Since 1912