



Browder's Convergence for One-Parameter Nonexpansive Semigroups

Shigeki Akiyama and Tomonari Suzuki

Abstract. We give the sufficient and necessary conditions of Browder's convergence theorem for one-parameter nonexpansive semigroups which was proved by Suzuki. We also discuss the perfect kernels of topological spaces.

1 Introduction

Let C be a closed convex subset of a Banach space E . A family of mappings $\{T(t) : t \geq 0\}$ is called a *one-parameter strongly continuous semigroup of nonexpansive mappings* (*one-parameter nonexpansive semigroup*, for short) on C if the following are satisfied:

- (i) For each $t \geq 0$, $T(t)$ is a nonexpansive mapping on C , that is,

$$\|T(t)x - T(t)y\| \leq \|x - y\|$$

holds for all $x, y \in C$.

- (ii) $T(s + t) = T(s) \circ T(t)$ for all $s, t \geq 0$.
(iii) For each $x \in C$, the mapping $t \mapsto T(t)x$ from $[0, \infty)$ into C is strongly continuous.

There are six papers concerning the existence of common fixed points of $\{T(t) : t \geq 0\}$; see [1, 2, 4, 5, 9, 11]. Recently, Suzuki [11] proved that $\bigcap_{t \geq 0} F(T(t))$ is nonempty provided every nonexpansive mapping on C has a fixed point, where $F(T(t))$ is the set of all fixed points of $T(t)$. He also proved a semigroup version of Browder's [3] convergence theorem in [10, 12].

Theorem 1.1 ([12]) *Let τ be a nonnegative real number. Let $\{\alpha_n\}$ and $\{t_n\}$ be real sequences satisfying*

- (i) $0 < \alpha_n < 1$ and $0 < t_n$ for $n \in \mathbb{N}$;
(ii) $\lim_n t_n = \tau$;
(iii) $t_n \neq \tau$ for $n \in \mathbb{N}$ and $\lim_n \alpha_n / (t_n - \tau) = 0$.

Let C be a weakly compact convex subset of a Banach space E . Assume that either of the following holds:

Received by the editors January 10, 2009.

Published electronically April 15, 2011.

The authors were supported in part by grants-in-Aid for Scientific Research from the Japanese Ministry of Education, Culture, Sports, Science and Technology (No. 18540020, No. 18740075).

AMS subject classification: 47H20.

Keywords: nonexpansive semigroup, common fixed point, Browder's convergence, perfect kernel.

- E is uniformly convex with uniformly Gâteaux differentiable norm.
- E is uniformly smooth.
- E is a smooth Banach space with the Opial property and the duality mapping J of E is weakly sequentially continuous at zero.

Let $\{T(t) : t \geq 0\}$ be a one-parameter nonexpansive semigroup on C . Fix $u \in C$ and define a sequence $\{u_n\}$ in C by

$$(1.1) \quad u_n = (1 - \alpha_n) T(t_n)u_n + \alpha_n u$$

for $n \in \mathbb{N}$. Then $\{u_n\}$ converges strongly to Pu , where P is the unique sunny nonexpansive retraction from C onto $\bigcap_{t \geq 0} F(T(t))$.

See [6, 7, 15] for the notions such as the Opial property, etc.

In this paper, we give the sufficient and necessary conditions on $\{\alpha_n\}$ and $\{t_n\}$.

2 Sufficiency

Throughout this paper we denote by \mathbb{N} the set of all positive integers and by \mathbb{R} the set of all real numbers.

In this section, we generalize Theorem 1.1.

Theorem 2.1 Let $\{\alpha_n\}$ and $\{t_n\}$ be real sequences satisfying

- $0 < \alpha_n < 1$ and $0 \leq t_n$ for $n \in \mathbb{N}$;
- $\{t_n\}$ is bounded;
- $\lim_n \alpha_n / (t_n - \tau) = 0$ for all $\tau \in [0, \infty)$, where $1/0 = \infty$.

Let $E, C, \{T(t) : t \geq 0\}, P, u$ and $\{u_n\}$ be as in Theorem 1.1. Then $\{u_n\}$ converges strongly to Pu .

Proof Let $\{f(n)\}$ be an arbitrary subsequence of $\{n\}$. Since $\{t_n\}$ is bounded, so is $\{t_{f(n)}\}$. Hence there exists a cluster point $\tau \in [0, \infty)$ of $\{t_{f(n)}\}$. From (iii), there exists $\nu \in \mathbb{N}$ such that $t_{f(n)} \neq \tau$ and $t_{f(n)} \neq 0$ for $n \in \mathbb{N}$ with $n \geq \nu$. We choose a subsequence $\{g(n)\}$ of $\{n\}$ such that $g(1) \geq \nu$ and $\{t_{f \circ g(n)}\}$ converges to τ . From (iii) again, we have

$$\lim_{n \rightarrow \infty} \frac{\alpha_{f \circ g(n)}}{t_{f \circ g(n)} - \tau} = 0.$$

By Theorem 1.1, $\{u_{f \circ g(n)}\}$ converges strongly to Pu . Since $\{f(n)\}$ is arbitrary, we obtain that $\{u_n\}$ converges strongly to Pu . ■

As a direct consequence of Theorem 2.1, we obtain the following.

Corollary 2.2 Let $\{\alpha_n\}$ and $\{t_n\}$ be real sequences satisfying conditions (i)–(iii) of Theorem 2.1. Let $\{T(t) : t \geq 0\}$ be a one-parameter nonexpansive semigroup on a bounded closed convex subset C of a Hilbert space E . Let P be the metric projection from C onto $\bigcap_{t \geq 0} F(T(t))$. Fix $u \in C$ and define a sequence $\{u_n\}$ in C by (1.1). Then $\{u_n\}$ converges strongly to Pu .

We note that we need condition (i) in order to define $\{u_n\}$. In the remainder of this paper, we discuss conditions (ii) and (iii).

3 Necessity

In this section, we shall show that conditions (ii) and (iii) of Theorem 2.1 are best possible, in the sense that we cannot relax these conditions on $\{\alpha_n\}$ and $\{t_n\}$ any more.

For real numbers s and t with $t > 0$, we define “mod” by

$$s \text{ mod } t = s - [s/t] t,$$

where $[s/t]$ is the maximum integer not exceeding s/t .

Lemma 3.1 *Let $\{\alpha_n\}$ and $\{t_n\}$ be real sequences satisfying condition (i) of Theorem 2.1. Assume $\limsup_{n \rightarrow \infty} t_n = \infty$. Then for every nonnegative real number v , there exists a positive real number τ such that*

$$\limsup_{n \rightarrow \infty} \frac{\alpha_n}{(t_n \text{ mod } \tau) - v} = \infty.$$

Proof We shall define two real sequences $\{\varepsilon_n\}$ and $\{\tau_n\}$ and a subsequence $\{f(n)\}$ of $\{n\}$ satisfying the following.

- (i) $0 < \varepsilon_n < 1$ and $v + 1 + \varepsilon_n < \tau_n$.
- (ii) $\alpha_{f(n)} / ((t_{f(n)} \text{ mod } \tau) - v) \geq n$ for $\tau \in [\tau_n - \varepsilon_n, \tau_n]$.
- (iii) $[\tau_n - \varepsilon_n, \tau_n] \supset [\tau_{n+1} - \varepsilon_{n+1}, \tau_{n+1}]$.

We denote $t_{f(n)}$ by s_n and $\alpha_{f(n)}$ by β_n for $n \in \mathbb{N}$. We choose $f(1)$ satisfying $s_1 > 2v + 2$. We put

$$\varepsilon_1 := \beta_1/2 \in (0, 1) \quad \text{and} \quad \tau_1 := s_1 - v > v + 2 > v + 1 + \varepsilon_1.$$

If $\tau \in [\tau_1 - \varepsilon_1, \tau_1]$, then since

$$1 \leq \frac{s_1}{s_1 - v} = \frac{s_1}{\tau_1} \leq \frac{s_1}{\tau} \leq \frac{s_1}{\tau_1 - \varepsilon_1} < \frac{s_1}{s_1/2} = 2,$$

we have $0 \leq (s_1 \text{ mod } \tau) - v = s_1 - \tau - v = \tau_1 - \tau \leq \varepsilon_1 \leq \beta_1$, which implies (ii). We assume that ε_n, τ_n and $f(n)$ are defined for some $n \in \mathbb{N}$. We choose $f(n + 1)$ satisfying $f(n + 1) > f(n)$ and $s_{n+1} \geq 2\tau_n(\tau_n - \varepsilon_n)/\varepsilon_n$. Then we have

$$s_{n+1} > \tau_n \quad \text{and} \quad \frac{s_{n+1}}{\tau_n - \varepsilon_n} \geq \frac{s_{n+1}}{\tau_n} + 2.$$

Hence there exist real numbers p and q such that

$$\tau_n - \varepsilon_n \leq p < q \leq \tau_n \quad \text{and} \quad \frac{s_{n+1}}{p} = \frac{s_{n+1}}{q} + 1 \in \mathbb{N}.$$

We put

$$\tau_{n+1} = \frac{(s_{n+1} - v)q}{s_{n+1}} \quad \text{and} \quad \varepsilon_{n+1} = \frac{\beta_{n+1}q}{(n + 1)s_{n+1}}.$$

Then it is obvious that $\tau_{n+1} \leq q$. Since

$$p - v - \beta_{n+1}/(n+1) \geq p - v - 1 \geq \tau_n - \varepsilon_n - v - 1 > 0,$$

we have

$$\begin{aligned} \tau_{n+1} - \varepsilon_{n+1} &= q \frac{s_{n+1} - v - \beta_{n+1}/(n+1)}{s_{n+1}} = \frac{s_{n+1} p}{s_{n+1} - p} \frac{s_{n+1} - v - \beta_{n+1}/(n+1)}{s_{n+1}} \\ &= p \frac{s_{n+1} - v - \beta_{n+1}/(n+1)}{s_{n+1} - p} > p. \end{aligned}$$

Therefore,

$$\tau_n - \varepsilon_n \leq p < \tau_{n+1} - \varepsilon_{n+1} < \tau_{n+1} \leq q \leq \tau_n.$$

So we note

$$(s_{n+1} \bmod \tau) - v = s_{n+1} - \tau s_{n+1}/q - v$$

for $\tau \in [\tau_{n+1} - \varepsilon_{n+1}, \tau_{n+1}]$. Since

$$(s_{n+1} \bmod \tau_{n+1}) - v = 0 \quad \text{and} \quad (s_{n+1} \bmod (\tau_{n+1} - \varepsilon_{n+1})) - v = \beta_{n+1}/(n+1),$$

we have

$$0 \leq (s_{n+1} \bmod \tau) - v \leq \beta_{n+1}/(n+1)$$

for $\tau \in [\tau_{n+1} - \varepsilon_{n+1}, \tau_{n+1}]$. Therefore we have defined $\{\varepsilon_n\}$, $\{\tau_n\}$ and $\{f(n)\}$ which satisfy (i)–(iii). Cantor's intersection theorem yields that there exists $\tau \in \mathbb{R}$ such that $\tau \in \bigcap_{n=1}^{\infty} [\tau_n - \varepsilon_n, \tau_n]$. By (ii), we have

$$\limsup_{n \rightarrow \infty} \frac{\alpha_n}{(t_n \bmod \tau) - v} \geq \limsup_{n \rightarrow \infty} \frac{\beta_n}{(s_n \bmod \tau) - v} \geq \lim_{n \rightarrow \infty} n = \infty. \quad \blacksquare$$

Lemma 3.2 Let $\{\alpha_n\}$ and $\{t_n\}$ be real sequences with condition (i) of Theorem 2.1. Assume

$$(3.1) \quad \lim_{n \rightarrow \infty} \frac{\alpha_n}{|t_n - \tau|} > 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} t_n = \tau$$

for some $\tau \in (0, \infty)$. Then there exists a subsequence $\{f(n)\}$ of $\{n\}$ such that either

$$(3.2) \quad \lim_{n \rightarrow \infty} \frac{\alpha_{f(n)}}{t_{f(n)} \bmod \tau} > 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} (t_{f(n)} \bmod \tau) = 0$$

or

$$(3.3) \quad \lim_{n \rightarrow \infty} \frac{\alpha_{f(n)}}{(t_{f(n)} \bmod \tau) - \tau} < 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} (t_{f(n)} \bmod \tau) = \tau$$

holds.

Proof If there exists a subsequence $\{f(n)\}$ of $\{n\}$ such that $t_{f(n)} \geq \tau$ for all $n \in \mathbb{N}$, then

$$t_{f(n)} \bmod \tau = t_{f(n)} - \tau = |t_{f(n)} - \tau|$$

for sufficiently large $n \in \mathbb{N}$. Thus (3.2) holds. If there exists a subsequence $\{f(n)\}$ of $\{n\}$ such that $t_{f(n)} < \tau$ for all $n \in \mathbb{N}$, then

$$(t_{f(n)} \bmod \tau) - \tau = t_{f(n)} - \tau = -|t_{f(n)} - \tau|$$

for all $n \in \mathbb{N}$. Thus (3.3) holds. ■

Lemma 3.3 Let $\{\alpha_n\}$ and $\{t_n\}$ be real sequences with condition (i) of Theorem 2.1. Assume

$$\lim_{n \rightarrow \infty} \frac{\alpha_n}{t_n} > 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} t_n = 0.$$

Then (3.2) holds for every positive real number τ and every subsequence $\{f(n)\}$ of $\{n\}$.

Proof Obvious. ■

Lemma 3.4 Let $\{\alpha_n\}$ and $\{t_n\}$ be real sequences with condition (i) of Theorem 2.1. Assume that the conjunction of conditions (ii) and (iii) of Theorem 2.1 does not hold. Then there exist a positive real number τ and a subsequence $\{f(n)\}$ of $\{n\}$ such that either (3.2) or (3.3) holds.

Proof We consider the following four cases:

- $\limsup_n t_n = \infty$
- $\limsup_n t_n < \infty$ and $\limsup_n \alpha_n > 0$
- $\limsup_n t_n < \infty$, $\lim_n \alpha_n = 0$ and $\limsup_n \alpha_n/|t_n - \tau| > 0$ for some $\tau \in (0, \infty)$
- $\limsup_n t_n < \infty$, $\lim_n \alpha_n = 0$ and $\limsup_n \alpha_n/t_n > 0$.

In the first case, using Lemma 3.1, there exist a positive real number τ and a subsequence $\{f(n)\}$ of $\{n\}$ such that

$$\lim_{n \rightarrow \infty} \frac{\alpha_{f(n)}}{t_{f(n)} \bmod \tau} = \infty.$$

It is obvious that $\lim_n (t_{f(n)} \bmod \tau) = 0$. Thus (3.2) holds. Next we note that it is sufficient to show the existence of a subsequence $\{g(n)\}$ of $\{n\}$ such that we can apply either Lemma 3.2 or Lemma 3.3. In the second case, we can choose a subsequence $\{g(n)\}$ of $\{n\}$ such that $\lim_n \alpha_{g(n)} > 0$ and $\{t_{g(n)}\}$ converges to some nonnegative real number τ . Then $\{\alpha_{g(n)}\}$ and $\{t_{g(n)}\}$ satisfy (3.1). So we can apply either Lemma 3.2 or Lemma 3.3. In the third case, we can choose a subsequence $\{g(n)\}$ of $\{n\}$ such that $\lim_n \alpha_{g(n)}/|t_{g(n)} - \tau| > 0$. Then $\lim_n |t_{g(n)} - \tau| = 0$ holds. Hence we can apply Lemma 3.2. Similarly, in the fourth case, we can apply Lemma 3.3. ■

Example 1 Let $\{\alpha_n\}$ and $\{t_n\}$ be real sequences with condition (i) of Theorem 2.1. Let γ be a positive real number. Let E be the two-dimensional real Hilbert space and put $C = \{x \in E : \|x\| \leq 1\}$. For $t \geq 0$, define a 2×2 matrix $T(t)$ by

$$T(t) = \begin{bmatrix} \cos(\gamma t) & -\sin(\gamma t) \\ \sin(\gamma t) & \cos(\gamma t) \end{bmatrix}$$

We can consider that $\{T(t) : t \geq 0\}$ is a linear nonexpansive semigroup on C . Let P be the metric projection from C onto $\bigcap_{t \geq 0} F(T(t))$, that is, $Px = 0$ for all $x \in C$. Put $u = (1, 0)$ and define a sequence $\{u_n\}$ by (1.1). Assume that the conjunction of condition (ii) and condition (iii) of Theorem 2.1 does not hold. Then there exists γ such that $\{u_n\}$ does not converge strongly to Pu .

Proof By Lemma 3.4, there exist a positive real number τ and a subsequence $\{f(n)\}$ of $\{n\}$ such that either (3.2) or (3.3) holds. We note that both (3.2) and (3.3) do not hold simultaneously. We put

$$\gamma = 4\pi/\tau.$$

We also put

$$\eta := \begin{cases} \lim_n (t_{f(n)} \bmod \tau) / \alpha_{f(n)} \in [0, \infty) & \text{if (3.2) holds,} \\ \lim_n ((t_{f(n)} \bmod \tau) - \tau) / \alpha_{f(n)} \in (-\infty, 0] & \text{if (3.3) holds.} \end{cases}$$

In the case where (3.2) holds, since

$$\sin(\gamma t_{f(n)}) = \sin(\gamma t_{f(n)} \bmod 4\pi) = \sin(\gamma (t_{f(n)} \bmod \tau)),$$

we have

$$\lim_{n \rightarrow \infty} \frac{\sin(\gamma t_{f(n)})}{\alpha_{f(n)}} = \lim_{n \rightarrow \infty} \frac{\gamma (t_{f(n)} \bmod \tau)}{\alpha_{f(n)}} = \gamma \eta.$$

In the case where (3.3) holds, since

$$\sin(\gamma t_{f(n)}) = \sin((\gamma t_{f(n)} \bmod 4\pi) - 4\pi) = \sin(\gamma ((t_{f(n)} \bmod \tau) - \tau)),$$

we have

$$\lim_{n \rightarrow \infty} \frac{\sin(\gamma t_{f(n)})}{\alpha_{f(n)}} = \lim_{n \rightarrow \infty} \frac{\gamma ((t_{f(n)} \bmod \tau) - \tau)}{\alpha_{f(n)}} = \gamma \eta.$$

Similarly, $\lim_n \sin(\gamma t_{f(n)}/2) / \alpha_{f(n)} = \gamma \eta/2$ holds in both cases. For $n \in \mathbb{N}$, we put a 2×2 matrix P_n by

$$P_n = \frac{\alpha_n}{4(1 - \alpha_n) \sin^2(\gamma t_n/2) + \alpha_n^2} \begin{bmatrix} a_n & -b_n \\ b_n & a_n \end{bmatrix},$$

where $a_n = \alpha_n + 2(1 - \alpha_n) \sin^2(\gamma t_n/2)$ and $b_n = (1 - \alpha_n) \sin(\gamma t_n)$. It is easy to verify that $u_n = P_n u$ for $n \in \mathbb{N}$ (see [14]). We obtain

$$\lim_{n \rightarrow \infty} P_{f(n)} = \frac{1}{\gamma^2 \eta^2 + 1} \begin{bmatrix} 1 & -\gamma \eta \\ \gamma \eta & 1 \end{bmatrix} = \frac{1}{\sqrt{\gamma^2 \eta^2 + 1}} \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix},$$

where $\theta := \arctan(\gamma \eta) \in (-\pi/2, \pi/2)$. Therefore

$$\lim_{n \rightarrow \infty} u_{f(n)} = \frac{1}{\sqrt{\gamma^2 \eta^2 + 1}} \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix} u \neq 0 = Pu$$

holds. ■

From Corollary 2.2 and Example 1, we obtain the following.

Theorem 3.5 *Let E be a Hilbert space whose dimension is more than 1. Let $\{\alpha_n\}$ and $\{t_n\}$ be real sequences satisfying condition (i) of Theorem 2.1. Then the following are equivalent:*

- Conditions (ii) and (iii) of Theorem 2.1 hold.
- If $\{T(t) : t \geq 0\}$ is a one-parameter nonexpansive semigroup on a bounded closed convex subset C of E , $u \in C$, $\{u_n\}$ is a sequence defined by (1.1) and P is the metric projection from C onto $\bigcap_{t \geq 0} F(T(t))$, then $\{u_n\}$ converges strongly to Pu .

4 Additional Results

In [13], we improved Theorem 1.1 as follows. In this section, we first compare Theorem 2.1 with Theorem 4.1.

Theorem 4.1 ([13]) *Let $\{\alpha_n\}$ and $\{t_n\}$ be real sequences satisfying Conditions (i) and (ii) of Theorem 2.1 and*

- (iii) $s_n := \liminf_m |t_m - t_n| > 0$ for $n \in \mathbb{N}$ and $\lim_n \alpha_n/s_n = 0$.

Then the same conclusion of Theorem 2.1 holds.

Theorem 4.1(iii) is stronger than condition (iii) of Theorem 2.1 because condition (iii) of Theorem 2.1 is a sufficient and necessary condition. It is a natural question of whether Theorem 4.1(iii) is strictly stronger.

Example 2 Define functions f and g from \mathbb{N} into $\mathbb{N} \cup \{0\}$ and real sequences $\{\alpha_n\}$ and $\{t_n\}$ by

- $f(n) = \max \{k \in \mathbb{N} \cup \{0\} : k(k+1)/2 < n\}$
- $g(n) = n - f(n) (f(n) + 1) / 2$
- $t_n = 2^{-g(n)}$ if $n = g(n) (g(n) + 1)/2$, and $t_n = 2^{-g(n)} + 4^{-n}$ otherwise.
- $\alpha_n = 4^{-2n}$.

Then $\{\alpha_n\}$ and $\{t_n\}$ satisfy Conditions (i)–(iii) of Theorem 2.1, however, do not satisfy (iii) of Theorem 4.1.

Remark 1 The sequence $\{t_n\}$ is

$$\frac{1}{2}, \frac{1}{2} + \frac{1}{4^2}, \frac{1}{2^2}, \frac{1}{2} + \frac{1}{4^4}, \frac{1}{2^2} + \frac{1}{4^3}, \frac{1}{2^3}, \frac{1}{2} + \frac{1}{4^7}, \frac{1}{2^2} + \frac{1}{4^8}, \frac{1}{2^3} + \frac{1}{4^9}, \frac{1}{2^4}, \frac{1}{2} + \frac{1}{4^{11}}, \dots$$

Proof We note that if $n = m(m+1)/2$ for some $m \in \mathbb{N}$, then $g(n) = m$. It is obvious that conditions (i) and (ii) of Theorem 2.1 hold. Since $2^{-\nu}$ is a cluster point of $\{t_n\}$ for every $\nu \in \mathbb{N}$, we have

$$s_{m(m+1)/2} := \liminf_{j \rightarrow \infty} |t_j - t_{m(m+1)/2}| = \liminf_{j \rightarrow \infty} |t_j - 2^{-m}| = 0$$

for all $m \in \mathbb{N}$. Hence Theorem 4.1(iii) does not hold. Let us prove condition (iii) of Theorem 2.1. Fix $\tau \in [0, \infty)$. We consider the following three cases:

- $\tau = 0$
- $\tau = 2^{-\nu}$ for some $\nu \in \mathbb{N}$
- otherwise

In the first case, we have

$$\lim_{n \rightarrow \infty} \frac{\alpha_n}{t_n - \tau} = \lim_{n \rightarrow \infty} \frac{4^{-2n}}{t_n} \leq \lim_{n \rightarrow \infty} \frac{4^{-2n}}{2^{-g(n)}} \leq \lim_{n \rightarrow \infty} \frac{4^{-2n}}{2^{-n}} = 0.$$

In the second case, considering the two cases of $g(n) \leq \nu$ and $g(n) > \nu$, we have

$$|t_n - \tau| \geq \min \{4^{-n}, 2^{-\nu-1} - 4^{-n}\}$$

for $n \in \mathbb{N}$ with $n > \nu(\nu + 1)/2$. Hence

$$\lim_{n \rightarrow \infty} \frac{\alpha_n}{|t_n - \tau|} \leq \lim_{n \rightarrow \infty} \frac{4^{-2n}}{\min \{4^{-n}, 2^{-\nu-1} - 4^{-n}\}} = 0.$$

In the third case, we have

$$\lim_{n \rightarrow \infty} \frac{\alpha_n}{|t_n - \tau|} \leq \frac{\lim_n 4^{-2n}}{\liminf_n |t_n - \tau|} = \frac{0}{\liminf_n |t_n - \tau|} = 0.$$

Therefore condition (iii) of Theorem 2.1 holds. ■

Finally we study Condition (iii) of Theorem 2.1 more deeply.

For an arbitrary set A , we denote by $\sharp A$ the cardinal number of A . For a subset A of a topological space, we also denote by A^d the derived set of A . That is, $x \in A^d$ if and only if x belongs to the closure of $A \setminus \{x\}$. We recall that A is *dense in itself* if $A \subset A^d$. We define A^p by

$$A^p = \bigcup \{B \subset A : B \text{ is dense in itself}\}.$$

A^p is called the *perfect kernel* of A . A is called *scattered* if $A^p = \emptyset$. We know that A^p is perfect under the relative topology for A . We also know that $A \setminus A^p$ is scattered, that is, A can be written as the union of two disjoint sets, one perfect, the other scattered. See [8, 16].

Let α be an ordinal number. We denote by α^+ and α^- the successor and the predecessor of α , respectively. We recall that α is *isolated* if α^- exists. α is *limit* if α^- does not exist.

Proposition 4.2 *Let A be a subset of a topological space. Let γ be an ordinal number with $\sharp \gamma > \sharp A$ and $\sharp \gamma \geq \sharp \mathbb{N}$. Put $D = \{\alpha : \alpha \leq \gamma\}$. Define a net $\{A_\alpha\}_{\alpha \in D}$ of subsets of A by*

$$A_\alpha = \begin{cases} A & \text{if } \alpha = 0, \\ A_{\alpha^-} \cap (A_{\alpha^-})^d & \text{if } \alpha \text{ is isolated,} \\ \bigcap \{A_\beta : \beta < \alpha\} & \text{if } \alpha \text{ is limit.} \end{cases}$$

Then $A_\gamma = A^p$ holds.

Proof It is obvious that $\alpha \leq \beta$ implies $A_\beta \subset A_\alpha$. We can easily show by transfinite induction $A^p \subset A_\alpha$ because $A^p \subset B$ implies $A^p \subset B \cap B^d$. Arguing by contradiction, we assume $A^p \not\subset A_\gamma$. Since $A^p \subset B \subset A$ implies $B \cap B^d \subset B$, we have $A_{\alpha^+} \subset A_\alpha$. Thus

$$\begin{aligned} \# \gamma &= \#\{\alpha : \alpha < \gamma\} = \#\{\alpha : \alpha \leq \gamma\} \\ &= \#\{\alpha : \alpha \leq \gamma, \alpha \text{ is isolated}\} \\ &\leq \#\bigsqcup\{A_{\alpha^-} \setminus A_\alpha : \alpha \leq \gamma, \alpha \text{ is isolated}\} \\ &= \#(A \setminus A_\gamma) \leq \#A, \end{aligned}$$

which contradicts $\#A < \# \gamma$. Therefore we obtain $A_\gamma = A^p$. ■

Proposition 4.3 Let $\{t_n\}$ be a real sequence and put $A = \{t_n : n \in \mathbb{N}\}$. Then the following are equivalent:

- (i) There exists a sequence $\{\alpha_n\}$ of positive real numbers satisfying $\lim_n \alpha_n / (t_n - \tau) = 0$ for all $\tau \in \mathbb{R}$.
- (ii) A is scattered, and $\#\{n : t_n = \tau\} < \infty$ for all $\tau \in \mathbb{R}$.

Remark 2 If $\{t_n\}$ satisfies the assumption of Theorem 4.1, then A is obviously scattered.

Proof In order to show (i) implies (ii), we assume that (ii) does not hold and let $\{\alpha_n\}$ be a sequence of positive real numbers. In the case where $\#\{n : t_n = \tau\} = \infty$ for some $\tau \in \mathbb{R}$, it is obvious $\limsup_n \alpha_n / (t_n - \tau) = \infty$. So we consider the other case where $A^p \neq \emptyset$. We first choose $f(1) \in \mathbb{N}$ such that $t_{f(1)} \in A^p$, and put $B_1 = (t_{f(1)} - \alpha_{f(1)}, t_{f(1)} + \alpha_{f(1)})$. Then from $t_{f(1)} \in (A^p)^d$, we have $\#(A^p \cap B_1) = \infty$. So we can choose $f(2) \in \mathbb{N}$ such that $f(2) > f(1)$ and $t_{f(2)} \in A^p \cap B_1$. We put

$$B_2 = B_1 \cap (t_{f(2)} - \alpha_{f(2)}, t_{f(2)} + \alpha_{f(2)}).$$

Then since $t_{f(2)} \in (A^p)^d$, we have $\#(A^p \cap B_2) = \infty$. So we can choose $f(3) \in \mathbb{N}$ such that $f(3) > f(2)$ and $t_{f(3)} \in A^p \cap B_2$. Continuing this argument, we have a subsequence $\{f(n)\}$ of $\{n\}$ and a sequence $\{B_n\}_{n=1}^\infty$ of nonempty open intervals satisfying

- $B_1 \supset B_2 \supset B_3 \supset \dots$;
- $B_n \subset [t_{f(n)} - \alpha_{f(n)}, t_{f(n)} + \alpha_{f(n)}]$ for all $n \in \mathbb{N}$.

So $\{[t_{f(n)} - \alpha_{f(n)}, t_{f(n)} + \alpha_{f(n)}]\}$ has the finite intersection property. Hence there exists $\tau \in \mathbb{R}$ such that $\tau \in \bigcap_{n=1}^\infty [t_{f(n)} - \alpha_{f(n)}, t_{f(n)} + \alpha_{f(n)}]$. Then we have

$$\limsup_{n \rightarrow \infty} \frac{\alpha_n}{|t_n - \tau|} \geq \limsup_{n \rightarrow \infty} \frac{\alpha_{f(n)}}{|t_{f(n)} - \tau|} \geq 1.$$

Therefore (i) does not hold in both cases. We have shown (i) implies (ii). Let us prove (ii) implies (i). We assume (ii). Let γ be an ordinal number with $\# \gamma = \# \mathbb{R}$ and put $D = \{\alpha : \alpha \leq \gamma\}$. Define a net $\{A_\alpha\}_{\alpha \in D}$ of subsets of A as in Proposition 4.2. By

Proposition 4.2, $A_\gamma = \emptyset$ holds. So we can define a function κ from \mathbb{N} into D such that

$$t_n \in A_{\kappa(n)} \quad \text{and} \quad t_n \notin A_{\kappa(n)^+}.$$

Define a function δ from \mathbb{N} into $(0, \infty]$ by

$$\delta(n) = \inf\{|t_n - s| : s \in A_{\kappa(n)} \setminus \{t_n\}\},$$

where $\inf \emptyset = \infty$. We note $\delta(n) > 0$ because $t_n \notin A_{\kappa(n)^+}$. We choose a real sequence $\{\alpha_n\}$ satisfying

$$0 < \alpha_n < \delta(n)/n \quad \text{and} \quad \alpha_{n+1} < \alpha_n.$$

Fix $\tau \in \mathbb{R}$ and $\varepsilon > 0$. Then there exists $\nu \in \mathbb{N}$ such that $2/\nu < \varepsilon$. It is obvious that $n \geq \nu$ implies $2\alpha_n/\varepsilon < \delta(n)$. We shall show

- $m > n \geq \nu$, $\alpha_n/|t_n - \tau| > \varepsilon$, $\alpha_m/|t_m - \tau| > \varepsilon$ and $t_m \neq t_n$ imply $\kappa(m) < \kappa(n)$.

Arguing by contradiction, we assume $\kappa(m) \geq \kappa(n)$. Then since $t_m \in A_{\kappa(n)} \setminus \{t_n\}$, we have

$$|t_n - t_m| \geq \delta(n) > 2\alpha_n/\varepsilon.$$

Since $\alpha_m < \alpha_n$, we have

$$2\alpha_n/\varepsilon < |t_n - t_m| \leq |t_n - \tau| + |t_m - \tau| < \alpha_n/\varepsilon + \alpha_m/\varepsilon < 2\alpha_n/\varepsilon,$$

which is a contradiction. Therefore we have shown $\kappa(m) < \kappa(n)$. Since there does not exist a strictly decreasing infinite sequence of ordinal numbers, we have

$$\#\{n \in \mathbb{N} : \alpha_n/|t_n - \tau| > \varepsilon\} < \infty.$$

Since $\varepsilon > 0$ is arbitrary, we obtain $\lim_n \alpha_n/|t_n - \tau| = 0$. ■

References

- [1] L. P. Belluce and W. A. Kirk, *Nonexpansive mappings and fixed-points in Banach spaces*. Illinois J. Math. **11**(1967), 474–479.
- [2] F. E. Browder, *Nonexpansive nonlinear operators in a Banach space*. Proc. Nat. Acad. Sci. U.S.A. **54**(1965), 1041–1044. doi:10.1073/pnas.54.4.1041
- [3] ———, *Convergence of approximants to fixed points of nonexpansive non-linear mappings in Banach spaces*. Arch. Rational Mech. Anal. **24** (1967), 82–90.
- [4] R. E. Bruck, *A common fixed point theorem for a commuting family of nonexpansive mappings*. Pacific J. Math. **53**(1974), 59–71.
- [5] R. DeMarr, *Common fixed points for commuting contraction mappings*. Pacific J. Math. **13**(1963), 1139–1141.
- [6] K. Goebel and W. A. Kirk, *Topics in Metric Fixed Point Theory*. Cambridge Studies in Advanced Mathematics 28, Cambridge University Press/Cambridge, 1990.
- [7] W. A. Kirk and B. Sims, (eds.) *Handbook of Metric Fixed Point Theory*. Kluwer Academic Publishers, dordrecht, 2001.
- [8] K. Kuratowski, *Topology*. I. Academic Press, New York, 1966.
- [9] T. C. Lim, *A fixed point theorem for families on nonexpansive mappings*. Pacific J. Math. **53**(1974), 487–493.
- [10] T. Suzuki, *On strong convergence to common fixed points of nonexpansive semigroups in Hilbert spaces*. Proc. Amer. Math. Soc. **131**(2003), no. 7, 2133–2136. doi:10.1090/S0002-9939-02-06844-2

- [11] ———, *Common fixed points of one-parameter nonexpansive semigroups*. Bull. London Math. Soc. **38**(2006), no. 6, 1009–1018. doi:10.1112/S0024609306018893
- [12] ———, *Browder's type convergence theorems for one-parameter semigroups of nonexpansive mappings in Banach spaces*. Israel J. Math. **157**(2007), 239–257. doi:10.1007/s11856-006-0010-6
- [13] ———, *Some comments about recent results on one-parameter nonexpansive semigroups*. Bull. Kyushu Inst. Technol. Pure Appl. Math. **54**(2007), 13–26.
- [14] ———, *Browder convergence and Mosco convergence for families of nonexpansive mappings*. Cubo **10**(2008), no. 4, 101–108.
- [15] W. Takahashi, *Nonlinear Functional Analysis. Fixed Point Theory and its Applications*. Yokohama Publishers, Yokohama, 2000.
- [16] S. Willard, *General Topology*. Dover, Mineola, NY, 2004.

Department of Mathematics, Faculty of Science, Niigata University, Niigata 950-2181, Japan
e-mail: akiyama@math.sc.niigata-u.ac.jp

Department of Basic Sciences, Kyushu Institute of Technology, Tobata, Kitakyushu 804-8550, Japan
e-mail: suzuki-t@mns.kyutech.ac.jp