



Only two Betchov homogeneity constraints exist for isotropic turbulence

Maurizio Carbone^{1,†} and Michael Wilczek^{1,2,†}

¹Max Planck Institute for Dynamics and Self-Organization, Am Faßberg 17, 37077 Göttingen, Germany

²Theoretical Physics I, University of Bayreuth, Universitätsstr. 30, 95447 Bayreuth, Germany

(Received 23 January 2022; revised 1 June 2022; accepted 27 July 2022)

Statistically homogeneous flows obey exact kinematic relations. The Betchov homogeneity constraints (Betchov, *J. Fluid Mech.*, vol. 1, 1956, pp. 497–504) for the average principal invariants of the velocity gradient are among the most well-known and extensively employed homogeneity relations. These homogeneity relations have far-reaching implications for the coupled dynamics of strain and vorticity, as well as for the turbulent energy cascade. Whether the Betchov homogeneity constraints are the only possible ones or whether additional homogeneity relations exist has not been proven yet. Here we show that the Betchov homogeneity constraints are the only homogeneity constraints for incompressible and statistically isotropic velocity gradient fields. Our analysis also applies to compressible/perceived velocity gradients, and it allows the derivation of homogeneity relations involving the velocity gradient and other dynamically relevant quantities, such as the pressure Hessian and viscous stresses.

Key words: turbulence theory, isotropic turbulence, homogeneous turbulence

1. Introduction

The velocity gradients, i.e. spatial derivatives of the velocity field, $\mathbf{A} = \nabla \mathbf{u}$, contain a wealth of information about small-scale turbulence, including the topology of vorticity and strain (Meneveau 2011). The moments of the velocity gradients of an incompressible and statistically homogeneous field obey exact kinematic relations (Betchov 1956). The two so-called Betchov constraints for the velocity gradient principal invariants, namely the matrix traces $\text{Tr}(\mathbf{A}^2)$ and $\text{Tr}(\mathbf{A}^3)$, are of central importance for a statistical description of the turbulent dynamics (Davidson 2004). The first Betchov constraint states that the second principal invariant of the velocity gradient is on average zero, $\langle \text{Tr}(\mathbf{A}^2) \rangle = 0$, which implies

† Email addresses for correspondence: maurizio.carbone@ds.mpg.de,
michael.wilczek@uni-bayreuth.de

the proportionality between the mean dissipation rate and the mean squared vorticity:

$$\varepsilon = \nu \langle \boldsymbol{\omega}^2 \rangle. \tag{1.1}$$

Here ν is the kinematic viscosity of the fluid, $\varepsilon = 2\nu \langle \text{Tr}(\mathbf{S}^2) \rangle$ is the mean dissipation rate, \mathbf{S} is the strain rate and $\boldsymbol{\omega} = \nabla \times \mathbf{u}$ is the vorticity. The second Betchov relation $\langle \text{Tr}(\mathbf{A}^3) \rangle = 0$ connects strain self-amplification and vortex stretching:

$$4 \langle \text{Tr}(\mathbf{S}^3) \rangle = -3 \langle \boldsymbol{\omega} \cdot (\mathbf{S}\boldsymbol{\omega}) \rangle. \tag{1.2}$$

While (1.1) constrains the strain-rate and vorticity magnitudes, (1.2) constrains their production rates. The latter relation was derived first by Townsend & Taylor (1951) and then rederived and extensively used by Betchov (1956). It allows one to characterize the average turbulent energy cascade in physical space (Davidson 2004; Carbone & Bragg 2020; Johnson 2020) and to predict the preferential configuration of the strain-rate eigenvalues (Betchov 1956). It also implies that vortex stretching has a positive average in the presence of an average forward energy cascade, related to the negative skewness of the longitudinal velocity increment and gradient statistics. This positive average has implications, for example, on the vorticity magnitude and orientation relative to the strain rate (Tsinober 2009; Tom, Carbone & Bragg 2021) and on the attenuation of extreme velocity gradients (Buaria, Pumir & Bodenschatz 2020).

Additionally, relations (1.1) and (1.2) have their analogues for the velocity structure functions (Hill 1997). The relations for the velocity structure functions are related to those for the velocity gradients through a simple Taylor expansion at small scales, and at larger scales through a filtered/perceived velocity gradient corrected for compressible effects (Carbone & Bragg 2020). The compressible/perceived gradients in homogeneous turbulence also follow analogous relations (Yang, Pumir & Xu 2020; Yang *et al.* 2022), which, for example, constrain the dynamics of velocity gradients as perceived by Lagrangian tetrads (Chertkov, Pumir & Shraiman 1999; Xu, Pumir & Bodenschatz 2011).

Applications of the homogeneity relations (1.1) and (1.2) are not limited to the theoretical understanding of turbulence, but they carry over to the modelling of turbulent flows. For example, stochastic models for the velocity gradient should in principle obey the constraints (1.1) and (1.2) (Johnson & Meneveau 2016), which help to reduce the number of free parameters in such models (Leppin & Wilczek 2020). The homogeneity relations can also be used to improve the performance of neural networks designed for machine learning of turbulent flows by including them into the training (Tian, Livescu & Chertkov 2021; Momenifar *et al.* 2022).

The Betchov relations (1.1) and (1.2) follow by writing the matrix traces $\text{Tr}(\mathbf{A}^2)$ and $\text{Tr}(\mathbf{A}^3)$ as the divergence of a vector field. Then, because of statistical homogeneity, the average of such traces is zero since a spatial derivative can be factored out, and it acts on an average that does not depend on space explicitly. For example, due to incompressibility, the second principal invariant of the velocity gradient can be rewritten as

$$\text{Tr}(\mathbf{A}^2) = \nabla_j u_i \nabla_i u_j = \nabla_i (u_j \nabla_j u_i), \tag{1.3}$$

so that its average vanishes, and relation (1.1) follows. An analogous procedure applies to retrieve (1.2), since the third principal invariant can be expressed as

$$\text{Tr}(\mathbf{A}^3) = \nabla_j u_i \nabla_k u_j \nabla_i u_k = \nabla_i (u_k \nabla_j u_i \nabla_k u_j - \frac{1}{2} u_i \nabla_k u_j \nabla_j u_k). \tag{1.4}$$

However, while it is straightforward to check the validity of the Betchov homogeneity relations for the velocity gradient, it is more complicated to show whether those

homogeneity relations are the only possible ones or if additional constraints exist. If there existed higher-order constraints, we could, for example, improve the current reduced-order models of the velocity gradient dynamics just by imposing these additional homogeneity constraints.

Most of the previous attempts to find higher-order homogeneity relations were based on swapping the spatial derivatives: a scalar contraction of powers of the velocity gradient is manipulated by factoring out the spatial derivative, in order to rewrite the contraction as the divergence of some quantity (if possible). Attempts to obtain relations for the fourth-order moments of the velocity increments/gradients through this derivative-swapping procedure include Hill (1997), Hierro & Dopazo (2003) and Bragg *et al.* (2022). However, it is very difficult to show the completeness of the homogeneity constraints for the velocity gradient through this approach. One would need to consider linear combinations of infinitely many contractions of the velocity gradients and try to recast them into the spatial derivative of some field. In this framework, Siggia (1981) showed that no homogeneity constraints exist on polynomials of fourth-order velocity gradient invariants.

Interestingly, the scenario is analogous to the search for inviscid invariants of the Navier–Stokes equations (Majda & Bertozzi 2001), which are central for the occurrence of cascades (Alexakis & Biferale 2018). While it is straightforward to check the conservation of kinetic energy and helicity in the incompressible three-dimensional Euler equations by a derivative-swapping procedure (Majda & Bertozzi 2001), it is much more involved to show whether those conserved quantities are the only possible ones or if additional ones exist. This completeness question has been answered by Serre (1984) for the incompressible Euler equations and by Enciso, Peralta-Salas & de Lizaaur (2016) for volume-preserving diffeomorphisms.

In this work, we investigate the existence of higher-order homogeneity constraints for the velocity gradient using tensor function representation theory (Zheng 1994; Itskov 2015). The analysis allows one to identify the homogeneity relations as the solutions of a system of partial differential equations, and it shows that no additional homogeneity constraints for the incompressible velocity gradient exist other than those already known from Betchov (1956). We furthermore extend the analysis to the velocity gradients as perceived by Lagrangian tetrads in turbulence (Chertkov *et al.* 1999; Naso & Pumir 2005; Xu *et al.* 2011); in particular, we show the completeness of the homogeneity relations for compressible/perceived gradients derived in Yang *et al.* (2020, 2022).

2. An equation encoding the homogeneity constraints on the velocity gradient

We consider a three-dimensional, incompressible and statistically homogeneous and isotropic velocity field $\mathbf{u}(\mathbf{x}, t)$, together with its spatial gradient $\mathbf{A} = \nabla \mathbf{u}$ ($A_{ij} = \nabla_j u_i$ in Cartesian component notation). Incompressibility implies that $\text{Tr}(\mathbf{A}) = 0$; we will extend the analysis to the compressible case afterwards.

We search for homogeneity relations for scalar single-point statistics of the velocity gradient only. By ‘homogeneity constraint’ we mean the vanishing of an average of a scalar quantity due to homogeneity. Statistical homogeneity means that all statistical quantities are invariant under spatial translations. In particular, single-point statistics do not explicitly depend on the spatial coordinate. As a consequence, any scalar field ϕ that is the divergence of a vector field, $\phi = \nabla \cdot \mathbf{F}$, has zero ensemble/spatial average:

$$\langle \phi \rangle = \langle \nabla \cdot \mathbf{F} \rangle = \nabla \cdot \langle \mathbf{F} \rangle = 0. \quad (2.1)$$

For example, in the first Batchov relation $\langle \text{Tr}(\mathbf{A}^2) \rangle = 0$, the vector field is $\mathbf{F} = \mathbf{A}\mathbf{u}$ ($F_i = A_{ij}u_j$ in component notation, see (1.3)). To generalize this, we search for scalar functions of the velocity gradient $\phi(\mathbf{A})$ that are the divergence of a vector field \mathbf{F} .

The vector field \mathbf{F} is in general a *functional* of the velocity field $\mathbf{u}(\mathbf{x}, t)$. We restrict the analysis to *functions* of the velocity and its spatial derivatives because we search for homogeneity relations on the single-point statistics of the velocity gradient. By restricting the analysis to functions of the velocity and its spatial derivatives, we are implicitly assuming isotropy. Indeed, in a statistically isotropic flow, the governing equations and associated boundary conditions do not introduce any characteristic direction. Therefore, in that statistically isotropic situation, the velocity and velocity gradients are all the possible variables upon which the vector \mathbf{F} can depend. We thereby exclude, for example, rotations of the frame of the flow, anisotropic forcing, boundary layers, etc.

Focusing on single-point statistics of isotropic flows, we have $\mathbf{F} = \mathbf{F}(\mathbf{u}, \mathbf{A}, \nabla\mathbf{A}, \nabla(\nabla\mathbf{A}), \dots)$ so that the corresponding $\phi(\mathbf{A})$ is, by the chain rule and in component notation,

$$\phi(\mathbf{A}) = \nabla_i F_i(\mathbf{u}(\mathbf{x}, t), \dots) = \frac{\partial F_i}{\partial u_p} A_{pi} + \frac{\partial F_i}{\partial A_{pq}} \nabla_i A_{pq} + \frac{\partial F_i}{\partial (\nabla_k A_{pq})} \nabla_i (\nabla_k A_{pq}) + \dots \tag{2.2}$$

The fact that the left-hand side of (2.2) depends only on the velocity gradient strongly constrains the functional form of the vector field \mathbf{F} . Namely, the right-hand side of (2.2) should explicitly involve neither the velocity \mathbf{u} nor the gradients of the velocity gradient, $\nabla\mathbf{A}$, $\nabla(\nabla\mathbf{A})$, etc. This implies that all the terms on the right-hand side of (2.2) featuring gradients of the velocity gradient should identically cancel, while only $(\partial F_i / \partial u_p) A_{pi}$ can contribute to ϕ . Moreover, the part of $(\partial F_i / \partial u_p) A_{pi}$ that contributes to ϕ can depend only on \mathbf{A} .

Therefore, we just need to consider vector functions of the velocity and velocity gradient, $\mathbf{F}(\mathbf{u}, \mathbf{A})$, that are linear in the velocity. Based on this, (2.2) splits into

$$\phi(\mathbf{A}) = \frac{\partial F_i}{\partial u_p}(\mathbf{u}, \mathbf{A}) A_{pi}, \tag{2.3a}$$

$$\frac{\partial F_i}{\partial A_{pq}}(\mathbf{u}, \mathbf{A}) \nabla_i A_{pq} = 0. \tag{2.3b}$$

Equation (2.3b) yields the main differential equation to determine \mathbf{F} . The gradient of the gradient, $\nabla_i A_{pq} = \nabla_i \nabla_q u_p$, is symmetric in i and q , so that only the part of $\partial F_i / \partial A_{pq}$ that is symmetric in i and q contributes to (2.3b). Additionally, the contractions i, p and q, p of $\nabla_i A_{pq}$ are zero by incompressibility. Therefore, \mathbf{F} solves (2.3b) only if, for some vector \mathbf{v} ,

$$\frac{\partial F_i}{\partial A_{pq}} + \frac{\partial F_q}{\partial A_{pi}} = v_i \delta_{pq} + v_q \delta_{pi}. \tag{2.4}$$

Here, δ_{ij} denotes the Kronecker delta, and the vector \mathbf{v} is easily determined by contracting two of the free indices, e.g. $v_p = \partial F_k / \partial A_{pk}$.

Equations (2.3) and (2.4) allow the search for homogeneity constraints to be more systematic: instead of attempting to factor out the spatial derivatives in tensor contractions of velocity gradients, we need to solve a system of partial differential equations. Solving (2.4) for vectors \mathbf{F} that are linear in \mathbf{u} yields all possible vectors $\mathbf{F}(\mathbf{u}, \mathbf{A})$, whose divergence depends only on the velocity gradient, $\phi(\mathbf{A}) = \nabla \cdot \mathbf{F}(\mathbf{u}, \mathbf{A})$, as in (2.3a). Therefore, finding all solutions of (2.4) that are linear in the velocity amounts to deriving all possible homogeneity constraints on scalar functions of an incompressible and statistically isotropic velocity gradient.

3. Tensor function representation of the homogeneity constraints

In the following, we construct the general isotropic tensor function $F(\mathbf{u}, \mathbf{A})$. Tensor function representation theory (Weyl 1946; Rivlin & Ericksen 1955; Pennisi & Trovato 1987; Zheng 1994; Itskov 2015) allows one to write down all possible vector functions F of the generating vector \mathbf{u} and tensor \mathbf{A} that transform consistently under any change of basis: when the arguments \mathbf{u} and \mathbf{A} undergo a rotation, F rotates accordingly (Itskov 2015). The vector field F will be finally determined by requiring that it is linear in \mathbf{u} and that it solves (2.4).

In general, F can depend separately on the symmetric and antisymmetric parts of the velocity gradient (Rivlin & Ericksen 1955):

$$\mathbf{S} = \frac{1}{2}(\mathbf{A} + \mathbf{A}^T), \quad \mathbf{W} = \frac{1}{2}(\mathbf{A} - \mathbf{A}^T), \quad (3.1a,b)$$

with \mathbf{A}^T denoting the matrix transpose of \mathbf{A} . Therefore, we consider all the vector functions $F(\mathbf{u}, \mathbf{S}, \mathbf{W})$ constructed through the velocity and velocity gradients that, due to (2.3a), are linear in the velocity:

$$\mathbf{F} = \sum_{n=0}^8 f_n(\mathcal{I}) \mathbf{B}^n \mathbf{u}. \quad (3.2)$$

Here \mathbf{B}^n are the basis tensors that can be formed through \mathbf{S} and \mathbf{W} (Pennisi & Trovato 1987):

$$\left. \begin{aligned} \mathbf{B}^0 &= \mathbf{I}, & \mathbf{B}^3 &= \mathbf{SS}, & \mathbf{B}^6 &= \mathbf{WW}, \\ \mathbf{B}^1 &= \mathbf{S}, & \mathbf{B}^4 &= \mathbf{SW} - \mathbf{WS}, & \mathbf{B}^7 &= \mathbf{SWW} + \mathbf{WWS}, \\ \mathbf{B}^2 &= \mathbf{W}, & \mathbf{B}^5 &= \mathbf{SW} + \mathbf{WS}, & \mathbf{B}^8 &= \mathbf{SSW} + \mathbf{WSS}, \end{aligned} \right\} \quad (3.3)$$

with \mathbf{I} denoting the identity matrix and the standard matrix product is implied. Two additional tensors would be necessary to fix degeneracies of the basis (3.3), which occur when the vorticity is an eigenvector of the strain-rate tensor or the strain rate has two identical eigenvalues (Rivlin & Ericksen 1955). We ignore that zero-measure configuration of the gradients. Also, note that the superscript of \mathbf{B}^n serves to number the basis tensors rather than indicating powers of the tensor.

The components f_n in (3.2) are functions of the set \mathcal{I} of independent invariants that can be formed through the velocity gradients (Pennisi & Trovato 1987):

$$\left. \begin{aligned} \mathcal{I}_1 &= \text{Tr}(\mathbf{SS}), & \mathcal{I}_3 &= \text{Tr}(\mathbf{SSS}), & \mathcal{I}_5 &= \text{Tr}(\mathbf{SSWW}), \\ \mathcal{I}_2 &= \text{Tr}(\mathbf{WW}), & \mathcal{I}_4 &= \text{Tr}(\mathbf{SWW}), \end{aligned} \right\} \quad (3.4)$$

with standard matrix product implied. A sixth invariant would be necessary to fix the orientation/handedness of the vorticity with respect to the strain-rate eigenvectors. We do not consider the sixth invariant as an independent variable, since it is determined by the invariants (3.4) up to a sign (Lund & Novikov 1992).

The number of basis tensors and independent invariants is finite due to the Cayley–Hamilton theorem, which plays a major role in the theory of tensor representation (Rivlin & Ericksen 1955). Betchov (1956) showed that the average of the velocity gradient principal invariants is zero in a homogeneous flow, thus constraining all the independent traces of powers of \mathbf{A} . However, one can take contractions of combinations of the strain rate and rotation rate, and pose the question whether the average of those scalar contractions vanishes due to homogeneity.

4. Solution of the equation encoding the homogeneity constraints

We use the general expression (3.2) combined with (2.4) in order to determine the components $f_n(\mathcal{I})$ of \mathbf{F} . This will yield a vector field \mathbf{F} associated with the homogeneity constraints for the velocity gradient through $\phi(\mathbf{A}) = \nabla \cdot \mathbf{F}(\mathbf{u}, \mathbf{A})$ and (2.1).

Inserting the general expression (3.2) of \mathbf{F} into (2.4) gives, in component notation,

$$\left[\frac{\partial f_l}{\partial \mathcal{I}_k} \frac{\partial \mathcal{I}_k}{\partial A_{pq}} B_{ij}^l + \frac{\partial f_l}{\partial \mathcal{I}_k} \frac{\partial \mathcal{I}_k}{\partial A_{pi}} B_{qj}^l + f_l \frac{\partial B_{ij}^l}{\partial A_{pq}} + f_l \frac{\partial B_{qj}^l}{\partial A_{pi}} \right] u_j = v_i \delta_{pq} + v_q \delta_{pi}. \quad (4.1)$$

Here and throughout, repeated indices imply summation, unless otherwise specified. As shown in Appendix A, the derivatives of the invariants (3.4) can be written as

$$\frac{\partial \mathcal{I}_k}{\partial A_{pq}} = M_{km} B_{pq}^m, \quad (4.2)$$

while the derivatives of the basis tensors (3.3) can be expressed as

$$\frac{\partial B_{ij}^n}{\partial A_{pq}} = \Gamma_{lm}^{1,n} B_{ip}^l B_{qj}^m + \Gamma_{lm}^{2,n} B_{iq}^l B_{pj}^m + \Gamma_{lm}^{3,n} B_{ij}^l B_{pq}^m, \quad (4.3)$$

with $0 \leq l, m, n \leq 8$ and $1 \leq k \leq 5$.

The matrix entries M_{km} featured in (4.2) are specified in (A1). The symbols $\Gamma_{lm}^{P,n}$ in (4.3) play the role of Christoffel symbols (Grinfeld 2013) and their components are listed in (A9), (A10) and (A11). Inserting the expressions (4.2) and (4.3) for the derivatives into (4.1) yields the following independent equations:

$$f_n (\Gamma_{lm}^{2,n} B_{iq}^l B_{pj}^m + \Gamma_{lm}^{2,n} B_{qi}^l B_{pj}^m) u_j = 0, \quad (4.4a)$$

$$\left[\frac{\partial f_l}{\partial \mathcal{I}_k} M_{km} B_{ij}^l B_{pq}^m + f_n (\Gamma_{lm}^{3,n} B_{ij}^l B_{pq}^m + \Gamma_{lm}^{1,n} B_{qp}^l B_{ij}^m) \right] u_j = v_i B_{pq}^0. \quad (4.4b)$$

Equations (4.4) should hold for all \mathbf{u} and \mathbf{A} , so that, separating out the basis tensors, we have the following equations for the components:

$$\sum_{n=0}^8 [\Gamma_{lm}^{2,n} + t(l) \Gamma_{lm}^{2,n}] f_n = 0, \quad \forall 0 \leq l, m \leq 8, \quad (4.5a)$$

$$\sum_{k=1}^5 \frac{\partial f_l}{\partial \mathcal{I}_k} M_{km} = - \sum_{n=0}^8 [\Gamma_{lm}^{3,n} + t(m) \Gamma_{ml}^{1,n}] f_n, \quad \forall 0 \leq l \leq 8, 1 \leq m \leq 8, \quad (4.5b)$$

where indices l, m are not contracted, and $t(l) = 1$ if \mathbf{B}^l is symmetric and $t(l) = -1$ if \mathbf{B}^l is antisymmetric. In the steps from (4.4b) to (4.5b), the components at $m = 0$ have been absorbed into the generic right-hand side $v_i B_{pq}^0$ of (4.4b), and therefore (4.5b) only concerns components with $m \geq 1$.

The linear system (4.5a) of 81 equations in the nine variables f_n , $0 \leq n \leq 8$, can be solved using symbolic calculus (Meurer *et al.* 2017). This yields

$$f_1 = f_2, \quad f_3 = f_5 = f_6, \quad f_4 = f_7 = f_8 = 0. \quad (4.6a-c)$$

Next, (4.5b) has a solution only if, for all l , the right-hand side is orthogonal to the kernel of \mathbf{M} , but this condition imposes no further constraints on f_n . Finally, with this orthogonality

Only two Betchov constraints for isotropic turbulence

condition ensured, the derivatives $\partial f_i/\partial \mathcal{I}_k$ are obtained by multiplying (4.5b) by the Moore–Penrose inverse of \mathbf{M} , with components $M_{mk'}^{-1}$, thus yielding

$$\frac{\partial f_0}{\partial \mathcal{I}_1} = \frac{\partial f_0}{\partial \mathcal{I}_2} = -\frac{1}{2}f_3, \tag{4.7a}$$

$$\frac{\partial f_0}{\partial \mathcal{I}_k} = 0, \quad \forall 3 \leq k \leq 5, \tag{4.7b}$$

$$\frac{\partial f_n}{\partial \mathcal{I}_k} = 0, \quad \forall 1 \leq k \leq 5, 1 \leq n \leq 8. \tag{4.7c}$$

By solving the straightforward linear system (4.7) with the conditions (4.6a–c), we obtain the components f_n of \mathbf{F} that solves (2.4) and is linear in \mathbf{u} :

$$\mathbf{F} = \bar{f}_1 \mathbf{u} + \bar{f}_2 \mathbf{A}\mathbf{u} + \bar{f}_3 (\mathbf{A}^2 - \frac{1}{2} \text{Tr}(\mathbf{A}^2)\mathbf{I})\mathbf{u}, \tag{4.8}$$

where the \bar{f}_n are arbitrary constants. The solution (4.8) of (2.4) encodes all the Betchov constraints, since its divergence yields the gradient principal invariants

$$\nabla \cdot \mathbf{F} = \bar{f}_2 \text{Tr}(\mathbf{A}^2) + \bar{f}_3 \text{Tr}(\mathbf{A}^3), \tag{4.9}$$

thus retrieving (1.3) and (1.4) and, by homogeneity, $(\bar{f}_2 \text{Tr}(\mathbf{A}^2) + \bar{f}_3 \text{Tr}(\mathbf{A}^3)) = 0$.

The Betchov homogeneity relations, obtained by averaging (4.9), are all the possible homogeneity constraints on the single-point statistics of an incompressible and isotropic gradient since they follow from all the independent solutions of (2.4). In other words, no scalar function of the velocity gradient invariants can be written as the divergence of a vector field, other than the principal invariants $\text{Tr}(\mathbf{A}^2) = \mathcal{I}_1 + \mathcal{I}_2$ and $\text{Tr}(\mathbf{A}^3) = \mathcal{I}_3 + 3\mathcal{I}_4$.

5. Homogeneity constraints for the velocity gradient and additional quantities

Equation (4.8) shows that the homogeneity relations for the velocity gradient alone consist only of the two Betchov constraints. However, generalizations of (4.8) easily generate homogeneity constraints concerning the velocity gradient together with additional variables. Indeed, the divergence of \mathbf{F} in (4.8) does not depend on the gradient of the velocity gradient even when \mathbf{u} in (4.8) is replaced by any scalar, vector or tensor quantity \mathbf{q} that does not explicitly depend on the velocity gradient itself. This is because \mathbf{F} solves (2.4), which features only derivatives with respect to \mathbf{A} , while the dependence of \mathbf{F} upon \mathbf{u} is parametric. Therefore, for any vector \mathbf{q} , one can construct homogeneity relations for the scalar quantities

$$\psi(\mathbf{A}, \nabla \mathbf{q}) = \nabla \cdot [\bar{f}_1 \mathbf{q} + \bar{f}_2 \mathbf{A}\mathbf{q} + \bar{f}_3 (\mathbf{A}^2 - \frac{1}{2} \text{Tr}(\mathbf{A}^2)\mathbf{I})\mathbf{q}], \tag{5.1}$$

where the standard matrix–vector product is implied and the left-hand side depends neither on \mathbf{q} nor on $\nabla \mathbf{A}$. For example, using (5.1) with the pressure gradient divided by the fluid density, $\mathbf{q} = \nabla P/\rho$, yields homogeneity relations for the pressure Hessian in incompressible flows, namely $\langle A_{ij} \nabla_i \nabla_j P \rangle = 0$ and $\langle A_{ik} A_{kj} \nabla_i \nabla_j P \rangle = -\rho \langle (A_{ij} A_{ji})^2 \rangle / 2$. Analogously, employing (5.1) with the velocity Laplacian, $\mathbf{q} = \nabla^2 \mathbf{u}$, gives homogeneity relations for the Laplacian of a traceless gradient, $\langle A_{ij} \nabla^2 A_{ji} \rangle = 0$ and $\langle A_{ik} A_{kj} \nabla^2 A_{ji} \rangle = 0$. Such relations play an important role in the budget equation for the turbulent dissipation rate (Buaria, Pumir & Bodenschatz 2022), and turn out to be useful in the Lagrangian modelling of velocity gradients (Leppin & Wilczek 2020).

6. Homogeneity constraints for compressible and perceived velocity gradients

The analysis presented in the previous sections easily generalizes, e.g. to compressible or two-dimensional flows. In this section we consider a compressible velocity gradient $\tilde{\mathbf{A}}$, with non-zero trace $\tilde{\mathcal{L}}_0 = \text{Tr}(\tilde{\mathbf{A}})$ (the tilde denoting quantities relative to the compressible velocity field). The presented homogeneity analysis allows one to prove the completeness of the homogeneity constraints recently derived in Yang *et al.* (2020) for a compressible gradient, which hold approximately even in strongly non-homogeneous compressible turbulence (Yang *et al.* 2022). Additionally, they apply one-to-one to the perceived velocity gradient (or tetrad) dynamics in homogeneous flows (Chertkov *et al.* 1999; Xu *et al.* 2011).

As done above for the incompressible gradient, we search for a scalar function of the compressible gradient only, $\tilde{\phi}(\tilde{\mathbf{A}})$, that has zero average due to statistical homogeneity. In the compressible case, the components \tilde{f}_n of the vector field $\tilde{\mathbf{F}}$ associated with the homogeneity constraints through $\tilde{\phi}(\tilde{\mathbf{A}}) = \nabla \cdot \tilde{\mathbf{F}}$ depend on the additional invariant $\tilde{\mathcal{L}}_0$. The procedure to derive the vector field $\tilde{\mathbf{F}}$, detailed in § 4 for the incompressible gradient, applies to the compressible gradient with the modifications listed in Appendix B. With those modifications, the same procedure as outlined in § 4 yields for the compressible/perceived gradient

$$\tilde{\mathbf{F}} = \bar{f}_1 \tilde{\mathbf{u}} + \bar{f}_2 (\tilde{\mathbf{A}} - \text{Tr}(\tilde{\mathbf{A}}) \mathbf{I}) \tilde{\mathbf{u}} + \bar{f}_3 [\tilde{\mathbf{A}}^2 - \text{Tr}(\tilde{\mathbf{A}}) \tilde{\mathbf{A}} - \frac{1}{2} (\text{Tr}(\tilde{\mathbf{A}}^2) - \text{Tr}(\tilde{\mathbf{A}})^2) \mathbf{I}] \tilde{\mathbf{u}}. \quad (6.1)$$

The average of the divergence of the vector field (6.1),

$$\tilde{\phi}(\tilde{\mathbf{A}}) = \bar{f}_1 \text{Tr}(\tilde{\mathbf{A}}) + \bar{f}_2 (\text{Tr}(\tilde{\mathbf{A}}^2) - \text{Tr}(\tilde{\mathbf{A}})^2) + \bar{f}_3 (\text{Tr}(\tilde{\mathbf{A}}^3) - \frac{3}{2} \text{Tr}(\tilde{\mathbf{A}}) \text{Tr}(\tilde{\mathbf{A}}^2) + \frac{1}{2} \text{Tr}(\tilde{\mathbf{A}})^3), \quad (6.2)$$

vanishes by homogeneity for all the arbitrary constants \bar{f}_i , thus recovering all the homogeneity relations derived in Yang *et al.* (2020, 2022) and proving their completeness.

7. Conclusions

We have shown that the Betchov homogeneity relations are all the possible homogeneity constraints for the velocity gradient in incompressible and statistically isotropic turbulence. Our conclusions apply to the single-point statistics of scalar functions of the velocity gradient. Our approach also allows the derivation of homogeneity relations involving additional quantities, like the pressure Hessian and velocity gradient Laplacian. The presented methodology furthermore generalizes to derive homogeneity constraints for the compressible/perceived gradient, in two- and three-dimensional isotropic flows, as well as in less idealized (e.g. axisymmetric) flows. More generally, the outcome of these calculations will help to deal with high-dimensional tensor equations, which are ubiquitous in fluid dynamics.

Acknowledgements. We thank L. Bentkamp, A.D. Bragg, T.D. Drivas, M. Iovieno, P. Johnson, A. Pumir and H. Xu for insightful comments on the manuscript.

Declaration of interests. The authors report no conflict of interest.

Author ORCID.

✉ Maurizio Carbone <https://orcid.org/0000-0003-0409-6946>;

✉ Michael Wilczek <https://orcid.org/0000-0002-1423-8285>.

Appendix A. Tensor derivatives of the basis tensors and independent invariants

In this appendix, we compute the derivatives with respect to the incompressible gradient \mathbf{A} of the basis tensors \mathbf{B}^n (3.3) and invariants \mathcal{I}_k (3.4).

We start with the derivatives of the invariants (3.4), which can be expressed as linear combinations of the basis tensors, as in (4.2). The matrix \mathbf{M} featuring the components of the derivatives of the invariants (3.4) in the employed basis (3.3) is computed by contracting (4.2) with the basis tensors,

$$M_{km} = Z_{ml}^{-1} B_{pq}^l \frac{\partial \mathcal{I}_k}{\partial A_{pq}} = \begin{bmatrix} 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -2 & 0 & 0 & 0 & 0 & 0 & 0 \\ -\mathcal{I}_1 & 0 & 0 & 3 & 0 & 0 & 0 & 0 & 0 \\ -\frac{1}{3}\mathcal{I}_2 & 0 & 0 & 0 & 0 & -1 & 1 & 0 & 0 \\ -\frac{2}{3}\mathcal{I}_4 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & -1 \end{bmatrix}, \tag{A1}$$

where $Z^{lm} = B_{pq}^l B_{pq}^m$ is the metric tensor and \mathbf{Z}^{-1} denotes its matrix inverse.

Next, we compute the derivatives of the basis tensors, for which we first introduce some notation. The basis tensors (3.3) are products of the symmetric and antisymmetric parts of the velocity gradient (3.1a,b), which can in turn be expressed through the fourth-order tensors

$$Q_{ijpq}^{(t)} = \frac{1}{2} \left(\delta_{ip}\delta_{jq} + t\delta_{iq}\delta_{jp} - \frac{1+t}{3}\delta_{ij}\delta_{pq} \right) \tag{A2}$$

contracted with the gradient itself, $S_{ij} = Q_{ijpq}^{(+1)} A_{pq}$ and $W_{ij} = Q_{ijpq}^{(-1)} A_{pq}$. Then, any basis tensor (3.3) of degree d consists of a linear combination of the products

$$b_{ij}^{t_1 t_2 \dots t_d} = Q_{ik_1 p_1 q_1}^{(t_1)} Q_{k_1 k_2 p_2 q_2}^{(t_2)} \dots Q_{k_{d-1} j p_d q_d}^{(t_d)} (A_{p_1 q_1} A_{p_2 q_2} \dots A_{p_d q_d}), \tag{A3}$$

with $t_l = \pm 1$ and summation over repeated indices, e.g. $\mathbf{b}^{+1,-1} = \mathbf{SW}$. Also, the components of the basis tensors (3.3) with respect to the elementary products (A3) are constant:

$$B_{ij}^n = \sum_{t_1 t_2 \dots t_d} c_{t_1 t_2 \dots t_d}^n b_{ij}^{t_1 t_2 \dots t_d}, \tag{A4}$$

e.g. the non-zero components of $\mathbf{B}^4 = \mathbf{SW} - \mathbf{WS}$ are $c_{+1,-1}^4 = -c_{-1,+1}^4 = 1$.

Using (A3) and (A4) we can take the derivative of any basis tensor of degree d :

$$\begin{aligned} \frac{\partial B_{ij}^n}{\partial A_{pq}} &= \sum_{t_1 t_2 \dots t_d} c_{t_1 t_2 \dots t_d}^n Q_{ik_1 p_1 q_1}^{(t_1)} Q_{k_1 k_2 p_2 q_2}^{(t_2)} \dots Q_{k_{d-1} j p_d q_d}^{(t_d)} \frac{\partial}{\partial A_{pq}} (A_{p_1 q_1} A_{p_2 q_2} \dots A_{p_d q_d}) \\ &= \sum_{t_1 t_2 \dots t_d} c_{t_1 t_2 \dots t_d}^n \sum_{m=1}^d [b_{ik_{m-1}}^{t_1 t_2 \dots t_{m-1}} Q_{k_{m-1} k_m p q}^{(t_m)} b_{k_m j}^{t_{m+1} t_{m+2} \dots t_d}], \end{aligned} \tag{A5}$$

with $t_l = \pm 1$, $k_0 = i$, $k_d = j$ and contraction over repeated indices. As expected, the derivative of any basis tensor consists of a linear combination of lower-order tensors. The derivative $\partial(\cdot)/\partial A_{pq}$ is understood as a directional derivative in $\mathbb{R}^{3,3}$ (Itskov 2015) and it is traceless due to incompressibility. In the steps to (A5), this property is taken into account by $\mathbf{Q}^{(t)}$, since $Q_{ijpq}^{(t)} \delta_{pq} = 0$ (in fact, $\mathbf{Q}^{(t)}$ is the tensor derivative of \mathbf{b}^t with respect to \mathbf{A}).

Inserting the expression of the fourth-order tensors (A2) into (A5), we can write the derivatives of the basis tensors more explicitly. For the basis tensors (3.3) of degree one (i.e. for $1 \leq n \leq 2$), we have

$$\frac{\partial B_{ij}^n}{\partial A_{pq}} = \frac{1}{2} \sum_{t_1} c_{t_1}^n \left(\delta_{ip} \delta_{qj} + t_1 \delta_{iq} \delta_{pj} - \frac{1+t_1}{3} \delta_{ij} \delta_{pq} \right). \tag{A6}$$

The derivatives of the basis tensors (3.3) of degree two (i.e. for $3 \leq n \leq 6$) read

$$\begin{aligned} \frac{\partial B_{ij}^n}{\partial A_{pq}} = \frac{1}{2} \sum_{t_1 t_2} c_{t_1 t_2}^n \left(\delta_{ip} b_{qj}^{t_2} + t_1 \delta_{iq} b_{pj}^{t_2} - \frac{1+t_1}{3} b_{ij}^{t_2} \delta_{pq} \right. \\ \left. + b_{ip}^{t_1} \delta_{qj} + t_2 b_{iq}^{t_1} \delta_{pj} - \frac{1+t_2}{3} b_{ij}^{t_1} \delta_{pq} \right). \end{aligned} \tag{A7}$$

Finally, the derivatives of the basis tensors (3.3) of degree three (i.e. for $7 \leq n \leq 8$) read

$$\begin{aligned} \frac{\partial B_{ij}^n}{\partial A_{pq}} = \frac{1}{2} \sum_{t_1 t_2 t_3} c_{t_1 t_2 t_3}^n \left(\delta_{ip} b_{qj}^{t_2 t_3} + t_1 \delta_{iq} b_{pj}^{t_2 t_3} - \frac{1+t_1}{3} b_{ij}^{t_2 t_3} \delta_{pq} + b_{ip}^{t_1} b_{qj}^{t_3} \right. \\ \left. + t_2 b_{iq}^{t_1} b_{pj}^{t_3} - \frac{1+t_2}{3} b_{ij}^{t_1 t_3} \delta_{pq} + b_{ip}^{t_1 t_2} \delta_{qj} + t_3 b_{iq}^{t_1 t_2} \delta_{pj} - \frac{1+t_3}{3} b_{ij}^{t_1 t_2} \delta_{pq} \right). \end{aligned} \tag{A8}$$

After replacing the symmetric and antisymmetric parts of \mathbf{b}^{t_1} and $\mathbf{b}^{t_1 t_2}$ with the corresponding \mathbf{B}^n , the tensor derivatives (A6), (A7) and (A8) can be compactly rewritten as contractions of the basis tensors with the Christoffel symbols $\Gamma_{lm}^{p,n}$, as in (4.3) in the main text. For the tensor basis (3.3) employed here, the non-zero elements of $\Gamma_{lm}^{1,n}$ are

$$\left. \begin{aligned} \Gamma_{00}^{1,1} = \frac{1}{2}, \quad \Gamma_{02}^{1,4} = \frac{1}{2}, \quad \Gamma_{10}^{1,5} = \frac{1}{2}, \quad \Gamma_{05}^{1,7} = \frac{1}{4}, \quad \Gamma_{50}^{1,7} = \frac{1}{4}, \quad \Gamma_{12}^{1,8} = \frac{1}{2}, \\ \Gamma_{00}^{1,2} = \frac{1}{2}, \quad \Gamma_{10}^{1,4} = \frac{1}{2}, \quad \Gamma_{20}^{1,5} = \frac{1}{2}, \quad \Gamma_{06}^{1,7} = \frac{1}{2}, \quad \Gamma_{60}^{1,7} = \frac{1}{2}, \quad \Gamma_{21}^{1,8} = \frac{1}{2}, \\ \Gamma_{01}^{1,3} = \frac{1}{2}, \quad \Gamma_{20}^{1,4} = -\frac{1}{2}, \quad \Gamma_{02}^{1,6} = \frac{1}{2}, \quad \Gamma_{12}^{1,7} = \frac{1}{2}, \quad \Gamma_{03}^{1,8} = \frac{1}{2}, \quad \Gamma_{30}^{1,8} = \frac{1}{2}, \\ \Gamma_{10}^{1,3} = \frac{1}{2}, \quad \Gamma_{01}^{1,5} = \frac{1}{2}, \quad \Gamma_{20}^{1,6} = \frac{1}{2}, \quad \Gamma_{21}^{1,7} = \frac{1}{2}, \quad \Gamma_{04}^{1,8} = \frac{1}{4}, \quad \Gamma_{40}^{1,8} = -\frac{1}{4}, \\ \Gamma_{01}^{1,4} = -\frac{1}{2}, \quad \Gamma_{02}^{1,5} = \frac{1}{2}, \quad \Gamma_{04}^{1,7} = -\frac{1}{4}, \quad \Gamma_{40}^{1,7} = \frac{1}{4}, \quad \Gamma_{05}^{1,8} = \frac{1}{4}, \quad \Gamma_{50}^{1,8} = \frac{1}{4}, \end{aligned} \right\} \tag{A9}$$

the non-zero elements of $\Gamma_{lm}^{2,n}$ are

$$\left. \begin{aligned} \Gamma_{00}^{2,1} = \frac{1}{2}, \quad \Gamma_{02}^{2,4} = \frac{1}{2}, \quad \Gamma_{10}^{2,5} = -\frac{1}{2}, \quad \Gamma_{05}^{2,7} = -\frac{1}{4}, \quad \Gamma_{50}^{2,7} = -\frac{1}{4}, \quad \Gamma_{12}^{2,8} = \frac{1}{2}, \\ \Gamma_{00}^{2,2} = -\frac{1}{2}, \quad \Gamma_{10}^{2,4} = -\frac{1}{2}, \quad \Gamma_{20}^{2,5} = \frac{1}{2}, \quad \Gamma_{06}^{2,7} = \frac{1}{2}, \quad \Gamma_{60}^{2,7} = \frac{1}{2}, \quad \Gamma_{21}^{2,8} = \frac{1}{2}, \\ \Gamma_{01}^{2,3} = \frac{1}{2}, \quad \Gamma_{20}^{2,4} = -\frac{1}{2}, \quad \Gamma_{02}^{2,6} = -\frac{1}{2}, \quad \Gamma_{12}^{2,7} = -\frac{1}{2}, \quad \Gamma_{03}^{2,8} = -\frac{1}{2}, \quad \Gamma_{30}^{2,8} = -\frac{1}{2}, \\ \Gamma_{10}^{2,3} = \frac{1}{2}, \quad \Gamma_{01}^{2,5} = -\frac{1}{2}, \quad \Gamma_{20}^{2,6} = -\frac{1}{2}, \quad \Gamma_{21}^{2,7} = -\frac{1}{2}, \quad \Gamma_{04}^{2,8} = \frac{1}{4}, \quad \Gamma_{40}^{2,8} = -\frac{1}{4}, \\ \Gamma_{01}^{2,4} = \frac{1}{2}, \quad \Gamma_{02}^{2,5} = \frac{1}{2}, \quad \Gamma_{04}^{2,7} = \frac{1}{4}, \quad \Gamma_{40}^{2,7} = -\frac{1}{4}, \quad \Gamma_{05}^{2,8} = \frac{1}{4}, \quad \Gamma_{50}^{2,8} = \frac{1}{4}, \end{aligned} \right\} \tag{A10}$$

and the non-zero elements of $\Gamma_{lm}^{3,n}$ are

$$\Gamma_{00}^{3,1} = -\frac{1}{3}, \quad \Gamma_{10}^{3,3} = -\frac{2}{3}, \quad \Gamma_{20}^{3,5} = -\frac{2}{3}, \quad \Gamma_{60}^{3,7} = -\frac{2}{3}, \quad \Gamma_{50}^{3,8} = -\frac{2}{3}. \tag{A11}$$

To obtain (4.3) from (A6), (A7) and (A8), we first computed the Christoffel symbols relative to $\{\mathbf{b}^{t_1}, \mathbf{b}^{t_1 t_2}, \mathbf{b}^{t_1 t_2 t_3}\}$ and then changed the basis to the $\{\mathbf{B}^n\}$ in (3.3). Indeed, any

set of basis tensors can be expressed through the transformation $\hat{\mathbf{B}}^{n'} = T_n^{n'}(\mathcal{I})\mathbf{B}^n$, with T an invertible matrix function of the invariants. Then, the Christoffel symbols transform as

$$\left. \begin{aligned} \hat{\Gamma}_{l'm'}^{1,n'} &= T_n^{n'} \Gamma_{lm}^{1,n} (T^{-1})_{l'}^l (T^{-1})_{m'}^m, & \hat{\Gamma}_{l'm'}^{2,n'} &= T_n^{n'} \Gamma_{lm}^{2,n} (T^{-1})_{l'}^l (T^{-1})_{m'}^m, \\ \hat{\Gamma}_{l'm'}^{3,n'} &= \left(T_n^{n'} \Gamma_{lm}^{3,n} + \frac{\partial T_n^{n'}}{\partial \mathcal{I}_k} M_{km} \right) (T^{-1})_{l'}^l (T^{-1})_{m'}^m. \end{aligned} \right\} \quad (\text{A12})$$

Appendix B. Modifications for compressible or perceived velocity gradients

As for the incompressible gradient, all the homogeneity relations for a compressible or perceived gradient $\tilde{\mathbf{A}}$ follow from (2.4), with $v_i \equiv 0$ in the compressible/perceived case. This is because $\nabla \tilde{\mathbf{A}}$ has no identically zero contractions. To derive homogeneity constraints for the compressible gradient, we follow the same steps as outlined in § 4, now employing the appropriate Christoffel symbols $\tilde{\Gamma}_{lm}^{P,n}$ and derivative components \tilde{M}_{km} , characterizing the tensor derivatives of the compressible/perceived basis tensors and invariants. In this appendix we compute $\tilde{\Gamma}_{lm}^{P,n}$ and \tilde{M}_{km} .

The basis tensors (3.3) are the same for the incompressible and compressible/perceived gradient, while the independent invariants $\tilde{\mathcal{I}}_k$ (3.4) feature the additional invariant $\tilde{\mathcal{I}}_0 = \text{Tr}(\tilde{\mathbf{A}})$ in the compressible/perceived case. The contributions from incompressibility to the tensor derivatives (4.2) and (4.3) vanish for a compressible/perceived gradient. This implies that $\tilde{\Gamma}_{lm}^{3,n} = 0$, while $\tilde{\Gamma}_{lm}^{1,n} = \Gamma_{lm}^{1,n}$ and $\tilde{\Gamma}_{lm}^{2,n} = \Gamma_{lm}^{2,n}$. The matrix of the derivatives of the invariants is

$$\tilde{M}_{km} = Z_{ml}^{-1} B_{pq}^l \frac{\partial \tilde{\mathcal{I}}_k}{\partial \tilde{\mathcal{A}}_{pq}} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 3 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & -1 \end{bmatrix}, \quad (\text{B1})$$

where the first row refers to $\tilde{\mathcal{I}}_0$, and the first column is now zero for $k > 0$ since the tensor derivatives of the invariants $\partial \tilde{\mathcal{I}}_k / \partial \tilde{\mathbf{A}}$ are no longer traceless.

REFERENCES

- ALEXAKIS, A. & BIFERALE, L. 2018 Cascades and transitions in turbulent flows. *Phys. Rep.* **767–769**, 1–101.
- BETCHOV, R. 1956 An inequality concerning the production of vorticity in isotropic turbulence. *J. Fluid Mech.* **1** (5), 497–504.
- BRAGG, A.D., HAMMOND, A.L., DHARIWAL, R. & MENG, H. 2022 Hydrodynamic interactions and extreme particle clustering in turbulence. *J. Fluid Mech.* **933**, A31.
- BUARIA, D., PUMIR, A. & BODENSCHATZ, E. 2020 Self-attenuation of extreme events in Navier–Stokes turbulence. *Nat. Commun.* **11**, 5852.
- BUARIA, D., PUMIR, A. & BODENSCHATZ, E. 2022 Generation of intense dissipation in high Reynolds number turbulence. *Phil. Trans. R. Soc. A* **380** (2218), 20210088.
- CARBONE, M. & BRAGG, A.D. 2020 Is vortex stretching the main cause of the turbulent energy cascade? *J. Fluid Mech.* **883**, R2.
- CHERTKOV, M., PUMIR, A. & SHRAIMAN, B.I. 1999 Lagrangian tetrad dynamics and the phenomenology of turbulence. *Phys. Fluids* **11** (8), 2394–2410.
- DAVIDSON, P.A. 2004 *Turbulence: An Introduction for Scientists and Engineers*. Oxford University Press.
- ENCISO, A., PERALTA-SALAS, D. & DE LIZAUR, F.T. 2016 Helicity is the only integral invariant of volume-preserving transformations. *Proc. Natl Acad. Sci. USA* **113** (8), 2035–2040.

- GRINFELD, P. 2013 *Introduction to Tensor Analysis and the Calculus of Moving Surfaces*. Springer.
- HIERRO, J. & DOPAZO, C. 2003 Fourth-order statistical moments of the velocity gradient tensor in homogeneous, isotropic turbulence. *Phys. Fluids* **15** (11), 3434–3442.
- HILL, R.J. 1997 Applicability of Kolmogorov's and Monin's equations of turbulence. *J. Fluid Mech.* **353**, 67–81.
- ITSKOV, M. 2015 *Tensor Algebra and Tensor Analysis for Engineers*. Springer.
- JOHNSON, P.L. 2020 Energy transfer from large to small scales in turbulence by multiscale nonlinear strain and vorticity interactions. *Phys. Rev. Lett.* **124**, 104501.
- JOHNSON, P.L. & MENEVEAU, C. 2016 A closure for Lagrangian velocity gradient evolution in turbulence using recent-deformation mapping of initially Gaussian fields. *J. Fluid Mech.* **804**, 387–419.
- LEPPIN, L.A. & WILCZEK, M. 2020 Capturing velocity gradients and particle rotation rates in turbulence. *Phys. Rev. Lett.* **125**, 224501.
- LUND, T.S. & NOVIKOV, E.A. 1992 Parameterization of subgrid-scale stress by the velocity gradient tensor. Center for Turbulence Research (Stanford University and NASA).
- MAJDA, A.J. & BERTOZZI, A.L. 2001 *Vorticity and Incompressible Flow*. Cambridge Texts in Applied Mathematics. Cambridge University Press.
- MENEVEAU, C. 2011 Lagrangian dynamics and models of the velocity gradient tensor in turbulent flows. *Annu. Rev. Fluid Mech.* **43** (1), 219–245.
- MEURER, A., *et al.* 2017 Sympy: symbolic computing in Python. *PeerJ Comput. Sci.* **3**, e103.
- MOMENIFAR, M., DIAO, E., TAROKH, V. & BRAGG, A.D. 2022 Dimension reduced turbulent flow data from deep vector quantisers. *J. Turbul.* **23** (4–5), 232–264.
- NASO, A. & PUMIR, A. 2005 Scale dependence of the coarse-grained velocity derivative tensor structure in turbulence. *Phys. Rev. E* **72**, 056318.
- PENNISI, S. & TROVATO, M. 1987 On the irreducibility of professor G.F. Smith's representations for isotropic functions. *Intl J. Engng Sci.* **25** (8), 1059–1065.
- RIVLIN, R.S. & ERICKSEN, J.L. 1955 Stress-deformation relations for isotropic materials. *J. Rat. Mech. Anal.* **4**, 323–425.
- SERRE, D. 1984 Les invariants du premier ordre de l'équation d'Euler en dimension trois. *Physica D* **13** (1), 105–136.
- SIGGIA, E.D. 1981 Invariants for the one-point vorticity and strain rate correlation functions. *Phys. Fluids* **24** (11), 1934–1936.
- TIAN, Y., LIVESCU, D. & CHERTKOV, M. 2021 Physics-informed machine learning of the Lagrangian dynamics of velocity gradient tensor. *Phys. Rev. Fluids* **6**, 094607.
- TOM, J., CARBONE, M. & BRAGG, A.D. 2021 Exploring the turbulent velocity gradients at different scales from the perspective of the strain-rate eigenframe. *J. Fluid Mech.* **910**, A24.
- TOWNSEND, A.A. & TAYLOR, G.I. 1951 On the fine-scale structure of turbulence. *Proc. Math. Phys. Engng Sci.* **208** (1095), 534–542.
- TSINOBER, A. 2009 *An Informal Conceptual Introduction to Turbulence*. Springer.
- WEYL, H. 1946 *The Classical Groups: Their Invariants and Representations*. Princeton Landmarks in Mathematics and Physics, Nr. 1, Teil 1. Princeton University Press.
- XU, H., PUMIR, A. & BODENSCHATZ, E. 2011 The pirouette effect in turbulent flows. *Nat. Phys.* **7**, 709–712.
- YANG, P.-F., FANG, J., FANG, L., PUMIR, A. & XU, H. 2022 Low-order moments of the velocity gradient in homogeneous compressible turbulence. *J. Fluid Mech.* **947**, R1.
- YANG, P.-F., PUMIR, A. & XU, H. 2020 Dynamics and invariants of the perceived velocity gradient tensor in homogeneous and isotropic turbulence. *J. Fluid Mech.* **897**, A9.
- ZHENG, Q.-S. 1994 Theory of representations for tensor functions—a unified invariant approach to constitutive equations. *Appl. Mech. Rev.* **47** (11), 545–587.