

CLASSICAL SOLUTIONS OF THE THIRD PAINLEVÉ EQUATION

YOSHIHIRO MURATA

1. Introduction and main results

The big problem “Do Painlevé equations define new functions? ”, what is called the problem of irreducibilities of Painlevé equations, was essentially solved by H. Umemura [16], [17] and K. Nishioka [9].

Umemura [16] analyzed Painlevé’s Stockholm Lessons [15] and extracted the concept of “classical functions”. To define “classical functions”, Umemura introduced the permissible operations to construct new known functions from already known functions. First, we note that we identify a holomorphic function f on an open set $U \subset \mathbf{C}$ with its restriction $f|_V$ onto an open subset $V \subset U$. Let S be a certain set of meromorphic functions on a domain $D \subset \mathbf{C}$. We assume that all the elements in S are already known functions. Permissible operations to construct new known functions from the set S are as follows.

DEFINITION I [16, Part II §2]. (O) Let $f(t) \in S$. Then the derived function $f'(t)$ is a new known function.

(P1) If $f_1, f_2 \in S$, then the sum $f_1 + f_2$ and the product $f_1 f_2$ are new known functions. Moreover if $f_2 \neq 0$, then the quotient f_1/f_2 is a new known function.

(P2) Let $a_1, \dots, a_n \in S$. Then any solution f of an algebraic equation $f^n + a_1 f^{n-1} + \dots + a_n = 0$ is a new known function.

(P3) Let $f(t) \in S$. Then the quadrature $\int f(t) dt$ is a new known function.

(P4) Let $a_1, \dots, a_n \in S$. Then any solution f of a linear differential equation $d^n f/dt^n + a_1 d^{n-1} f/dt^{n-1} + \dots + a_n f = 0$ is a new known function.

(P5) Let $\Gamma \subset \mathbf{C}^n$ be a lattice such that the quotient \mathbf{C}^n/Γ is an abelian variety. Let $\pi: \mathbf{C}^n \rightarrow \mathbf{C}^n/\Gamma$ be the projection. Let $f_1, \dots, f_n \in S$ be holomorphic functions on a domain $D \subset \mathbf{C}$ and ϕ be a meromorphic function on \mathbf{C}^n/Γ . Then the function $\phi \cdot \pi \cdot (f_1, \dots, f_n)$ is a new known function if it is not the constant function taking

Received March 28, 1994.

infinity.

In any operation of (P2), \dots , (P5), we consider that we take an appropriate subdomain $D' \subset D$ such that the newly constructed function is meromorphic and single valued on D' . Using these permissible operations, Umemura defined “classical functions” as follows:

DEFINITION II [16, Part II §2, Definition (2.27)]. Let f be a meromorphic function on a domain $D \subset \mathbf{C}$, \mathcal{M}_D be the set of all meromorphic functions on D and $\mathbf{C}(t)$ be the field of rational functions in a variable t . The function f is called classical if and only if there exists a tower of differential subfields $K_0 = \mathbf{C}(t)$, K_1, \dots, K_m of \mathcal{M}_D such that

(i) For any $j = 1, \dots, m$, $K_j = K_{j-1}\langle g_j \rangle = K_{j-1}(g_j, g_j', g_j'', \dots)$, where g_j is a meromorphic function obtained by one of permissible operations (P2), \dots , (P5) from the field K_{j-1} .

(ii) $f \in K_m$.

In this sense, rational functions in one variable, e^t , $\log t$, elliptic functions, the hypergeometric function and confluent hypergeometric functions are examples of classical functions. It is non-classical functions that are essentially new functions. We call a non-classical function an irreducible function.

Using the idea of Nishioka [9] and the fact that Painlevé I does not have algebraic solutions, Umemura [17] showed the theorem that every solution of Painlevé I is irreducible. After that, by the same idea, M. Noumi [10] clarified the distribution of classical solutions and irreducible solutions of Painlevé II, K. Okamoto [14] solved the case of Painlevé IV. They slightly generalized the techniques of Umemura [17] and used the facts on rational solutions of Painlevé II and IV and on solutions of Riccati equations contained in Painlevé II and IV [7]. Then our next target is Painlevé III. In this paper, we investigate the distribution of classical solutions of Painlevé III', which is equivalent to Painlevé III, in connection with the transformation group of solutions. Particularly, we completely determine all algebraic solutions. In the forthcoming paper [8], we prove a theorem that except for classical solutions derived in this paper, any solution of Painlevé III' is irreducible.

1.1. Two expressions of the third Painlevé equation

Painlevé III has an equivalent equation Painlevé III' [13, Introduction];

$$(1.1) \quad P_{III} : \frac{d^2 y}{dx^2} = \frac{1}{y} \left(\frac{dy}{dx} \right)^2 - \frac{1}{x} \frac{dy}{dx} + \frac{1}{x} (\alpha y^2 + \beta) + \gamma y^3 + \frac{\delta}{y}$$

$$(1.2) \quad P_{III'} : \frac{d^2 q}{dt^2} = \frac{1}{q} \left(\frac{dq}{dt} \right)^2 - \frac{1}{t} \frac{dq}{dt} + \frac{q^2}{4t^2} (\gamma q + \alpha) + \frac{\beta}{4t} + \frac{\delta}{4q}$$

In fact, a solution of P_{III} corresponds to that of $P_{III'}$ by the change of the variables:

$$(1.3) \quad t = x^2, \quad q = xy.$$

In this paper, we mainly treat $P_{III'}$, because it has a transformation group with good structure (See 1.2).

As is well known [13, Proposition 1.1], by the change of the variables: $t = t_1$, $q = t/q_1$, $P_{III'}(\alpha, \beta, \gamma, \delta)$ is transformed into $P_{III'}(-\beta, -\alpha, -\delta, -\gamma)$. In the same way, by the change of the variables: $t = t_1^2$, $q = q_1^2$, $P_{III'}(\alpha, \beta, 0, 0)$ is transformed into $P_{III'}(0, 0, 2\alpha, 2\beta)$ [11 II, Remark 1]. The correspondence (1.3) implies that $P_{III}(\alpha, \beta, \gamma, \delta)$ also has similar transformations to the above ones.

From these facts, we may consider that the values of the complex parameters $\alpha, \beta, \gamma, \delta$ of $P_{III'}$ and P_{III} satisfy one of the three cases:

- (A) $\alpha = \gamma = 0$ (or $\beta = \delta = 0$)
- (1.4) (B) $\gamma = 0, \alpha\delta \neq 0$ (or $\delta = 0, \beta\gamma \neq 0$)
- (C) $\gamma\delta \neq 0$.

In the case (A), $P_{III'}$ and P_{III} are solvable by quadratures [13, Proposition 1.5]. In the case (B), V. I. Gromak [1], [4, Theorem 2], [5, 2] showed that P_{III} has 3-sheeted algebraic solutions for special values of β (or α). In the case (C), any solution of $P_{III'}$ (resp. P_{III}) governs the isomonodromic deformation of a second order linear differential equation $L_{III'}$ (resp. L_{III}) which has irregular singularities of Poincaré rank 1 at the origin and at infinity, and a nonlogarithmic singularity at q [11 II, Proposition 1], [12, 4.3]. Then the case (C) is essential for $P_{III'}$ and P_{III} . In this case, Gromak [5, Theorem 9] obtained the necessary and sufficient conditions for $P_{III}(\alpha, \beta, 1, -1)$ to have rational solutions by the use of transformations of solutions of P_{III} . But, as we will mention in 1.2, the transformation group of solutions of $P_{III'}$ is isomorphic to the Affine Weyl group of the type B_2 , and so, we can treat $P_{III'}$ more successfully than P_{III} . Therefore, in this paper, by the help of this transformation group of $P_{III'}$ and by different approaches from those of Gromak [5], in this case (C), we investigate algebraic solutions and solutions expressible by Riccati solutions of $P_{III'}$ in detail. In the process of studying

algebraic solutions of $P_{III'}$, we obtain a result on algebraic solutions of P_{III} which contains the theorem by Gromak [5, Theorem 9].

Let parameters $(\alpha, \beta, \gamma, \delta)$ satisfy the case (C). Then $(\alpha, \beta, \gamma, \delta)$ can be replaced by other parameters $(\eta_0, \eta_\infty, \theta_0, \theta_\infty)$ [11 II, 2];

$$(1.5) \quad \alpha = -4\eta_\infty\theta_\infty, \beta = 4\eta_0(\theta_0 + 1), \gamma = 4\eta_\infty^2, \delta = -4\eta_0^2.$$

In addition, by the change of variables:

$$(1.6) \quad t = \lambda t_1, \quad q = \mu q_1, \quad (\lambda\mu \neq 0),$$

$P_{III'}(\eta_0, \eta_\infty, \theta_0, \theta_\infty)$ is transformed into $P_{III'}((\lambda/\mu)\eta_0, \mu\eta_\infty, \theta_0, \theta_\infty)$. Then, if the values of λ and μ are chosen appropriately, $P_{III'}(\eta_0, \eta_\infty, \theta_0, \theta_\infty)$ is transformed into a canonical type equation $P_{III'}(1, 1, \theta_0, \theta_\infty)$, which we express by $P_{III'}(\theta_0, \theta_\infty)$. From now on, we mainly consider this canonical type equation

$$P_{III'}(\theta_0, \theta_\infty) : \frac{d^2q}{dt^2} = \frac{1}{q} \left(\frac{dq}{dt} \right)^2 - \frac{1}{t} \frac{dq}{dt} + \frac{q^2}{t^2} (q - \theta_\infty) + \frac{\theta_0 + 1}{t} - \frac{1}{q}.$$

$P_{III}(\eta_0, \eta_\infty, \theta_0, \theta_\infty)$ is also transformed into a canonical type equation

$$P_{III}(\theta_0, \theta_\infty) : \frac{d^2y}{dx^2} = \frac{1}{y} \left(\frac{dy}{dx} \right)^2 - \frac{1}{x} \frac{dy}{dx} + \frac{4}{x} (-\theta_\infty y^2 + \theta_0 + 1) + 4y^3 - \frac{4}{y}$$

by a similar transformation to (1.6).

1.2. Transformation groups of $P_{III'}$ and P_{III}

As Okamoto [13, Introduction] pointed out, $P_{III'}(\eta_0, \eta_\infty, \theta_0, \theta_\infty)$ is transformed into a Hamiltonian system. In fact, putting

$$(1.7) \quad \begin{cases} q = q \\ p = \frac{t dq/dt + \eta_\infty q^2 + \theta_0 q - \eta_0 t}{2q^2}, \end{cases}$$

then we get a Hamiltonian system

$$(1.8) \quad \begin{cases} \frac{dq}{dt} = \frac{1}{t} [2q^2 p - (\eta_\infty q^2 + \theta_0 q - \eta_0 t)] = \frac{\partial H}{\partial p} \\ \frac{dp}{dt} = -\frac{1}{t} [2qp^2 - (2\eta_\infty q + \theta_0)p + \frac{1}{2} \eta_\infty (\theta_0 + \theta_\infty)] = -\frac{\partial H}{\partial q}, \end{cases}$$

where $H = (1/t)[q^2 p^2 - (\eta_\infty q^2 + \theta_0 q - \eta_0 t)p + (1/2)\eta_\infty(\theta_0 + \theta_\infty)q]$. Let $H_{III'}(\theta_0, \theta_\infty)$ denote the Hamiltonian system which corresponds to $P_{III'}(\theta_0, \theta_\infty)$, i.e.,

the system (1.8) with parameters $(1, 1, \theta_0, \theta_\infty)$. Okamoto obtained a transformation group \mathbf{G}'_H of $H_{III'}(\theta_0, \theta_\infty)$ and showed that it is isomorphic to the Affine Weyl group of the type B_2 [13, Theorem 1]. In this paper, we consider the transformation group \mathbf{G}' of $P_{III'}(\theta_0, \theta_\infty)$ which is derived from \mathbf{G}'_H by (1.7).

Generators of \mathbf{G}' and transformations which we use later are as follows:

$$\begin{aligned}
 s_0 & \begin{cases} (t, q) \rightarrow (s, Q) = (t, -t/q) \\ (\theta_0, \theta_\infty) \rightarrow (\bar{\theta}_0, \bar{\theta}_\infty) = (-\theta_\infty - 1, -\theta_0 - 1) \end{cases} \\
 s_1 & \begin{cases} (t, q) \rightarrow (s, Q) = \left(t, q \frac{t dq/dt - q^2 + \theta_\infty q - t}{t dq/dt - q^2 + \theta_0 q - t} \right) \\ (\theta_0, \theta_\infty) \rightarrow (\bar{\theta}_0, \bar{\theta}_\infty) = (\theta_\infty, \theta_0) \end{cases} \\
 s_2 & \begin{cases} (t, q) \rightarrow (s, Q) = (-t, -q) \\ (\theta_0, \theta_\infty) \rightarrow (\bar{\theta}_0, \bar{\theta}_\infty) = (\theta_0, -\theta_\infty) \end{cases} \\
 l & \begin{cases} (t, q) \rightarrow (s, Q) = \left(t, -\frac{t}{q} \frac{t dq/dt - q^2 - (\theta_0 + 2)q + t}{t dq/dt - q^2 + \theta_\infty q + t} \right) \\ (\theta_0, \theta_\infty) \rightarrow (\bar{\theta}_0, \bar{\theta}_\infty) = (\theta_0 + 1, \theta_\infty + 1) \end{cases} \\
 l^{-1} & \begin{cases} (t, q) \rightarrow (s, Q) = \left(t, -\frac{t}{q} \frac{t dq/dt + q^2 + \theta_0 q - t}{t dq/dt + q^2 - \theta_\infty q - t} \right) \\ (\theta_0, \theta_\infty) \rightarrow (\bar{\theta}_0, \bar{\theta}_\infty) = (\theta_0 - 1, \theta_\infty - 1) \end{cases} \\
 m & \begin{cases} (t, q) \rightarrow (s, Q) = \left(t, \frac{t}{q} \frac{t dq/dt + q^2 - (\theta_0 + 2)q + t}{t dq/dt + q^2 - \theta_\infty q + t} \right) \\ (\theta_0, \theta_\infty) \rightarrow (\bar{\theta}_0, \bar{\theta}_\infty) = (\theta_0 + 1, \theta_\infty - 1) \end{cases} \\
 m^{-1} & \begin{cases} (t, q) \rightarrow (s, Q) = \left(t, \frac{t}{q} \frac{t dq/dt - q^2 + \theta_0 q - t}{t dq/dt - q^2 + \theta_\infty q - t} \right) \\ (\theta_0, \theta_\infty) \rightarrow (\bar{\theta}_0, \bar{\theta}_\infty) = (\theta_0 - 1, \theta_\infty + 1) \end{cases} \\
 h & \begin{cases} (t, q) \rightarrow (s, Q) = (t, -q) \\ (\theta_0, \theta_\infty) \rightarrow (\bar{\theta}_0, \bar{\theta}_\infty) = (-\theta_0 - 2, -\theta_\infty). \end{cases}
 \end{aligned}$$

Each transformation of the above all is applicable only when a solution $q(t)$ does not vanish the numerator and the denominator of Q . \mathbf{G}' is generated by s_j 's

($j = 0, 1, 2$). These transformations combine together like

$$\begin{aligned} s_j^2 &= \text{id} \quad (j = 0, 1, 2), \\ l &= (s_1 s_2)^2 s_0 s_1, \quad m = s_2 \ell s_2, \quad h = s_1 m s_0, \\ lm &= ml, \quad hl = l^{-1} h, \quad hm = m^{-1} h. \end{aligned}$$

We also introduce the transformation group \mathbf{G} of $P_{\text{III}}(\theta_0, \theta_\infty)$, which is derived from \mathbf{G}' by (1.3). Let S_j denote the corresponding transformation to s_j ($j = 0, 1$):

$$\begin{aligned} S_0 & \begin{cases} (x, y) \rightarrow (u, Y) = (x, -1/y) \\ (\theta_0, \theta_\infty) \rightarrow (\bar{\theta}_0, \bar{\theta}_\infty) = (-\theta_\infty - 1, -\theta_0 - 1) \end{cases} \\ S_1 & \begin{cases} (x, y) \rightarrow (u, Y) = \left(x, y \frac{(x/2)(dy/dx) - xy^2 + (\theta_\infty + 1/2)y - x}{(x/2)(dy/dx) - xy^2 + (\theta_0 + 1/2)y - x} \right) \\ (\theta_0, \theta_\infty) \rightarrow (\bar{\theta}_0, \bar{\theta}_\infty) = (\theta_\infty, \theta_0). \end{cases} \end{aligned}$$

S_1 is applicable under the conditions $(x/2)(dy/dx) - xy^2 + (\theta_\infty + 1/2)y - x \neq 0$ and $(x/2)(dy/dx) - xy^2 + (\theta_0 + 1/2)y - x \neq 0$. The transformation s_2 corresponds to the transformation

$$S_\xi \quad \begin{cases} (x, y) \rightarrow (u, Y) = (\xi x, \xi y) \\ (\theta_0, \theta_\infty) \rightarrow (\bar{\theta}_0, \bar{\theta}_\infty) = (\theta_0, -\theta_\infty), \end{cases}$$

where ξ is i or $-i$ ($i = \sqrt{-1}$). S_0, S_1 and S_ξ generate \mathbf{G}' . Correctly speaking, $\mathbf{G}' = \langle S_0, S_1, S_i \rangle = \langle S_0, S_1, S_{-i} \rangle$ holds, because $(S_{\pm i})^3 = S_{(\pm i)^3} = S_{\mp i}$. Here, we have $S_0^2 = S_1^2 = \text{id}$ and $S_i S_{-i} = S_{-i} S_i = \text{id}$. The transformation s_j ($j = 0, 1$) actually corresponds to both of S_j and $(S_i)^2 S_j$. The transformation m corresponds to the transformation

$$M \quad \begin{cases} (x, y) \rightarrow (u, Y) = \left(x, \frac{1}{y} \frac{(x/2)(dy/dx) + xy^2 - (\theta_0 + 3/2)y + x}{(x/2)(dy/dx) + xy^2 - (\theta_\infty - 1/2)y + x} \right) \\ (\theta_0, \theta_\infty) \rightarrow (\bar{\theta}_0, \bar{\theta}_\infty) = (\theta_0 + 1, \theta_\infty - 1). \end{cases}$$

The transformation m^{-1} corresponds to the transformation

$$M^{-1} \quad \begin{cases} (x, y) \rightarrow (u, Y) = \left(x, \frac{1}{y} \frac{(x/2)(dy/dx) - xy^2 + (\theta_0 + 1/2)xy - x}{(x/2)(dy/dx) - xy^2 + (\theta_\infty - 1/2)xy - x} \right) \\ (\theta_0, \theta_\infty) \rightarrow (\bar{\theta}_0, \bar{\theta}_\infty) = (\theta_0 - 1, \theta_\infty + 1). \end{cases}$$

Here each transformation of the above two is applicable only when a solution

$y(x)$ does not vanish the numerator and the denominator of Y . The transformation m (resp. m^{-1}) actually corresponds to both of M and $(S_\xi)^2 M$ (resp. M^{-1} and $(S_\xi)^2 M^{-1}$). Furthermore, whichever value ξ takes, we have $M = S_\xi(S_1 S_\xi)^2 S_0(S_1 S_\xi)$. Lastly, the transformation h corresponds to the transformation

$$H \quad \begin{cases} (x, y) \rightarrow (u, Y) = (x, -y) \\ (\theta_0, \theta_\infty) \rightarrow (\bar{\theta}_0, \bar{\theta}_\infty) = (-\theta_\infty - 2, -\theta_\infty). \end{cases}$$

and the transformation $(S_\xi)^2 H$.

The Hamiltonian expression (1.8) of $P_{III'}$ and the transformation groups \mathbf{G}'_H , \mathbf{G}' also play important roles in the forthcoming paper [8].

1.3. Main results

In this paper, if we do not comment especially, “a rational function” means a single-valued algebraic function defined on a Riemann sphere \mathbf{P} , and “an algebraic function” means a many-valued algebraic function defined on \mathbf{P} .

THEOREM 1. $P_{III'}(\alpha, \beta, \gamma, \delta)$ has a rational solution if and only if $\alpha = \gamma = 0$ or $\beta = \delta = 0$.

THEOREM 2. Assume that $\gamma\delta \neq 0$.

- (1) $P_{III'}(\theta_0, \theta_\infty)$ ($(\theta_0, \theta_\infty) \in \mathbf{C}^2$) does not have rational solutions.
- (2) $P_{III'}(\theta_0, \theta_\infty)$ ($(\theta_0, \theta_\infty) \in \mathbf{C}^2$) has algebraic solutions if and only if there exists an integer I such that $\theta_\infty - \theta_0 - 1 = 2I$ or $\theta_\infty + \theta_0 + 1 = 2I$.
- (3) If $P_{III'}(\theta_0, \theta_\infty)$ has algebraic solutions, then the number of algebraic solutions is one or two. $P_{III'}(\theta_0, \theta_\infty)$ has two algebraic solutions if and only if there exist two integers I and J such that $\theta_\infty - \theta_0 - 1 = 2I$ and $\theta_\infty + \theta_0 + 1 = 2J$.
- (4) Let

$$D_+ = \{(-5/2, 1/2), (1/2, -1/2)\}, D_- = \{(-5/2, -1/2), (1/2, 1/2)\},$$

$$\alpha_+ = 1, \alpha_- = i,$$

$$\Delta_\pm = \{((L - 2K - 1)/2, \pm (L + 2K + 1)/2) \mid K \text{ and } L \text{ are integers such that } K \geq 2, L = -2, -4, \dots, -2K\}$$

$$\cup \{((L - 2K - 1)/2, \pm (L + 2K + 1)/2) \mid K \text{ and } L \text{ are integers such that } K \leq -2, L = 0, 2, \dots, 2(-K - 1)\},$$

where double signs correspond with each other. Then, algebraic solutions are classified as in the table I. In the types III_\pm, IV_\pm , the values of N, ϵ_j, b_j ($j = 1, \dots, N$) depend

on the values of θ_0 , θ_∞ and α_\pm .

Type	Conditions of $(\theta_0, \theta_\infty)$	Forms of algebraic solutions
I_\pm	$\theta_\infty \mp \theta_0 \mp 1 = 0, (\theta_0, \theta_\infty) \in \mathbf{C}^2$	$q = \alpha_\pm \sqrt{t}$
II_\pm	$(\theta_0, \theta_\infty) \in D_\pm$	$q = \alpha_\pm \sqrt{t} + \theta_\infty$
III_\pm	$\theta_\infty \mp \theta_0 \mp 1 = 2I, (\theta_0, \theta_\infty) \in \mathbf{C}^2$ $I \in \mathbf{Z} - \{0\}, (\theta_0, \theta_\infty) \notin D_\pm \cup \Delta_\pm$	$q = \sqrt{t} \left(\alpha_\pm + \frac{1}{2} \sum_{j=1}^N \frac{\varepsilon_j}{\sqrt{t} - b_j} \right)$ $\left(\begin{array}{l} N \in \mathbf{Z}, N > 0 \\ \varepsilon_j = 1 \text{ or } -1 \\ b_j \in \mathbf{C} - \{0\} \\ \text{if } j \neq j', \text{ then } b_j \neq b_{j'} \\ I = \sum_{j=1}^N \varepsilon_j \end{array} \right) \dots (*)$
IV_\pm	$(\theta_0, \theta_\infty) \in \Delta_\pm$	$q = \sqrt{t} \left(\alpha_\pm + \frac{1}{2} \sum_{j=1}^N \frac{\varepsilon_j}{\sqrt{t} - b_j} \right) + \theta_\infty$ under the same conditions as in (*)

Table I. Algebraic solutions of $P_{III'}(\theta_0, \theta_\infty)$.

Remark 1.1. (1) Every algebraic solution is actually calculable from the algebraic solution of the type I_+ by means of transformation group \mathbf{G}' . (Refer to Propositions 3.4, 3.10 and 3.11.)

(2) When the values of θ_0 and θ_∞ are restricted to real numbers, algebraic solutions are distributed as in the Figure 1.

As is well known [13, 4.1], when $\theta_0 + \theta_\infty = 0$ ($(\theta_0, \theta_\infty) \in \mathbf{C}^2$), $P_{III'}(\theta_0, \theta_\infty)$ contains all solutions of a Riccati equation:

$$(1.9) \quad \frac{dq}{dt} = -\frac{1}{t}q^2 - \frac{\theta_0}{t}q + 1.$$

By the change of variables: $q = (s/2)(d/ds) \log u - \theta_0/2$ ($u \neq 0$), $t = -s^2/4$, (1.9) is transformed into a Bessel equation:

$$\frac{d^2u}{ds^2} + \frac{1}{s} \frac{du}{ds} + \left(1 - \frac{\theta_0^2}{s^2}\right)u = 0 \quad (u \neq 0).$$

Let $\phi_\sigma(t)$ ($\sigma \in \mathbf{C}$) denote the one-parameter family of solutions of (1.9).




Figure 1. The distribution of algebraic solutions of $P_{III'}(\theta_0, \theta_\infty)$ in the case $(\theta_0, \theta_\infty) \in \mathbb{R}^2$.
 On the line --- , type I_+ .
 On a line --- , type III_+ .
 On a point \circ , type II_+ .
 On a point \bullet , type IV_+ .
 On the wavy line ~ , type I_- .
 On a wavy line ~ , type III_- .
 On a point \triangle , type II_- .
 On a point \blacktriangle , type IV_- .
 --- is a fundamental cell of the Affine Weyl group of the type B_2 .

THEOREM 3. Let I and J be any integers. If $\theta_\infty + \theta_0 = 2I$ (resp. $\theta_\infty - \theta_0 = 2J$), $P_{III'}(\theta_0, \theta_\infty)$ has a one-parameter family of solutions of the form

$$q_I(t) = \mathcal{Q}_I(\theta_0, t, \phi_\sigma(t)) \quad (\sigma \in \mathbb{C})$$

$$\text{(resp. } \bar{q}_J(t) = \bar{\mathcal{Q}}_J(\theta_0, t, \phi_\sigma(-t)) \text{) } (\sigma \in \mathbb{C}).$$

Here, \mathcal{Q}_I and $\bar{\mathcal{Q}}_J$ are rational functions in three variables with integer coefficients, forms of which depend on the values of I and J respectively. If $\theta_\infty + \theta_0 = 2I$ and $\theta_\infty - \theta_0 = 2J$, then $P_{III'}(\theta_0, \theta_\infty)$ has two one-parameter families $q_I(t)$ and $\bar{q}_J(t)$ which do not have common solutions.

Remark 1.2. (1) Theorem 3 itself does not guarantee that $q_I(t)$ and $\bar{q}_J(t)$ are all the solutions which can be expressible by solutions of Riccati equations. But, as we will show in the next paper [8], it is actually true.

(2) Okamoto [13, 4.1, 4.2, 4.3] obtained the τ -function of the solutions $q_I(t)$ and investigated its properties, relating it to the Toda lattice equation.

In the forthcoming paper [8], by the theory of differential fields, we prove the theorem that any solution of $P_{\text{III}'}(\theta_0, \theta_\infty)$ is irreducible except for algebraic solutions of Theorem 2 and one-parameter families of solutions $q_I(t)$ and $\bar{q}_J(t)$ of Theorem 3.

2. Proof of Theorem 1

First we note that the equation $P_{\text{III}'}(\alpha, \beta, \gamma, \delta)$ (See (1.2)) is equivalent to the equation

$$(2.1) \quad 4t^2 q \frac{d^2 q}{dt^2} = 4t^2 \left(\frac{dq}{dt} \right)^2 - 4tq \frac{dq}{dt} + q^3(\gamma q + \alpha) + \beta tq + \delta t^2 \quad (q \neq 0).$$

Proof of Theorem 1. Let $q(t)$ be a rational solution of $P_{\text{III}'}(\alpha, \beta, \gamma, \delta)$. We suppose that $q(t)$ has a pole at $t = \infty$. Putting the Laurent expansion of $q(t)$ at $t = \infty$ into (2.1) and comparing the coefficients of the terms of the highest degree with respect to t , we obtain the condition $\alpha = \gamma = 0$. Similarly, if $q(t)$ is holomorphic at $t = \infty$, the condition $\beta = \delta = 0$ is derived. Conversely $P_{\text{III}'}(\alpha, 0, \gamma, 0)$ has a general solution $q = 2\lambda^2 F / [(F - 1)\{(\alpha/4 - \varepsilon)F - (\alpha/4 + \varepsilon)\}]$, where $F = \mu t^\lambda$, $\varepsilon = \sqrt{\gamma} \lambda / 2$, $(\lambda, \mu) \in (\mathbf{C} - \{0\}) \times \mathbf{C}$ (Okamoto [13, Proposition 1.5]). Then, if λ is an integer, q is a rational solution of $P_{\text{III}'}(\alpha, 0, \gamma, 0)$. $P_{\text{III}'}(0, \beta, 0, \delta)$ also has rational solutions. \square

3. Proof of Theorem 2

3.1. Possible rational solutions of $P_{\text{III}'}(\theta_0, \theta_\infty)$

From now on, we assume that $\gamma\delta \neq 0$, and we consider the canonical equation $P_{\text{III}'}(\theta_0, \theta_\infty)$.

Let $q(t)$ be an algebraic solution of $P_{\text{III}'}(\theta_0, \theta_\infty)$, \mathcal{R} the Riemann surface of $q(t)$, T a Riemann sphere $\mathbf{P}, \text{pr}; \mathcal{R} \rightarrow T$ the canonical projection. By simple calculations, we can check that any algebraic singularities of $q(t)$ on $t = b \in \mathbf{C} -$

{0} is a pole. Then, \mathcal{R} has branching points only on $t = 0, \infty \in T$. By the Riemann-Hurwitz's formula, \mathcal{R} has one branching point on each point of $t = 0$ and $t = \infty$, and the two branching points have the same multiplicity. Let $n (\geq 1)$ denote this multiplicity. If we put $X = \mathbf{P}$, a Riemann sphere, and $\phi_n : X \rightarrow T, x \rightarrow t = x^n$, then $\text{pr}^{-1}\phi_n$ uniformize \mathcal{R} . Hence $z(x) = (q\phi_n)(x)$ is a rational function on $X = \mathbf{P}$. Conversely if $z(x)$ is a rational function on X , then $q(t) = (z\phi_n^{-1})(t)$ is a d -sheeted algebraic function, where d is a divisor of n .

PROPOSITION 3.1. *Let $q(t)$ be an algebraic solution of $P_{\text{III}}(\theta_0, \theta_\infty)$. Then $q(t)$ is two-sheeted and is expanded at $t = \infty$ as*

$$q(t) = C_1x + C_0 + C_{-1}x^{-1} + \dots + C_{-k}x^{-k} + \dots,$$

where $x = \sqrt{t}$, $(C_1, C_0) = (\pm 1, (\theta_\infty - \theta_0 - 1)/4)$ or $(C_1, C_0) = (\pm i, (\theta_\infty + \theta_0 + 1)/4)$. For any integer $k \geq 1$, $C_{-k} = C_1C_0R_k(\theta_0, \theta_\infty, C_1, C_0)$ holds, where $R_k(X, Y, Z, W)$ is a polynomial in four variables with complex coefficients.

Proof. Suppose $q(t)$ is an n -sheeted algebraic solution with an expansion at $t = \infty$ as

$$q(t) = C_mx^m + C_{m-1}x^{m-1} + \dots,$$

where $x = t^{1/n}$, $m \in \mathbf{Z}$, $C_m \neq 0$. We substitute this expansion into (2.1) and compare the coefficients. If $m \leq 0$, then we cannot cancel the coefficient of x^{2m} . Therefore, m must be a positive integer. We also find that $m = n/2$ and $C_m = \sqrt[4]{1} = \pm 1, \pm i$. Thus we have

$$\begin{aligned} & \text{(the right hand side of (2.1))} \\ & - \text{(the left hand side of (2.1))} \\ & = 4 \sum_{4m > p \geq 3m} \left(\sum_{k+l+v+w=p} C_k C_l C_v C_w \right) x^p + 4(-\theta_\infty C_m^2 + \theta_0 + 1) C_m x^{3m} \\ & + 4 \sum_{3m > p \geq 2m} \left\{ \sum_{k+l+v+w=p} C_k C_l C_v C_w - \theta_\infty \sum_{k+l+v=p} C_k C_l C_v + (\theta_0 + 1) C_{p-2m} \right\} x^p \\ & + 4 \sum_{2m > p} \left\{ \sum_{k+l+v+w=p} C_k C_l C_v C_w - \theta_\infty \sum_{k+l+v=p} C_k C_l C_v \right. \\ & \left. + (\theta_0 + 1) C_{p-2m} + \frac{1}{4m^2} \sum_{k+l=p} (k - 2m)(l - k) C_k C_l \right\} x^p = 0. \end{aligned}$$

Comparing the coefficients of x^{4m-d} ($1 \leq d < m$), we inductively obtain $C_{m-1} = \dots = C_1 = 0$. Next, from the coefficients of x^{3m} , we have $4(4C_0C_m^2 - \theta_\infty C_m^2 + \theta_0 + 1)C_m = 0$. Then $C_m = \pm 1$ and $C_m = \pm i$ imply $C_0 = (\theta_\infty - \theta_0 - 1)/4$ and $C_0 = (\theta_\infty + \theta_0 + 1)/4$ respectively. Continuing this process succes-

sively, from the coefficients of $x^{3m-(em+d)}$ ($e \geq 0, 1 \leq d < m$), we obtain $C_{-em-1} = \dots = C_{-(e+1)m+1} = 0$. From the coefficients of $x^{3m-(e+1)m}$ ($e \geq 1$), we find that

$$\begin{aligned} C_{-(e+1)m} &= - (C_m/4)C_0 \cdot (\text{a polynomial of } \theta_0, \theta_\infty, C_m, C_0, \dots, C_{-em} \\ &\hspace{15em} \text{with complex coefficients}) \\ &= C_m C_0 \cdot (\text{a polynomial of } \theta_0, \theta_\infty, C_m, C_0 \text{ with complex coefficients}). \end{aligned}$$

By the above arguments, it follows that the expansion of $q(t)$ at $t = \infty$ must be $q(t) = \sum_{k \leq 1} C_{mk} x^{mk} = \sum_{k \leq 1} C_{mk} (\sqrt{t})^k$, where $x = t^{1/2m}$. Hence the multiplicity $2m$ is equal to 2. □

The arguments above and (1.3) in 1.1 lead us to the following result.

PROPOSITION 3.2. *If $q(t)$ is an algebraic solution of $P_{III'}(\theta_0, \theta_\infty)$, then $y(x) = q(x^2)/x$ is a rational solution of $P_{III}(\theta_0, \theta_\infty)$. Conversely, if $y(x)$ is a rational solution of $P_{III}(\theta_0, \theta_\infty)$, then $q(t) = \sqrt{t}y(\sqrt{t})$ is an algebraic solution of $P_{III'}(\theta_0, \theta_\infty)$.*

By this proposition, what we should do to find all algebraic solutions of $P_{III'}(\theta_0, \theta_\infty)$ is to find all rational solutions of $P_{III}(\theta_0, \theta_\infty)$. In Proposition 3.3, we determine all possible forms of rational solutions of $P_{III}(\theta_0, \theta_\infty)$. From Propositions 3.1, 3.2, we first obtain the following lemma.

LEMMA 3.1. *Let $y(x)$ be a rational solution of $P_{III}(\theta_0, \theta_\infty)$. Then $y(x)$ is expanded at $x = \infty$ as*

$$y(x) = D_0 + D_{-1}/x + \dots + D_{-k-1}/x^{k+1} + \dots,$$

where $(D_0, D_1) = (\pm 1, (\theta_\infty - \theta_0 - 1)/4)$ or $(\pm i, (\theta_\infty + \theta_0 + 1)/4)$. For any integer $k \geq 1$, $D_{-k-1} = D_0 D_{-1} \cdot R_k(\theta_0, \theta_\infty, D_0, D_{-1})$ holds, where R_k is the same polynomial as in Proposition 3.1.

By simple calculation, we obtain the next lemma.

LEMMA 3.2. (1) *If $y(x)$ is a rational solution of $P_{III}(\theta_0, \theta_\infty)$ with a pole at $x = 0$, then it is expanded at $x = 0$ as*

$$y(x) = \theta_\infty x^{-1} + E_0 + E_1 x + \dots,$$

where $\theta_\infty \neq 0$.

(2) *If $y(x)$ is a rational solution of $P_{III}(\theta_0, \theta_\infty)$ with a pole at $x = b (\in \mathbf{C} -$*

{0}), then it is expanded at $x = b$ as

$$y(x) = (\pm 1/2)(x - b)^{-1} + F_0 + F_1(x - b) + \dots$$

The above two lemmas give us the information on the possible forms of rational solutions of $P_{III}(\theta_0, \theta_\infty)$.

PROPOSITION 3.3. *If $y(x)$ is a rational solution of $P_{III}(\theta_0, \theta_\infty)$, then it must belong to one of the eight types in the table II, where $\alpha_+ = 1$ or -1 , $\alpha_- = i$ or $-i$, and double signs correspond with each other.*

Type	Conditions of $(\theta_0, \theta_\infty)$	Forms of $y(x)$'s
I_\pm	$\theta_\infty \mp \theta_0 \mp 1 = 0, (\theta_0, \theta_\infty) \in \mathbb{C}^2$	$y = \alpha_\pm$
II_\pm	$3\theta_\infty \pm \theta_0 \pm 1 = 0, \theta_\infty \neq 0$ $(\theta_0, \theta_\infty) \in \mathbb{C}^2$	$y = \alpha_\pm + \theta_\infty/x$
III_\pm	$\theta_\infty \mp \theta_0 \mp 1 = 2I, (\theta_0, \theta_\infty) \in \mathbb{C}^2$	$y = \alpha_\pm + \frac{1}{2} \sum_{j=1}^N \frac{\varepsilon_j}{x - b_j}$ $\left(\begin{array}{l} N \in \mathbb{Z}, N > 0 \\ \varepsilon_j = 1 \text{ or } -1 \\ b_j \in \mathbb{C} - \{0\} \\ \text{if } j \neq j', \text{ then } b_j \neq b_{j'} \\ I = \sum_{j=1}^N \varepsilon_j \end{array} \right) \dots (*)$
IV_\pm	$3\theta_\infty \pm \theta_0 \pm 1 = -2I, \theta_\infty \neq 0$ $(\theta_0, \theta_\infty) \in \mathbb{C}^2$	$y = \alpha_\pm + \frac{\theta_\infty}{x} + \frac{1}{2} \sum_{j=1}^N \frac{\varepsilon_j}{x - b_j}$ under the same conditions as in (*)

Table II. Possible types of rational solutions of $P_{III}(\theta_0, \theta_\infty)$.

Proof. By Lemma 3.1 and Lemma 3.2, the possible decompositions of $y(x)$ into partial fractions are classified into the eight types I_\pm, \dots, IV_\pm . For each type, we can derive the conditions which θ_0 and θ_∞ should fulfill. Here we show the case IV_\pm only.

If $|b_j| < |x|$, then we have

$$\frac{\varepsilon_j}{x - b_j} = \frac{\varepsilon_j}{x} \frac{1}{1 - b_j/x} = \frac{\varepsilon_j}{x} (1 + b_j/x + b_j^2/x^2 + \dots).$$

Hence, in the neighborhood of $x = \infty$, we obtain an expansion

$$y(x) = \alpha_{\pm} + \theta_{\infty}/x + (1/2)\{I/x + (\sum_{j=1}^N \varepsilon_j b_j)/x^2 + \dots\}.$$

From Lemma 3.1, it follows that $\theta_{\infty} + I/2 = (\theta_{\infty} \mp \theta_0 \mp 1)/4$. Then, we have $3\theta_{\infty} \pm \theta_0 \pm 1 = -2I$. \square

Here we introduce new notations $R(\theta_0, \theta_{\infty})$ and $r(\theta_0, \theta_{\infty})$. Let $R(\theta_0, \theta_{\infty})$ denote the set of all rational solutions of $P_{\text{III}}(\theta_0, \theta_{\infty})$. If $P_{\text{III}}(\theta_0, \theta_{\infty})$ does not have a rational solution, we consider $R(\theta_0, \theta_{\infty}) = \emptyset$. We express an element of $R(\theta_0, \theta_{\infty})$ by a form (x, y) , where x is the independent variable of $P_{\text{III}}(\theta_0, \theta_{\infty})$ and y is a rational solution of $P_{\text{III}}(\theta_0, \theta_{\infty})$ expressed by the variable x . Next, let $r(\theta_0, \theta_{\infty})$ denote the set of all rational solutions of $P_{\text{III}}(\theta_0, \theta_{\infty})$ of the types $\text{I}_+, \dots, \text{IV}_+$. $r(\theta_0, \theta_{\infty})$ is a subset of $R(\theta_0, \theta_{\infty})$. By Proposition 3.3, we must study eight types of rational solutions. But the following result decreases our labor.

PROPOSITION 3.4. (1) *The mapping*

$$\begin{aligned} S_i: R(\theta_0, \theta_{\infty}) &\rightarrow R(\bar{\theta}_0, \bar{\theta}_{\infty}) = R(\theta_0, -\theta_{\infty}), \\ (x, y) &\rightarrow S_i(x, y) = (u, Y) = (ix, iy) \end{aligned}$$

is bijective, where $i = \sqrt{-1}$.

(2) If (x, y) is a rational solution of the type J_{\pm} ($J = \text{I}, \dots, \text{IV}$), then $S_i(x, y) = (u, Y)$ is of the type J_{\mp} , where double signs correspond with each other.

We omit the proof. By this proposition, we may concentrate on the study of rational solutions of the types $\text{I}_+, \dots, \text{IV}_+$ in the following.

3.2. Existence and uniqueness

First, we note that $P_{\text{III}}(\theta_0, \theta_{\infty})$ is equivalent to the equation

$$(3.1) \quad xy \frac{d^2 y}{dx^2} = x \left(\frac{dy}{dx} \right)^2 - y \frac{dy}{dx} + 4(-\theta_{\infty} y^2 + \theta_0 + 1)y + 4xy^4 - 4x \quad (y \neq 0).$$

PROPOSITION 3.5. (1) (Existence and uniqueness of the type I_+) If $y(x) = 1$ or $y(x) = -1$ belongs to $r(\theta_0, \theta_{\infty})$, then $(\theta_0, \theta_{\infty}) \in \mathbf{C}^2$ satisfies the condition $\theta_{\infty} - \theta_0 - 1 = 0$. Conversely, if $\theta_{\infty} - \theta_0 - 1 = 0$ ($(\theta_0, \theta_{\infty}) \in \mathbf{C}^2$), then $r(\theta_0, \theta_{\infty}) = \{y(x) = \pm 1\}$.

(2) (Existence and uniqueness of the type II_+) If $y(x) = 1 + \theta_{\infty}/x$ or $y(x) =$

$-1 + \theta_\infty/x$ belongs to $r(\theta_0, \theta_\infty)$, then $(\theta_0, \theta_\infty) = (-5/2, 1/2)$ or $(1/2, -1/2)$. Conversely, if $(\theta_0, \theta_\infty) = (-5/2, 1/2)$ or $(1/2, -1/2)$, then $r(\theta_0, \theta_\infty) = \{y(x) = \pm 1 + \theta_\infty/x\}$.

Proof. We prove (2) only. The proof of (1) is the same as that of (2). If $P_{III}(\theta_0, \theta_\infty)$ has $y(x) = 1 + \theta_\infty/x$ or $y(x) = -1 + \theta_\infty/x$ as a solution, it follows from Proposition 3.3 that $3\theta_\infty + \theta_0 + 1 = 0$ and $\theta_\infty \neq 0$. On the other hand, substituting $y(x)$ into (3.1) and comparing the coefficients of x^{-2} , we obtain $4\theta_\infty^2 = 1$. Therefore $(\theta_0, \theta_\infty) = (-5/2, 1/2)$ or $(1/2, -1/2)$. Conversely, if $(\theta_0, \theta_\infty) = (-5/2, 1/2)$ or $(1/2, -1/2)$, then we have $3\theta_\infty + \theta_0 + 1 = 0$ and $4\theta_\infty^2 = 1$. At this time, $y(x) = \pm 1 + \theta_\infty/x$ satisfy (3.1). Moreover, Lemma 3.1 ensure that $r(\theta_0, \theta_\infty)$ does not contain rational solutions of other types. \square

PROPOSITION 3.6 (Existence and uniqueness of the type III_+ and the type IV_+ (1)). (1) If $P_{III}(\theta_0, \theta_\infty)$ has a rational solution of the type III_+ , then it must be that $r(\theta_0, \theta_\infty) = \{y(x) = \pm 1 + (1/2) \sum_{j=1}^N \epsilon_j/(x \mp b_j)\}$, where double signs correspond with each other.

(2) If $P_{III}(\theta_0, \theta_\infty)$ has a rational solution of the type IV_+ , then it must be that $r(\theta_0, \theta_\infty) = \{y(x) = \mp 1 + \theta_\infty/x + (1/2) \sum_{j=1}^N \epsilon_j/(x \mp b_j)\}$, where double signs correspond with each other.

Proof. We prove (2) only. Since the transformation $(S_i)^2: R(\theta_0, \theta_\infty) \rightarrow R(\theta_0, \theta_\infty)$, $(x, y) \rightarrow (u, Y) = (-x, -y)$ is bijective, if $y(x) = 1 + \theta_\infty/x + (1/2) \sum_{j=1}^N \epsilon_j/(x - b_j)$ is a solution of $P_{III}(\theta_0, \theta_\infty)$, then $Y(u) = -1 + \theta_\infty/u + (1/2) \sum_{j=1}^N \epsilon_j/(u + b_j)$ is also a solution of $P_{III}(\theta_0, \theta_\infty)$, and vice versa. Next, Lemma 3.1 ensure that for a fixed $(\theta_0, \theta_\infty)$ a rational solution expressed as $y = 1 + D_{-1}/x + D_{-2}/x^2 + \dots$ or $y = -1 + D_{-1}/x + D_{-2}/x^2 + \dots$ is uniquely determined. \square

Remark 3.1. From Propositions 3.4, 3.5 and 3.6, we see that $R(\theta_0, \theta_\infty)$ has only the following four possibilities:

- (i) $R(\theta_0, \theta_\infty) = \phi$.
- (ii) $R(\theta_0, \theta_\infty) = \{\text{two rational solutions of the type } J_+\}$
 $(J_+ = I_+, \dots, IV_+)$.
- (iii) $R(\theta_0, \theta_\infty) = \{\text{two rational solutions of the type } K_-\}$
 $(K_- = I_-, \dots, IV_-)$.

$$(iv) \quad R(\theta_0, \theta_\infty) = \{\text{two rational solutions of the type } J_+\} \\ \cup \{\text{two rational solutions of type } K_-\} \\ (J_+ = I_+, \dots, IV_+, K_- = I_-, \dots, IV_-).$$

From now on, we study the transformations S_0, H and M in order to determine necessary and sufficient conditions for rational solutions of the types III_+, IV_+ to exist. The next lemma is prepared for the proofs of Proposition 3.7 and Proposition 3.9. We omit the proof of it.

LEMMA 3.3. *Let $y(x) = \pm 1 + P(x)/Q(x)$ be a rational solution of the type J_+ ($J_+ = II_+, \dots, IV_+$). Here $P(x)$ and $Q(x)$ are polynomials with complex coefficients. We do not assume that $P(x)$ and $Q(x)$ are coprime.*

- (1) *If $Q(0) \neq 0$, then $y(x)$ is of the type III_+ .*
- (2) *If $Q(0) = 0$ and $P(0) \neq 0$, then $y(x)$ is of the type II_+ or the type IV_+ .*

PROPOSITION 3.7. (1) *The mapping*

$$S_0 : r(\theta_0, \theta_\infty) \rightarrow r(\bar{\theta}_0, \bar{\theta}_\infty) = r(-\theta_\infty - 1, -\theta_0 - 1), \\ (x, y) \rightarrow S_0(x, y) = (x, Y) = (x, -1/y)$$

is bijective.

(2) *Let y be a type IV_+ rational solution of $P_{III}(\theta_0, \theta_\infty)$, where $3\theta_\infty + \theta_0 + 1 = -2I$ ($I \in \mathbf{Z}$), $\theta_\infty \neq 0$, and let $S_0(x, y) = (x, Y)$. Then Y is a type III_+ rational solution of $P_{III}(\bar{\theta}_0, \bar{\theta}_\infty)$ with the condition $Y(0) = 0$, where $\bar{\theta}_\infty - \bar{\theta}_0 - 1 = 2\bar{I}$ ($\bar{I} \in \mathbf{Z} - \{0\}$).*

Proof. (1) It is obvious.

(2) Substituting $y = \pm 1 + \theta_\infty/x + (1/2) \sum_{j=1}^N \epsilon_j/(x \mp b_j)$ into $Y = -1/y = \mp 1 + (\pm y - 1)/y$, we obtain $Y(x) = \mp 1 + P(x)/Q(x)$, where

$$P(x) = (\pm y - 1)x \prod_{j=1}^N (x \mp b_j) \in \mathbf{C}[x] \\ Q(x) = yx \prod_{j=1}^N (x \mp b_j) \in \mathbf{C}[x] \\ \deg P \leq N, \deg Q = N + 1.$$

Since $Q(0) = \theta_\infty \prod_{j=1}^N (\mp b_j) \neq 0$, by Lemma 3.3, $Y(x)$ is of the type III_+ . Then, we have $\bar{\theta}_\infty - \bar{\theta}_0 - 1 = 2\bar{I}$, where $\bar{I} \in \mathbf{Z} - \{0\}$ because of Proposition 3.5 (1). Next, if we assume $Y(0) \neq 0$, from $S_0(x, Y) = (x, y)$ and the similar argument as the above, we obtain that y is of the type III_+ . This is a contradiction. Then $Y(0) = 0$ must hold. □

COROLLARY 3.1. *We consider a set*

$$\tilde{D} = \{((L - 2K - 1)/2, (L + 2K + 1)/2) \in \mathbf{R}^2 \mid K, L \in \mathbf{Z}, K \neq 0\}.$$

If $P_{\text{III}}(\theta_0, \theta_\infty)$ has a rational solution of the type IV_+ , then $(\theta_0, \theta_\infty) \in \tilde{D}$.

Proof. By Proposition 3.7 (2), if (x, y) is a type IV_+ rational solution of $P_{\text{III}}(\theta_0, \theta_\infty)$, then $S_0(x, y) = (x, Y)$ is a type III_+ rational solution of $P_{\text{III}}(\bar{\theta}_0, \bar{\theta}_\infty)$, and $(\bar{\theta}_0, \bar{\theta}_\infty)$ satisfies a relation $\bar{\theta}_\infty - \bar{\theta}_0 - 1 = 2\bar{I}$ ($\bar{I} \in \mathbf{Z} - \{0\}$). Hence,

$$\theta_\infty - \theta_0 - 1 = (-\bar{\theta}_0 - 1) - (-\bar{\theta}_\infty - 1) - 1 = \bar{\theta}_\infty - \bar{\theta}_0 - 1 = 2\bar{I}.$$

On the other hand, we have $3\theta_\infty + \theta_0 + 1 = -2I$ ($I \in \mathbf{Z}$). By the both equalities, we obtain $\theta_0 = (-I - 3\bar{I} - 2)/2$, $\theta_\infty = (-I + \bar{I})/2$. Putting $-I - \bar{I} - 1 = L$ and $\bar{I} = K (\neq 0)$, we obtain $\theta_0 = (L - 2K - 1)/2$, $\theta_\infty = (L + 2K + 1)/2$. □

Remark 3.2. A point $((L - 2K - 1)/2, (L + 2K + 1)/2)$ is an intersection of two lines $\theta_\infty - \theta_0 - 1 = 2K$ and $\theta_\infty + \theta_0 = L$ in the real plane \mathbf{R}^2 .

From the table II of Proposition 3.3, Proposition 3.5 (2) and the Corollary 3.1, we obtain

COROLLARY 3.2. *Let $\Sigma_\pm = \{(\theta_0, \theta_\infty) \in \mathbf{C}^2 \mid \text{there exists an integer } I \text{ such that } \pm I \geq 0, \theta_\infty - \theta_0 - 1 = 2I\}$. If $r(\theta_0, \theta_\infty) = \phi$, then $(\theta_0, \theta_\infty) \in \Sigma_+ \cup \Sigma_-$.*

PROPOSITION 3.8. (1) *The mapping*

$$\begin{aligned} H : r(\theta_0, \theta_\infty) &\rightarrow r(\bar{\theta}_0, \bar{\theta}_\infty) = r(-\theta_0 - 2, -\theta_\infty), \\ (x, y) &\rightarrow H(x, y) = (x, Y) = (x, -y) \end{aligned}$$

is bijective.

(2) *If (x, y) is a rational solution of the type J_+ ($J_+ = \text{I}_+, \dots, \text{IV}_+$), then $H(x, y) = (x, Y)$ is also a rational solution of the type J_+ .*

We omit the proof of this proposition. This proposition implies that the set $r(\theta_0, \theta_\infty)$ ($(\theta_0, \theta_\infty) \in \Sigma_+$) are uniquely determined by the set $r(\theta_0, \theta_\infty)$ ($(\theta_0, \theta_\infty) \in \Sigma_-$), because $\bar{\theta}_\infty - \bar{\theta}_0 - 1 = (-\theta_\infty) - (-\theta_0 - 2) - 1 = -(\theta_\infty - \theta_0 - 1)$. Therefore, we may investigate the set $r(\theta_0, \theta_\infty)$ only in the case $(\theta_0, \theta_\infty) \in \Sigma_-$.

PROPOSITION 3.9. (1) *The mapping*

$$\begin{aligned} M : r(\theta_0, \theta_\infty) &\rightarrow r(\bar{\theta}_0, \bar{\theta}_\infty) = r(\theta_0 + 1, \theta_\infty - 1), \\ (x, y) &\rightarrow M(x, y) = (x, Y) \\ &= \left(x, \frac{1}{y} \frac{(x/2)(dy/dx) + xy^2 - (\theta_0 + 3/2)y + x}{(x/2)(dy/dx) + xy^2 - (\theta_\infty - 1/2)y + x} \right) \end{aligned}$$

is bijective.

(2) If (x, y) is a rational solution of the type J_+ ($J_+ = I_+, \dots, IV_+$), then $M(x, y) = (x, Y)$ is a rational solution as in the table III. Here double signs correspond with each other, and $(\theta_0, \theta_\infty) \in \Sigma_-$.

Proof. (1) By Lemma 3.1, any rational solution $y(x)$ of the type J_+ ($J_+ = I_+, \dots, IV_+$) of $P_{III}(\theta_0, \theta_\infty)$ is developed as

$$y(x) = \pm 1 + D_{-1}/x + \dots + D_{-k-1}/x^{k+1} + \dots$$

at $x = \infty$. Then it follows that

$$(x/2)(dy/dx) + xy^2 - Ay + x = 2x + O(1) \neq 0,$$

at $x = \infty$, whichever $A = \theta_0 + 3/2$ or $A = \theta_\infty - 1/2$. Therefore we can apply M to any solution in $r(\theta_0, \theta_\infty)$. By a similar reason, the inverse transformation M^{-1} is applicable to any solution in $r(\bar{\theta}_0, \bar{\theta}_\infty)$. Therefore M is bijective.

(2) We prove only the case $J_+ = III_+$. Proofs of the other cases are done in the same ways. First, by the assumption $(\theta_0, \theta_\infty) \in \Sigma_-$ and Proposition 3.5 (1), we have $\theta_\infty - \theta_0 - 1 = 2I \leq -2$, and $\bar{\theta}_\infty - \bar{\theta}_0 - 1 = (\theta_\infty - 1) - (\theta_0 + 1) - 1 = 2(I - 1) \leq -4$. Therefore, from Proposition 3.5, Y must be of the type III_+ or the type IV_+ . Next, we substitute $y = \pm 1 + (1/2) \sum_{j=1}^N \varepsilon_j / (x \mp b_j)$ into

$$\begin{aligned} (3.2) \quad Y &= \frac{1}{y} \frac{(x/2)(dy/dx) + xy^2 - (\theta_0 + 3/2)y + x}{(x/2)(dy/dx) + xy^2 - (\theta_\infty - 1/2)y + x} \\ &= \frac{1}{y} + \frac{\theta_\infty - \theta_0 - 2}{(x/2)(dy/dx) + xy^2 - (\theta_\infty - 1/2)y + x}. \end{aligned}$$

Here we note that

$$\frac{1}{y} = \pm 1 + \frac{\mp y + 1}{y} = \pm 1 + \frac{P_1(x)}{Q_1(x)},$$

where

$$P_1(x) = (\mp y + 1) \prod_{j=1}^N (x \mp b_j) \in \mathbf{C}[x]$$

$$Q_1(x) = y \prod_{j=1}^N (x \mp b_j) \in \mathbf{C}[x]$$

$$\deg P_1 = N - 1, \quad \deg Q_1 = N.$$

Type of $y(x)$	Conditions of $(\theta_0, \theta_\infty) \in \Sigma_-$ Form of $y(x)$	Conditions of $(\bar{\theta}_0, \bar{\theta}_\infty) \in \Sigma_-$ Type of $Y(x)$
I_+	$\theta_\infty - \theta_0 - 1 = 0$ $y = \pm 1$	$\bar{\theta}_\infty - \bar{\theta}_0 - 1 = -2$ (i) If $(\bar{\theta}_0, \bar{\theta}_\infty) \neq (1/2, -1/2)$, then Y is of the type III_+ . Here $Y(0) = 0$ if and only if $(\bar{\theta}_0, \bar{\theta}_\infty) = (-1/2, -3/2)$. (ii) If $(\bar{\theta}_0, \bar{\theta}_\infty) = (1/2, -1/2)$, then Y is of the type II_+ .
II_+	$(\theta_0, \theta_\infty) = (1/2, -1/2)$ $y = \pm 1 + (-1/2)/x$	$(\bar{\theta}_0, \bar{\theta}_\infty) = (3/2, -3/2)$ Y is of the type IV_+ .
III_+	$\theta_\infty - \theta_0 - 1 = 2I$ $(I = \sum_{j=1}^N \varepsilon_j < 0)$ $y = \pm 1 + (1/2) \sum_{j=1}^N \varepsilon_j / (x \mp b_j)$	$\bar{\theta}_\infty - \bar{\theta}_0 - 1 = 2(I - 1)$ (i) If $\theta_0 \neq 1/2$ and $y(0) \neq 0$, then Y is of the type III_+ . $Y(0) = 0$ if and only if $\theta_0 = -3/2$. (ii) If $\theta_\infty = 1/2$ and $y(0) \neq 0$, then Y is of the type IV_+ . (iii) If $y(0) = 0$, then Y is of the type III_+ or the type IV_+ .
IV_+	$(\theta_0, \theta_\infty) = ((L - 2K - 1)/2, (L + 2K + 1)/2) \in \bar{D}$ $K < 0$ $y = z + \theta_\infty/x$ ($\theta_\infty \neq 0$), where $z = \pm 1 + (1/2) \sum_{j=1}^N \varepsilon_j / (x \mp b_j)$	$(\bar{\theta}_0, \bar{\theta}_\infty) = ((\bar{L} - 2\bar{K} - 1)/2, (\bar{L} + 2\bar{K} + 1)/2) \in \bar{D}$ $\bar{K} = K - 1 < 0, \bar{L} = L$ (i) If $\theta_\infty \neq -1/2$ and $z(0) \neq 0$, then Y is of the type III_+ . (ii) If $\theta_\infty = -1/2$ or $z(0) = 0$, then Y is of the type IV_+ .

Table III. Actions of the mapping M .

Particularly, we have

$$(3.3) \quad P_1(0) = (\mp y(0) + 1) \prod_{j=1}^N (\mp b_j), \quad Q_1(0) = y(0) \prod_{j=1}^N (\mp b_j).$$

Furthermore, we note that

$$\frac{\theta_\infty - \theta_0 - 2}{(x/2)(dy/dx) + xy^2 - (\theta_\infty - 1/2)y + x} = \frac{\theta_\infty - \theta_0 - 2}{x\{(1/2)(dy/dx) + y^2 + 1\} - (\theta_\infty - 1/2)y} = \frac{P_2(x)}{Q_2(x)},$$

where

$$P_2(x) = (\theta_\infty - \theta_0 - 2) \prod_{j=1}^N (x \mp b_j)^2 \in \mathbf{C}[x]$$

$$Q_2(x) = x\{(1/2)(dy/dx) + y^2 + 1\} \prod_{j=1}^N (x \mp b_j)^2 - (\theta_\infty - 1/2)y \prod_{j=1}^N (x \mp b_j)^2 \in \mathbf{C}[x]$$

$$\deg P_2 = 2N, \quad \deg Q_2 = 2N + 1.$$

In particular, we obtain

$$(3.4) \quad \begin{aligned} P_2(0) &= (\theta_\infty - \theta_0 - 2) \prod_{j=1}^N (\mp b_j)^2 = (2I - 1) \prod_{j=1}^N b_j^2 \neq 0 \\ Q_2(0) &= -(\theta_\infty - 1/2)y(0) \prod_{j=1}^N b_j^2. \end{aligned}$$

Since $Y = \pm 1 + P_1(x)/Q_1(x) + P_2(x)/Q_2(x)$, it follows from (3.3), (3.4) and Lemma 3.3 that

- (i) When $\theta_\infty \neq 1/2$ and $y(0) \neq 0$, Y is of the type III₊.
- (ii) When $\theta_\infty = 1/2$ and $y(0) \neq 0$, Y is of the type IV₊.
- (iii) When $y(0) = 0$, Y is either of the type III₊ or of the type IV₊.

In the case (i), (3.2) imply that

$$Y(0) = \frac{1}{y(0)} \frac{-(\theta_0 + 3/2)y(0)}{-(\theta_\infty - 1/2)y(0)} = \frac{(\theta_0 + 3/2)}{(\theta_\infty - 1/2)} \frac{1}{y(0)} = 0$$

if and only if $\theta_0 = -3/2$. □

Here we prepare a simple lemma for Proposition 3.10.

LEMMA 3.4. *Let $y = z + \theta_\infty/x$ ($\theta_\infty \neq 0$) be a rational solution of the type IV₊ of $P_{\text{III}}(\theta_0, \theta_\infty)$, where $z = \pm 1 + (1/2) \sum_{j=1}^N \varepsilon_j/(x \mp b_j)$. If $\theta_\infty \neq \pm 1/2$, then we have $z(0) = 0$.*

Proof. The solution $y(x)$ is developed at $x = 0$ as $y = \theta_\infty/x + z(0) + O(x)$. We substitute this expansion into the equation (3.1). Comparing the coefficients of x^{-2} , we obtain $(4\theta_\infty^2 - 1)z(0) = 0$. Therefore, if $\theta_\infty \neq \pm 1/2$, then $z(0) = 0$ must hold. □

From Propositions 3.5, \dots , 3.9 and Lemma 3.4, we can derive the following final result about the existence of rational solutions of the type III_+ and the type IV_+ .

PROPOSITION 3.10 (Existence and uniqueness of the type III_+ and the type IV_+ (2)). *We set*

$$E_+ = \{((L - 2K - 1)/2, (L + 2K + 1)/2) \mid K \text{ and } L \text{ are integers such that } K \geq 2, L = -2, -4, \dots, -2K\}$$

$$E_- = \{((L - 2K - 1)/2, (L + 2K + 1)/2) \mid K \text{ and } L \text{ are integers such that } K \leq -2, L = 0, 2, \dots, 2(-K - 1)\}$$

$$E = E_+ \cup E_- \cup \{(-5/2, 1/2), (1/2, -1/2)\}$$

$$\Sigma_* = \{(\theta_0, \theta_\infty) \in \mathbb{C}^2 \mid \text{there exists a nonzero integer } I \text{ such that } \theta_\infty - \theta_0 - 1 = 2I\}.$$

- (1) *If $r(\theta_0, \theta_\infty)$ contains a rational solution of the type III_+ , then it must be that $(\theta_0, \theta_\infty) \in \Sigma_* - E$. Conversely, if $(\theta_0, \theta_\infty) \in \Sigma_* - E$, then $r(\theta_0, \theta_\infty) = \{\text{two rational solutions of the type } \text{III}_+\}$.*
- (2) *If $r(\theta_0, \theta_\infty)$ contains a rational solution of the type IV_+ , then it must be that $(\theta_0, \theta_\infty) \in E_+ \cup E_-$. Conversely, if $(\theta_0, \theta_\infty) \in E_+ \cup E_-$, then $r(\theta_0, \theta_\infty) = \{\text{two rational solutions of the type } \text{IV}_+\}$.*

Proof. We prove (1) and (2) simultaneously. Refer to the figure 1 in the Remark 1.1 after Theorem 2. By Proposition 3.5 and Corollary 3.2, we first note that if $r(\theta_0, \theta_\infty)$ contains rational solutions of the type III_+ or the type IV_+ , then $(\theta_0, \theta_\infty) \in \Sigma_*$.

Step 1. For any integer I , we define a set

$$\Pi_I = \{(\theta_0, \theta_\infty) \in \mathbb{C}^2 \mid \theta_\infty - \theta_0 - 1 = 2I\}.$$

By their definitions, $\Sigma_* \cup \Pi_0 = \Sigma_+ \cup \Sigma_-, E_+ \cup \{(-5/2, 1/2)\} \subset \Sigma_+, E_- \cup \{(1/2, -1/2)\} \subset \Sigma_-$ hold. Let $P_{K,L}$ denote a point $((L - 2K - 1)/2, (L + 2K + 1)/2)$ in \mathbb{C}^2 , where K and L are integers. As we mentioned in the Remark 3.2 after Corollary 3.1, $P_{K,L}$ is an intersection of a plane Π_K and a plane $\theta_\infty + \theta_0 = L$. When $(\theta_0, \theta_\infty) = P_{K,L}$, we express $r(\theta_0, \theta_\infty)$ by $r(P_{K,L})$.

Step 2. By Proposition 3.5, Proposition 3.9 (2) I_+ and Proposition 3.6 (1), it turns out that when $(\theta_0, \theta_\infty) \in \Pi_{-1} - \{P_{-1,-2}, P_{-1,0}\}$, $r(\theta_0, \theta_\infty) = \{\text{two rational}$

solutions of the type III_+ , which satisfy $y(0) \neq 0$ }, and that $r(P_{-1,-2}) = \{\text{two rational solutions of the type } \text{III}_+, \text{ which satisfy } y(0) = 0\}$, $r(P_{-1,0}) = \{\text{two rational solutions of the type } \text{II}_+\}$.

Step 3. From now, we prove the following facts (F1), (F2), (F3) by induction. Here we assume $-K = k \geq 2$.

(F1) $r(\theta_0, \theta_\infty) = \{\text{two rational solutions of the type } \text{III}_+, \text{ which satisfy } y(0) \neq 0\}$ for any $(\theta_0, \theta_\infty) \in \Pi_{-k} - \{P_{-k,L} \mid L = -2k, -2(k-1), \dots, 2(k-1)\}$

(F2) $r(P_{-k,L}) = \{\text{two rational solutions of the type } \text{III}_+, \text{ which satisfy } y(0) = 0\}$ for $L = -2k, -2(k-1), \dots, -2$.

(F3) $r(P_{-k,L}) = \{\text{two rational solutions of the type } \text{IV}_+\}$ for $L = 0, 2, \dots, 2(k-1)$.

By proposition 3.9 (1), we note that $M(r(\theta_0, \theta_\infty)) = r(\bar{\theta}_0, \bar{\theta}_\infty)$, where $\bar{\theta}_\infty - \bar{\theta}_0 - 1 = (\theta_\infty - \theta_0 - 1) - 2$, and that $M(r(P_{-k,L})) = r(P_{-(k+1),L})$.

(i) Let $k = 2$. From the results in step 2 and Proposition 3.9 (2) II_+ , III_+ (ii), we see that if $L = 0$ or 2 , then $r(P_{-2,L}) = \{\text{two rational solutions of the type } \text{IV}_+\}$. By Proposition 3.7 (2), we see that if $L = -4$ or -2 , then $r(P_{-2,L}) = \{\text{two rational solutions of the type } \text{III}_+, \text{ which satisfy } y(0) = 0\}$. Hence, Proposition 3.9 (2) III_+ (i) tells us that for any $(\theta_0, \theta_\infty) \in \Pi_{-2} - \{P_{-2,L} \mid L = -4, -2, 0, 2\}$, $r(\theta_0, \theta_\infty) = \{\text{two rational solutions of the type } \text{III}_+, \text{ which satisfy } y(0) \neq 0\}$ holds.

(ii) We suppose that $k \geq 2$ and that (F1), (F2), (F3) hold for k . By Proposition 3.9 (2) III_+ (ii), IV_+ (ii) and Lemma 3.4, we see that if $L = 0, 2, \dots, 2k$, then $r(P_{-(k+1),L}) = \{\text{two rational solutions of the type } \text{IV}_+\}$. By Proposition 3.7 (2), we see that if $L = -2(k+1), -2k, \dots, -2$, then $r(P_{-(k+1),L}) = \{\text{two rational solutions of the type } \text{III}_+, \text{ which satisfy } y(0) = 0\}$. Therefore, by Proposition 3.9 (2) III_+ (i), we find that if $(\theta_0, \theta_\infty) \in \Pi_{-(k+1)} - \{P_{-(k+1),L} \mid L = -2(k+1), -2k, \dots, 2k\}$, $r(\theta_0, \theta_\infty) = \{\text{two rational solutions of the type } \text{III}_+, \text{ which satisfy } y(0) \neq 0\}$ holds.

(iii) From (i) and (ii), we obtain the desired results (F1), (F2), (F3) for any $k (\geq 2)$.

Step 4. We have proved (1) and (2) of the present proposition in the case that $(\theta_0, \theta_\infty) \in \Sigma_-$. By Proposition 3.8, we can also obtain the same result in the case that $(\theta_0, \theta_\infty) \in \Sigma_+$. Hence we have finished the proof. \square

3.3. Rational solutions of $P_{III}(\theta_0, \theta_\infty)$ and proof of Theorem 2

Using the results in previous subsections, we give a proof of Theorem 2. We

prove the following proposition first, and then prove Theorem 2.

PROPOSITION 3.11. (1) $P_{III}(\theta_0, \theta_\infty)$ does not have algebraic solutions.

(2) $P_{III}(\theta_0, \theta_\infty)$ has rational solutions if and only if there exists an integer I such that $\theta_\infty - \theta_0 - 1 = 2I$ or $\theta_\infty + \theta_0 + 1 = 2I$, where $(\theta_0, \theta_\infty) \in \mathbb{C}^2$.

(3) If $P_{III}(\theta_0, \theta_\infty)$ has rational solutions, then the number of rational solutions is two or four. $P_{III}(\theta_0, \theta_\infty)$ has four rational solutions if and only if there exist two integers I and J such that $\theta_\infty - \theta_0 - 1 = 2I$ and $\theta_\infty + \theta_0 + 1 = 2J$.

(4) Rational solutions are classified as in the table IV. Here the sets D_\pm, Δ_\pm and the numbers α_\pm are the same ones as in Theorem 2, and double signs correspond with each other. In the types III_\pm, IV_\pm , the values of N, ε_j, b_j ($j = 1, \dots, N$) depend on the values of θ_0, θ_∞ and α_\pm .

Type	Conditions of $(\theta_0, \theta_\infty)$	Forms of rational solutions
I_\pm	$\theta_\infty \mp \theta_0 \mp 1 = (\theta_0, \theta_\infty) \in \mathbb{C}^2$	$y = \alpha_\pm, y = -\alpha_\pm$
II_\pm	$(\theta_0, \theta_\infty) \in D_\pm$	$y = \alpha_\pm + \theta_\infty/x, y = -\alpha_\pm + \theta_\infty/x$
III_\pm	$\theta_\infty \mp \theta_0 \mp 1 = 2I, (\theta_0, \theta_\infty) \in \mathbb{C}^2$ $I \in \mathbb{Z} - \{0\}, (\theta_0, \theta_\infty) \notin D_\pm \cup \Delta_\pm$	$y = \alpha_\pm + \frac{1}{2} \sum_{j=1}^N \frac{\varepsilon_j}{x - b_j}$ $y = -\alpha_\pm + \frac{1}{2} \sum_{j=1}^N \frac{\varepsilon_j}{x + b_j}$ $\left(N, \varepsilon_j, b_j, I \text{ satisfy} \right.$ $\left. \begin{array}{l} \text{the same conditions} \\ \text{as in Theorem 2, } III_\pm(*) \end{array} \right) \dots (**)$
IV_\pm	$(\theta_0, \theta_\infty) \in \Delta_\pm$	$y = \alpha_\pm + \frac{\theta_\infty}{x} + \frac{1}{2} \sum_{j=1}^N \frac{\varepsilon_j}{x - b_j}$ $y = -\alpha_\pm + \frac{\theta_\infty}{x} + \frac{1}{2} \sum_{j=1}^N \frac{\varepsilon_j}{x + b_j}$ under the same conditions as in (**)

Table IV. Rational solutions of $P_{III}(\theta_0, \theta_\infty)$.

Proof. (1) If $y(x)$ is a k -sheeted ($k \geq 1$) algebraic solution of $P_{III}(\theta_0, \theta_\infty)$, then $q(t) = \sqrt{t} y(\sqrt{t})$ is an algebraic solution of $P_{III'}(\theta_0, \theta_\infty)$. Since $q(x^2)/x = \sqrt{x^2} y(\sqrt{x^2})/x = y(x)$ is a rational solution by Proposition 3.2, it turns out that $k = 1$. Therefore, $P_{III}(\theta_0, \theta_\infty)$ does not have a many-valued algebraic solution.

(2) (3) (4) From Propositions 3.5, 3.6 and 3.10, we obtain the results for rational

solutions of the type J_+ ($J_+ = I_+, \dots, IV_+$). Next, from these results and Proposition 3.4, we obtain the results for rational solutions of the type J_- ($J_- = I_-, \dots, IV_-$). By Proposition 3.3, we thus exhausted all rational solutions of $P_{III}(\theta_0, \theta_\infty)$. \square

Remark 3.3. Gromak's result [5, Theorem 9] corresponds to (2) in this proposition.

Proof of Theorem 2. From Propositions 3.2 and Proposition 3.11, we obtain the desired results. Here, by the following calculation, we can check that two rational solutions of the type J_\pm ($J = I, \dots, IV$) give the same algebraic solution of $P_{III'}(\theta_0, \theta_\infty)$. For example, we assume $J_+ = IV_+$, and take

$$\begin{aligned} y_1(x) &= 1 + \theta_\infty/x + (1/2) \sum_{j=1}^N \varepsilon_j/(x - b_j) \\ y_2(x) &= -1 + \theta_\infty/x + (1/2) \sum_{j=1}^N \varepsilon_j/(x + b_j). \end{aligned}$$

Since

$$\begin{aligned} q_1(t) &= \sqrt{t}y_1(\sqrt{t}) = \sqrt{t}\{1 + \theta_\infty/\sqrt{t} + (1/2) \sum_{j=1}^N \varepsilon_j/(\sqrt{t} - b_j)\} \\ q_2(t) &= \sqrt{t}y_2(\sqrt{t}) = \sqrt{t}\{-1 + \theta_\infty/\sqrt{t} + (1/2) \sum_{j=1}^N \varepsilon_j/(\sqrt{t} + b_j)\}, \end{aligned}$$

if we analytically continue $q_1(t)$ around $t = 0$,

$$q_1(t) = (-\sqrt{t})\{1 + \theta_\infty/(-\sqrt{t}) + (1/2) \sum_{j=1}^N \varepsilon_j/(-\sqrt{t} - b_j)\} = q_2(t). \quad \square$$

4. Proof of Theorem 3

In this section, we suppose that parameters θ_0 and θ_∞ of $P_{III'}(\theta_0, \theta_\infty)$ always satisfy the condition $\theta_\infty + \theta_0 = 2I$ or $\theta_\infty - \theta_0 = 2J$, where I and J are any integers. To clarify the condition which parameters θ_0 and θ_∞ fulfill, we will often use notations $P_{III'}(\theta_\infty + \theta_0 = 2I)$, $P_{III'}(\theta_\infty - \theta_0 = 2J)$ (or $P_{III'}(\theta_\infty + \theta_0 + 1 = 2I + 1)$, $P_{III'}(\theta_\infty - \theta_0 - 1 = 2J - 1)$). Let us consider the following four Riccati equations:

$$\begin{aligned} R_1 : dq/dt &= (1/t)(-q^2 - \theta_0 q + t) \\ R_{-1} : dq/dt &= (1/t)(q^2 - \theta_\infty q - t) \\ \bar{R}_1 : dq/dt &= (1/t)(-q^2 + \theta_\infty q - t) \\ \bar{R}_{-1} : dq/dt &= (1/t)(q^2 - \theta_0 q + t). \end{aligned}$$

R_1 coincides with the equation (1.9) and is contained in $P_{III'}(\theta_\infty + \theta_0 + 1 = 1)$. R_{-1} , \bar{R}_1 and \bar{R}_{-1} are contained in $P_{III'}(\theta_\infty + \theta_0 + 1 = -1)$, $P_{III'}(\theta_\infty - \theta_0 - 1 = 1)$

and $P_{III'}(\theta_\infty - \theta_0 - 1 = -1)$ respectively. We express the Riccati equation R_1 contained in $P_{III'}(\theta_\infty + \theta_0 + 1 = 1)$ by $R_1(\theta_\infty + \theta_0 + 1 = 1)$. Similarly we use notations $R_{-1}(\theta_\infty + \theta_0 + 1 = -1)$, $\bar{R}_1(\theta_\infty - \theta_0 - 1 = 1)$ and $\bar{R}_{-1}(\theta_\infty - \theta_0 - 1 = -1)$. If we apply the transformation h to solutions of R_1 , then we get solutions of R_{-1} . Conversely, if we apply h to solutions of R_{-1} , then we get solutions of R_1 . We express this relation by the symbol

$$(4.1) \quad R_1 \xleftrightarrow{h} R_{-1}.$$

In the same meanings, we have the following relations:

$$(4.2) \quad \begin{aligned} \bar{R}_1 &\xleftrightarrow{h} \bar{R}_{-1} \\ R_1 &\xleftrightarrow{s_2} \bar{R}_{-1}, \quad R_{-1} \xleftrightarrow{s_2} \bar{R}_1 \end{aligned}$$

PROPOSITION 4.1. Transformation $l^{\pm 1}$ (resp. $m^{\pm 1}$) is applicable to a solution $q(t)$ of $P_{III'}(\theta_0, \theta_\infty)$ if and only if $q(t)$ is not a solution of R_{\mp} (resp. $\bar{R}_{\pm 1}$). Here double signs correspond with each other, and R_{+1}, \bar{R}_{+1} denote R_1, \bar{R}_1 respectively.

Proof. We prove only the case of l . Let $q(t)$ be a solution of $P_{III'}(\theta_0, \theta_\infty)$. We can not apply l to $q(t)$ if and only if $q(t)$ satisfies the equation

$$(4.3) \quad dq/dt = (1/t)(q^2 - Aq - t),$$

where $A = -(\theta_0 + 2)$ or $A = \theta_\infty$.

Suppose $q(t)$ satisfies (4.3). Differentiating (4.3) by t , we obtain

$$(4.4) \quad \frac{d^2q}{dt^2} = \frac{2q^3 - (3A + 1)q^2 + (-2t + A^2 + A)q + At}{t^2}.$$

Substituting (4.3) and (4.4) into

$$t^2 q \frac{d^2q}{dt^2} = \left(t \frac{dq}{dt}\right)^2 - tq \frac{dq}{dt} + q^3(q - \theta_\infty) + (\theta_0 + 1)tq - t^2,$$

we obtain

$$(4.5) \quad (\theta_\infty - A)q^2 - (\theta_0 + 2 + A)t = 0.$$

Whichever A may be, (4.5) induces $\theta_\infty + \theta_0 = -2$, and (4.3) coincides with $R_{-1}(\theta_\infty + \theta_0 + 1 = -1)$. Conversely, if $q(t)$ is a solution of $R_{-1}(\theta_\infty + \theta_0 + 1 = -1)$, then $q(t)$ satisfies (4.3). □

We put

$$\begin{aligned}\Pi_{\pm 1} &= \{(\theta_0, \theta_\infty) \in \mathbf{C}^2 \mid \theta_\infty + \theta_0 + 1 = \pm 1\} \\ \bar{\Pi}_{\pm 1} &= \{(\theta_0, \theta_\infty) \in \mathbf{C}^2 \mid \theta_\infty - \theta_0 - 1 = \pm 1\}.\end{aligned}$$

Apparently, we have

$$\begin{aligned}\Pi_1 \cap \bar{\Pi}_1 &= \{(-1, 1)\} & \Pi_1 \cap \bar{\Pi}_{-1} &= \{(0, 0)\}, \\ \Pi_{-1} \cap \bar{\Pi}_1 &= \{(-2, 0)\}, & \Pi_{-1} \cap \bar{\Pi}_{-1} &= \{(-1, -1)\}.\end{aligned}$$

In connection with these facts, we have the following proposition.

PROPOSITION 4.2. $P_{\text{III}'}(-1, 1)$ contains two Riccati equations R_1, \bar{R}_1 and these equations do not have common solutions. Similar results hold for $P_{\text{III}'}(0, 0)$ and $(R_1, \bar{R}_{-1}), P_{\text{III}'}(-2, 0)$ and $(R_{-1}, \bar{R}_1), P_{\text{III}'}(-1, -1)$ and (R_{-1}, \bar{R}_1) respectively.

We omit the proof. By simple calculations, we obtain the following result.

PROPOSITION 4.3. (1) If $q(t)$ is a solution of $R_{\pm 1}(\theta_\infty + \theta_0 + 1 = \pm 1)$, then $m(q(t))$ (resp. $m^{-1}(q(t))$) is a solution of $R_{\pm 1}((\theta_\infty - 1) + (\theta_0 + 1) + 1 = \pm 1)$ (resp. $R_{\pm 1}((\theta_\infty + 1) + (\theta_0 - 1) + 1 = \pm 1)$).
(2) If $q(t)$ is a solution of $\bar{R}_{\pm 1}(\theta_\infty - \theta_0 - 1 = \pm 1)$, then $l(q(t))$ (resp. $l^{-1}(q(t))$) is a solution of $\bar{R}_{\pm 1}((\theta_\infty + 1) - (\theta_0 + 1) - 1 = \pm 1)$ (resp. $\bar{R}_{\pm 1}((\theta_\infty - 1) - (\theta_0 - 1) - 1 = \pm 1)$).

Using the above results, we can prove Theorem 3.

Proof of Theorem 3. Let I be any integer, t be the independent variable of $P_{\text{III}'}(\theta_\infty + \theta_0 = 2I)$. Let $\phi_\sigma(t)$ (resp. $\psi_\sigma(t)$) ($\sigma \in \mathbf{C}$) be a general solution of $R_1(\theta_\infty + \theta_0 = 0)$ (resp. $R_{-1}(\theta_\infty + \theta_0 = -2)$). By Proposition 4.1, $l^I(\phi_\sigma(t))$ ($\sigma \in \mathbf{C}$) is a one-parameter family of solutions of $P_{\text{III}'}(\bar{\theta}_\infty + \bar{\theta}_0 = 2I)$, where $I \geq 0$, $\bar{\theta}_0 = \theta_0 + I$, $\bar{\theta}_\infty = \theta_\infty + I$. When $I = 1$, $l(\phi_\sigma(t))$ and $(d/dt)\{l(\phi_\sigma(t))\}$ are rational functions of $\bar{\theta}_0, t$ and $\phi_\sigma(t)$ with integer coefficients. By induction, we find that $l^I(\phi_\sigma(t))$ is a rational function of $\bar{\theta}_0, t$ and $\phi_\sigma(t)$ with integer coefficients. By similar arguments, we see that $P_{\text{III}'}(\bar{\theta}_\infty + \bar{\theta}_0 = -2(I + 1))$ has a one-parameter family of solutions $(l^{-1})^I(\psi_\sigma(t))$, which is a rational function of $\bar{\theta}_0, t$ and $\psi_\sigma(t)$ with integer coefficients, where $I \geq 0$, $\bar{\theta}_0 = \theta_0 - I$, $\bar{\theta}_\infty = \theta_\infty - I$. Here let us recall the correspondence of R_1 and R_{-1} by the transformation h (See (4.1)). By this correspondence, we see that $P_{\text{III}'}(\bar{\theta}_0 + \bar{\theta}_\infty = 2I)$ has a one-parameter family of solutions of the form

$$(4.6) \quad q(t) = \mathcal{Q}_I(\bar{\theta}_0, t, \phi_\sigma(t)),$$

where \mathcal{Q}_I is a rational function of three variables with integer coefficients. The form of \mathcal{Q}_I depends on the value of I . From (4.2) and the same arguments as in the above, we find that $P_{III'}(\bar{\theta}_\infty - \bar{\theta}_0 = -2J)$ has a one-parameter family of solution of the form

$$(4.7) \quad \bar{q}(t_1) = \bar{\mathcal{Q}}_{-J}(\bar{\theta}_0, t_1, \phi_\sigma(-t_1)),$$

where J is any integer, $t_1 = -t$ is the independent variable of $P_{III'}(\bar{\theta}_\infty - \bar{\theta}_0 = -2J)$, $\bar{\mathcal{Q}}_{-J}$ has the same property as \mathcal{Q}_I in (4.6).

Next, we suppose that $(\bar{\theta}_0, \bar{\theta}_\infty)$ satisfy the conditions $\bar{\theta}_\infty + \bar{\theta}_0 = 2I$ and $\bar{\theta}_\infty - \bar{\theta}_0 = -2J$, where I, J are nonnegative integers. We use a variable t as the independent variable of $P_{III'}(\bar{\theta}_0, \bar{\theta}_\infty)$. Then, $P_{III'}(\bar{\theta}_0, \bar{\theta}_\infty)$ has a one-parameter family of solutions $q(t)$ of the form (4.6) and a one-parameter family of solutions $\bar{q}(t)$ of the form (4.7). Noting Proposition 4.3 and the fact that transformations l and m do not change the independent variable of $P_{III'}(\theta_0, \theta_\infty)$, and that $lm = ml$, we see that

$$\begin{aligned} q(t) &= l^I(\phi_\sigma(t; \theta_\infty + \theta_0 = 0)) = l^I(m^J(\phi_\sigma(t; 0 + 0 = 0))) \\ &= m^J(l^I(\phi_\sigma(t; 0 + 0 = 0))), \\ \bar{q}(t) &= m^J(\bar{\phi}_\sigma(t; \theta_\infty - \theta_0 = 0)) = m^J(l^I(\bar{\phi}_\sigma(t; 0 - 0 = 0))), \end{aligned}$$

where $\phi_\sigma(t; \theta_\infty + \theta_0 = 0)$, $\bar{\phi}_\sigma(t; \theta_\infty - \theta_0 = 0)$ are general solutions of the Riccati equations $R_1(\theta_\infty + \theta_0 = 0)$, $\bar{R}_{-1}(\theta_\infty - \theta_0 = 0)$ respectively (Refer to Figure 2). By Proposition 4.2, $R_1(0 + 0 = 0)$, $\bar{R}_{-1}(0 - 0 = 0)$ do not have common solutions. Therefore one-parameter families $q(t)$ and $\bar{q}(t)$ can not have common solutions. When I and J satisfy other conditions, for example, $I \leq -1, J \geq 0$, we can prove in the same way. □




Figure 2. The relation $m^j l^i = l^i m^j$ holds.

REFERENCES

- [1] V. I. Gromak, Solutions of the third Painlevé equation, *Differential Equations*, **9** (1973), 1599–1600.
- [2] —, Theory of Painlevé's equation, *Differential Equations*, **11** (1975), 285–287.
- [3] —, One-parameter systems of solutions of Painlevé's equations, *Differential Equations*, **14** (1978), 1510–1513.
- [4] V. I. Gromak and N. A. Lukashevich, Special classes of solutions of Painlevé's equations, *Differential Equations*, **18** (1982), 317–326.
- [5] V. I. Gromak, Reducibility of Painlevé equations, *Differential Equations*, **20** (1984), 1191–1198.
- [6] N. A. Lukashevich, On the theory of the third Painlevé equation, *Differential Equations*, **3** (1967), 994–999.

- [7] Y. Murata, Rational solutions of the second and the fourth Painlevé equations, *Funkcial. Ekvac.*, **28** (1985), 1–32.
- [8] —, On the irreducibility of the third Painlevé equation, in preparation.
- [9] K. Nishioka, A note on the transcendency of Painlevé's first transcendent, *Nagoya Math. J.*, **109** (1988), 63–67.
- [10] M. Noumi, private communication.
- [11] K. Okamoto, Polynomial Hamiltonians associated with Painlevé equations. I, *Proc. Japan Acad. Ser. A Math. Sci.*, **56** (1980), 264–268; II, *ibid.*, 367–371.
- [12] —, Isomonodromic deformation and Painlevé equations, and the Garnier system, *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, **33** (1986), 575–618.
- [13] —, Studies on the Painlevé equations IV. Third Painlevé equation P_{III} , *Funkcial. Ekvac.*, **30** (1987), 305–332.
- [14] —, private communication.
- [15] P. Painlevé, *Leçons de Stockholm*, in “Œuvres de P. Painlevé I”, Editions du C. N. R. S., Paris, 1972, 199–818.
- [16] H. Umemura, Birational automorphism groups and differential equations, *Nagoya Math. J.*, **119** (1990), 1–80.
- [17] —, On the irreducibility of the first differential equation of Painlevé, in “Algebraic geometry and commutative algebra in honor of Masayoshi Nagata”, Kinokuniya, Tokyo, 1987, 771–789.

Faculty of Economics
Nagasaki University
4-2-1, Katafuchi, Nagasaki-shi 850
Japan