

Diet quality in young adults and its association with food-related behaviours

Maree G Thorpe, Mark Kestin, Lynn J Riddell, Russell SJ Keast and Sarah A McNaughton*

School of Exercise and Nutrition Sciences, Deakin University, 221 Burwood Highway, Burwood, Victoria 3125, Australia

Submitted 31 July 2012: Final revision received 7 April 2013: Accepted 8 May 2013: First published online 18 July 2013

Abstract

Objective: To determine the diet quality of a group of young adults and explore its associations with two food-related behaviours (involvement in meal preparation and consumption of commercially prepared meals).

Design: Cross-sectional study of young adults. Sample characteristics, food-related behaviours and dietary intake were assessed using a self-administered questionnaire including an FFQ. Diet quality was measured using the fifteen-item Dietary Guideline Index (DGI) designed to assess adherence to Australian dietary guidelines. One-way ANOVA, *t* tests and multiple linear regression analyses were used to explore the relationships between DGI scores, sample characteristics and food-related behaviours.

Setting: University students enrolled in an undergraduate nutrition class, Melbourne, Australia.

Subjects: Students (*n* 309) aged 18–36 years.

Results: The DGI score was normally distributed, with a mean score of 93·4 (SD 17·1) points (range 51·9–127·4 points), out of a possible score of 150 points. In multivariate analyses adjusted for age, sex, nationality, BMI and maternal education, cooking meals for oneself was positively associated with DGI score ($\beta = 0\cdot15$; 95% CI 1·15, 10·03; $P = 0\cdot01$); frequency of takeaway and frequency of convenience meal consumption were inversely associated with DGI score ($\beta = -0\cdot21$; 95% CI -9·96, -2·32; $P = 0\cdot002$ and $\beta = -0\cdot16$; 95% CI -7·40, -0·97; $P < 0\cdot01$, respectively).

Conclusions: Cooking meals for oneself was linked to higher diet quality among young adults, while consumption of commercially prepared meals was associated with poorer diet quality. Maintaining education programmes that promote cooking skills within young adults has the potential to improve DGI scores.

Keywords
Diet quality
Young adults
Cooking
Takeaway

It is widely acknowledged that poor diet quality contributes to an increased risk of chronic diseases such as CVD, type 2 diabetes and some cancers^(1,2). In order to improve dietary intakes, it is first necessary to explore the determinants of dietary behaviours and their influence on diet quality. Young adulthood presents as an ideal period to explore dietary habits as it is an important period of transition from adolescence to adulthood where independent influences on food behaviours and greater responsibility for food choices become established⁽³⁾.

Promoting good nutrition and health is a primary focus of dietary guidelines⁽⁴⁾. Diet quality can be measured by comparing dietary intakes and dietary behaviours with these existing guidelines or recommendations using diet quality indices⁽⁵⁾. These tools utilise a holistic approach, providing a comprehensive alternative measurement to traditional approaches based on individual nutrients^(5,6).

Diet quality indices provide a single numerical value that represents overall diet quality based on current scientific evidence and dietary guidelines. In most cases a higher score indicates better diet quality or better adherence to recommendations.

Although diet quality generally increases with age⁽⁷⁾, a decline is often observed during the transition period from adolescence to adulthood⁽⁸⁾. Alarmingly, some food-related behaviours developed by young adults including irregular meal patterns, such as meal skipping and frequent snacking^(9–11), and frequent consumption of commercially prepared meals, such as takeaway food, pre-packaged or restaurant meals, are often associated with a poorer diet quality^(12–14). An additional concern is that these potentially negative behaviours developed in earlier life are likely to track into later life, having a lasting impact on the long-term health of individuals⁽³⁾.

*Corresponding author: Email sarah.mcnaughton@deakin.edu.au

Given the poor diet quality observed among many young adults⁽¹⁵⁾, it is important to explore the determinants or correlates of diet quality. Food-related lifestyle factors such as consumers' attitudes to food quality, purchasing, cooking methods, ways of shopping and consumption situations and other food-related behaviours have been used to examine the potential determinants of food intake and obesity^(16–18). However, food-related behaviours, such as food purchasing and meal preparation, may be a more proximal determinant of diet quality and have been investigated previously in young adults^(19–21).

Previous studies of adolescents (11–18 years) and young adults (18–23 years) in the USA have suggested that the level of involvement in food purchasing and preparation influences diet quality^(19,20). However, an Australian study of 26–36-year-olds exploring associations between adherence to dietary recommendations and involvement in meal preparation showed little evidence to support this⁽²¹⁾. This may be a reflection of the slightly older age group in the Australian study compared with the US studies. Young adults often perceive their cooking skills as a barrier to preparing meals, along with available time and funds^(19,22). As a result, they might choose to purchase commercially prepared meals^(22–25). In previous research, consumption of commercially prepared meals has been inversely associated with diet quality^(13,26–29). Individuals who consume high levels of commercially prepared meals are more likely to report less favourable dietary patterns, such as lower intakes of fruit, vegetables, dietary fibre and some vitamins and minerals, as well as report higher intakes of energy, fat, meat and confectionery^(13,26–29). However, much of this evidence is based on adult populations. Research that focuses on young adult populations is sparse.

Since the diet quality of young adults is often poor⁽⁸⁾, it is important to explore the determinants of diet quality among this population group in order to inform future health initiatives. Given the limited research in the area, the purpose of the present study was to determine the diet quality of a sample of young adults and to assess the associations of young adults' diet quality with food-related behaviours such as meal preparation and consumption of commercially prepared meals.

Experimental methods

Participants and procedures

The participants were selected from a convenience sample of 369 students enrolled in a first-year university food and nutrition class in 2011. A total of 337 students agreed to take part in the study (91% response rate). Inclusion was restricted to young adults, defined as those aged 18–36 years^(19,21). Participants who were outside the defined age range ($n = 5$) or who had substantial missing responses on the FFQ ($>10\%$ missing responses; $n = 23$)

were excluded, leading to a sample of 309 young adults (84% of the initial sample). Further exclusions were made for the final multiple linear regression analysis due to participants with missing data ($n = 54$). Therefore, a total of 255 (69% of initial sample) participants were included in the multiple linear regression analysis. The socio-demographic characteristics (age, sex, BMI, nationality, maternal education, living arrangements, and study and work commitments) of this sample were not significantly different from those of the total sample. Participants completed a self-assessed food and diet questionnaire that included an FFQ^(30–32). The study was conducted according to ethical guidelines laid down in the Declaration of Helsinki and all procedures were approved by Deakin University Human Research Ethics Committee (ethics number EC2009-163). Written informed consent was obtained from all participants.

Participant characteristics

The self-administered questionnaire was used to obtain participant characteristics (age, sex, BMI (calculated as weight (in kilograms) divided by the square of height (in metres)), nationality, maternal education, living arrangements, and study and work commitments). The questionnaire was similar to one used by Riddell *et al.*⁽³³⁾ based on a questionnaire previously used to investigate the eating habits of young adults^(34,35).

Dietary assessment

Dietary intake was measured by a 107-item FFQ previously used in the 1995 National Nutrition Survey^(30,31) and based on an existing validated FFQ developed for Australian populations⁽³²⁾. The FFQ asked the participants to recall the average number of times each food or beverage was consumed during the previous month. Participants were able to respond using nine categories ranging from 'never or less than once a month' to 'six or more times per day'.

Measurement of diet quality

Diet quality was measured using data from the FFQ and the previously developed Dietary Guideline Index (DGI)⁽¹⁵⁾. The DGI has been shown to reflect intakes of key nutrients and was inversely associated with poor health outcomes in previous research^(15,36), suggesting it is a valid measure of diet quality among Australians. The DGI is a 150-point diet quality index, comprised of fifteen components that reflect the Dietary Guidelines for Australian Adults⁽⁴⁾. Each component contributed 10 points to the total score (Table 1). Criteria for the scores were devised from age- and sex-specific recommendations found in the Australian Guide to Healthy Eating⁽³⁷⁾. The fifteen components include diet variety, intakes of fruit, vegetables, cereal, meat/protein, dairy, alcohol, saturated fat, added sugar, added salt, fluids and 'extra' foods (foods not essential in providing nutrient requirements, such as high-energy and nutrient-poor

Table 1 Components and scoring methods of the Dietary Guideline Index (DGI) score

DGI component	Component scoring criteria*		
	Minimum score (0)	Maximum score (10)	
Diet variety			
Proportion of foods from each core food group† consumed at least once per week	0%	100%	
Fruit intake			
Servings per day	0	≥2	
Vegetable intake			
Servings per day	0	≥5	
Cereal intake			
Servings per day			
	Male	0	≥6
	Female	0	≥4
Wholegrain cereals			
Proportion of wholegrain cereal relative to total cereal	0%	100%	
Protein intake (lean meat and protein alternatives)			
Servings per day	0	≥1	
Lean protein source			
Proportion of lean meats and alternatives relative to total meat and alternatives	0%	100%	
Dairy intake			
Servings per day	0		
Reduced-fat dairy choices			
Type of milk 'usually' consumed	Whole-fat milk	Low-fat milk	
Alcohol intake			
Servings per day			
	Male	≥4	≤2
	Female	≥2	≤1
Saturated fat intake‡			
Type of milk 'usually' consumed	Whole-fat milk	Low-fat milk	
Trim fat from meat	Never/rarely	Usually	
Added sugars			
Intake of high-sugar foods§,			
Servings per day			
	Male	>1.5	<1.5
	Female	>1.25	<1.25
Added salt intake‡			
Salt added in cooking	Never/rarely	Usually	
Salt added the table	Never/rarely	Usually	
Fluid intake‡			
Beverage servings per day	≥8	0	
Proportion of water consumed relative to total beverages	50%	0%	
'Extra' food intake			
Servings per day			
	Male	>3	<3
	Female	>2.5	<2.5
Total DGI score			
Sum of all components	0	150	

*Intakes in between the maximum and minimum intake are assigned a score proportionately.

†Core food groups consist of fruit, vegetables, protein, cereal and dairy.

‡There are two indicators for these components; each is allocated 50% of the component score.

§High-sugar foods include soft drink, cordial, fruit juice drink, jam, chocolate and confectionery.

||'Extra foods' include foods not essential in providing nutrient requirements such as high-sugar foods, chips, hot chips, hamburgers, cakes, muffins, pastry and desserts.

sweets and snacks). The remaining three components reflect healthy choices made by the individual, including usually choosing wholegrain cereal, lean protein and reduced-fat dairy products. Appropriate adjustments in the score calculation were made for those who indicated that they were vegetarian or vegan. For all components, participants in between the criteria for maximum and minimum scores were assigned scores proportionately. The sum of the fifteen components represents the DGI score, where a higher score indicates better adherence to dietary recommendations. Full details of the DGI are described elsewhere⁽¹⁵⁾.

Measurement of food-related behaviours

Involvement in meal preparation was self-reported using two questions: 'Do you do your own food shopping?' and 'Do you cook meals for yourself?'⁽³³⁾. Participants who indicated that they do cook meals for themselves also reported the frequency of cooking using five response categories ranging from 'daily' to 'rarely'. Consumption of commercially prepared meals was measured by asking: 'In an average week, how often would your main meal of the day be...?' (i) 'takeaway'; (ii) 'convenience or pre-packaged' (hereafter called a 'convenience meal'); or (iii) 'from a restaurant, café or pub'. The six response

categories ranged from 'never' to '6–7 meals per week or more'. These questions were based on a similar question on takeaway consumption used in previous studies^(28,38). As few participants reported consumption of commercially prepared meals greater than two times per week, the frequency categories were collapsed into 'never', '<1 meal/week', 'about 1 meal/week' and '≥2 meals/week'. These variables were used as continuous variables (frequencies) for the multiple linear regression analyses.

Statistical analysis

Data analysis was conducted using the statistical software package IBM SPSS Statistics Version 18. The normality of the distribution of DGI score was assessed before further analysis. As it was found to be approximately normal, no transformation was necessary. One-way ANOVA and *t* test analyses were used to examine associations between DGI score and the categorical variables. A multiple linear regression model was developed and used to assess the association between DGI score and the food-related behaviours. The model was adjusted for the potential

confounders age, sex, BMI and maternal education as these are commonly associated with diet quality^(1,7,8). Adjustments were also made for nationality as this was shown to be associated with DGI score within the bivariate analysis of this sample. The low sample size and the further reduction in the sample size for the multivariate analysis was a limitation. However, with *n* 255, 80% power and α set at 0.05, it was estimated that a correlation of 0.16 could be detected, which was only minimally reduced to a correlation of 0.14 with the sample size of 337 that originally agreed to participate. Significance level was set at 5%.

Results

Table 2 shows the distribution of the participants' key characteristics and the mean DGI scores. The majority of participants were aged 18–21 years with a mean age of 20.4 (SD 2.9) years. The mean DGI score was 93.4 (SD 17.1) points, with a range of 51.9–127.4 points. The

Table 2 Mean Dietary Guideline Index (DGI) score according to key characteristics of a sample of university students aged 18–36 years in Melbourne, Australia, 2011

	<i>n</i> *	%	DGI score			<i>P</i> value†
			Mean	SE	Range	
Age (years)						
18–19	159	51.5	94.1	1.37	53.5–123.7	0.93
20–21	91	29.4	92.0	1.76	52.0–124.0	
22–23	25	8.1	93.3	3.49	56.0–123.3	
24–25	13	4.2	93.4	4.21	72.7–118.1	
26–36	21	6.8	93.7	4.18	51.9–127.4	
Sex						
Male	58	19.5	93.1	2.45	52.0–123.3	0.51‡
Female	239	80.5	94.8	1.02	54.8–127.4	
BMI						
Underweight (BMI <18.5 kg/m ²)	19	6.5	87.1	4.90	56.9–127.4	0.25
Normal weight (BMI ≥ 18.5 to <25.0 kg/m ²)	226	77.7	93.4	1.16	52.0–124.0	
Overweight/obese (BMI ≥ 25.0 kg/m ²)	46	15.8	94.8	2.18	63.4–120.4	
Nationality						
Australian	250	84.2	94.2	1.07	51.9–124.0	0.02
Chinese	15	5.1	82.5	3.93	62.7–120.8	
Other Asian	14	4.7	94.9	4.81	57.7–121.9	
Other	18	6.0	86.6	4.12	56.5–127.4	
Mother's highest education						
High school or less	111	37.9	91.1	18.3	53.5–124.0	0.26
Trade or certificate	68	23.2	94.4	15.8	51.9–123.7	
Tertiary	114	38.9	94.5	16.7	52.0–127.4	
Living arrangements						
Parents	177	57.5	93.7	17.8	52.0–124.0	0.67
Flatmates	92	29.9	92.2	16.5	56.5–120.8	
Partner or spouse	26	8.4	92.4	13.2	62.7–117.5	
On own	13	4.2	98.1	20.2	72.6–127.4	
Study and work commitments						
Study part time & work full time	8	2.6	89.7	22.2	56.0–117.5	0.71
Study part time & work part time	15	4.9	97.5	16.7	70.9–118.4	
Study full time & work part time	214	69.7	93.5	16.7	53.5–124.0	
Study full time with no work	70	22.8	92.6	18.1	51.9–127.4	

*Total *n* varies between measures due to missing responses.

†One-way ANOVA, except as indicated.

‡*t* Test.

Table 3 Mean Dietary Guideline Index (DGI) score according to food-related behaviours of a sample of university students aged 18–36 years in Melbourne, Australia, 2011

	n*	%	DGI score			P value†
			Mean	SE	Range	
Food shopping						
Do own food shopping	156	50.6	93.3	1.43	51.9–127.4	0.98‡
Don't do own food shopping	152	49.4	93.3	1.33	52.0–123.7	
Meal cooking						
Cook own meals	220	71.4	94.7	1.13	51.9–127.4	0.04‡
Don't cook own meals	88	28.6	90.2	1.89	53.5–121.1	
Frequency of meal cooking						
Daily	106	48.6	95.2	1.59	51.8–127.4	0.86
3 times/week	70	32.1	94.9	2.05	54.8–122.9	
1 time/week	28	12.8	94.5	2.85	56.5–188.8	
Rarely (once per month/special occasion)	14	6.4	91.1	5.81	52.0–119.6	
Frequency of takeaway as a main meal						
Never	40	13.0	102.2	2.68	51.9–127.4	<0.001
<1 meal/week	124	40.3	95.2	1.51	52.0–123.7	
About 1 meal/week	105	34.1	89.8	1.61	54.8–124.0	
≥2 meals/week§	37	12.6	87.6	2.56	56.5–123.3	
Frequency of a convenience meal as a main meal						
Never	101	32.7	96.4	1.66	51.9–127.4	0.04
<1 meal/week	115	37.2	93.7	1.57	52.0–124.0	
About 1 meal/week	49	15.9	91.0	2.55	59.0–122.9	
≥2 meals/week§	44	14.2	88.1	2.52	54.8–117.5	
Frequency of eating at a restaurant, café or pub for a main meal						
Never	20	6.5	103.9	3.29	70.4–127.4	0.04
<1 meal/week	136	44.2	92.8	1.53	52.0–121.1	
About 1 meal/week	114	37.0	92.4	1.52	51.9–124.0	
≥2 meals/week§	38	12.3	92.3	2.79	56.0–123.3	

*Total *n* varies between behaviour measures due to missing responses.

†One-way ANOVA, except as indicated.

‡*t* Test.

§Variable categories were collapsed because very few participants reported 4–5 meals/week or 6–7 meals/week (<2% each).

majority of participants were female, had a BMI within the normal weight range, identified themselves as Australian, had a mother with a tertiary education, were living with their parents and were studying full time. As there was evidence to suggest that nationality was associated with DGI score, Tukey's Honestly Significant Difference *post hoc* test was performed (not shown in table) and it indicated that participants of Australian nationality had a significantly higher diet quality than those who were of Chinese nationality ($P=0.02$). There was little evidence to suggest that age, sex, BMI, maternal education, living arrangements, or study and work commitments were associated with DGI score ($P>0.05$).

Approximately half of the sample did their own food shopping (50.6%), while the majority (71.4%) of participants indicated that they cook meals for themselves (Table 3). There was no evidence ($P=0.98$) to indicate a difference in diet quality between students who participated in shopping for food and those who did not. Cooking meals for oneself was associated with a higher mean DGI score compared with those who did not cook for themselves (mean 94.7 (SD 1.13) *v.* 90.2 (SD 1.89) points; $P=0.04$). While 71.4% of young adults reported cooking meals, only 48.6% of these cooked daily, 32.1% cooked three times weekly and 19.2% cooked once weekly or less. No evidence was found to suggest an

association between diet quality and cooking frequency ($P=0.86$). There was evidence to support associations between DGI score and the frequency of takeaway ($P<0.001$), convenience meals ($P=0.04$) and restaurant meals ($P=0.04$; Table 3).

Those food-related behaviours significantly associated with DGI score in the preliminary bivariate analyses were included in the multivariate linear regression model (involvement in cooking of meals, consumption frequency of takeaway, consumption frequency of convenience meals and consumption frequency of restaurant meals; Table 4). In the multivariate model adjusted for age, sex, BMI, nationality and maternal education, cooking meals for oneself was positively associated with DGI score ($\beta=0.15$; 95% CI 1.15, 10.03; $P=0.01$), while the frequency of takeaway consumption ($\beta=-0.21$; 95% CI -9.96, -2.32; $P=0.002$) and the frequency of convenience meal consumption ($\beta=-0.16$; 95% CI -7.40, -0.97; $P=0.01$) were both inversely associated with DGI score. The frequency of restaurant meal consumption was not associated with DGI score in the model ($P=0.89$; Table 4). We also examined the relationships after excluding all nationalities other than Australian, since we did not have a sufficient number of participants to stratify the analyses for nationality, and the results were similar (results not shown).

Table 4 Relationship of food-related behaviours to the Dietary Guideline Index (DGI) score of a sample of university students aged 18–36 years in Melbourne, Australia, 2011: results of bivariate and multivariate linear regression analyses (*n* 255)

	Bivariate				Multivariate*			
	β †	95% CI		<i>P</i> value	β †	95% CI		<i>P</i> value
		Lower	Upper			Lower	Upper	
Cook (yes/no)	0.12	−0.20	8.55	0.06	0.15	1.15	10.03	0.01
Takeaway meal frequency	−0.29	−11.59	−4.90	<0.001	−0.21	−9.96	−2.32	0.002
Convenience meal frequency	−0.22	−8.87	−2.63	<0.001	−0.16	−7.40	−0.97	0.01
Restaurant meal frequency	−0.06	−5.74	1.85	0.31	0.01	−3.84	4.44	0.89

*Model $R^2 = 0.16$ and $P < 0.001$, adjusted for age, sex, nationality, BMI and maternal education.
 † β is the standard deviation change in DGI score per unit change in predictor variable.

Discussion

The present study suggested that participation in food preparation is associated with diet quality in young adults. Involvement in cooking and avoidance of commercially prepared meals were independently associated with higher diet quality. These results are consistent with previous research, which indicates that diet quality is positively associated with involvement in meal preparation^(19–21,25) and negatively associated with the consumption of commercially prepared meals^(13,26–29). However, much of the previous research is based on either adult or adolescent population samples and few studies were within Australian populations. A number of key cross-country variations in food behaviours have been demonstrated^(39,40) which is why it is important to consider the local context when conducting research and developing interventions specific for a population. Literature regarding the association of food-related behaviours and diet quality specifically among young adults is sparse^(19,21,28,33). While diet quality was higher among those who reported never consuming restaurant meals in the bivariate analyses, the association was not supported in the multiple regression analyses after adjustment for confounders.

In line with two US studies, one of young adults aged 18–23 years (mean age 20.4 years)⁽¹⁹⁾ and another of adolescents aged 11–18 years (mean age 14.9 years)⁽²⁰⁾, the current study indicated that involvement in cooking meals is associated with increased diet quality among young adults. Engaging in meal preparation may lead to increased skills and variety in food items consumed⁽²²⁾ which, in turn, positively influences diet quality⁽²⁵⁾. Conversely, those who do not cook their own meals may be more inclined to purchase commercially prepared meals. This behaviour has previously been associated with the consumption of foods with poor nutrient content and consequently linked to lower diet quality^(26–28,41). However, the results from a previous study of Australian young adults (mean age 31.6 years) were not consistent with this⁽²¹⁾. That study did not find the level of involvement in cooking meals to be associated with diet quality⁽²¹⁾. The inconsistencies between studies could reflect the varying age range of the participants in the study samples. It might be that the diet

quality of adolescents and the younger age bracket of young adults is more likely to be influenced by involvement in meal preparation than the diet quality of an older age group. The different measures used to measure involvement in meal preparation as well as the different contextual factors of the samples in these studies may also have contributed to the different results.

In previous research, young adults have identified cooking as a potential barrier to preparing healthy food^(19,22,42). For young adults who do not cook or cannot cook their own meals, their reliance on commercially prepared and processed foods is likely to be greater than for those who do cook^(22,23,25). As previously mentioned, this is concerning because foods and meals purchased from commercial providers are generally associated with a poorer diet quality compared with meals prepared at home^(22,41). In line with our results, a study of young adults in the USA (mean age 19.7 years) suggested that increased ability to cook is associated with increased diet quality⁽²⁵⁾. Limited research is available regarding diet quality and ability to cook as it poses many methodological challenges, such as defining cooking abilities⁽⁴³⁾. Future research might benefit from further exploration of the associations between diet quality and cooking among young adults.

Within the present study only a small proportion of the participants reported that they regularly consume commercially prepared meals as their main meal of the day. A previous study from the USA found that 27.3% of young adults (mean age 20.5 years) had takeaway three or more times in the week prior to the survey⁽⁴⁴⁾. Similarly, in an Australian study 37.9% of male and 17.7% of female young adults (mean age 31.6 years) reported consuming takeaway at least two times weekly⁽²⁸⁾. In comparison, only 12.6% of the current sample reported that they consumed takeaway meals as a main meal two or more times in an average week. It is surprising that the university students in our sample reported a low frequency of eating commercially prepared meals. A difficulty with interpreting this is that the survey question did not specify whether the main meal in which the commercially prepared meal was consumed was lunch or dinner. The survey also did not measure the frequency of

takeaway food as a snack, which may be important since young adults have a tendency to skip meals and frequently snack^(9–11,45).

Although low frequencies were reported, our results indicate that consumption of commercially prepared meals was associated with lower diet quality. The effect of takeaway consumption on diet quality has previously been reported in Australian adolescent and young adult populations^(12,28) as well as in adolescent and adult populations in other countries^(13,26,27). These studies along with a systematic review⁽⁴⁶⁾ consistently demonstrate that eating commercially prepared meals is associated with poorer diet quality. Analysis of data from a large US sample of adults (n 13 429; age >20 years) indicated that for every one meal purchased or prepared away from home, there was an additional energy intake of 544 kJ and a 2-point decrease in a diet quality index score⁽²⁶⁾. A likely contributor to the negative relationship between commercially prepared food and diet quality is the poorer nutrient content of the food. Commercially prepared food typically contains more total and saturated fat, and less fibre and important micronutrients than meals prepared and consumed at home^(47,48). This is concerning as it is common to find an increase in takeaway consumption in young adults^(11,12), particularly following university enrolment⁽³⁾. Furthermore, the social context of eating commercially prepared meals encourages the consumption of larger portion sizes⁽⁴⁹⁾ and higher energy-dense foods compared with eating meals cooked at home⁽⁵⁰⁾. It may be important to investigate motivators for young adults to choose commercially prepared meals over home cooked meals.

There are limited data available that explore diet quality and its relationship specifically with restaurant, café or pub meals^(29,51). A Japanese study found that the frequency of young women eating out at full-service restaurants, takeaway pizza delivery shops or cafés was associated with lower consumption of fruit, vegetables, rice and fibre, along with increased intakes of energy, meat, fat, bread and confectionery⁽²⁹⁾. Research on US young adults (mean age 25.3 years) has identified that the type of restaurant chosen influences dietary intake patterns⁽⁵¹⁾. Eating food from restaurants that served primarily burgers and fries was associated with negative dietary patterns⁽⁵¹⁾ while eating at restaurants that served sandwiches or full-service restaurants was unrelated to poor dietary patterns⁽⁵¹⁾. Since eating out is increasing in popularity^(47,52), this may be an important area for further investigation.

Previous research has indicated that students living independently or out of their family home may have poorer diet quality^(3,11,53). However existing research is mixed, with one UK study suggesting that living independently does not necessarily lead to a poorer diet quality in young adults⁽⁵⁴⁾. Consistent with the UK findings, in the current study, 58% of the sample lived with their family but living arrangements were not associated with

diet quality. This may reflect the different circumstances of the young adults studied.

There are several limitations in the present study that should be considered. The study's cross-sectional design cannot provide evidence of causality and the evidence would be strengthened by future longitudinal study designs. Although the FFQ has been validated previously^(32,34,35), the retrospective nature of this dietary data collection method may result in recall bias and measurement error. To reduce the cognitive burden and to capture the transition period as young adults begin university, participants were asked to report on their average dietary intake in the previous month, as opposed to the past 12 months, which was in the original questionnaire^(30,31). Furthermore, the low sample size, the relatively narrow age range and the large proportion of females within the sample made it difficult to examine associations between population characteristics and DGI score. In general, it is understood that diet quality increases with age and differs according to sex^(1,7). As a result of the disproportions in these population characteristics, the study was limited in its ability to generalise the findings to the broader young adult population. Furthermore, since all of the study participants were university students undertaking a unit of study in nutrition, they may be more health and nutritionally conscious and therefore are unlikely to represent the behaviours of all young adults⁽³⁴⁾. Consequently, the diet quality of other young adults may be worse than reported herein. There are limited studies of comparable nature to compare the frequency of commercially prepared meal consumption in Australian young adults. Data from a national sample of young Australians aged 26–36 years old indicated that 37.9% of men and 17.7% of women ate takeaway two or more times weekly⁽²⁸⁾, which is more than the 12.6% of the young adults in the current study. Furthermore, many of the participants (38.9%) had mothers who had completed tertiary education. This is relatively high compared with national data for Australians, where approximately 23% of 45–64-year-olds have completed a bachelor degree or higher⁽⁵⁵⁾.

The utilisation of the comprehensive dietary guideline index (DGI) and a detailed FFQ was a significant strength of the current study. As mentioned, both the FFQ^(30–32) and the DGI⁽¹⁵⁾ were designed for use in Australian populations and therefore were able to suitably measure the dietary intake and diet quality of the participants. In addition to the consumption frequency of individual food items, the FFQ used for the study included several questions relating to eating habits, such as the type of milk typically consumed and the frequency of adding salt to a meal. These questions assisted in the calculation of the DGI and its ability to represent the degree of adherence to dietary guidelines by participants. However, comparisons of the DGI with other indices such as the Healthy Eating Index⁽⁵⁶⁾ is difficult since the latter and

most other dietary indices have been developed in the USA. Furthermore, other indices use varying methods of calculation and can have a different range of values compared with the DGI. It is also difficult to define a reference range to indicate 'good' or 'bad' diet quality as the DGI provides a continuous score in which an optimal score is one that is the highest possible. A strength of the present study was the ability to capture data from a unique population of young adults experiencing a life stage transition as they enter tertiary education. This may be an important time in which food-related behaviours are established among young adults.

Conclusion

The present study suggests that participation in cooking meals and avoiding commercially prepared meals may be important correlates of diet quality of young adults. These food-related behaviours warrant further investigation in longitudinal studies and in general young adult populations.

Acknowledgements

Sources of funding: S.A.M. is supported by an Australian Research Council Future Fellowship. *Conflicts of interest:* None of the authors declared a conflict of interest. *Authors' contributions:* M.G.T. assisted with the study design, performed the statistical analysis, interpreted the results, prepared the tables and prepared the manuscript. S.A.M. assisted with the study design, in particular the calculation of the DGI scores, interpretation of the results, preparing the tables and writing and editing the manuscript draft. M.K. conceived the study design, oversaw the project, managed data collection, and assisted with the statistical analysis, interpreting the results and editing the manuscript draft. L.J.R. and R.S.J.K. conceived the study design, oversaw development of the data collection tools, secured ethical approval and assisted with editing the manuscript. All authors read and approved the final manuscript.

References

1. Kant AK (2004) Dietary patterns and health outcomes. *J Am Diet Assoc* **104**, 615–635.
2. World Health Organization (2003) *Diet, Nutrition and the Prevention of Chronic Diseases. Joint FAO/WHO Expert Consultation. WHO Technical Report Series* no. 916. Geneva: WHO; available at http://whqlibdoc.who.int/trs/WHO_TRS_916.pdf
3. Papadaki A, Hondros G, Scott JA *et al.* (2007) Eating habits of university students living at, or away from home in Greece. *Appetite* **49**, 169–176.
4. National Health and Medical Research Council (2003) *Dietary Guidelines for Australians*. Canberra: NHMRC; available at <http://www.nhmrc.gov.au/publications/synopses/dietsyn.htm>
5. Waijers PM, Feskens EJ & Ocke MC (2007) A critical review of predefined diet quality scores. *Br J Nutr* **97**, 219–231.

6. Kant AK (1996) Indexes of overall diet quality: a review. *J Am Diet Assoc* **96**, 785–791.
7. Thiele S, Mensink GB & Beitz R (2004) Determinants of diet quality. *Public Health Nutr* **7**, 29–37.
8. Forshee R & Storey M (2006) Demographics, not beverage consumption, is associated with diet quality. *Int J Food Sci Nutr* **57**, 494–511.
9. Al-Rethaiaa AS, Fahmy AE & Al-Shwaiyat NM (2010) Obesity and eating habits among college students in Saudi Arabia: a cross sectional study. *Nutr J* **9**, 39.
10. Satalic Z, Baric IC & Keser I (2007) Diet quality in Croatian university students: energy, macronutrient and micronutrient intakes according to gender. *Int J Food Sci Nutr* **58**, 398–410.
11. Kremmyda LS, Papadaki A, Hondros G *et al.* (2008) Differentiating between the effect of rapid dietary acculturation and the effect of living away from home for the first time, on the diets of Greek students studying in Glasgow. *Appetite* **50**, 455–463.
12. Burns C, Jackson M, Gibbons C *et al.* (2002) Foods prepared outside the home: association with selected nutrients and body mass index in adult Australians. *Public Health Nutr* **5**, 441–448.
13. French SA, Story M, Neumark-Sztainer D *et al.* (2001) Fast food restaurant use among adolescents: associations with nutrient intake, food choices and behavioral and psychosocial variables. *Int J Obes Relat Metab Disord* **25**, 1823–1833.
14. Nicklas TA, Myers L, Reger C *et al.* (1998) Impact of breakfast consumption on nutritional adequacy of the diets of young adults in Bogalusa, Louisiana: ethnic and gender contrasts. *J Am Diet Assoc* **98**, 1432–1438.
15. McNaughton SA, Ball K, Crawford D *et al.* (2008) An index of diet and eating patterns is a valid measure of diet quality in an Australian population. *J Nutr* **138**, 86–93.
16. Pérez-Cueto FJA, Verbeke W, de Barcellos MD *et al.* (2010) Food-related lifestyles and their association to obesity in five European countries. *Appetite* **54**, 156–162.
17. Brunsø K, Scholderer J & Grunert KG (2004) Testing relationships between values and food-related lifestyle: results from two European countries. *Appetite* **43**, 195–205.
18. Saba A, Cupellaro E & Vassallo M (2013) Which dimensions of food-related lifestyle are likely to be associated with obesity in Italy? *Public Health Nutr* (Epublication ahead of print version).
19. Larson N, Perry C, Story M *et al.* (2006) Food preparation by young adults is associated with better diet quality. *J Am Diet Assoc* **106**, 2001–2007.
20. Larson N, Story M, Eisenberg M *et al.* (2006) Food preparation and purchasing roles among adolescents: associations with sociodemographic characteristics and diet quality. *J Am Diet Assoc* **106**, 211–218.
21. Smith KJ, McNaughton SA, Gall SL *et al.* (2010) Involvement of young Australian adults in meal preparation: cross-sectional associations with sociodemographic factors and diet quality. *J Am Diet Assoc* **110**, 1363–1367.
22. Soliah L, Walter J & Antosh D (2006) Quantifying the impact of food preparation skills among college women. *Coll Student J* **40**, 729–739.
23. Candel MJJM (2001) Consumers' convenience orientation towards meal preparation: conceptualization and measurement. *Appetite* **36**, 15–28.
24. van der Horst K, Brunner TA & Siegrist M (2011) Ready-meal consumption: associations with weight status and cooking skills. *Public Health Nutr* **14**, 239–245.
25. Levy J & Auld G (2004) Cooking classes outperform cooking demonstrations for college sophomores. *J Nutr Educ Behav* **36**, 197–203.
26. Mancino L, Todd J & Lin B-H (2009) Separating what we eat from where: measuring the effect of food away from home on diet quality. *Food Policy* **34**, 557–562.

27. Schroder H, Fito M & Covas MI (2007) Association of fast food consumption with energy intake, diet quality, body mass index and the risk of obesity in a representative Mediterranean population. *Br J Nutr* **98**, 1274–1280.
28. Smith KJ, McNaughton SA, Gall SL *et al.* (2009) Takeaway food consumption and its associations with diet quality and abdominal obesity: a cross-sectional study of young adults. *Int J Behav Nutr Phys Act* **6**, 29.
29. Murakami K, Sasaki S, Takahashi Y *et al.* (2011) Neighborhood restaurant availability and frequency of eating out in relation to dietary intake in young Japanese women. *J Nutr Sci Vitaminol (Tokyo)* **57**, 87–94.
30. McLennan W & Podger A (1999) *National Nutrition Survey: Foods Eaten, Australia, 1995. ABS Catalogue* no. 4804.0. Canberra: Australian Government Publishing Service.
31. McLennan W & Podger A (1998) *National Nutrition Survey Users' Guide, 1995. ABS Catalogue* no. 4801.0. Canberra: Australian Government Publishing Service.
32. Ireland P, Jolley D, Giles G *et al.* (1994) Development of the Melbourne FFQ: a food frequency questionnaire for use in an Australian prospective study involving and ethnically diverse cohort. *Asia Pac J Clin Nutr* **3**, 19–31.
33. Riddell LJ, Ang B, Keast RS *et al.* (2011) Impact of living arrangements and nationality on food habits and nutrient intakes in young adults. *Appetite* **56**, 726–731.
34. Georgiou CC, Betts NM, Hoerr SL *et al.* (1997) Among young adults, college students and graduates practiced more healthful habits and made more healthful food choices than did nonstudents. *J Am Diet Assoc* **97**, 754–759.
35. Soriano JM, Moltó JC & Mañes J (2000) Dietary intake and food pattern among university students. *Nutr Res* **20**, 1249–1258.
36. McNaughton SA, Dunstan DW, Ball K *et al.* (2009) Dietary quality is associated with diabetes and cardio-metabolic risk factors. *J Nutr* **139**, 734–742.
37. Kellett E, Smith A & Schmerlaib Y (1998) *The Australian Guide to Healthy Eating*. Canberra: Commonwealth Department of Health and Family Services.
38. Smith KJ, Blizzard L, McNaughton SA *et al.* (2012) Takeaway food consumption and cardio-metabolic risk factors in young adults. *Eur J Clin Nutr* **66**, 577–584.
39. Ball K, Timperio AF & Crawford DA (2006) Understanding environmental influences on nutrition and physical activity behaviors: where should we look and what should we count? *Int J Behav Nutr Phys Act* **3**, 33.
40. El Ansari W, Stock C & Mikolajczyk R (2012) Relationships between food consumption and living arrangements among university students in four European countries – a cross-sectional study. *Nutr J* **11**, 28.
41. Lin B & Guthrie J (1998) Popularity of dining out presents barrier to dietary improvements. *Food Rev* **21**, 2–10.
42. Marquis M (2005) Exploring convenience orientation as a food motivation for college students living in residence halls. *Int J Consum Stud* **29**, 55–63.
43. Engler-Stringer R (2010) Food, cooking skills, and health: a literature review. *Can J Diet Pract Res* **71**, 141–145.
44. Larson N, Neumark-Sztainer D, Story M *et al.* (2008) Fast food intake: longitudinal trends during the transition to young adulthood and correlates of intake. *J Adolesc Health* **43**, 79–86.
45. Zizza C, Siega-Riz AM & Popkin BM (2001) Significant increase in young adults' snacking between 1977–1978 and 1994–1996 represents a cause for concern! *Prev Med* **32**, 303–310.
46. Lachat C, Nago E, Verstraeten R *et al.* (2012) Eating out of home and its association with dietary intake: a systematic review of the evidence. *Obes Rev* **13**, 329–346.
47. Guthrie JF, Lin BH & Frazao E (2002) Role of food prepared away from home in the American diet, 1977–78 versus 1994–96: changes and consequences. *J Nutr Educ Behav* **34**, 140–150.
48. Todd J, Mancino L & Lin B-H (2010) *The Impact of Food Away from Home on Adult Diet Quality. Economic Research Report* no. ERR-90. Washington, DC: US Department of Agriculture, Economic Research Service.
49. de Castro JM (1997) Socio-cultural determinants of meal size and frequency. *Br J Nutr* **77**, Suppl. 1, S39–S54.
50. Nielsen SJ & Popkin BM (2003) Patterns and trends in food portion sizes, 1977–1998. *JAMA* **289**, 450–453.
51. Larson N, Neumark-Sztainer D, Laska MN *et al.* (2011) Young adults and eating away from home: associations with dietary intake patterns and weight status differ by choice of restaurant. *J Am Diet Assoc* **111**, 1696–1703.
52. Stewart H, Blisard N, Bhuyan S *et al.* (2004) *The Demand for Food Away From Home: Full-Service or Fast Food. Agricultural Economic Report* no. AER-829. Washington, DC: US Department of Agriculture, Economic Research Service.
53. Sharma B, Harker M, Harker D *et al.* (2008) Living independently and the impact on young adult eating behaviour in Germany. *Br Food J* **111**, 436–451.
54. Beasley IJ, Hackett AF & Maxwell SM (2004) The dietary and health behaviour of young people aged 18–25 years living independently or in the family home in Liverpool, UK. *Int J Consum Stud* **28**, 315–316.
55. Australian Bureau of Statistics (2010) 6227.0 – Education and work, Australia, May 2010. <http://www.abs.gov.au/ausstats/abs@.nsf/mf/6227.0> (accessed October 2011).
56. Kennedy ET, Ohls J, Carlson S *et al.* (1995) The Healthy Eating Index: design and applications. *J Am Diet Assoc* **95**, 1103–1108.