

# Differences between Perfect Powers

Michael A. Bennett

*Abstract.* We apply the hypergeometric method of Thue and Siegel to prove that if  $a$  and  $b$  are positive integers, then the inequality  $0 < |a^x - b^y| < \frac{1}{4} \max\{a^{x/2}, b^{y/2}\}$  has at most a single solution in positive integers  $x$  and  $y$ . This essentially sharpens a classic result of LeVeque.

## 1 Introduction

In 1950, LeVeque [10] proved, given fixed positive integers  $a$  and  $b$ , that the Diophantine equation  $a^x - b^y = 1$  has at most a single solution in positive integers  $x$  and  $y$ , unless  $(a, b) = (3, 2)$ , in which case two such solutions accrue. Nowadays, this might be regarded as a very special case of the profound work of Mihailescu [11] on Catalan's conjecture, but, in fairness, one should note that [10] inspired the work of Cassels [4, 5] which, in turn, proved crucial to Mihailescu.

If one considers more general equations of the shape

$$(1.1) \quad a^x - b^y = c$$

where  $c > 1$  is fixed, then no conclusion of even remotely comparable strength to those in [11] is available to us. If, in analogy to LeVeque [10], we assume that  $a$  and  $b$  are fixed, however, then equation (1.1) has at most two solutions in positive integers  $(x, y)$  (see [2] and earlier work of Herschfeld [7] and Pillai [12–15]). Recently, this result has been extended to equations of the shape  $|a^x \pm b^y| = c$  by Scott and Styer [17].

The goal of this paper is a broad generalization of the main theorem of [10], where, instead of a Diophantine equation, we consider a corresponding Diophantine inequality.

**Theorem 1.1** *Let  $a$  and  $b$  be positive integers. Then there exists at most one pair of positive integers  $(x, y)$  for which*

$$(1.2) \quad 0 < |a^x - b^y| < \frac{1}{4} \max\{a^{x/2}, b^{y/2}\}.$$

It should be noted that lower bounds for linear forms in logarithms may be used to show that there are in fact no solutions whatsoever to (1.2), provided  $x \geq x_0(a, b)$  (see Ellison [6]; more recent work of Laurent, Mignotte and Nesterenko [9] may be used to sharpen this result), which leads to an alternative proof of Theorem 1.1, for

---

Received by the editors April 4, 2006.

The author was supported in part by a grant from NSERC

AMS subject classification: Primary: 11D61; secondary: 11D45.

©Canadian Mathematical Society 2008.

sufficiently large  $a$  and  $b$ . Our proof, in contrast, will rely upon the hypergeometric method of Thue–Siegel which, to our knowledge, has not been applied previously in this context.

Theorem 1.1 leads rather easily to a sharpening of the results of [2, 17]; we will not undertake this here.

## 2 Elementary Preliminaries

We will suppose, here and henceforth, that  $a$  and  $b$  are positive integers, and that  $(x_1, y_1)$  and  $(x_2, y_2)$  are two solutions in positive integers to inequality (1.2) with, say,  $x_2 > x_1$ . Without loss of generality, we may assume that neither  $a$  nor  $b$  is a perfect power. Let us write

$$(2.1) \quad a^{x_i} - b^{y_i} = c_i,$$

where, by symmetry, we may assume that  $c_1 > 0$ . For future use, it will prove convenient to note that

$$(2.2) \quad \min\{a^{x_i}, b^{y_i}\} > \frac{15}{16} \max\{a^{x_i}, b^{y_i}\}.$$

To see this, observe that the inequality  $\min\{a^{x_i}, b^{y_i}\} \leq \frac{15}{16} \max\{a^{x_i}, b^{y_i}\}$  implies

$$|c_i| \geq \frac{1}{16} \max\{a^{x_i}, b^{y_i}\},$$

whence from (1.2),  $\frac{1}{16} \max\{a^{x_i}, b^{y_i}\} < \frac{1}{4} \max\{a^{x_i}, b^{y_i}\}^{1/2}$ , and so  $\max\{a^{x_i}, b^{y_i}\} < 16$ , contradicting (1.2) and the fact that  $|c_i| \geq 1$ .

Next, let us show that necessarily  $x_i$  and  $y_i$  are coprime. If we suppose

$$\gcd(x_i, y_i) = d > 1$$

and write  $x_i = x_0 d$ ,  $y_i = y_0 d$ , then, from (2.1) and the fact that  $a^{x_i} \neq b^{y_i}$  (whereby  $|a^{x_0} - b^{y_0}| \geq 1$ ), we have

$$|c_i| \geq d \min\{a^{x_0(d-1)}, b^{y_0(d-1)}\} = d \min\{a^{x_i}, b^{y_i}\}^{(d-1)/d},$$

and so  $d \min\{a^{x_i}, b^{y_i}\}^{(d-1)/d} < \frac{1}{4} \max\{a^{x_i}, b^{y_i}\}^{1/2}$ . Applying inequality (2.2), it follows that  $d \min\{a^{x_i}, b^{y_i}\}^{(d-1)/d} < 1/\sqrt{15} \min\{a^{x_i}, b^{y_i}\}^{1/2}$ , whereby

$$\min\{a^{x_i}, b^{y_i}\}^{\frac{1}{2} - \frac{1}{d}} < \frac{1}{d\sqrt{15}},$$

contradicting  $d \geq 2$ .

### 3 A Gap Principle

As is rather standard when counting solutions to Diophantine equations or inequalities, we will require a result which guarantees that the putative solutions  $(x_1, y_1)$  and  $(x_2, y_2)$  to (1.2) are of very different size. To derive this, we will begin with equation (2.1) which, after dividing by  $b^{y_i}$  becomes  $a^{x_i} b^{-y_i} - 1 = c_i b^{-y_i}$ . Examination of the Maclaurin series for  $e^z$  thus shows that

$$t|x_1 \log a - y_1 \log b| < c_1 b^{-y_1} \quad \text{and} \quad |x_2 \log a - y_2 \log b| < 2|c_2| b^{-y_2}$$

(recall that  $c_1 > 0$ ). Thus

$$(3.1) \quad \left| \frac{\log b}{\log a} - \frac{x_i}{y_i} \right| < \frac{2^{i-1}|c_i|}{y_i b^{y_i} \log a},$$

whereby we may conclude that  $x_i/y_i$  is a convergent in the simple continued fraction expansion to  $\log b/\log a$ , provided, say,

$$(3.2) \quad \frac{b^{y_i} \log a}{|c_i| y_i} > 4 \geq 2^i.$$

Now, from (1.2) and (2.2), we have that

$$\frac{b^{y_i} \log a}{|c_i| y_i} > \frac{\sqrt{15} b^{y_i/2} \log a}{y_i}.$$

If  $a = 2$ , then  $b \geq 3$  and hence  $b^{y_i/2}/y_i \geq 3/2$ , while, if  $a \geq 3$ ,  $b^{y_i/2}/y_i \geq 2\sqrt{2}/3$ . In both cases inequality (3.2) holds.

It follows, therefore, that  $x_i/y_i$  is a convergent in the simple continued fraction expansion to  $\log b/\log a$  for both  $i = 1$  and  $i = 2$ . On the other hand, if  $p_n/q_n$  is the  $n$ -th such convergent, then

$$(3.3) \quad \left| \frac{\log b}{\log a} - \frac{p_n}{q_n} \right| > \frac{1}{(a_{n+1} + 2) q_n^2},$$

where  $a_{n+1}$  is the  $(n + 1)$ -st partial quotient to  $\log b/\log a$  (see [8]). Since

$$\gcd(x_1, y_1) = \gcd(x_2, y_2) = 1,$$

it follows, if  $x_1/y_1 = p_r/q_r$  and  $x_2/y_2 = p_s/q_s$ , that  $x_1 = p_r$ ,  $y_1 = q_r$ ,  $x_2 = p_s$ , and  $y_2 = q_s$ . Combining (3.1) and (3.3) thus yields

$$a_{r+1} > \frac{b^{y_1} \log a}{c_1 y_1} - 2,$$

and, since  $p_s \geq p_{r+1} > a_{r+1} p_r$ ,

$$(3.4) \quad x_2 > \left( \frac{b^{y_1} \log a}{c_1 y_1} - 2 \right) x_1.$$

From (1.2) and (2.2), we thus have that

$$(3.5) \quad x_2 > \left( \frac{\sqrt{15} b^{y_1/2} \log a}{y_1} - 2 \right) x_1.$$

Similarly, we obtain the inequality

$$(3.6) \quad a_{s+1} > \frac{b^{y_2} \log a}{2|c_2|y_2} - 2 > \frac{\sqrt{15} b^{q_s/2} \log a}{2q_s} - 2.$$

### 4 Some Useful Polynomials

Our main tool in proving Theorem 1.1 will be (off-diagonal) Padé approximants to binomial functions of the shape  $(1 - z)^k$ . We will generate these as in [1] (see also [3]). Let  $A, B$  and  $C$  be positive integers and define

$$(4.1) \quad \begin{aligned} P_{A,B,C}(z) &= \frac{(A + B + C + 1)!}{A! B! C!} \int_0^1 u^A (1 - u)^B (z - u)^C du, \\ Q_{A,B,C}(z) &= \frac{(-1)^C (A + B + C + 1)!}{A! B! C!} \int_0^1 u^B (1 - u)^C (1 - u + zu)^A du, \\ E_{A,B,C}(z) &= \frac{(A + B + C + 1)!}{A! B! C!} \int_0^1 u^A (1 - u)^C (1 - zu)^B du. \end{aligned}$$

Arguing as in [1, §2], we find that

$$(4.2) \quad P_{A,B,C}(z) - (1 - z)^{B+C+1} Q_{A,B,C}(z) = z^{A+C+1} E_{A,B,C}(z).$$

It is worth observing that if  $A = C$ , then  $P_{A,B,C}(z)$  and  $Q_{A,B,C}(z)$  correspond to the diagonal Padé approximants to  $(1 - z)^{B+C+1}$  with error term  $E_{A,B,C}(z)$ . The following results are given in [1, 3].

**Lemma 4.1** *The expressions  $P_{A,B,C}(z)$ ,  $Q_{A,B,C}(z)$ , and  $E_{A,B,C}(z)$  satisfy*

$$\begin{aligned} P_{A,B,C}(z) &= \sum_{r=0}^C \binom{A + B + C + 1}{r} \binom{A + C - r}{A} (-z)^r, \\ Q_{A,B,C}(z) &= (-1)^C \sum_{r=0}^A \binom{A + C - r}{C} \binom{B + r}{r} z^r, \\ E_{A,B,C}(z) &= \sum_{r=0}^B \binom{A + r}{r} \binom{A + B + C + 1}{A + C + r + 1} (-z)^r. \end{aligned}$$

**Lemma 4.2** *There is a non-zero integer  $D = D(A, B)$  for which*

$$P_{A,B,A}(z) Q_{A+1,B-1,A+1}(z) - Q_{A,B,A}(z) P_{A+1,B-1,A+1}(z) = Dz^{2A+1}.$$

In summary, Lemma 4.1 implies that  $P_{A,B,C}(z)$ ,  $Q_{A,B,C}(z)$  and  $E_{A,B,C}(z)$  are polynomials in  $z$  with integer coefficients, while Lemma 4.2 ensures that

$$(P_{A,B,A}(z), P_{A+1,B-1,A+1}(z)) \quad \text{and} \quad (Q_{A,B,A}(z), Q_{A+1,B-1,A+1}(z))$$

are pairs of relatively prime polynomials.

### 5 Bounding the Approximants

For our purposes, we will need to find reasonably sharp upper bounds on the approximating polynomials defined in the previous section.

**Lemma 5.1** *If  $n = m - \delta$  for  $\delta \in \{0, 1\}$  and  $0 < z < 1/2$ , then*

$$|P_n(z)| < \frac{4\sqrt{2}}{3\pi} \cdot 4^m \quad \text{and} \quad |E_n(z)| < \frac{4}{3\sqrt{2}\pi} \cdot 16^m.$$

**Proof** We take  $A = C = n = m - \delta$  and  $B = 3m - n - 1 = 2m + \delta - 1$  and begin by noting that a routine application of Stirling’s formula yields the inequality

$$\frac{(4m)!}{(m!)^2(2m)!} < \frac{1}{\sqrt{2\pi m}} \cdot 64^m,$$

valid for all positive integers  $m$ . It follows from (4.1), if we define

$$u_1 = \frac{1}{8} (3z + 2 + \sqrt{4 - 4z + 9z^2}) \quad \text{and} \quad P(z) = u_1 (1 - u_1)^2 (z - u_1),$$

that

$$|P_n(z)| < \frac{\sqrt{2}}{8^\delta \pi} \cdot 64^m |P(z)|^{m-1} \left| \int_0^1 u^{1-\delta} (1-u)^{1+\delta} (z-u)^{1-\delta} du \right|.$$

Via calculus, it is easy to show that  $|P(z)| < 1/16$ , for  $0 < z < 1/2$ . Also

$$\left| \int_0^1 u(1-u)(z-u) du \right| = \frac{1}{12} - \frac{z}{6} < \frac{1}{12} \quad \text{and} \quad \int_0^1 (1-u)^2 du = \frac{1}{3},$$

and hence the bound for  $|P_n(z)|$  follows.

Similarly, if we define

$$u_2 = \frac{1}{8z} (3z + 2 - \sqrt{4 - 4z + 9z^2}) \quad \text{and} \quad E(z) = u_2(1 - u_2)(1 - zu_2)^2,$$

then

$$|E_n(z)| < \frac{\sqrt{2}}{8^\delta \pi} \cdot 64^m |E(z)|^{m-1} \left| \int_0^1 u^{1-\delta} (1-u)^{1-\delta} (1-zu)^{1+\delta} du \right|.$$

Once again, it is easy to show that  $|E(z)| < 1/4$ , for  $0 < z < 1/2$ , and that

$$\left| \int_0^1 u(1-u)(1-zu) du \right| = \frac{1}{6} - \frac{z}{12} < \frac{1}{6}$$

and

$$\int_0^1 (1-zu)^2 du = 1 - z + \frac{z^2}{3} < 1,$$

which leads to the desired result. ■

Lemma 5.1 provides us with archimedean bounds for our approximants. Regarding non-archimedean information, let us define

$$G(n) = \gcd_{r \in \{0,1,\dots,n\}} \left( \binom{2n-r}{n} \binom{3m-n-1+r}{r} \right).$$

If we take  $n = m$  or  $m - 1$ , it follows from [1, Lemma 7] that

$$\lim_{m \rightarrow \infty} \frac{1}{m} \log G(n) = \frac{\pi}{2} - 3 \log 2,$$

and hence there exists a constant  $c$  such that, for  $n = m$  or  $m - 1$ , and  $m \geq 1$ ,

$$G(n) > c \cdot 1.663^m.$$

For our purposes, we will have need of a completely explicit result along these lines; the proof of this follows arguments sketched in [1, p. 200] and relies upon Chebyshev-type estimates for primes in intervals.

**Proposition 5.2** *If  $m$  is a positive integer and  $n = m$  or  $m - 1$ , then*

$$G(n) > 0.00279 \cdot 1.5498^m.$$

We note that we could avoid use of this proposition if we were prepared to treat certain “small” cases of Theorem 1.1 via lower bounds for linear forms in logarithms.

### 6 The Proof of Theorem 1.1

To proceed with the proof of Theorem 1.1, let us begin by writing  $x_2 = 3x_1m + \alpha$  and  $y_2 = 3y_1m' + \beta$ , where  $0 \leq \alpha < 3x_1$  and  $0 \leq \beta < 3y_1$ , so that  $c_2 = a^{3x_1m}M_1 - b^{3y_1m'}M_2$ , with  $M_1 = a^\alpha$  and  $M_2 = b^\beta$ . We claim that  $m' \geq m$ . If not, then

$$a^{x_2} - b^{y_2} \geq a^{3x_1m_2} \cdot a^{3x_1+\alpha} - b^{3y_1m'} \cdot b^\beta > a^{3x_1m'} \cdot a^{3x_1} - b^{3y_1m'} \cdot b^{3y_1}$$

and so

$$a^{x_2} - b^{y_2} > b^{3y_1} \left( (a^{x_1})^{3m'} - (b^{y_1})^{3m'} \right).$$

It follows that either  $m' = 0$  (so that  $0 \leq y_2 < 3y_1$ , contradicting the combination of (2.2) and (3.5)) or that  $3m' \geq 3$ . In the latter case, we have

$$(a^{x_1})^{3m'} - (b^{y_1})^{3m'} > c_1 \cdot 3m' \cdot (b^{y_1})^{3m'-1}$$

whence  $c_2 = a^{x_2} - b^{y_2} > c_1 \cdot 3m' \cdot (b^{y_1})^{3m'+1} > b^{y_2}$ , a contradiction. It follows that we may write

$$(6.1) \quad a^{3x_1m}M_1 - b^{3y_1m}M_3 = c_2,$$

with  $M_3 = b^{\beta+3y_1(m'-m)}$ .

We take  $n = m$  or  $m - 1$ . Here and subsequently, let  $A = C = n, B = 3m - n - 1$  and write, suppressing various dependencies,

$$P_n(z) = P_{n,3m-n-1,n}(z), \quad Q_n(z) = Q_{n,3m-n-1,n}(z), \quad E_n(z) = E_{n,3m-n-1,n}(z).$$

Fixing once and for all  $z = z_0 = c_1/a^{x_1}$  and substituting this into (4.2), we find that

$$(6.2) \quad a^{3x_1m}P - b^{3y_1m}Q = E,$$

where

$$P = \frac{1}{G(n)}a^{x_1n}P_n(z_0), \quad Q = \frac{1}{G(n)}a^{x_1n}Q_n(z_0), \quad E = \frac{1}{G(n)}(a^{x_1})^{3m-n-1}c_1^{2n+1}E_n(z_0).$$

It follows that  $P, Q$  and  $E$  are all integers. Multiplying (6.1) by  $P$  and (6.2) by  $M_1$ , we deduce the inequality  $b^{3y_1m}|M_3P - M_1Q| \leq |c_2||P| + |E|M_1$ . We claim that for at least one of  $n = m$  or  $n = m - 1$ , say  $n = m - \delta$ , we have  $M_3P \neq M_1Q$ . Indeed, if this fails to be the case, then  $M_3P_{m-1}(z_0) = M_1Q_{m-1}(z_0)$  and  $M_3P_m(z_0) = M_1Q_m(z_0)$ , whereby  $P_{m-1}(z_0)Q_m(z_0) = Q_{m-1}(z_0)P_m(z_0)$ , contradicting Lemma 4.2. For this  $n = m - \delta$ , we therefore have

$$(6.3) \quad b^{3y_1m} \leq |c_2||P| + |E|M_1.$$

To proceed, we will show that each of  $|P|$  and  $|E|$  is not too large, whereby we may employ (6.3) to obtain a (typically contradictory) lower bound on  $M_1$ .

Let us begin by showing that

$$(6.4) \quad b^{3y_1m} > 31|c_2||P|.$$

We will first assume  $b^{y_1} \geq 86$ . From (3.4) and (3.5), this enables us to suppose that

$$(6.5) \quad x_2 \geq 43x_1.$$

Applying Lemma 5.1 and the trivial inequality  $G(n) \geq 1$ , we have

$$|P| < a^{x_1(m-\delta)} \frac{4\sqrt{2}}{3\pi} \cdot 4^m$$

and hence

$$\frac{b^{3y_1m}}{|c_2||P|} > \frac{3\pi}{\sqrt{2}} \left( \frac{b^{3y_1}}{4a^{x_1}} \right)^m \max\{a^{x_2}, b^{y_2}\}^{-1/2}.$$

Since

$$m = \frac{x_2 - \alpha}{3x_1},$$

it follows that

$$(6.6) \quad m > \frac{x_2}{3x_1} - 1,$$

and so, together with  $b^{y_1} > \frac{15}{16}a^{x_1}$ , we have

$$\frac{b^{3y_1m}}{|c_2||P|} > \frac{3\pi}{\sqrt{2}} (15^3 2^{-14} a^{2x_1})^{\frac{x_2}{3x_1}-1} \max\{a^{x_2}, b^{y_2}\}^{-1/2},$$

whence

$$\frac{b^{3y_1m}}{|c_2||P|} > \frac{8192}{1125} \pi \sqrt{2} (15^3 2^{-14})^{\frac{x_2}{3x_1}} a^{2x_2/3-2x_1} \max\{a^{x_2}, b^{y_2}\}^{-1/2}.$$

From (2.2) and the fact that  $15^3 2^{-14} > \frac{1}{5}$ , we thus have

$$\frac{b^{3y_1m}}{|c_2||P|} > \frac{2048}{1125} \pi \sqrt{30} \cdot \left( a^{\frac{1}{2} - \frac{6x_1}{x_2}} \right)^{x_2/3}.$$

Inequality (6.5) and the fact that  $b^{y_1} \geq 86$  (whereby  $a^{x_1} \geq 87$ ) thus imply

$$(6.7) \quad a^{\frac{1}{2} - \frac{6x_1}{x_2}} < 5^{1/x_1},$$

and so

$$\frac{b^{3y_1m}}{|c_2||P|} > \frac{2048}{1125} \pi \sqrt{30}$$

which yields (6.4).

To treat the cases where  $b^{y_1} \leq 85$ , we note that inequality (6.7) (and hence (6.4)) follows as before, from (3.4), unless we have either  $16 \leq b^{y_1} \leq 36$  and  $a^{x_1} = b^{y_1} + 1$ , or

$$(a, x_1, b, y_1) = (2, 6, 63, 1), (65, 1, 2, 6), (66, 1, 2, 6), (83, 1, 3, 4).$$

If  $b^{y_1} \geq 25$ , then we have in each case (6.7) and hence (6.4), unless  $x_2 \leq 996$ . For each  $(a, b)$  under consideration, we compute the initial terms in the simple continued fraction expansion to  $\frac{\log a}{\log b}$  and check that in each case convergents  $p_s/q_s$  with  $x_1 < p_s \leq 996$  have corresponding partial quotients  $a_{s+1}$  violating (3.6).

To treat the cases  $16 \leq b^{y_1} \leq 24$ , we argue as previously only with the trivial lower bound upon  $G(n)$  replaced by that of Proposition 5.2. After a little work, we deduce the inequality

$$\frac{b^{3y_1m}}{|c_2||P|} > 0.087 (0.319)^{\frac{x_2}{3x_1}} a^{x_2/6-2x_1}.$$

In every case, this implies (6.4), unless  $x_2 \leq 158$ . Again, examining the simple continued fraction expansions to  $\frac{\log a}{\log b}$  for  $a = b + 1$  and  $17 \leq b \leq 23$ , and  $(a, b) = (17, 2), (5, 24)$ , we find that all convergents  $p_s/q_s$  with  $x_1 < p_s \leq 158$  have corresponding partial quotients  $a_{s+1}$  which contradict (3.6).

From inequalities (6.3) and (6.4), we thus have

$$\frac{30 b^{3y_1m}}{31 |E|} < M_1 = a^\alpha \leq a^{3x_1-1}.$$

Since

$$|E| < \frac{4}{3\sqrt{2}\pi} G(n)^{-1} c_1^{1-2\delta} a^{x_1(\delta-1)} (16c_1^2 a^{2x_1})^m,$$

it follows from Proposition 5.2 that

$$\left( \frac{1.5498 b^{3y_1}}{16c_1^2 a^{2x_1}} \right)^m < 112 a^{(2+\delta)x_1-1} c_1^{1-2\delta}.$$

Now  $c_1 = \frac{1}{4} a^{\theta x_1}$  where  $0 < \theta < 1/2$  and hence we have

$$\left( \frac{1.5498 b^{3y_1}}{a^{(2+2\theta)x_1}} \right)^m < 112 \cdot 2^{4\delta-2} a^{(2+\delta+\theta-2\theta\delta)x_1-1} < 448 a^{(3-\theta)x_1-1}.$$

Again the fact that  $b^{y_1} > \frac{15}{16} a^{x_1}$  yields  $(1.2769 a^{(1-2\theta)x_1})^m < 448 a^{(3-\theta)x_1-1}$  and so, since  $0 < \theta < 1/2$  and  $a^{x_1} < \frac{16}{15} b^{y_1}$ ,

$$(6.8) \quad m < 4.1 \cdot \log(448 a^{3x_1-1}) < 25.9 + 12.3 \log(b^{y_1}) - 4.1 \log a.$$

On the other hand, from (3.5) and (6.6),

$$m > \frac{\sqrt{15} b^{y_1/2} \log a}{3y_1} - \frac{5}{3},$$

whence, with (6.8),

$$(6.9) \quad \frac{b^{y_1/2} \log a}{y_1} < 21.4 + 9.6 \log(b^{y_1}) - 3.1 \log a.$$

This inequality provides an immediate contradiction for suitably large  $b^{y_1}$  (and hence for all but finitely many quadruples  $(a, x_1, b, y_1)$ ). We will treat these exceptions in the next section, completing the proof of Theorem 1.1.

## 7 Computations

Let us first dispense with the possibility that  $\min\{x_1, y_1\} > 1$ . A short computation reveals that there are exactly 122 quadruples  $(a, x_1, b, y_1)$  with  $\min\{x_1, y_1\} \geq 2$  and

$$(7.1) \quad b^{y_1} < a^{x_1} \leq 10^8,$$

satisfying (1.2).

From inequality (6.9), since  $a \geq 2$ , we may check that if  $y_1 = 2$ , then necessarily  $b \leq 385$ , and, more generally

$y_1 = 2$	$b \leq 385$	$y_1 = 7$	$b \leq 7$
$y_1 = 3$	$b \leq 72$	$y_1 = 8$	$b \leq 6$
$y_1 = 4$	$b \leq 29$	$y_1 = 9$	$b \leq 5$
$y_1 = 5$	$b \leq 16$	$10 \leq y_1 \leq 15$	$b \leq 3$
$y_1 = 6$	$b \leq 11$	$16 \leq y_1 \leq 25$	$b = 2$

while, if  $y_1 \geq 26$ , we have  $b < 2$ , a contradiction. From (1.2), the inequalities in (7.1) thus hold and it is therefore easy to check that the only quadruples satisfying the above bounds upon  $y_1$  and  $b$ , together with (1.2), are

$$(a, x_1, b, y_1) = (13, 3, 3, 7), (56, 2, 5, 5), (15, 3, 58, 2), (2, 15, 181, 2), (2, 17, 362, 2).$$

To treat these remaining quadruples, in each case we begin by noting that from (6.8)  $m \leq 167$ . Inequality (6.6) and the fact that  $x_1 \leq 17$  thus imply that  $x_2 \leq 8567$ . For each of our five cases, as in the preceding section, we compute some initial terms in the infinite simple continued fraction expansion to  $\frac{\log b}{\log a}$  via Maple 9.5. Since  $x_2$  and  $y_2$  are coprime,  $x_2$  is the numerator of a convergent in such an expansion, say  $x_2 = p_s$ . In each case, there are fewer than 5 convergents for which  $x_1 < p_s \leq 8567$ ; in no case does  $a_{s+1}$  satisfy (3.6).

We may thus suppose  $\min\{x_1, y_1\} = 1$ . Let us begin by assuming that  $x_1 = 1$ . It follows that  $a > b^{y_1}$  and hence we may replace (6.9) with the simpler

$$b^{y_1/2} \log b < 21.4 + 96.5 \log(b^{y_1}),$$

which implies the inequalities

$y_1 = 1$	$b \leq 120$	$y_1 = 4$	$b \leq 6$
$y_1 = 2$	$b \leq 20$	$5 \leq y_1 \leq 7$	$b \leq 3$
$y_1 = 3$	$b \leq 7$	$8 \leq y_1 \leq 13$	$b = 2$

We consider  $a = b^{y_1} + t$  where, from (1.2),

$$1 \leq t < \frac{\sqrt{1 + 64b^{y_1}} + 1}{32}.$$

Since we omit perfect powers for  $a$  and  $b$ , this leaves us with precisely 306 triples  $(a, b, y_1)$ . Combining (6.8) and (6.6), we thus have that  $x_2 = p_s, y_2 = q_s$  for a convergent  $p_s/q_s$  in the simple continued fraction expansion to  $\frac{\log b}{\log a}$ , satisfying

$$1 < p_s < 77.1 + 24.6 \log a.$$

A simple calculation reveals that none of these convergents have corresponding  $a_{s+1}$  satisfying (3.6).

Finally, let us suppose that  $y_1 = 1$  (and, from the preceding work, that  $x_1 \geq 2$ ). If  $a = 2$ , then from (6.9), we have  $b \leq 28913$  and so, via (1.2),  $x_1 \leq 14$ . Similarly, for larger values of  $a$ , we may conclude as follows:

$a = 2$	$x_1 \leq 14$	$a = 6$	$x_1 \leq 4$
$a = 3$	$x_1 \leq 8$	$a = 7, 10$	$x_1 \leq 3$
$a = 5$	$x_1 \leq 5$	$11 \leq a \leq 22$	$x_1 = 2$

If  $a \geq 23$ , we contradict  $x_1 \geq 2$ . For each pair  $(a, x_1)$ , we consider  $b = a^{x_1} - t$ , where

$$1 \leq t < \frac{1}{4}a^{x_1/2}.$$

Once again, (6.8) and (6.6) imply the existence of a convergent  $p_s/q_s$  in the simple continued fraction expansion to  $\frac{\log b}{\log a}$  with  $x_1 < p_s < 12.3 \cdot \log(a^{3x_1-1}) + 3x_1 - 1$  and, via (3.6), corresponding partial quotient  $a_{s+1}$  satisfying

$$a_{s+1} > \frac{\sqrt{15} b^{q_s/2} \log a}{2q_s} - 2.$$

A short calculation with Maple 9.5 verifies that this does not occur, completing the proof of Theorem 1.1.

**Acknowledgments** The author would like to thank Michael Filaseta for numerous helpful discussions.

## References

- [1] M. Bennett, *Fractional parts of powers of rational numbers*. Math. Proc. Cambridge Philos. Soc. **114**(1993), no. 2, 191–201.
- [2] ———, *On some exponential equations of S. S. Pillai*. Canad. J. Math. **53**(2001), no. 5, 897–922.
- [3] F. Beukers, *Fractional parts of powers of rationals*. Math. Proc. Cambridge Philos. Soc. **90**(1981), no. 1, 13–20.
- [4] J. W. S. Cassels, *On the equation  $a^x - b^y = 1$* . Amer. J. Math. **75**(1953), 159–162.
- [5] ———, *On the equation  $a^x - b^y = 1$ . II*. Proc. Cambridge Philos. Soc. **56**(1960), 97–103.
- [6] W. J. Ellison, *On a theorem of S. Sivasankaranarayana Pillai*. In: Séminaire de théorie des nombres, 1970–71, no. 12, Lab. Théorie des Nombres, Centre Nat. Recherche Sci., Talence, 1971.
- [7] A. Herschfeld, *The equation  $2^x - 3^y = d$* . Bull. Amer. Math. Soc. **42**(1936), no. 4, 231–234.
- [8] A. Y. Khinchin, *Continued Fractions*. Third edition. P. Noordhoff Ltd., Groningen, 1963.
- [9] M. Laurent, M. Mignotte, and Y. Nesterenko, *Formes linéaires en deux logarithmes et déterminants d'interpolation*. J. Number Theory **55**(1995), no. 2, 285–321.
- [10] W. J. LeVeque, *On the equation  $a^x - b^y = 1$* . Amer. J. Math. **74**(1952), 325–331.
- [11] P. Mihăilescu, *Primary cyclotomic units and a proof of Catalan's conjecture*. J. Reine Angew. Math. **572**(2004), 167–195.
- [12] S. S. Pillai, *On the inequality  $0 < a^x - b^y \leq n$* . J. Indian Math. Soc. **19**(1931), 1–11.
- [13] ———, *On  $a^x - b^y = c$* . J. Indian Math. Soc. (N.S.) **2**(1936), 119–122, 215.
- [14] ———, *On  $a^x - b^y = b^y \pm a^x$* . J. Indian Math. Soc. (N.S.) **8**(1944), 10–13.
- [15] ———, *On the equation  $2^x - 3^y = 2^x + 3^y$* . Bull. Calcutta Math. Soc. **37**(1945), 18–20.
- [16] P. Ribenboim, *Catalan's Conjecture*. Academic Press, Boston, MA, 1994.
- [17] R. Scott and R. Styer, *On the generalized Pillai equation  $\pm a^x \pm b^y = c$* . J. Number Theory **118**(2006), no. 2, 236–265.

Department of Mathematics, University of British Columbia, Vancouver, BC, V6T 1Z2  
e-mail: bennett@math.ubc.ca