

LOCAL TIMES OF SELF-INTERSECTION FOR MULTIDIMENSIONAL BROWNIAN MOTION*

SHENG-WU HE, WEN-QIANG YANG, RONG-QIN YAO
AND JIA-GANG WANG

We will define local times of self-intersection for multidimensional Brownian motion as generalized Wiener functionals under the framework of white noise analysis as in H. Watanabe ([6]). By making use of the chaotic representation of δ -function and precise computation we get a deep insight into the problem. In the section 1 multiple Wiener integrals with respect to multidimensional Brownian motion and chaotic representations for square-integrable Wiener functionals are given. They are indispensable, but seem not to be formulated clearly and correctly before. The useful concepts and results of white noise analysis are illustrated in the section 2. Section 3 is the main part of the paper. The applications to local times are introduced in the section 4 briefly.

1. Multiple Wiener integrals

Let $\mathbf{B} = \{(B_t^1, \dots, B_t^d), -\infty < t < \infty\}$ be a d -dimensional Brownian motion defined on a probability space $(\Omega, \mathcal{F}, \mathbf{P})$ such that $\mathcal{F} = \sigma\{(B_t^1, \dots, B_t^d), -\infty < t < \infty\}$. We will define the multiple Wiener integrals with respect to \mathbf{B} . The procedure will be sketched, and all proofs omitted, since they are completely similar to that for one-dimensional Brownian motion (cf. [3]).

Let $n = n_1 + \dots + n_d$, $n_j \geq 0$, $j = 1, \dots, d$. For each $f \in L^2(\mathbf{R}^n)$ define

$$(1.1) \quad \tilde{f}(t_1, \dots, t_n) = \frac{1}{n_1! \dots n_d!} \sum_{\pi^1, \dots, \pi^d} f(t_{\pi_1^1}, \dots, t_{\pi_{n_1}^1}, \dots, t_{\pi_1^d}, \dots, t_{\pi_{n_d}^d}),$$

where π^j , $j = 1, \dots, d$, are the permutations of $(n_1 + \dots + n_{j-1} + 1, \dots, n_1 + \dots + n_j)$, $j = 1, \dots, d$, respectively, and the summation is over all permutations $\{\pi^1, \dots, \pi^d\}$. Write

Received October 29, 1993.

*) The project supported by National Natural Science Foundation of China

$$\tilde{L}^2(\mathbf{R}^{n_1}) \otimes \cdots \otimes \tilde{L}^2(\mathbf{R}^{n_d}) = \{\tilde{f} : f \in L^2(\mathbf{R}^n)\}.$$

For each $f \in L^2(\mathbf{R}^n)$ we can define

$$(1.2) \quad I_{n_1, \dots, n_d}(f) = \int f(t_1, \dots, t_n) dB_{t_n}^1 \cdots dB_{t_{n_1}}^1 \cdots dB_{t_{n_1+\dots+t_{n_{d-1}+1}}}^d \cdots dB_{t_n}^d,$$

such that for any $f \in L^2(\mathbf{R}^n)$

$$(1.3) \quad 1) \quad I_{n_1, \dots, n_d}(f) = I_{n_1, \dots, n_d}(\tilde{f}),$$

$$(1.4) \quad 2) \quad E[I_{n_1, \dots, n_d}(f)] = 0,$$

3) for any $g \in L^2(\mathbf{R}^m)$, $m = m_1 + \cdots + m_d$, $m_j \geq 0$, $j = 1, \dots, d$,

$$(1.5) \quad E[I_{n_1, \dots, n_d}(f)I_{m_1, \dots, m_d}(g)] = n_1! \cdots n_d! \langle \tilde{f}, \tilde{g} \rangle \delta_{(n_1, \dots, n_d)(m_1, \dots, m_d)}$$

4) for any $g \in L^2(\mathbf{R}^n)$, $a, b \in \mathbf{R}$

$$(1.6) \quad I_{n_1, \dots, n_d}(af + bg) = aI_{n_1, \dots, n_d}(f) + bI_{n_1, \dots, n_d}(g),$$

5) for $f(t_1, \dots, t_n) = f_1(t_1, \dots, t_{n_1}) \cdots f_d(t_{n_1+\dots+n_{d-1}+1}, \dots, t_n)$, $f_j \in L^2(\mathbf{R}^{n_j})$,

$$(1.7) \quad j = 1, \dots, d,$$

$$I_{n_1, \dots, n_d}(f) = \int f_1 dB_{t_1}^1 \cdots dB_{t_{n_1}}^1 \cdots \int f_d dB_{t_1}^d \cdots dB_{t_{n_d}}^d.$$

Thus $I_{n_1, \dots, n_d}(\tilde{f})$ is an isometric mapping from $\tilde{L}^2(\mathbf{R}^{n_1}) \otimes \cdots \otimes \tilde{L}^2(\mathbf{R}^{n_d})$ into $(L^2) = L^2(\Omega, \mathcal{F}, \mathbf{P})$. The most important result is the following chaotic representation for square-integrable Wiener functionals:

To any $\phi \in (L^2)$ corresponds a unique sequence $\{\phi_{n_1, \dots, n_d} \in \tilde{L}^2(\mathbf{R}^{n_1}) \otimes \cdots \otimes \tilde{L}^2(\mathbf{R}^{n_d}), n_j \geq 0, j = 1, \dots, d\}$ such that

$$(1.8) \quad \phi = \sum_{n=0}^{\infty} \sum_{n_1+\dots+n_d=n} I_{n_1, \dots, n_d}(\phi_{n_1, \dots, n_d}),$$

and

$$(1.9) \quad \|\phi\|_2^2 = \sum_{n=0}^{\infty} \sum_{n_1+\dots+n_d=n} n_1! \cdots n_d! \|\phi_{n_1, \dots, n_d}\|_2^2,$$

where $I_0(f) = E[f]$. Later we will denote (1.8) simply by $\phi \sim (\phi_{n_1, \dots, n_d})$.

2. White noise space

We adopt the framework of white noise analysis set by T. Hida (cf. [1] or [2]).

Let $\mathcal{S}(\mathbf{R})$ be the Schwartz space of rapidly decreasing functions on \mathbf{R} . Denote by A the self-adjoint extension of the harmonic oscillator operator on $L^2(\mathbf{R})$:

$$(2.1) \quad Af(t) = -f''(t) + (1 + t^2)f(t), \quad f \in \mathcal{S}(\mathbf{R}).$$

Let $H_n(x)$, $n \geq 0$, be the Hermite polynomial of order n , and

$$(2.2) \quad e_n(x) = (n!2^n)^{-1/2}(\pi)^{-1/4}H_n(x)e^{-x^2/2}, \quad n \geq 0.$$

Then $e_n \in \mathcal{S}(\mathbf{R})$, and $\{e_n, n \geq 0\}$ is an orthogonal normal basis of $L^2(\mathbf{R})$ and

$$(2.3) \quad Ae_n = (2n + 2)e_n, \quad n \geq 0.$$

Put

$$(2.4) \quad \|f\|_{2,p}^2 = \|A^p f\|_2^2 = \sum_{n=0}^{\infty} (2n + 2)^{2p} |\langle f, e_n \rangle|^2, \quad f \in L^2(\mathbf{R}),$$

$$(2.5) \quad \mathcal{S}_p(\mathbf{R}) = \mathcal{D}(A^p) = \{f \in L^2(\mathbf{R}) : \|f\|_{2,p}^2 < \infty\}, \quad p \geq 0.$$

With $\{\|\cdot\|_{2,p}, p \geq 0\}$ $\mathcal{S}(\mathbf{R})$ is a nuclear space. Let $\mathcal{S}'(\mathbf{R})$ be its dual. Set

$$(2.6) \quad \mathcal{S}_p'(\mathbf{R}) = \left\{ f \in \mathcal{S}'(\mathbf{R}) : \|f\|_{2,p}^2 = \sum_{n=0}^{\infty} (2n + 2)^{2p} |\langle f, e_n \rangle|^2 < \infty \right\}, \quad p \in \mathbf{R},$$

where $\langle \cdot, \cdot \rangle$ denotes the pairing between $\mathcal{S}(\mathbf{R})$ and $\mathcal{S}'(\mathbf{R})$. Then

$$(2.7) \quad \mathcal{S}(\mathbf{R}) = \bigcap_{p \in \mathbf{R}} \mathcal{S}_p(\mathbf{R}), \quad \mathcal{S}'(\mathbf{R}) = \bigcup_{p \in \mathbf{R}} \mathcal{S}_p'(\mathbf{R}).$$

By Minlos theorem there exists a unique probability measure μ on $\mathcal{B}(\mathcal{S}'(\mathbf{R}))$, the σ -field generated by cylinder sets, such that

$$(2.8) \quad \int_{\mathcal{S}'(\mathbf{R})} e^{i\langle x, \xi \rangle} \mu(dx) = \exp\left\{-\frac{1}{2} \|\xi\|_2^2\right\}, \quad \xi \in \mathcal{S}(\mathbf{R}).$$

The measure μ is called the white noise measure, and the probability space $(\mathcal{S}'(\mathbf{R}), \mathcal{B}(\mathcal{S}'(\mathbf{R})), \mu)$ is called the white noise space. Define

$$(2.9) \quad (\Omega_d, \mathcal{F}_d, \mu_d) = (\mathcal{S}'(\mathbf{R}), \mathcal{B}(\mathcal{S}'(\mathbf{R})), \mu)^d.$$

It will be our fundamental probability space, and denote by $(L^{2,d})$ the L^2 -space on it. Set

$$(2.10) \quad X_\xi^j(x) = \langle x_j, \xi \rangle, \quad x = (x_1, \dots, x_d) \in \Omega_d, \quad \xi \in \mathcal{S}(\mathbf{R}), \quad j = 1, \dots, d.$$

For each j , $\xi \rightarrow X_\xi^j$ is an isometric mapping from $\mathcal{S}(\mathbf{R})$ into $(L^{2,d})$, and can be extended as an isometric mapping from $L^2(\mathbf{R})$ into $(L^{2,d})$, i.e., X_ξ^j is well defined for

$\xi \in L^2(\mathbf{R})$. Put

$$f_t = \begin{cases} 1_{(0,t)}, & t \geq 0, \\ 1_{(t,0)}, & t < 0, \end{cases}$$

and

$$(2.11) \quad B_i^j(\mathbf{x}) = \langle x_j, f_i \rangle, \quad -\infty < t < \infty, j = 1, \dots, d.$$

Then $\mathbf{B} = \{(B_i^1, \dots, B_i^d), -\infty < t < \infty\}$ is a d -dimensional Brownian motion, and $\mathcal{F}_d = \sigma\{(B_i^1, \dots, B_i^d), -\infty < t < \infty\}$. Let $\phi \sim (\phi_{n_1, \dots, n_d}) \in (L^{2,d})$. For $p \geq 0$, define

$$(2.12) \quad (\mathcal{S}^d)_p = \left\{ \phi \sim (\phi_{n_1, \dots, n_d}) \in (L^{2,d}) : \|\phi\|_{2,p}^2 = \sum_{n=0}^{\infty} \sum_{n_1+\dots+n_d=n} n_1! \cdots n_d! | (A^p)^{\otimes n} \phi_{n_1, \dots, n_d} |^2 < \infty \right\}.$$

Denote by $(\mathcal{S}^d)_{-p}$ the dual of $(\mathcal{S}^d)_p$. Define

$$(\mathcal{S}^d) = \bigcap_{p \geq 0} (\mathcal{S}^d)_p.$$

With $\{\|\cdot\|_{2,p}, p \geq 0\}$ (\mathcal{S}^d) is a nuclear space, and its dual

$$(\mathcal{S}^d)^* = \bigcup_{p \geq 0} (\mathcal{S}^d)_{-p}.$$

Each element of (\mathcal{S}^d) is called a test functional, and each element of $(\mathcal{S}^d)^*$ is called a generalized Wiener functional or Hida distribution.

For $\xi = (\xi_1, \dots, \xi_d) \in \mathcal{S}^d(\mathbf{R})$, exponential functional

$$(2.13) \quad \mathcal{E}(\xi)(\mathbf{x}) = \exp\left\{ \sum_{j=1}^d \left(\langle x_j, \xi_j \rangle - \frac{1}{2} |\xi_j|_2^2 \right) \right\} \sim \left(\frac{1}{n_1! \cdots n_d!} \xi_1^{\otimes n_1} \otimes \cdots \otimes \xi_d^{\otimes n_d} \right),$$

$$\mathbf{x} = (x_1, \dots, x_d) \in \Omega_d.$$

is a test functional. For any $\Phi \in (\mathcal{S}^d)^*$ the S -transform of Φ is defined as

$$(2.14) \quad (S\Phi)(\xi) = \langle\langle \Phi, \mathcal{E}(\xi) \rangle\rangle, \quad \xi \in \mathcal{S}^d(\mathbf{R}),$$

where $\langle\langle \cdot, \cdot \rangle\rangle$ denotes the pairing between (\mathcal{S}^d) and $(\mathcal{S}^d)^*$. If $\Phi \in (L^{2,d})^*$, then

$$(2.15) \quad (S\Phi)(\xi) = \int_{\Omega_d} \Phi(\mathbf{x} + \xi) \mu_d(d\mathbf{x}).$$

A functional U on $\mathcal{S}^d(\mathbf{R})$ is called a U -functional. If

1) for each $\xi = (\xi_1, \dots, \xi_d) \in \mathcal{S}^d(\mathbf{R})$ the mapping

$$(\lambda_1, \dots, \lambda_d) \rightarrow U(\lambda_1 \xi_1, \dots, \lambda_d \xi_d)$$

has analytic continuation, denoted by $u(z_1, \dots, z_d; \xi_1, \dots, \xi_d)$, on \mathbf{C}^n ;

2) for any $n = n_1 + \dots + n_d$, $n_j \geq 0$, $j = 1, \dots, d$,

$$U_{n_1, \dots, n_d}(\xi_1, \dots, \xi_d) = \frac{1}{n_1! \dots n_d!} \sum_{j_1=1}^{n_1} \dots \sum_{j_d=1}^{n_d} (-1)^{n-(j_1+\dots+j_d)}$$

$$\sum_{i_1^d < \dots < i_1^d} \dots \sum_{i_1^d < \dots < i_1^d} \frac{\partial^n}{\partial^{n_1} z_1 \dots \partial^{n_d} z_d} u(0, \dots, 0; \xi_{i_1^1} + \dots + \xi_{i_1^1}, \dots, \xi_{i_1^d} + \dots + \xi_{i_1^d})$$

is multilinear in (ξ_1, \dots, ξ_d) ;

3) there exist constants $C_1 > 0$, $C_2 > 0$, $p \in \mathbf{R}$ such that for all $(z_1, \dots, z_d) \in \mathbf{C}^n$ and $(\xi_1, \dots, \xi_d) \in \mathcal{D}^d(\mathbf{R})$

$$|u(z_1, \dots, z_d; \xi_1, \dots, \xi_d)| \leq C_1 \exp\left\{C_2 \sum_{j=1}^d |z_j|^2 |\xi_j|_{2,p}^2\right\}.$$

Pothoff-Streit characterization theorem states that a functional on $\mathcal{D}^d(\mathbf{R})$ is the S -transform of a Hida distribution if and only if it is a U -functional (cf. [5], only the case of $d = 1$ is discussed there). Every Hida distribution is uniquely determined by its S -transform. In particular, if

$$(2.16) \quad U(\xi) = \sum_{n=0}^{\infty} \sum_{n_1+\dots+n_d=n} \langle f_{n_1, \dots, n_d}, \xi_1^{\otimes n_1} \otimes \dots \otimes \xi_d^{\otimes n_d} \rangle$$

and for some $p \geq 0$

$$(2.17) \quad \sum_{n=0}^{\infty} \sum_{n_1+\dots+n_d=n} n_1! \dots n_d! |A^{-p}|^{\otimes n} f_{n_1, \dots, n_d}|_2^2 < \infty,$$

where $f_{n_1, \dots, n_d} \in \tilde{L}^2(\mathbf{R}^{n_1}) \otimes \dots \otimes \tilde{L}^2(\mathbf{R}^{n_d})$, then there exists $\Phi \in (\mathcal{D}^d)_{-p}$ such that $(S\Phi)(\xi) = U(\xi)$, and the expression in (2.17) is just equal to $\|\Phi\|_{2,-p}^2$. In this case, we also denote $\Phi \sim (f_{n_1, \dots, n_d})$.

3. Local times of self-intersection

Let f be a bounded Borel function on \mathbf{R}^d . Then it is well-known by (2.15) that for any $0 \leq s \leq t$

$$[Sf(\mathbf{B}_t - \mathbf{B}_s)](\xi) = [T_{t-s}^{(d)} f] \left(\int_s^t \xi(r) dr \right),$$

where $\{T_t^{(d)}, t \geq 0\}$ is the transition semigroup of d -dimensional Brownian mo-

tion:

$$(3.1) \quad [T_{t-s}^{(d)} f](\mathbf{x}) = \int f(y_1, \dots, y_d) \frac{1}{[2\pi(t-s)]^{d/2}} \exp\left\{-\frac{1}{2(t-s)} \sum_{j=1}^d (y_j - x_j)^2\right\} dy_1 \cdots dy_d$$

Let $\delta_{\mathbf{a}}, \mathbf{a} = (a_1, \dots, a_d)$, be the δ -function at point \mathbf{a} . From (3.1) we have formally

$$(3.2) \quad [S\delta_{\mathbf{a}}(\mathbf{B}_t - \mathbf{B}_s)](\xi) = \frac{1}{[2\pi(t-s)]^{d/2}} \exp\left\{-\frac{1}{2(t-s)} \sum_{j=1}^d \left(a_j - \int_s^t \xi_j(r) dr\right)^2\right\}.$$

It is not difficult to see that the right hand of (3.2) is a U -functional. In fact, by the formula of the generating function for Hermite polynomials we have

$$\begin{aligned} \exp\left\{-\frac{1}{2(t-s)} \left(\int_s^t \xi_j(r) dr - a_j\right)^2\right\} &= e^{-x_j^2} \sum_{n=0}^{\infty} \frac{1}{n!} H_n(x_j) \left(\frac{\int_s^t \xi_j(r) dr}{\sqrt{2(t-s)}}\right)^n \\ [2\pi(t-s)]^{-d/2} \exp\left\{-\frac{1}{2(t-s)} \sum_{j=1}^d \left(\int_s^t \xi_j(r) dr - a_j\right)^2\right\} \\ &= [2\pi(t-s)]^{-d/2} e^{-\sum_{j=1}^d x_j^2} \sum_{n=0}^{\infty} \sum_{n_1+\dots+n_d=n} \prod_{j=1}^d \left\{\frac{1}{n_j!} H_{n_j}(x_j) \left(\frac{\int_s^t \xi_j(r) dr}{\sqrt{2(t-s)}}\right)^{n_j}\right\}, \end{aligned}$$

where $x_j = \frac{a_j}{\sqrt{2(t-s)}}$. Put

$$(3.3) \quad \begin{aligned} \phi_{n_1, \dots, n_d}^{(\mathbf{a})}(s, t) &= [2\pi(t-s)]^{-d/2} e^{-\sum_{j=1}^d x_j^2} \prod_{j=1}^d \frac{1}{n_j!} H_{n_j}(x_j) \left(\frac{1_{(s,t)}}{\sqrt{2(t-s)}}\right)^{\otimes n_j} \\ &= [2(t-s)]^{-d/2} \pi^{-d/4} e^{-\sum_{j=1}^d x_j^2/2} \prod_{j=1}^d \frac{1}{\sqrt{n_j!}} e_{n_j}(x_j) \left(\frac{1_{(s,t)}}{\sqrt{t-s}}\right)^{\otimes n_j}. \end{aligned}$$

Note that $\{e_n(x), n \geq 0\}$ is uniformly bounded: $C = \sup_{n,x} |e_n(x)| < \infty$. Now for any $p > 0$ we have

$$(3.4) \quad \begin{aligned} |(A^{-p})^{\otimes n} \phi_{n_1, \dots, n_d}^{(\mathbf{a})}(s, t)|_2^2 &\leq 2^{-2np} |\phi_{n_1, \dots, n_d}^{(\mathbf{a})}(s, t)|_2^2 \\ &\leq 2^{-2np} [2(t-s)]^{-d} \pi^{-d/2} \prod_{j=1}^d \left(\frac{1}{n_j!} e_{n_j}^2(x_j)\right) \left|\frac{1_{(s,t)}}{\sqrt{t-s}}\right|_2^{2n} \\ &\leq C^2 2^{-2np} [2(t-s)]^{-d} \pi^{-d/2} \prod_{j=1}^d \frac{1}{n_j!}, \end{aligned}$$

where $n = n_1 + \dots + n_d$, and

$$\begin{aligned} & \sum_{n=0}^{\infty} \sum_{n_1+\dots+n_d=n} n_1! \cdots n_d! | (A^{-p})^{\otimes n} \phi_{n_1, \dots, n_d}^{(\mathbf{a})}(s, t) |^2 \\ & \leq C^2 [2(t-s)]^{-d} \pi^{-d/2} \sum_{n=0}^{\infty} \sum_{n_1+\dots+n_d=n} 2^{-2np} \\ & = C^2 [2(t-s)]^{-d} \pi^{-d/2} \sum_{n=0}^{\infty} 2^{-2np} C_{n+d-1}^{d-1} < \infty. \end{aligned}$$

According to (2.16) and (2.17), we obtain the following

LEMMA 1. For any $0 \leq s < t$, $p > 0$ and $\mathbf{a} \in \mathbf{R}^d$

$$\delta_{\mathbf{a}}(\mathbf{B}_t - \mathbf{B}_s) \sim (\phi_{n_1, \dots, n_d}^{(\mathbf{a})}(s, t)) \in (\mathcal{S}^d)_{-p}$$

is a Hida distribution.

THEOREM 1. If $\mathbf{a} \neq \mathbf{0}$, then for any $t \geq 0$, $p > 0$

$$G_{\mathbf{a}}^{\alpha}(t) = \int_{0 \leq u < v \leq t} \delta_{\mathbf{a}}(\mathbf{B}_v - \mathbf{B}_u) dudv \in (\mathcal{S}^d)_{-p}$$

is a Hida distribution.

Proof. Noting that

$$C_{\mathbf{a}} = \pi^{-d/4} \sup_{u>0} \{ (2u)^{-d/2} e^{-|\mathbf{a}|^2/4u} \} < \infty,$$

(3.4) can be modified as

$$| (A^{-p})^{\otimes n} \phi_{n_1, \dots, n_d}^{(\mathbf{a})}(s, t) |^2 \leq C^2 C_{\mathbf{a}}^2 2^{-2np} \prod_{j=1}^d \frac{1}{n_j!}.$$

Then we have

$$\begin{aligned} & \| \delta_{\mathbf{a}}(\mathbf{B}_t - \mathbf{B}_s) \|_{2, -p}^2 \leq C^2 C_{\mathbf{a}}^2 \sum_{n=0}^{\infty} 2^{-2np} C_{n+d-1}^{d-1}, \\ & \int_{0 \leq u < v \leq t} \| \delta_{\mathbf{a}}(\mathbf{B}_v - \mathbf{B}_u) \|_{2, -p} dudv < \infty. \end{aligned}$$

Hence $G_{\mathbf{a}}^{(\mathbf{a})}(t) \in (\mathcal{S}^d)_{-p}$. □

THEOREM 2. For any $t \geq 0$

(3.5)

$$G_d(t) = \int_{0 \leq u < v \leq t} [\delta(\mathbf{B}_v - \mathbf{B}_u) - \sum_{n=0}^{d-2} \sum_{n_1+\dots+n_d=n} I_{n_1, \dots, n_d}(\phi_{n_1, \dots, n_d}(u, v))] dudv$$

is a Hida distribution, called local times of self-intersection for d -dimensional Brownian motion, where $\delta = \delta_0$.

Proof. At first, observe that let $p > 0$

$$(3.6) \quad \begin{aligned} |A^{-p} \mathbf{1}_{(s,t)}|_2^2 &= \sum_{k=0}^{\infty} (2k+2)^{-2p} |\langle \mathbf{1}_{(s,t)}, e_k \rangle|^2 \\ &\leq \sum_{k=0}^{\infty} (2k+2)^{-2p} (t-s)^2 \sup_x e_k^2(x) \end{aligned}$$

Since $\sup_x e_k^2(x) = O(k^{-\frac{1}{6}})$, take p large enough such that

$$(3.7) \quad \sum_{k=0}^{\infty} (2k+2)^{-2p} \sup_x e_k^2(x) < 1.$$

Then by (3.6) and (3.7) we get

$$(3.8) \quad \begin{aligned} |(A^{-p})^{\otimes n} \phi_{n_1, \dots, n_d}(s, t)|_2^2 &= [2\pi(t-s)]^{-d} \prod_{j=1}^d \left(\frac{H_{n_j}(0)}{n_j!} \right)^2 \left| A^{-p} \frac{\mathbf{1}_{(s,t)}}{\sqrt{2(t-s)}} \right|_2^{2n} \\ &\leq \frac{1}{n_1! \cdots n_d!} (2\pi)^{-d} (t-s)^{n-d}, \end{aligned}$$

noting that $H_{2k+1}(0) = 0$, $H_{2k}(0) = (-1)^k \frac{(2k)!}{k!}$, $k \geq 0$, for any $k \geq 0$, $H_k^2(0) \leq k!2^k$. Let $\alpha > 0$, from (3.8) we have

$$(3.9) \quad \begin{aligned} &\left\| \delta(\mathbf{B}_t - \mathbf{B}_s) - \sum_{n=0}^{d-2} \sum_{n_1+\dots+n_d=n} I_{n_1, \dots, n_d}(\phi_{n_1, \dots, n_d}(s, t)) \right\|_{2, -(p+\alpha)}^2 \\ &= \left\| \sum_{n=d-1}^{\infty} \sum_{n_1+\dots+n_d=n} I_{n_1, \dots, n_d}(\phi_{n_1, \dots, n_d}(s, t)) \right\|_{2, -(p+\alpha)}^2 \\ &= \sum_{n=d-1}^{\infty} \sum_{n_1+\dots+n_d=n} n_1! \cdots n_d! |(A^{-(p+\alpha)})^{\otimes n} \phi_{n_1, \dots, n_d}(s, t)|_2^2 \\ &\leq \sum_{n=d-1}^{\infty} \sum_{n_1+\dots+n_d=n} n_1! \cdots n_d! 2^{-2n\alpha} |(A^{-p})^{\otimes n} \phi_{n_1, \dots, n_d}(s, t)|_2^2 \\ &\leq \sum_{n=d-1}^{\infty} \sum_{n_1+\dots+n_d=n} 2^{-2n\alpha} (2\pi)^{-d} (t-s)^{n-d} \\ &= (2\pi)^{-d} \sum_{n=d-1}^{\infty} 2^{-2n\alpha} C_{n+d-1}^{d-1} (t-s)^{n-d}. \end{aligned}$$

Take α large enough such that $2^{-\alpha}\sqrt{t} < 1$. Then

$$\int_{0 \leq u < v \leq t} \left\| \delta(\mathbf{B}_v - \mathbf{B}_u) - \sum_{n=0}^{d-2} \sum_{n_1+\dots+n_d=n} I_{n_1, \dots, n_d}(\psi_{n_1, \dots, n_d}(u, v)) \right\|_{2, -(p+\alpha)} dudv < \infty.$$

Hence, the integral in the right hand of (3.5) exists in $(\mathcal{S}^d)_{-(p+\alpha)}$ as Bochner integral, i.e., $G_d(t) \in (S^d)_{-(p+\alpha)}$ is a Hida distribution. □

From (3.5) we have

$$(3.10) \quad G_1(t) = \int_{0 \leq u < v \leq t} \delta(\mathbf{B}_v - \mathbf{B}_u) dudv,$$

$$(3.11) \quad G_2(t) = \int_{0 \leq u < v \leq t} \{\delta(\mathbf{B}_v - \mathbf{B}_u) - [2\pi(v - u)]^{-1}\} dudv,$$

$$(3.12) \quad G_3(t) = \int_{0 \leq u < v \leq t} \{\delta(\mathbf{B}_v - \mathbf{B}_u) - [2\pi(v - u)]^{-3/2}\} dudv,$$

$$(3.13) \quad G_4(t) = \int_{0 \leq u < v \leq t} \left\{ \delta(\mathbf{B}_v - \mathbf{B}_u) - \frac{1}{[2\pi(v - u)]^2} + \frac{1}{(2\pi)^2(v - u)^3} \sum_{j=1}^4 \int_u^v (B_r^j - B_u^j) dB_r^j \right\} dudv,$$

$$(3.14) \quad G_5(t) = \int_{0 \leq u < v \leq t} \left\{ \delta(\mathbf{B}_v - \mathbf{B}_u) - \frac{1}{[2\pi(v - u)]^{5/2}} + \frac{1}{(2\pi)^{5/2}(v - u)^{7/2}} \sum_{j=1}^5 \int_u^v (B_r^j - B_u^j) dB_r^j \right\} dudv,$$

.....

For $d \geq 2$, the integral

$$\int_{0 \leq u < v \leq t} \delta(\mathbf{B}_v - \mathbf{B}_u) dudv$$

has no meaning even in generalized sense, and must be renormalized according to (3.5). The renormalization parts in $G_4(t)$ and $G_5(t)$ in Watanabe ([6]) are mistaken, caused by his small error in computation.

LEMMA 2. Set

$$a_n^{(d)}(t) = \left| \int_{0 \leq u < v \leq t} (v - u)^{-(n+\frac{d}{2})} 1_{(u,v)}^{\otimes 2n} dudv \right|_2^2.$$

Then as $n \rightarrow \infty$, for $t > 0$ we have

$$(3.15) \quad a_n^{(d)}(t) \begin{cases} \sim \frac{2t^3}{3n^2}, & d = 1, \\ \sim \frac{2t^2}{n^2}, & d = 2, \\ = \infty, & d \geq 3. \end{cases}$$

Proof. By Fubini theorem on interchanging the order of integration, we have

$$(3.16) \quad \begin{aligned} a_n^{(d)}(t) &= \int_{\substack{0 \leq u < v \leq t \\ 0 \leq r < s \leq t}} \frac{|\langle 1_{(u,v)}, 1_{(r,s)} \rangle|^{2n}}{(v-u)^{n+d/2}(s-r)^{n+d/2}} dudvdrds \\ &= \int_{\substack{0 \leq u < v \leq t, 0 \leq r < s \leq t, \\ r < u < s \text{ or } u < r < v}} \frac{(v \wedge s - u \vee r)^{2n}}{(v-u)^{n+d/2}(s-r)^{n+d/2}} dudvdrds \\ &= 2 \int_{0 \leq r < u < s < v \leq t} \frac{(s-u)^{2n}}{(v-u)^{n+d/2}(s-r)^{n+d/2}} dudvdrds \\ &\quad + 2 \int_{0 \leq r < u < v < s \leq t} \frac{(v-u)^{n-d/2}}{(s-r)^{n+d/2}} dudvdrds \\ &= 2(a_{n1}^{(d)}(t) + a_{n2}^{(d)}(t)). \end{aligned}$$

$$(3.17) \quad \begin{aligned} a_{n1}^{(d)}(t) &= \int_{0 \leq r < s \leq t} (s-r)^{-(n+d/2)} drds \int_{r < u < v < s} (v-u)^{n-d/2} dudv \\ &= [(n-d/2+1)(n-d/2+2)]^{-1} \int_{0 \leq r < s \leq t} (s-r)^{2-d} drds \\ &= C_t^{(d)} [(n-d/2+1)(n-d/2+2)], \end{aligned}$$

where

$$(3.18) \quad C_t^{(d)} = \int_{0 \leq r < s \leq t} (s-r)^{-d+2} drds = \begin{cases} t^3/6, & d = 1, \\ t^2/2, & d = 2, \\ \infty, & d \geq 3. \end{cases}$$

Below we consider only the case of $d = 1$ or 2 .

$$(3.19) \quad a_{n1}^{(d)}(t) = 2 \int_{0 \leq u < s < v \leq t} \frac{(s-u)^{2n}}{(v-u)^{n+d/2}} dudvds \int_0^u \frac{dr}{(s-r)^{n+d/2}}$$

$$\begin{aligned}
 &= (n + d/2 - 1)^{-1} \int_{0 \leq u < s < v \leq t} \left[\frac{(s - u)^{n-d/2+1}}{(v - u)^{n+d/2}} - \frac{(s - u)^{2n}}{(v - u)^{n+d/2} s^{n+d/2-1}} \right] dudvds \\
 &= (n + d/2 - 1)^{-1} [a_{n11}^{(d)}(t) - a_{n12}^{(d)}(t)]. \quad (0 \leq a_{n12}^{(d)}(t) \leq a_{n11}^{(d)}(t).) \\
 (3.20) \quad a_{n11}^{(d)}(t) &= \int_{0 \leq u < v \leq t} (v - u)^{-(n+d/2)} dudv \int_u^v (s - u)^{n-d/2+1} ds \\
 &= C_t^{(d)} (n - d/2 + 2)^{-1}. \\
 a_{n12}^{(d)}(t) &= \int_{0 \leq u < s \leq t} \frac{(s - u)^{2n}}{s^{n+d/2-1}} duds \int_s^t \frac{dv}{(v - u)^{n+d/2}} \\
 &= (n + d/2 - 1)^{-1} \int_{0 \leq u < s \leq t} \left[\frac{(s - u)^{n-d/2+1}}{s^{n+d/2-1}} - \frac{(s - u)^{2n}}{s^{n+d/2-1} (t - u) s^{n+d/2-1}} \right] duds \\
 &= (n + d/2 - 1)^{-1} [a_{n121}^{(d)}(t) - a_{n122}^{(d)}(t)]. \quad (0 \leq a_{n122}^{(d)}(t) \leq a_{n121}^{(d)}(t).) \\
 a_{n121}^{(d)}(t) &= \int_0^t s^{-(n+d/2-1)} ds \int_0^s (s - u)^{n+d/2+1} du \\
 &= (n + d/2 + 1)^{-1} \int_0^t s^{-d+3} ds = o(1).
 \end{aligned}$$

Thus $a_{n122}^{(d)}(t) = o(1)$, $a_{n12}^{(d)}(t) = o(1)$, and by (3.19), (3.20) we have

$$(3.21) \quad a_{n1}^{(d)}(t) = \frac{1}{n^2} C_t^{(d)} (1 + o(1)), \quad d = 1, 2.$$

At last, (3.15) follows from (3.16), (3.17), (3.18) and (3.21). □

In Lemma 2 one needs only that $2n$ is an integer.

THEOREM 3. For any $t \geq 0$ and $\mathbf{a} \in \mathbf{R}$

$$G_d^{(\mathbf{a})}(t) \in (L^{2,d}), \quad d = 1, 2.$$

Proof. From (3.3) we have

$$G_d^{(\mathbf{a})}(t) \sim \left(\int_{0 \leq u < v \leq t} \frac{\exp(-\sum_{j=1}^d x_j^2)}{(2\pi)^{2/d}} \prod_{j=1}^d \frac{e_{n_j}(x_j)}{\sqrt{n_j!}} \frac{1_{(u,v)}^{\otimes n}}{(v - u)^{(n+d)/2}} dudv \right),$$

where $x_j = \frac{a_j}{\sqrt{2(v - u)}}$ and the first term is considered as zero when $\mathbf{a} = \mathbf{0}$ and $d = 2$. Hence,

$$\begin{aligned}
 & \| G_d^{(a)}(t) \|_2^2 \\
 &= \sum_{n=n_0}^{\infty} \sum_{n_1+\dots+n_d=n} \left| \int_{0 \leq u < v \leq t} \frac{e^{-\sum_{j=1}^d x_j^2}}{(2\pi)^{d/2}} \prod_{j=1}^d e^{n_j(x_j)} \frac{1_{(u,v]}^{\otimes n}}{(v-u)^{(n+d)/2}} dudv \right|_2^2 \\
 &\leq C \left(1 + \sum_{n=1}^{\infty} \sum_{n_1+\dots+n_d=n} (2\pi)^{-d} \prod_{j=1}^d (n_j)^{-1/6} a_{n/2}^{(d)}(t) \right) \\
 &\leq C \left(1 + \sum_{n=1}^{\infty} \sum_{n_1+\dots+n_d=n} n^{-2} \prod_{j=1}^d n_j^{-1/6} \right) \\
 &\leq C \left(1 + \sum_{n=1}^{\infty} \sum_{n_1+\dots+n_d=n} \prod_{j=1}^d n_j^{-1/6} n_j^{-1} \right) \\
 &\leq C \left(1 + \sum_{n=1}^{\infty} n^{-7/6} \right)^d < \infty,
 \end{aligned}$$

where

$$n_0 = \begin{cases} 1, & a = 0 \text{ and } d = 2, \\ 0, & \text{otherwise} \end{cases}$$

and C is a constant depending on only d and t , but may vary in different expressions. Thus $G_d^{(a)}(t) \in (L^{2,d})$ for $d = 1, 2$. □

Based on Lemma 2, it is plausible to reason that Theorem 3 does not hold for $d \geq 3$.

THEOREM 4. *For any bounded Borel function f and $t > 0$*

$$(3.22) \quad \int_{0 \leq u < v \leq t} f(\mathbf{B}_v - \mathbf{B}_u) dudv = \begin{cases} \int_{\mathbf{R}} f(\mathbf{a}) G_1^{(a)}(t) d\mathbf{a}, & d = 1, \\ \int_{\mathbf{R}^2} f(\mathbf{a}) G_2^{(a)}(t) d\mathbf{a}, & d = 2. \end{cases}$$

(3.22) is the so-called Tanaka’s formula.

Proof. We only give the proof for $d = 2$. We show the S -transforms of the two sides of (3.22) are the same. Let $\xi = (\xi_1, \xi_2) \in \mathcal{L}(\mathbf{R}) \times \mathcal{L}(\mathbf{R})$. Then

$$\begin{aligned}
 & \left[S \left(\int_{\mathbf{R}^2} f(\mathbf{a}) G_2^{(a)}(t) d\mathbf{a} \right) \right] (\xi) = \int_{\mathbf{R}^2} f(\mathbf{a}) [S(G_2^{(a)}(t))] (\xi) d\mathbf{a} \\
 &= \int_{\mathbf{R}^2} \int_{0 \leq u < v \leq t} \frac{f(\mathbf{a})}{2\pi(v-u)} \exp \left\{ -\frac{1}{2(v-u)} \sum_{j=1}^2 \left(a_j - \int_u^v \xi_j(r) dr \right)^2 \right\} dudv d\mathbf{a} \\
 &= \int_{0 \leq u < v \leq t} [T_{v-u}^{(2)} f] \left(\int_u^v \xi(r) dr \right) dudv.
 \end{aligned}$$

On the other hand,

$$\begin{aligned} \left[S\left(\int_{0 \leq u < v \leq t} f(\mathbf{B}_v - \mathbf{B}_u) dudv\right) \right](\xi) &= \int_{0 \leq u < v \leq t} [S(f(\mathbf{B}_v - \mathbf{B}_u))](\xi) dudv \\ &= \int_{0 \leq u < v \leq t} [T_{v-u}^{(2)} f]\left(\int_u^v \xi(r) dr\right) dudv. \end{aligned}$$

Hence (3.22) follows. □

Even for $d \geq 3$ the Tanaka's formula (3.22) holds, but the integrals in the right side of (3.22) should be understood in the sense of the Bochner integral in $(S^d)_{-p}$ for any $p > 0$.

4. Local times

Compared with the results and proofs in the above section, we can easily obtain the following results:

1) For any $d \geq 1, t \geq 0$

$$L_d(t) = \int_0^t \left[\delta(\mathbf{B}_u) - \sum_{n=0}^{d-2} \sum_{n_1+\dots+n_d=n} I_{n_1, \dots, n_d}(\phi_{n_1, \dots, n_d}(0, u)) \right] du \in (\mathcal{L}^d)^*.$$

Naturally, $L_d(t)$ may be considered as the local times at 0.

2) For any $t > 0, L_1(t) \in (L^{2,1})$. In fact, note that

$$\left| \int_0^t u^{-(n+1/2)} \mathbf{1}_{(0,u]}^{\otimes 2n} du \right|_2 = \begin{cases} \frac{4t}{2n+1}, & d = 1, \\ \infty, & d \geq 2. \end{cases}$$

So even for $d = 2, L_2(t), t > 0$, are impossible to be ordinary Wiener functionals.

3) If $\mathbf{a} \neq \mathbf{0}$, then for any $t \geq 0, p > 0$

$$L_d^{(\mathbf{a})}(t) = \int_0^t \delta_{\mathbf{a}}(\mathbf{B}_u) du \in (\mathcal{L}^d)_{-p}$$

is a Hida distribution. $L_d^{(\mathbf{a})}(t)$ may also be considered as the local time at \mathbf{a} .

4) For any $t > 0, \mathbf{a} \in \mathbf{R}, L_1^{(\mathbf{a})}(t) \in (L^{2,1})$. In fact, noting that

$$\int_{0 \leq u, v \leq t} \left| \left\langle \frac{\mathbf{1}_{(0,u]}}{\sqrt{u}}, \frac{\mathbf{1}_{(0,v]}}{\sqrt{v}} \right\rangle \right|^n dudv = \frac{2t^2}{n+2},$$

it seems that for $\mathbf{a} \neq \mathbf{0}, L_2^{(\mathbf{a})}(t)$ may not be ordinary Wiener functionals.

For the case of $d = 1$ we can give another treatment. Kubo has established

the following generalized Ito's formula (cf. [4]): for all $f \in \mathcal{B}'(\mathbf{R})$, $0 < s < t$,

$$(4.1) \quad f(\mathbf{B}_t) - f(\mathbf{B}_s) = \int_s^t \partial_u^* f'(\mathbf{B}_u) du + \frac{1}{2} \int_s^t f''(\mathbf{B}_u) du.$$

Take $f(u) = u \mathbf{1}_{(a, \infty)}(u) = (u - a)^+$, then $f' = \mathbf{1}_{(a, \infty)}$, $f'' = \delta_a$. Substituting them into (4.1) yields

$$(4.2) \quad (\mathbf{B}_t - a)^+ - (\mathbf{B}_s - a)^+ = \int_s^t \partial_u^* \mathbf{1}_{(a, \infty)}(\mathbf{B}_u) du + \frac{1}{2} \int_s^t \delta_a(\mathbf{B}_u) du.$$

Since $\mathbf{1}_{(a, \infty)}(\mathbf{B}_u)$ is adapted, letting $s \rightarrow 0$ in (4.2) yields

$$L_1^{(a)}(t) = \int_s^t \delta_a(\mathbf{B}_u) du = 2[(\mathbf{B}_t - a)^+ - (-a)^+ - \int_0^t \mathbf{1}_{(a, \infty)}(\mathbf{B}_u) d\mathbf{B}_u] \in (L^{2,1}).$$

This is just the ordinary definition of local times for one-dimensional Brownian motion. Obviously, we provide indeed a white noise analysis treatment of local times for one-dimensional Brownian motion. This approach applies also to local times of self-intersection for one-dimensional Brownian motion. In fact, by using (4.1) it is easy to get

$$\begin{aligned} & \int_{0 \leq u < v \leq t} \delta_a(\mathbf{B}_v - \mathbf{B}_u) dudv \\ &= 2 \left[\int_0^t (\mathbf{B}_t - \mathbf{B}_u - a)^+ du - (-a)^+ t - \int_{0 \leq u < v \leq t} \mathbf{1}_{(a, \infty)}(\mathbf{B}_v - \mathbf{B}_u) dud\mathbf{B}_v \right] \in (L^{2,1}). \end{aligned}$$

REFERENCES

- [1] Hida, T., Analysis of Brownian Functionals, Carleton Math. Lecture, **13**, 1975.
- [2] Hida, T. and Potthoff, J., White noise analysis—an overview, In T. Hida et al. (eds.) White Noise Analysis—Mathematics and Applications, World Scientific, Singapore, 1990.
- [3] Ito, K., Multiple Wiener integrals, J. Math. Soc. Japan, **3** (1951), 157–169.
- [4] Kubo, I, Ito formula for generalized Brownian functionals, Lecture Notes in Control and Information Science, No. **49** (1983), 156–166, Springer-Verlag.
- [5] Potthoff, J. and Streit L., A characterization on Hida distributions, J. Funct. Anal., **101** (1991), 212–229.
- [6] Watanabe, H., The local time of self-intersections of Brownian motion as generalized Brownian functionals, Lett. Math. Phys., **23** (1991), 1–9.

S. He, W. Yang and R. Yao
Department of Mathematical Statistics
East China Normal University
Shanghai, 200062 China

J. Wang
Institute of Applied Mathematics
East China University of Technology
Shanghai, 200237 China