

The Journal of Laryngology and Otology

EDITED BY
WALTER HOWARTH

ASSISTANT EDITOR
G. H. BATEMAN

Contents

ACOUSTIC-VESTIBULAR AND HISTOLOGIC EXAMINATIONS IN GUINEA-PIGS AFTER INTERRUPTION OF MEMBRANOUS LABYRINTH IN SEMICIRCULAR CANALS OR COCHLEA	HARALD K. KRISTENSEN ✓
CYLINDROMAS OF THE UPPER RESPIRATORY TRACT	W. H. STRUBEN and J. F. HAMPE ✓
A METHOD FOR STUDYING THE VIBRATORY MOVEMENTS OF THE VOCAL CORDS	BERTIL SONESSON ✓
A NEW TECHNIQUE IN STAPES MOBILIZATION	BRUCE CORNISH ✓
SOME LONG-TERM RESULTS OF THE PATTERSON EXTERNAL ETHMOIDECTOMY	R. G. WILLIAMS ✓
CLINICAL RECORDS—	
INTRA-PHARYNGEAL HÆMORRHAGE FROM THE INTERNAL CAROTID ARTERY. REPORT OF A CASE AND ARTERIOGRAPHIC FINDINGS	T. B. HOGARTH ✓
A HUGE TRILOBED LARYNGEAL CYST	ALY EL MOFTY ✓
AN UNUSUAL CASE OF CARCINOMA OF THE LARYNX	M. FEINMESSER and I. GAY ✓
SOCIETIES' PROCEEDINGS—	
THE SCOTTISH OTOLARYNGOLOGICAL SOCIETY	✓
GENERAL NOTES	

London
Headley Brothers Ltd
109 Kingsway WC2

Annual Subscription £4/4/0 net, U.S.A. \$13

Monthly, 10/- net post free

The Journal of Laryngology and Otology

(FOUNDED in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN)

EDITED BY
WALTER HOWARTH

ASSISTANT EDITOR
G. H. BATEMAN

1. Original Articles are accepted on the condition that they have not been published elsewhere.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and sent at the same time as original photographs, etc.

The Harvard system of recording references should be used, e.g. GREEN, C., and BROWN, D. (1951) *J. Laryng.*, 65, 33. Abbreviations of Journals should follow the style recommended in *World Medical Periodicals*, published by World Health Organization, 1952.

It is most important that authors should verify *personally* the accuracy of every reference before submitting a paper for publication.

3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.

4. Illustration blocks are provided free up to the limit of £10 per article; beyond this authors are expected to pay half the cost. Coloured illustrations will be charged in full to authors.

Blocks will normally be held by the Printers for three years after which they will be destroyed. Any author who has borne a part of the cost of his blocks is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to HEADLEY BROTHERS LTD., 109 Kingsway, London, WC2.

5. Orders for reprints should be sent when returning galley proofs, and for this purpose special forms are supplied.

6. Authors of original communications on Oto-Laryngology in other journals are invited to send a copy, or two reprints, to the *Journal of Laryngology*. If they are willing, at the same time, to submit their own abstract (in English, French, Italian, or German) it will be welcomed.

7. Editorial communications may be addressed to THE EDITOR, *Journal of Laryngology*, c/o HEADLEY BROTHERS LTD., 109 Kingsway, London, WC2.

8. The annual subscription is four guineas sterling (U.S.A. \$13) post free, and is payable in advance.

9. Single copies will be on sale at 10s. od. each; copies of parts up to Vol. LXIII may be purchased at 4s. each.


10. All subscriptions, advertising and business communications should be sent to the publishers, HEADLEY BROTHERS LTD., 109 KINGSWAY, LONDON, WC2.

United States of America

Orders for this *Journal* may be sent through local bookseller, or to STECHERT-HAFNER, INC., 31-33 East 10th Street, New York, or direct to the publishers, HEADLEY BROTHERS LTD., 109 KINGSWAY, LONDON, WC2, England.

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

**When a
hearing
aid is
advised . .**



**. . you may confidently
recommend your patients
to S. C. INGRAM**

- ★ They can try the different makes of hearing aids
- ★ They get impartial advice on all makes of hearing aids
- ★ Their personal needs are given individual and continued attention
- ★ A full report on each fitting recommended is sent to the patient's ear specialist
- ★ They can compare at one consultation the different makes one with another

S. C. INGRAM Independent Hearing Aid Consultant

On the approved List of the National Institute for the Deaf

2 SHEPHERD STREET, SHEPHERD MARKET, LONDON, W.1

Hyde Park 9042

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

PERDILATAL

**in the treatment of sudden
perceptive deafness
and Ménière's syndrome.**

* Vascular spasm may well be responsible for the majority of these cases.¹

* Vascular changes in the labyrinth are an important factor in the physiopathology of Ménière's disease.²

* In the treatment of Ménière's disease emphasis must be put upon early restoration of blood flow to the inner ear.³

* Perdilatal in doses of at least 6 mg. 4 times a day is the most effective vasodilator for the treatment of these conditions.⁴

1 Fowler, E. P. *Ann. Otorhinolaryng.*, 1950, 59, 980.

2 Atkinson, M. *Proc. roy. Soc. Med.*, 1946, 39, 807

3 Wilmot, T. J. *Brit. med. J.*, 1957, 2, 1047.

4 Wilmot, T. J. Paper read on February 6th, 1959 at a Meeting of the Section of Otology, Roy. Soc. Med.

Perdilatal is available in tablets of 6 mg. as PERDILATAL FORTE in 50's, 250's and 1,000's.



Smith & Nephew Pharmaceuticals Limited

WELWYN GARDEN CITY · HERTFORDSHIRE

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

In schools for the DEAF— multitone Telesonic INDUCTION LOOP SYSTEM brings new advantages

- 1 Can be installed in adjoining classrooms without risk of 'spill over'.
- 2 Combines good hearing with complete freedom of movement. Unaffected by distance between pupil and teacher.
- 3 Clear speech at high amplification.
- 4 Any number of pupils may use the same installation.
- 5 The child can use the instrument as a normal hearing aid outside school.



The Kindergarten Class at the London Residential School for Jewish Deaf Children where all classrooms have been installed with the Multitone Telesonic Induction Loop System

Write for further details to:

multitone


ELECTRIC COMPANY LTD., 12-20 UNDERWOOD STREET, LONDON, N.1.

CRC M23

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

THE HALLPIKE—

BLACKMORE EAR MICROSCOPE*



USES

DIAGNOSIS

The excellent optical performance of the Hallpike Ear Microscope gives the user greatly increased confidence in the diagnosis of abnormal conditions; and many disorders may be recognized at an earlier stage than has hitherto been possible with "headlamp and speculum" techniques. Thus slight pulsation of swellings and vascular tumours—of high diagnostic significance and almost impossible to detect by conventional methods—can readily be detected under this microscope.

Use of the Siegle attachment permits observation of small bubbles present in fluid collections within the tympanum; it is also invaluable for demonstrating minute perforations. Where defects of the tympanic membrane are present, the middle-ear cavity can be scrutinized minutely; the detailed structure of the stapes, stapedius tendon, round window and Eustachian orifice—not normally visible by ordinary methods of examination—are seen with quite remarkable clarity when the Ear Microscope is used.

TREATMENT

Manipulative procedures may be undertaken with greater confidence and precision if the Ear Microscope and its special instruments are used.

The curettes enable adherent wax and debris to be removed with the greatest precision from the walls of the deep meatus or from the tympanic membrane itself. Cholesteatomatous masses or neoplasms may be examined critically and biopsy specimens obtained by means of the curettes. These may also be used as delicate probes for palpation of the incus and head of the stapes in certain cases of otosclerosis with retraction and atrophy of the tympanic membrane; in this way unmistakable evidence may be obtained of bony fixation of the footplate. Or again, it is sometimes possible to identify by palpation defects of the bony tympanic ring or meatal wall which may underlie granulations.

Myringotomy or removal of foreign bodies can be undertaken with surprisingly high precision because the optics of the Ear Microscope and the specially angled myringotome blade eliminate all shadows so that the surgeon can see clearly where and what he is cutting. The outfit is invaluable also for post-operative examinations and treatment of mastoid cavities. Granulations can be cauterized and exposed areas of dura mater carefully palpated.

**J. Laryng.* (1953) 67, 108.


39 WIGMORE STREET
LONDON W.1



5241 WHITBY AVENUE
PHILADELPHIA 43, Pa.,
U.S.A.

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

AMPLIVOX



F W N

MODEL 81 AUDIOMETER

For the first time it is now possible, with a compact and simple audiometer, to obtain accurate and reliable air and bone conduction measurements—even in cases of severe unilateral deafness.

This exclusive-featured audiometer is Amplivox Model 81, the outcome of many years' research and experience in the field of audiometry. In its development, considerable emphasis has been placed on simplicity of operation and long-term stability.

- * Filtered White Noise masking excludes all unwanted frequencies except the critical band around the pure tone frequency in use. This permits high masking intensities to be used without discomfort and fatigue to the subject under test.

Only Amplivox Model 81 Audiometer has these features:

- * Threshold calibration to British or American Standards.
- * Filtered White Noise masking tone to 5% band width above and below frequency being tested.
- * Insert-type earphone for masking increases inter-aural attenuation to 90dB.
- * Indicator light signals when controls are wrongly set for a particular test—also monitors speech level.

All enquiries to:

AMPLIVOX LIMITED

Medical Acoustic Division,
80 New Bond Street, London, W.1.
Telephone: HYD 9888

Please mention *The Journal of Laryngology and Otology* when replying to advertisements

The Journal of Laryngology and Otology

CONTENTS

	PAGE
ACOUSTIC-VESTIBULAR AND HISTOLOGIC EXAMINATIONS IN GUINEA-PIGS AFTER INTERRUPTION OF MEMBRANOUS LABYRINTH IN SEMICIRCULAR CANALS OR COCHLEA. Harald K. Kristensen (Copenhagen)	699
CYLINDROMAS OF THE UPPER RESPIRATORY TRACT. W. H. Struben and J. F. Hampe (Amsterdam)	722
A METHOD FOR STUDYING THE VIBRATORY MOVEMENTS OF THE VOCAL CORDS. Bertil Sonesson (Lund, Sweden)	732
A NEW TECHNIQUE IN STAPES MOBILIZATION. Bruce Cornish (Auckland, New Zealand)	738
SOME LONG-TERM RESULTS OF THE PATTERSON EXTERNAL ETHMOIDECTOMY. R. G. Williams (Wolverhampton)	745
CLINICAL RECORDS—	
Intra-Pharyngeal Hæmorrhage from the Internal Carotid Artery, Report of a Case and Arteriographic Findings. T. B. Hogarth (Leeds)	764
A Huge Trilobed Laryngeal Cyst. Aly El Mofty (Cairo)	768
An Unusual Case of Carcinoma of the Larynx. M. Feinmesser and I. Gay (Jerusalem)	772
SOCIETIES' PROCEEDINGS—	
The Scottish Otolaryngological Society	775
GENERAL NOTES	777

THE UNIVERSITY OF LIVERPOOL

Applications are invited for the post of Lecturer in Oto-Rhino-Laryngology. The initial salary will be within the range £1,400 to £1,600 per annum, according to qualifications and experience.

Applications, stating age, academic qualifications and experience, together with the names of three referees, should be received not later than Friday, February 12th, 1960, by the Registrar, from whom further particulars of the conditions of appointment may be obtained. Quoting Ref. CV/JL.