

it is important to acknowledge that hyperthyroidism can also lead to a rare but significant complication: psychosis. Psychosis in the context of hyperthyroidism refers to the presence of delusions, hallucinations, and disordered thinking, which can significantly impact an individual's mental health and overall well-being.

Objectives: This case report aims to describe a rare case of hyperthyroidism-related psychosis in a patient including the clinical presentation, diagnosis, and management. Additionally, we aim to increase awareness of and promote further research into this condition.

Methods: We present a comprehensive case report detailing the clinical course of a 29-year-old male patient with no previous medical or psychiatric history, who sought urgent psychiatric evaluation at the Razi Hospital La Manouba's emergency department due to escalating symptoms of agitation and paranoia persisting for three days. The patient, identified as Mr. S.O., a Tunisian male, presented with severe agitation and paranoia necessitating the use of restraints upon admission to the psychiatric emergency department. The initial physical examination revealed no notable abnormalities, except for the presence of tachycardia, which was subsequently confirmed on an electrocardiogram, arousing suspicion of a primary psychiatric illness.

Results: While the standard blood workup yielded unremarkable findings, the endocrine workup revealed decreased levels of thyroid-stimulating hormone (TSH) and elevated free thyroxine (FT4). Further laboratory investigations demonstrated elevated anti-thyroid-stimulating hormone receptor antibodies, leading to the diagnosis of Graves' disease. Collaborative consultation with an endocrinologist resulted in the initiation of a treatment regimen consisting of methimazole, propranolol, and risperidone. Notably, within three days of the initiated therapy, the patient exhibited significant improvement in terms of reduced agitation, coherent speech, and the development of self-reflection regarding the episode, ultimately leading to his discharge on the seventh day of hospitalization. This case report serves to highlight the complexity of psychiatric presentations associated with underlying endocrine disorders and underscores the importance of interdisciplinary collaboration in achieving optimal patient outcomes.

Conclusions: While mental health factors play a significant role in the development of psychosis, it is essential to recognize that underlying medical conditions may also contribute to its onset or exacerbation.

Disclosure of Interest: None Declared

EPP0156

Sociodemographic and Clinical Profile of Perpetrators of conjugal Homicides

W. Haouari*, S. Omri, A. Labyadh, R. Feki, I. Gassara, L. Zouari, J. Ben Thabet, N. Charfi, M. B. Maalej, N. Smaoui and M. Maalej
Psychiatrie C, Hedi Chaker university Hospital, Sfax, Tunisia

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.362

Introduction: Conjugal homicide refers to the act of killing a current or former intimate life partner, regardless of their marital status. This type of behavior is still inadequately addressed by prevention programs, as it is often regarded as exceptional compared to other forms of domestic violence.

Objectives: To describe the sociodemographic, clinical, and criminological data of individuals who commit spousal homicide.

Methods: This is a retrospective descriptive study of 21 psychiatric expertise files conducted between January 2002 and September 2023 in the psychiatric department of Hedi Chaker Hospital in Sfax. Sociodemographic, clinical, and criminological data were collected from criminal psychiatric expertise files and supplemented with information from medical records.

Results: The perpetrators of spousal homicide were predominantly male (85.7%), had an educational level above secondary school (57.1%), were married (85.7%), and had an average age at the time of the act of 40.3 years. Most of them had no psychiatric history (81%) or legal history (85%), and only 19% were using psychoactive substances.

Among the perpetrators, 66.7% had a history of violence against their partners, with threats of homicide in 19% of cases. The majority of homicides occurred during the day (42.9%), in a public place (28.6%), were perpetrated in isolation (95.2%), happened impulsively (47.6%), and employed a single method (81%), with knives being the primary weapon (42.9%). In 76.2% of cases, the perpetrators of spousal homicides were found criminally responsible, while 19% were hospitalized in a psychiatric setting as part of a judicial non-prosecution decision.

Conclusions: The study of specific characteristics of spousal homicides holds crucial importance for the early detection of domestic violence situations that carry a lethal risk. By highlighting these particularities, it enables the development of more targeted prevention strategies.

Disclosure of Interest: None Declared

EPP0157

Impact of social media on Working Memory and Academic Performance of Undergraduate Students- A Cross-sectional Study

N. E. Mgbeto¹, M. E. Musa^{1*}, P. Chhikara², R. Chaturvedi², N. Vyas³ and N. Zavrashvili¹

¹Medicine, University of Georgia; ²Medicine, Batumi State University and ³Medicine, Eastern European University, Tbilisi, Georgia

*Corresponding author.

doi: 10.1192/j.eurpsy.2024.363

Introduction: Over the course of the last decade social media has become a very important part of the human experience; it has become the main source of communication and entertainment for a lot of people young and old.

Objectives: This study aimed to examine the influence of social media on undergraduates working memory and academic performance. We assessed the association between the harmful use of social media on gender differences, physical activities, academic performance, and working memory.

Methods: This cross-sectional study was designed to examine the effect of social media on the working memory of undergraduate students from three different Universities in Georgia involving 722 participants. The collection survey form was distributed among Undergraduate students from the University of Georgia (UG), Eastern European University (EEU), and Batumi State University (BSU) through google forms from 14th June to 2nd July 2023. The