



A Note on p -adic Rankin–Selberg L -functions

David Loeffler

Abstract. We prove an interpolation formula for the values of certain p -adic Rankin–Selberg L -functions associated with non-ordinary modular forms.

1 Introduction

1.1 Background

Let f_1, f_2 be two modular eigenforms of weights $k_1 > k_2$. Then there is an associated Rankin–Selberg L -function $L(f_1, f_2, s)$, which is defined by a Dirichlet series $\sum c_n n^{-s}$ such that for ℓ prime we have $c_\ell = a_\ell(f) a_\ell(g)$.

If p is prime, and f_1 is *ordinary* at p , then a well-known construction due to Panichshkin [Pan82] and (independently) Hida [Hid85] gives rise to a p -adic Rankin–Selberg L -function $L_p(f_1, f_2, \sigma)$. This is a p -adic analytic function on the space \mathcal{W} of continuous characters of \mathbf{Z}_p^\times , with the property that if σ is a locally algebraic character $z \mapsto z^j \chi(z)$, with j in the critical range $k_2 \leq j \leq k_1 - 1$ and χ of finite order, then

$$L_p(f_1, f_2, \sigma) = (\star) \cdot L(f_1, f_2, \chi^{-1}, j),$$

where (\star) is an explicit factor. Hida [Hid88] subsequently showed that if f_2 is also ordinary, then $L_p(f_1, f_2, \sigma)$ extends to a 3-variable analytic function in which the forms f_1 and f_2 are allowed to vary in Hida families $\mathcal{F}_1, \mathcal{F}_2$. The existence of this p -adic L -function plays a major role in several recent works on arithmetic of Rankin–Selberg L -functions, in particular appearing in the explicit reciprocity law for the Euler system of Beilinson–Flach elements [BDR15a, BDR15b, KLZ17] (which is in turn crucial for several other recent works such as [BL16a, Cas15, Das16]).

It is natural to seek a generalisation of this construction to non-ordinary eigenforms, and variation in Coleman families. For fixed f_1 and f_2 of level prime to p and satisfying a suitable “small slope” hypothesis, such a construction was carried out by My [My91], but allowing variation in families has proved to be substantially more difficult. A construction of a 3-variable p -adic L -function with the expected interpolating property was initially announced in [Urb14], but an error in this construction was subsequently found, and (to the best of the author’s knowledge) this has not been fully resolved at the present time. (See “Note Added” at the end of this section.)

Received by the editors April 18, 2017; revised July 17, 2017.

Published electronically September 14, 2017.

The author’s research was supported by a Royal Society University Research Fellowship.

AMS subject classification: 11F85, 11F67, 11G40, 14G35.

Keywords: p -adic L -function, Iwasawa theory.

In the author's recent work with Zerbes [LZ16, Theorem 9.3.2], it was shown that there exists a 3-variable p -adic L -function with the expected interpolating property at *crystalline* points (i.e., where f_1 and f_2 are p -stabilisations of eigenforms of level prime to p , and χ is trivial). Moreover, this p -adic L -function is related by an explicit reciprocity law to the Euler system of Beilinson–Flach elements, as in the ordinary case. Unfortunately, we were not able to establish unconditionally that the p -adic L -function thus constructed also had the expected interpolation property at non-crystalline points, so our results fell short of giving a full proof of the results announced in [Urb14].

This gap in the published literature has become increasingly troublesome, since several papers have now been published that assume this stronger interpolation property; these include several papers making major contributions to famous open problems, such as the Iwasawa main conjecture for supersingular elliptic curves [BL16b, Wan15] and the Birch–Swinnerton–Dyer conjecture in analytic rank 1 [JSW15].

1.2 Aims of this Paper

The purpose of this note is to give a proof of an interpolation formula for the L -function of [LZ16] at all critical points, crystalline or otherwise, in a certain special case. The assumption we make is that the Coleman family \mathcal{F}_2 is ordinary, although \mathcal{F}_1 may not be; this suffices for the applications in the papers cited above (all of which correspond to the case where \mathcal{F}_2 is an ordinary family of CM-type). The author is cautiously optimistic that it might be possible to push these methods further in order to give a full proof of the results announced in [Urb14], but believes it is in the interests of the research community to release this partial proof without further delay, in order to place the already-published papers conditional on this result on a firm footing.

Our strategy will be to relate the 3-variable “geometric” p -adic L -function, constructed using Beilinson–Flach elements, with two families of “analytic” p -adic L -functions. These 2-variable functions, denoted here by superscripts \clubsuit and \diamond , are defined over 2-variable slices of the full 3-variable parameter space. Their construction involves nearly-overconvergent forms of a fixed degree, and therefore can be carried out using the methods of [Urb14] without the technical issues that arise when the degree of near-overconvergence is allowed to vary. The assumption that the second Coleman family \mathcal{F}_2 is ordinary implies that it is defined over an entire component of weight space; this gives sufficient “room” to move along \clubsuit and \diamond families from an arbitrary critical point to a crystalline one at which the results of [KLZ17] can be applied.

A secondary aim of this paper is to make the interpolation formula for the resulting p -adic L -function completely explicit, at least in the most important cases. This calculation is not new, but a precise statement of the formula seems to be difficult to find in the existing references (particularly in the non-crystalline cases); so we have given careful statements in Propositions 2.10 and 2.12 and an outline sketch of their proofs in an appendix.

Note Added During Review Since the initial version of this paper was released, the preprint [AI17] has appeared, which circumvents the problems with [Urb14] via a

new approach to nearly-overconvergent modular forms (as sections of a certain sheaf of Banach modules). This should in due course lead to a proof of an analogue of Theorem 6.3 of this paper for arbitrary pairs of Coleman families, without the restriction imposed here that \mathcal{F}_2 be ordinary. However, the author believes that there is still value in making this note available, since the preprint [AI17] has not yet been published, and the preliminary version of [AI17] seen by the author only considers families over the “centre” of weight space and thus does not cover most non-crystalline classical points.

2 Complex Rankin–Selberg L -functions and Period Integrals

2.1 The Complex L -function

Let k, k' be positive integers and let f_1, f_2 be two new, normalised cuspidal modular eigenforms of weights k_1, k_2 (and some levels N_1, N_2). We assume $k_1 \geq k_2$ without loss of generality.

Definition 2.1 The (imprimitive) Rankin–Selberg L -function of f_1 and f_2 is the Dirichlet series

$$L^{\text{imp}}(f_1, f_2, s) = L_{(N_1 N_2)}(\varepsilon_1 \varepsilon_2, 2s + 2 - k_1 - k_2) \cdot \sum_{n \geq 1} a_n(f_1) a_n(f_2) n^{-s}.$$

More generally, if χ is a Dirichlet character of conductor N_χ , we set

$$L^{\text{imp}}(f_1, f_2, \chi, s) = L_{(N_1 N_2 N_\chi)}(\varepsilon_1 \varepsilon_2 \chi^2, 2s + 2 - k_1 - k_2) \cdot \sum_{\substack{n \geq 1 \\ (n, N_\chi) = 1}} a_n(f_1) a_n(f_2) \chi(n) n^{-s}.$$

This L -function has an Euler product, in which the local factor for a prime $\ell \nmid N_1 N_2 N_\chi$ is given by $P_\ell(f_1, f_2, \chi(\ell) \ell^{-s})^{-1}$, where

$$P_\ell(f_1, f_2, X) = (1 - \alpha_1 \alpha_2 X)(1 - \alpha_1 \beta_2 X)(1 - \beta_1 \alpha_2 X)(1 - \beta_1 \beta_2 X).$$

Here, α_1, β_1 denote the roots of the polynomial $X^2 - a_\ell(f_1)X + \ell^{k_1-1} \varepsilon_1(\ell)$, and similarly for α_2, β_2 .

Remark 2.2 We refer to this L -function as an “imprimitive” L -function, since it differs by finitely many Euler factors from the L -function of the motive associated with $f_1 \otimes f_2 \otimes \chi$ (the “primitive” Rankin–Selberg L -function). The only primes ℓ at which the local Euler factors can differ are those ℓ dividing at least two of the three integers N_1, N_2, N_χ ; so if these are pairwise coprime, then the primitive and imprimitive L -functions coincide.

It is well known that $L^{\text{imp}}(f_1, f_2, \chi, s)$ has a meromorphic continuation to all $s \in \mathbf{C}$. It is entire unless $k_1 = k_2$ and $f_2 = f_1 \otimes \varepsilon_1^{-1} \chi^{-1}$, in which case there is a simple pole at $s = k_1$. The critical values are those in the interval $k_2 \leq s \leq k_1 - 1$.

2.2 A Petersson Product Formula

Now let p be prime and choose an embedding $\overline{\mathbf{Q}} \hookrightarrow \overline{\mathbf{Q}}_p$.

Definition 2.3 A locally algebraic character of \mathbf{Z}_p^\times is a homomorphism $\mathbf{Z}_p^\times \rightarrow \overline{\mathbf{Q}}_p^\times$ of the form $x \mapsto x^n \chi(x)$, where $n \in \mathbf{Z}$ and χ is a finite-order character (equivalently, a Dirichlet character of p -power conductor). We denote this character by $n + \chi$.

Definition 2.4 By a p -stabilised newform of tame level N , where N is an integer coprime to p , we shall mean a normalised cuspidal Hecke eigenform of level $\Gamma_1(Np^r)$, for some $r \geq 1$, such that either f is a newform, or f is a U_p -eigenform in the two-dimensional space of oldforms associated with some newform of level N . In the latter case, we say f is crystalline.

We define the weight-character of f to be the locally-algebraic character κ of \mathbf{Z}_p^\times defined by $\kappa = k + \varepsilon_p$, where k is the weight of f and ε_p is the p -part of the Nebentypus character of f .

If f is a p -stabilised newform, we denote by f^c the unique p -stabilised newform with the same weight-character as f satisfying

$$a_n(f^c) = \varepsilon_{N,f}(n)^{-1} a_n(f),$$

where $\varepsilon_{N,f}$ is the prime-to- p part of the Nebentypus of f , for all $(n, N) = 1$ (even if $p \mid n$).

Remark 2.5 Note that if f is a p -stabilised newform whose nebentypus is trivial at p , then f^c has the same Hecke eigenvalues away from p as the conjugate form f^* defined by $f^*(\tau) = \overline{f(-\bar{\tau})}$. However, f^c and f^* do not generally have the same U_p -eigenvalue; in particular, f^c is ordinary if f is (which is not true of f^*). On the other hand, if f has non-trivial character at p , then the Hecke eigenvalues of f^c and f^* away from p are different.

Let f_1, f_2 be p -stabilised newforms of some tame levels N_1, N_2 , and let $\kappa_1 = k_1 + \varepsilon_{1,p}, \kappa_2 = k_2 + \varepsilon_{2,p}$ be their weight-characters. We choose an integer N divisible by both N_1 and N_2 , and with the same prime factors as $N_1 N_2$. Given $\sigma = j + \chi$ a locally algebraic character, we consider the formal power series

$$\mathcal{E}_N(\kappa_1, \kappa_2, \sigma) := \sum_{\substack{n \geq 1 \\ p \nmid n}} \left(\sum_{d \mid n} d^{\sigma - \kappa_2} \left(\frac{n}{d}\right)^{\kappa_1 - \sigma - 1} \left[e^{2\pi i d/N} + (-1)^{\kappa_1 - \kappa_2} e^{-2\pi i d/N} \right] \right) q^n.$$

Lemma 2.6 If $1 \leq k_2 \leq j \leq k_1 - 1$, then $\mathcal{E}_N(\kappa_1, \kappa_2, \sigma)$ is the q -expansion of a nearly-holomorphic modular form of weight $k_1 - k_2$, level dividing Np^∞ , and degree at most $\min(k_1 - 1 - j, j - k_2)$, on which the diamond operators at p act via the character $\varepsilon_{1,p} - \varepsilon_{2,p}$.

Proof See [LLZ14, §5.3]. ■

If Π^{hol} denotes Shimura’s holomorphic projector, then the cuspidal modular form

$$\Pi^{\text{hol}}(f_2 \cdot \mathcal{E}_N(\kappa_1, \kappa_2, \sigma))$$

has level dividing Np^∞ , and its weight-character agrees with that of f_1 (and thus also of f_1^c).

Definition 2.7 Suppose f_1 has finite slope (that is, $a_p(f) \neq 0$). We let $\lambda_{f_1^c}$ denote the unique linear functional on $S_{k_1}(N_1 p^\infty, \bar{\epsilon}_{1,p})$, which factors through the Hecke eigenspace associated with f_1^c , and maps the normalised eigenform f_1^c itself to 1. We extend this to forms of tame level N by composing with the trace map.

Definition 2.8 We set

$$I(f_1, f_2, \sigma) = N^{\kappa_1 + \kappa_2 - 2\sigma - 2} \cdot \lambda_{f_1^c} \left(\Pi^{\text{hol}}(f_2 \cdot \mathcal{E}_N(\kappa_1, \kappa_2, \sigma)) \right).$$

Theorem 2.9 (Rankin–Selberg, Shimura) *If $1 \leq k_2 \leq j \leq k_1 - 1$, then we have*

$$I(f_1, f_2, j + \chi) = (\star) \cdot L^{\text{imp}}(f_1, f_2, \chi^{-1}, j),$$

where (\star) is an explicitly computable factor.

We shall not give the precise form of the factor (\star) in all possible cases, since this rapidly becomes messy, but we shall give a selection of useful cases. First, we treat the case where f_1 and f_2 are crystalline, hence p -stabilisations of forms f_1°, f_2° of levels N_1, N_2 coprime to p . We write α_i for the U_p -eigenvalue of f_i , so that α_i is a root of the Hecke polynomial of f_i° at p , and β_i for the other root of this polynomial. We assume¹ that $\alpha_1 \neq \beta_1$.

We define certain local Euler factors at p , as in [BDR15a] and [KLZ17, Theorem 2.7.4], by

$$\begin{aligned} \mathcal{E}(f_1) &= \left(1 - \frac{\beta_1}{p\alpha_1}\right), & \mathcal{E}^*(f_1) &= \left(1 - \frac{\beta_1}{\alpha_1}\right), \\ \mathcal{E}(f_1, f_2, j + \chi) &= \\ &\begin{cases} \left(1 - \frac{p^{j-1}}{\alpha_1\alpha_2}\right)\left(1 - \frac{p^{j-1}}{\alpha_1\beta_2}\right)\left(1 - \frac{\beta_1\alpha_2}{p^j}\right)\left(1 - \frac{\beta_1\beta_2}{p^j}\right) & \text{if } \chi = 1, \\ G(\chi)^2 \cdot \left(\frac{p^{2s-2}}{\alpha_1^2\alpha_2\beta_2}\right)^r & \text{if } \chi \text{ has conductor } p^r > 1. \end{cases} \end{aligned}$$

Here $G(\chi)$ is the Gauss sum $\sum_{a \in (\mathbb{Z}/p^r\mathbb{Z})^\times} \chi(a) e^{2\pi i a/p^r}$.

Proposition 2.10 *In the above setting, we have*

$$I(f_1, f_2, j + \chi) = \frac{\mathcal{E}(f_1, f_2, j)}{\mathcal{E}(f_1)\mathcal{E}^*(f_1)} \cdot \frac{(j-1)!(j-k_2)! i^{k_1-k_2}}{\pi^{2j+1-k_2} 2^{2j+k_1-k_2} \langle f_1^\circ, f_1^\circ \rangle_{N_1}} L^{\text{imp}}(f_1^\circ, f_2^\circ, \chi^{-1}, j).$$

Remark 2.11 For χ trivial, this formula is standard, and its derivation can be found in many references such as [BDR15a, LLZ14, LZ16]. For χ non-trivial, references are more scant; many sources, such as [Hid88], give more general but less explicit formulas, and the work involved in recovering a completely explicit form for all the local factors is routine but unpleasant. For the convenience of the reader, we give, in an appendix to this paper, an account of the main steps required in this case to evaluate $I(f_1, f_2, j + \chi)$.

¹This assumption is known to be true if $k_1 = 2$, and is known to follow from the Tate conjecture if $k_1 \geq 3$ [CE98].

The other case we shall consider is that where f_1 is still assumed crystalline, but f_2 has some non-trivial character $\varepsilon_{2,p}$ at p , and neither χ nor $\chi' = \chi\varepsilon_{2,p}^{-1}$ is trivial. We define $\beta_2 = p^{k_2-1}\varepsilon_{2,N}(p)/\alpha_2$, and we let the conductor of χ (resp. χ') be p^r (resp. $p^{r'}$).

Proposition 2.12 *In this setting, we have*

$$I(f_1, f_2, j + \chi) = \left(\frac{p^{j-1}}{\alpha_1\alpha_2}\right)^r G(\chi) \left(\frac{p^{j-1}}{\alpha_1\beta_2}\right)^{r'} G(\chi') \\ \times \frac{(j-1)!(j-k_2)!i^{k_1-k_2}}{\mathcal{E}(f_1)\mathcal{E}^*(f_1)\pi^{2j+1-k_2}2^{2j+k_1-k_2}\langle f_1^\circ, f_1^\circ \rangle_{N_1}} L^{\text{imp}}(f_1^\circ, f_2, \chi^{-1}, j).$$

3 Overconvergent Families

Let us fix a finite extension L/\mathbf{Q}_p (contained in our fixed choice of algebraic closure $\overline{\mathbf{Q}_p}$).

Definition 3.1 Let the *weight space*, \mathcal{W} , be the rigid-analytic space over L parametrising continuous characters of \mathbf{Z}_p^\times , so that for an affinoid L -algebra A , we have $\mathcal{W}(A) = \text{Hom}(\mathbf{Z}_p^\times, A^\times)$.

As in [KLZ17], we identify both \mathbf{Z} and the set of Dirichlet characters of p -power order with subsets of $\mathcal{W}(\bar{L})$ in the natural fashion, and we denote the group law on \mathcal{W} additively. If $\kappa = k + \chi$ is a locally algebraic character, we write $w(\kappa) := k$.

Now let N be an integer coprime to p . It will be convenient to assume that L contains the N -th roots of unity; let $\zeta_N \in L^\times$ denote the image of $e^{2\pi i/N} \in \overline{\mathbf{Q}}$ under our chosen embedding.

Lemma 3.2 *The power series in $E_{\mathbf{k}}^{[p]}$ and $F_{\mathbf{k}}^{[p]}$ in $\mathcal{O}(\mathcal{W})[[q]]$ given by*

$$E_{\mathbf{k}}^{[p]} := \sum_{\substack{n \geq 1 \\ p \nmid n}} \left(\sum_{d|n} d^{\mathbf{k}-1} (\zeta_N^d + (-1)^{\mathbf{k}} \zeta_N^{-d}) \right) q^n, \\ F_{\mathbf{k}}^{[p]} := \sum_{\substack{n \geq 1 \\ p \nmid n}} \left(\sum_{d|n} \binom{n}{d}^{\mathbf{k}-1} (\zeta_N^d + (-1)^{\mathbf{k}} \zeta_N^{-d}) \right) q^n$$

are both the q -expansions of families of overconvergent modular forms over \mathcal{W} of tame level $\Gamma_1(N)$ and weight \mathbf{k} (with radius of overconvergence bounded below over any affinoid in \mathcal{W}).

Lemma 3.3 *Let χ be a Dirichlet character of p -power conductor, with values in L . Then, for any family of overconvergent modular forms \mathcal{F} of tame level $\Gamma_1(N)$ and weight $\kappa : \mathbf{Z}_p^\times \rightarrow A^\times$, where A is an affinoid algebra, the power series defined by*

$$\theta^\chi \mathcal{F} := \sum_{\substack{n \geq 1 \\ p \nmid n}} a_n(\mathcal{F}) \chi(n) q^n$$

is the q -expansion of a family of overconvergent forms over A , of weight $\kappa + 2\chi$.

Sketch of proof Let χ have conductor p^r . Then there is a “twisting homomorphism” $t_j: X_1(Np^{2r}) \rightarrow X_1(N)$, given in terms of complex uniformizations by $\tau \mapsto \tau + \frac{j}{p^r}$, for any $j \in \mathbf{Z}/p^r\mathbf{Z}$. This preserves the component of the ordinary locus containing ∞ and extends to all sufficiently small overconvergent neighbourhoods of it, so it induces a pullback map on overconvergent modular (or cusp) forms. Since $\theta^\chi \mathcal{F}$ is equal to $\sum_{j \in (\mathbf{Z}/p^r\mathbf{Z})^\times} \chi(j)^{-1} t_j^*(\mathcal{F})$ up to a constant, it is overconvergent of level $\Gamma_1(Np^{2r})$ and weight-character κ ; and the diamond operators at p act on it via χ^2 , so it descends to an overconvergent form of level $\Gamma_1(N) \cap \Gamma_0(p^{2r})$ and weight $\kappa + 2\chi$. Via the canonical-subgroup map, we can regard it as an overconvergent form of level N . ■

In order to allow more general twists, we work with families of nearly-overconvergent modular forms (of some finite degree $r \geq 0$), in the sense of [Urb14, §3.3.2]. If τ is a locally algebraic weight with $w(\tau) \geq 0$, we can define $\theta^\tau(\mathcal{F})$ as a family of nearly-overconvergent forms of weight $\kappa + 2\tau$ and degree $w(\tau)$.

Lemma 3.4 *If \mathcal{F} is a Coleman family (a family of overconvergent normalised eigenforms of finite slope), new of some tame level N , defined over some affinoid $A \rightarrow \mathcal{W}$, then there is a unique tame level N Coleman family \mathcal{F}^c over A satisfying*

$$a_n(\mathcal{F}^c) = \varepsilon_N(n)^{-1} a_n(\mathcal{F})$$

for all $(n, N) = 1$ (including $n = p$). Here, $\varepsilon_N: (\mathbf{Z}/N\mathbf{Z})^\times \rightarrow L^\times$ is the prime-to- p nebentype of \mathcal{F} .

Proof This is proved in the same way as the previous lemma. ■

We now recall the construction of the universal object parametrising Coleman families: the eigencurve.

Definition 3.5 Let \mathcal{C}_N denote the Coleman–Mazur–Buzzard cuspidal eigencurve, of tame level N .

By definition, \mathcal{C}_N is a reduced rigid space, equidimensional of dimension 1, equipped with a morphism $\mathcal{C}_N \rightarrow \mathcal{W}$, and there is a universal eigenform over \mathcal{C}_N ; that is, \mathcal{C}_N comes equipped with a power series $\mathcal{F}^{\text{univ}} = \sum a_n q^n \in \mathcal{O}(\mathcal{C}_N)[[q]]$, with $a_1 = 1$ and a_p invertible on \mathcal{C}_N , with the following universal property:

For any affinoid X with a weight morphism $\kappa: X \rightarrow \mathcal{W}$, and any family of finite-slope eigenforms \mathcal{F}_X over X of tame level N and weight κ , there is a unique morphism $X \rightarrow \mathcal{C}_N$ lifting κ such that \mathcal{F}_X is the pullback of $\mathcal{F}^{\text{univ}}$.

4 Two-variable p -adic L -functions

Let U_1 and U_2 be two affinoid subdomains of \mathcal{W} . We write $\mathbf{k}_i: \mathbf{Z}_p^\times \rightarrow \mathcal{O}(U_i)^\times$ for the pullbacks of the canonical character \mathbf{k} . We suppose that we are given the following data:

- a finite flat covering $\tilde{U}_2 \rightarrow U_2$,

- an overconvergent family $\mathcal{F}_2 \in M_{\mathbf{k}_2}^\dagger(\Gamma_1(N); \tilde{U}_2)$ (not necessarily cuspidal or normalised),
- a locally analytic character $\tau \in \mathcal{W}(L)$, with $t = w(\tau) \geq 0$.

We define two families of nearly-overconvergent forms over $U_1 \times \tilde{U}_2$, both of weight \mathbf{k}_1 and degree of near-overconvergence $\leq t$, by

$$\begin{aligned} \Xi_\tau^\spadesuit &:= \mathcal{F}_2 \cdot \theta^\tau(E_{\mathbf{k}_1 - \mathbf{k}_2 - 2\tau}^{[p]}), \\ \Xi_\tau^\diamond &:= \mathcal{F}_2 \cdot \theta^\tau(F_{\mathbf{k}_1 - \mathbf{k}_2 - 2\tau}^{[p]}). \end{aligned}$$

We apply to both of these forms the overconvergent projector Π^{oc} of [Urb14, §3.3.4]. This gives elements

$$\Pi^{\text{oc}}(\Xi_\tau^\spadesuit), \Pi^{\text{oc}}(\Xi_\tau^\diamond) \in \frac{1}{\prod_{m=2}^{2t} (\nabla_1 - m)} S_{\mathbf{k}_1}^\dagger(\Gamma_1(N), U_1 \times \tilde{U}_2),$$

where $\nabla_1 \in \mathcal{O}(U_1)$ is the pullback to U_1 of the unique rigid-analytic function $\nabla \in \mathcal{O}(W)$ such that $\nabla(\kappa) = w(\kappa)$ for all locally-algebraic κ .

Proposition 4.1 *Let (κ_1, κ_2) be a locally-algebraic point of $U_1 \times U_2$ such that $1 \leq k_2 \leq k_1 - 1 - t$, where $k_i = w(\kappa_i)$, and with $k_1 \notin \{2, \dots, 2t\}$. Let $\tilde{\kappa}_2$ be a point of \tilde{U}_2 above κ_2 and let f_2 be the specialisation of \mathcal{F}_2 at $\tilde{\kappa}_2$. Let us suppose that f_2 is a classical modular form.*

Then the specialisations of $\Pi^{\text{oc}}(\Xi_\tau^\spadesuit)$ and $\Pi^{\text{oc}}(\Xi_\tau^\diamond)$ at $(\kappa_1, \tilde{\kappa}_2)$ are given by

$$\begin{aligned} \Pi^{\text{oc}}(\Xi_\tau^\spadesuit)(\kappa_1, \tilde{\kappa}_2) &= \Pi^{\text{hol}}(f_2 \cdot \mathcal{E}_N(\kappa_1, \kappa_2, \kappa_1 - 1 - \tau)), \\ \Pi^{\text{oc}}(\Xi_\tau^\diamond)(\kappa_1, \tilde{\kappa}_2) &= \Pi^{\text{hol}}(f_2 \cdot \mathcal{E}_N(\kappa_1, \kappa_2, \kappa_2 + \tau)). \end{aligned}$$

Proof An elementary computation shows that $\theta^\tau(E_{\mathbf{k}_1 - \mathbf{k}_2 - 2\tau}^{[p]}) = \mathcal{E}_N(\kappa_1, \kappa_2, \kappa_1 - 1 - \tau)$ and similarly that $\theta^\tau(F_{\mathbf{k}_1 - \mathbf{k}_2 - 2\tau}^{[p]}) = \mathcal{E}_N(\kappa_1, \kappa_2, \kappa_2 + \tau)$. The result now follows from the compatibility of the holomorphic and overconvergent projection operators. ■

Remark 4.2 We can consider the formal power series $\mathcal{F}_2 \cdot \mathcal{E}_N(\mathbf{k}_1, \mathbf{k}_2, \sigma)$ as a family of p -adic modular forms over $U_1 \times \tilde{U}_2 \times \mathcal{W}$. This is not overconvergent, or even nearly-overconvergent, in any reasonable sense, since the near-overconvergence degrees of its specialisations are not bounded above over any open affinoid in the parameter space $U_1 \times \tilde{U}_2 \times \mathcal{W}$. However, Proposition 4.1 gives two families of 2-dimensional “slices” of the parameter space for which the above family does become nearly-overconvergent, of bounded degree, over any given slice.

Let us now suppose that $k_1 \geq 2$ is a non-negative integer lying in U_1 , N_f is an integer dividing N , and $f_1 \in S_{k_1}(\Gamma_1(N_f) \cap \Gamma_0(p), L)$ is a “noble eigenform” in the sense of [LZ16, Definition 4.6.3]; that is, f_1 is a p -stabilisation of some normalised newform of level $\Gamma_1(N_f)$ whose Hecke polynomial at p has distinct roots, and a mild extra condition is satisfied in the case of critical-slope eigenforms.

Then, after possibly shrinking the affinoid neighbourhood $U_1 \ni k_1$, we can find a Coleman family of normalised eigenforms \mathcal{F}_1 over U_1 whose specialisation at k_1 is f_1 ;

and a continuous $\mathcal{O}(U_1)$ -linear functional

$$\lambda_{\mathcal{F}_1^c}: \mathcal{S}_{\mathbf{k}_1}^\dagger(\Gamma_1(N_f), U_1) \rightarrow \mathcal{O}(U_1)$$

factoring through the Hecke eigenspace associated with the dual family \mathcal{F}_1^c , and mapping the normalised eigenform \mathcal{F}_1^c itself to 1. We extend this to a linear functional on forms of level N by composing with the trace map. We can therefore define two meromorphic functions, both lying in the space $\frac{1}{\prod_{j=2}^{2w(\tau)}(\nabla_1-j)}\mathcal{O}(U_1 \times \tilde{U}_2)$, by the formulae

$$\begin{aligned} L_p^\star(\mathcal{F}_1, \mathcal{F}_2; \tau) &= N^{(-\mathbf{k}_1+\mathbf{k}_2+2\tau)} \lambda_{\mathcal{F}_1^c}[\Pi^{\text{oc}}(\Xi_\tau^\star)], \\ L_p^\diamond(\mathcal{F}_1, \mathcal{F}_2; \tau) &= N^{(\mathbf{k}_1-\mathbf{k}_2-2\tau-2)} \lambda_{\mathcal{F}_1^c}[\Pi^{\text{oc}}(\Xi_\tau^\diamond)]. \end{aligned}$$

By construction, L_p^\star interpolates the values $I(f_1, f_2, \kappa_1 - 1 - \tau)$, and L_p^\diamond the values $I(f_1, f_2, \kappa_2 + \tau)$, for varying f_1 and f_2 (but fixed τ).

Remark 4.3 Our eventual goal is to show that there is a 3-variable L -function on $U_1 \times \tilde{U}_2 \times \mathcal{W}$ interpolating all critical values of the Rankin L -function. The 2-variable L -functions L_p^\star and L_p^\diamond will turn out to be slices of this 3-variable L -function, along two different families of 2-dimensional subspaces of the parameter space.

Let us, finally, specialise to the case where \tilde{U}_2 is an affinoid subdomain of the eigen-curve \mathcal{C}_{N_2} , and \mathcal{F}_2 is the universal eigenform. One knows that \mathcal{C}_{N_2} is admissibly covered by affinoids \tilde{U}_2 with the property that \tilde{U}_2 is a finite flat covering of an admissible open in \mathcal{W} , as above, and the above construction is clearly compatible on overlaps, so we obtain two families of meromorphic functions on $U_1 \times \mathcal{C}_{N_2}$.

5 Compatibility of the Two Families

Definition 5.1 Given a locally algebraic τ with $w(\tau) \geq 0$, we define two 2-dimensional rigid-analytic subspaces of $U_1 \times \tilde{U}_2 \times \mathcal{W}$ by

$$\begin{aligned} \mathcal{W}^\star(\tau) &= \{(\kappa_1, \tilde{\kappa}_2, \kappa_1 - 1 - \tau) : \kappa_1 \in U_1, \tilde{\kappa}_2 \in U_2\}, \\ \mathcal{W}^\diamond(\tau) &= \{(\kappa_1, \tilde{\kappa}_2, \kappa_2 + \tau) : \kappa_1 \in U_1, \tilde{\kappa}_2 \in U_2\}. \end{aligned}$$

We set $\Sigma_{\text{crit}}^\star(\tau) = \Sigma_{\text{crit}} \cap \mathcal{W}^\star(\tau)$ and, similarly, $\Sigma_{\text{geom}}^\star(\tau), \Sigma_{\text{crit}}^\diamond(\tau), \Sigma_{\text{geom}}^\diamond(\tau)$.

We can then regard $L_p^\star(\mathcal{F}_1, \mathcal{F}_2; \tau)$ as a p -adic meromorphic function on $\mathcal{W}^\star(\tau)$ in a natural way, interpolating classical L -values at the points in $\Sigma_{\text{crit}}^\star(\tau)$, and similarly for \diamond .

We have the following technical lemma.

Lemma 5.2 Let τ, τ' be two locally-algebraic characters with $w(\tau) \geq 0, w(\tau') \geq 0$, and suppose that we have

$$\{\kappa - (1 + \tau + \tau') : \kappa \in U_1\} \subseteq U_2.$$

Then $L_p^\star(\mathcal{F}_1, \mathcal{F}_2; \tau)$ and $L_p^\diamond(\mathcal{F}_1, \mathcal{F}_2; \tau')$ coincide as functions on $\mathcal{W}^\star(\tau) \cap \mathcal{W}^\diamond(\tau')$.

Proof The intersection $\mathcal{W}^\bullet(\tau) \cap \mathcal{W}^\diamond(\tau')$ consists of those points of the form $(\kappa_1, \tilde{\kappa}_2, \kappa_1 - 1 - \tau)$ such that $\tilde{\kappa}_2$ lies above the point $\kappa_1 - (1 + \tau + \tau')$ of \mathcal{W} . In particular, under the assumptions of the lemma, this is simply a finite covering of U_1 .

Let $(\kappa_1, \tilde{\kappa}_2, \sigma)$ be a point in this intersection with κ_1 locally algebraic, and such that $w(\kappa_1) \geq 2 \max(w(\tau), w(\tau')) + 1$ in order to avoid singularities of the nearly-overconvergent projection operators. Then the two p -adic L -functions specialise to the image under $\lambda_{f_1^c}$ of the nearly-overconvergent modular forms with q -expansions

$$f_2 \theta^\tau (E_{\kappa_1 - \kappa_2 - 2\tau}^{[p]}) \quad \text{and} \quad f_2 \theta^{\tau'} (E_{\kappa_1 - \kappa_2 - 2\tau'}^{[p]}).$$

Since these two modular forms are identical, we deduce that the two L -functions agree at the given point. As the set of locally-algebraic $\kappa_1 \in U_1$ with $w(\kappa_1)$ greater than any given bound is clearly Zariski-dense, it follows that the two p -adic L -functions are identically equal on this intersection. ■

Lemma 5.3 *Let τ be a locally algebraic character with $w(\tau) \geq 0$. If U_2 is sufficiently large (depending on U_1 and τ), then the union of the intersections $\mathcal{W}^\bullet(t) \cap \mathcal{W}^\diamond(\tau)$, as t varies over integers ≥ 0 , is Zariski dense in $\mathcal{W}^\diamond(\tau)$.*

Proof This is easy to check. ■

6 The 3-variable Geometric L -function

We now turn from “ p -adic analytic” methods to “arithmetic” ones; that is, we invoke the existence of the Euler system of Beilinson–Flach elements.

Theorem 6.1 *Suppose \tilde{U}_2 is the preimage of U_2 in the ordinary locus of the eigencurve, and \mathcal{F}_2 the universal ordinary family over U_2 . Then there exists a p -adic meromorphic² function $L_p^{\text{geom}}(\mathcal{F}_1, \mathcal{F}_2)$ on $U_1 \times \tilde{U}_2 \times \mathcal{W}$ with the following property:*

(†) *For any crystalline character $\tau = t$ with $t \geq 0$, the 2-variable p -adic L -function $L_p^\bullet(\mathcal{F}_1, \mathcal{F}_2; \tau)$ is the restriction of L_p^{geom} to $\mathcal{W}^\bullet(\tau)$.*

Moreover, L_p^{geom} is related to the Euler system of Beilinson–Flach elements via the formula

$$L_p^{\text{geom}}(\mathcal{F}_1, \mathcal{F}_2) = (c^2 - \varepsilon_{N,1}(c)^{-1} \varepsilon_{N,2}(c)^{-1} c^{2s+2-k_1-k_2})^{-1} (-1)^s \lambda(\mathcal{F}_1)^{-1} \langle {}_c \mathcal{B} \mathcal{F}^{[\mathcal{F}_1, \mathcal{F}_2]}, \eta_{\mathcal{F}_1} \otimes \omega_{\mathcal{F}_2} \rangle$$

in the notation of [LZ16, §9.1], for any $c > 1$ coprime to $6pN_1N_2$.

Proof The only difference in our present statement is that we are allowing U_2 to be arbitrary, and permitting some finite flat covering $\tilde{U}_2 \rightarrow U_2$, whereas in our earlier work we assumed that both U_1 and U_2 were small neighbourhoods of some given

²It is analytic if the product of the prime-to- p nebentypus characters of \mathcal{F}_1 and \mathcal{F}_2 is non-trivial. Otherwise, it may have poles along the near-central points $(\kappa_1, \kappa_2, \sigma)$ such that $\kappa_1 + \kappa_2 = 2\sigma$. This is a consequence of the “smoothing factors” $c^2 - c^2$ appearing in the construction of the Beilinson–Flach elements. In particular, the restriction of L_p^{geom} to any \diamond or \clubsuit slice is well defined.

eigenforms f_1, f_2 . However, the latitude to shrink U_2 was only used in *op.cit.* at precisely two points:

- in the proof of Proposition 5.3.4 of *op.cit.*, in order to arrange that all specialisations of \mathcal{F}_2 at points of classical weight were classical; this is automatically satisfied for ordinary families.
- in Sections 6.3 and 6.4 of *op.cit.*, in order to find a triangulation of the (φ, Γ) -module associated with \mathcal{F}_2 , and canonical crystalline periods for the filtration steps; this can be carried out globally over an ordinary family, using Ohta’s results [Oht00], as in [KLZ17]. ■

In order to complete the proof, we shall manoeuvre from the rather weak interpolating property (\dagger) of L_p^{geom} into a much stronger one by repeatedly using the compatibility between the \spadesuit and \diamond slices.

Corollary 6.2 *Let τ be any locally-algebraic character (not necessarily crystalline) with $w(\tau) \geq 0$. If U_2 is sufficiently large (depending on U_1 and τ), then*

$$L_p^\diamond(\mathcal{F}_1, \mathcal{F}_2; \tau) = L_p^{\text{geom}}(\mathcal{F}_1, \mathcal{F}_2)|_{\mathcal{W}^\diamond(\tau)},$$

$$L_p^\spadesuit(\mathcal{F}_1, \mathcal{F}_2; \tau) = L_p^{\text{geom}}(\mathcal{F}_1, \mathcal{F}_2)|_{\mathcal{W}^\spadesuit(\tau)}.$$

Proof By Lemma 5.3, for the first equality, it suffices to show that L_p^\diamond and L_p^{geom} agree on the intersection $\mathcal{W}^\spadesuit(t) \cap \mathcal{W}^\diamond(\tau)$, for integers $t \geq 0$. However, we know that L_p^\diamond and L_p^\spadesuit coincide on these intersections and that L_p^{geom} in turn coincides with L_p^\spadesuit .

For the second equality, we consider the intersection of $\mathcal{W}^\spadesuit(\tau)$ with the slices $\mathcal{W}^\diamond(\tau')$, where τ' is an arbitrary locally-algebraic character of weight $w(\tau') \geq 0$. Using the previously-proved equality, we know that L_p^{geom} agrees with $L_p^\spadesuit(\mathcal{F}_1, \mathcal{F}_2; \tau)$ on each of these intersections. As before, the union of these is Zariski dense in $\mathcal{W}^\spadesuit(\tau)$, as required. ■

We conclude, finally, the following interpolation formula. Recall that we are assuming \mathcal{F}_2 to be an ordinary family.

Theorem 6.3 *Let $(\kappa_1, \tilde{\kappa}_2, \sigma)$ be a triple of locally-algebraic points in $U_1 \times \tilde{U}_2 \times \mathcal{W}$, with $1 \leq w(\kappa_2) \leq w(\sigma) \leq w(\kappa_1) - 1$. Let f_1, f_2 be the specialisations of $\mathcal{F}_1, \mathcal{F}_2$ at the weights κ_i , and suppose that these specialisations are classical.*

Then we have

$$L_p^{\text{geom}}(\mathcal{F}_1, \mathcal{F}_2)(\kappa_1, \tilde{\kappa}_2, \sigma) = I(f_1, f_2, \sigma).$$

Proof Given any such triple, let us write $\tau = \kappa_1 - 1 - \sigma$ and $\tau' = \sigma - \kappa_2$. Both of these are locally algebraic characters, and $w(\tau), w(\tau') \geq 0$.

Since $w(\tau) + w(\tau') = w(\kappa_1) - 1 - w(\kappa_2)$, at least one of the quantities $w(\tau)$ and $w(\tau')$ must be $\leq \frac{w(\kappa_1)-1}{2}$. If $w(\tau) \leq \frac{w(\kappa_1)-1}{2}$, then $(\kappa_1, \tilde{\kappa}_2, \sigma)$ lies in the interval in which $L_p^\spadesuit(\mathcal{F}_1, \mathcal{F}_2; \tau)$ interpolates the classical Rankin–Selberg period. Similarly, if $w(\tau')$ is smaller than this bound, we can invoke the interpolating property of L_p^\diamond .

Since \mathcal{F}_2 is an ordinary family, we can assume without loss of generality that U_2 is arbitrarily large, and via the previous theorem, we can conclude that L_p^\star or L_p^\diamond coincides with the appropriate specialisation of the 3-variable p -adic L -function. ■

A Evaluation of the Rankin–Selberg Period

For the convenience of the reader, we outline the derivation of the formula relating the period $I(f_1, f_2, \sigma)$ defined above to the Rankin–Selberg L -function. Our approach is closely based on that of [PR88]. We place ourselves in the setting of Proposition 2.10, and, since the case of trivial χ is covered in many references, we will assume that χ is non-trivial, of conductor p^r with $r \geq 1$.

Step 1 We express the linear functional $\lambda_{f_1^c}$ on $S_k(\Gamma_1(N) \cap \Gamma_0(p^n))$, for any $n \geq 1$, via the formula

$$\lambda_{f_1^c}(h) = \left(\frac{\varepsilon_1(p)}{\alpha_1} \right)^{n-1} \cdot \frac{\langle g_n, h \rangle_{N(p^n)}}{\langle g, f_1^c \rangle_{N_1(p)}}$$

where $g = W_{N_1 p}(f_{1,\beta})$ and $g_n = g|_k \left(\begin{smallmatrix} p^{n-1} & \\ & 1 \end{smallmatrix} \right)$. Here $f_{1,\beta}$ is the p -stabilisation of f_1° corresponding to the root β_1 of the Hecke polynomial; and the subscript $N(p^n)$ denotes the Petersson product at level $\Gamma_1(N) \cap \Gamma_0(Np^n)$. Cf. [Hid85, Proposition 4.5]. A computation closely analogous to the final step of [KLZ17, Proposition 10.1.1] shows that the denominator term is given by

$$\langle g, f_1^c \rangle_{N_1(p)} = \frac{\overline{\lambda(f_1^\circ)} \alpha \mathcal{E}(f_1) \mathcal{E}^*(f_1)}{\varepsilon_1(p)} \cdot \langle f_1^\circ, f_1^\circ \rangle_{N_1},$$

where $\lambda(f_1^\circ)$ denotes the Atkin–Lehner pseudo-eigenvalue of f_1° . This yields the formula

$$I(f_1, f_2, j + \chi) = \frac{\varepsilon_1(p)^{2r}}{\alpha_1^{2r} \overline{\lambda(f_1^\circ)} \mathcal{E}(f_1) \mathcal{E}^*(f_1) \langle f_1^\circ, f_1^\circ \rangle_{N_1}} \langle g_n, f_2 \cdot \mathcal{E}(k_1, k_2, j + \chi) \rangle_{N(p^{2r})}.$$

Step 2 We recognise the nearly-holomorphic Eisenstein series $\mathcal{E}(k_1, k_2, j + \chi)$ of level Np^{2r} as the twist by the character χ of a simpler Eisenstein series \tilde{E} of level Np^r and character χ^{-2} , whose q -expansion is

$$\sum_{n \geq 1} q^n \sum_{\substack{d|n \\ p \nmid \frac{n}{d}}} d^{j-k_2} (n/d)^{k_1-1-j} \chi(n/d)^{-2} (e^{2\pi i d/N} + (-1)^{k_1-k_2} e^{-2\pi i d/N}).$$

Since $a_n(g_{2r}) = 0$ unless $p^{2r-1} | n$, we can pull the twist through the Petersson product to write

$$\langle g_{2r}, f_2 \cdot \mathcal{E}(k_1, k_2, j + \chi) \rangle_{N(p^{2r})} = \chi(-1) \langle g_{2r}, f_{2,\chi} \cdot \tilde{E} \rangle_{N(p^{2r})}.$$

Step 3 We rewrite the last Petersson product using the local Atkin–Lehner operator $W_{p^{2r}}$ acting on forms of level Np^{2r} . We compute that

$$\tilde{E}|W_{p^{2r}} = p^{2r(k_1-2-j)} \chi(-1) \sum_{a \in (\mathbb{Z}/p^{2r}\mathbb{Z})^\times} \chi(a)^{-2} E_{1/N+a/p^{2r}}$$

where the nearly-holomorphic Eisenstein series $E_\gamma = E_\gamma^{k_1-k_2}(-, j-k_1+1)$ for $\gamma \in \mathbf{Q}/\mathbf{Z}$ is as in [LLZ14, §4–5]. On the other hand, the action on $f_{2,\chi}$ is given by

$$f_{2,\chi} | W_{p^{2r}} = p^{(k_2-3)r} \varepsilon_2(p)^r G(\chi)^2 f_{2,\chi^{-1}}.$$

Combining these formulae, we deduce

$$\begin{aligned} \langle g_{2r}, f_2 \cdot \mathcal{E}(k_1, k_2, j + \chi) \rangle_{N(p^{2r})} = \\ \left(\frac{p^{(2k_1+k_2-5-2j)r} G(\chi)^2 \chi(N^2)}{\varepsilon_1(p)^{2r} \varepsilon_2(p)^r} \right) \langle f_{1,\beta} |_{k_1} W_{N_1}, f_{2,\chi^{-1}} \cdot E_{1/Np^{2r}} \rangle_{Np^{2r}}. \end{aligned}$$

Step 4 Via the classical “unfolding” technique, integrating against the Eisenstein series $E_{1/Np^{2r}}$ gives the (imprimitive) Rankin–Selberg L -function at $s = j$; cf. [Kat04, Theorem 7.1]. That is, we have

$$\begin{aligned} \langle f_{1,\beta} |_{k_1} W_{N_1}, f_{2,\chi^{-1}} \cdot E_{1/Np^{2r}} \rangle_{Np^{2r}} = \\ \frac{(j-1)!(j-k_2)! i^{k_1-k_2} L^{\text{imp}}(\overline{f_{1,\beta} |_{k_1} W_{N_1}}, f_{2,\chi^{-1}}, j)}{N^{k_1+k_2-2j-2} p^{2r(k_1+k_2-2j-2)} \pi^{2j+1-k_2} 2^{2j+k_1-k_2}}. \end{aligned}$$

However, since all Fourier coefficients a_n of $f_{2,\chi^{-1}}$ with $p \mid n$ are zero, this formula is unchanged if we replace $f_{1,\beta} |_{k_1} W_{N_1}$ with any form having the same Fourier coefficients away from p ; one such form is $\lambda(f_1^\circ) f_1^\circ$, so this is

$$\begin{aligned} \langle f_{1,\beta} |_{k_1} W_{N_1}, f_{2,\chi^{-1}} \cdot E_{1/Np^{2r}} \rangle_{Np^{2r}} = \\ \frac{(j-1)!(j-k_2)! i^{k_1-k_2} \overline{\lambda(f_1^\circ)} \cdot L^{\text{imp}}(f_1^\circ, f_2^\circ, \chi^{-1}, j)}{N^{k_1+k_2-2j-2} p^{2r(k_1+k_2-2j-2)} \pi^{2j+1-k_2} 2^{2j+k_1-k_2}}. \end{aligned}$$

Combining Steps 1, 3, and 4 gives the formula stated in Proposition 2.10. A similar argument (using an Eisenstein series of level $Np^{r+r'}$) can be used to prove Proposition 2.12.

Acknowledgments I am grateful to Eric Urban and Xin Wan for helpful comments on the topic of this paper, and to Xin Wan in particular for encouraging me to write it up. Part of the work described in the paper was carried out during a visit to the Institute for Advanced Study in Princeton in the spring of 2016, and I am very grateful to the IAS for their hospitality.

References

- [AI17] F. Andreatta and A. Iovita (with an appendix by Eric Urban), *Triple product p -adic L -functions associated to finite slope p -adic families of modular forms*. arxiv:1708.02785
- [BDR15a] M. Bertolini, H. Darmon, and V. Rotger, *Beilinson–Flach elements and Euler systems I: syntomic regulators and p -adic Rankin L -series*. J. Algebraic Geom. 24(2015), no. 2, 355–378. <http://dx.doi.org/10.1090/S1056-3911-2014-00670-6>
- [BDR15b] ———, *Beilinson–Flach elements and Euler systems II: the Birch and Swinnerton-Dyer conjecture for Hasse–Weil–Artin L -functions*. J. Algebraic Geom. 24(2015), no. 3, 569–604. <http://dx.doi.org/10.1090/S1056-3911-2015-00675-0>
- [BL16a] K. Büyükboduk and A. Lei, *Anticyclotomic p -ordinary Iwasawa theory of elliptic modular forms*. arxiv:1602.07508
- [BL16b] ———, *Anticyclotomic Iwasawa theory of elliptic modular forms at non-ordinary primes*. arxiv:1605.05310

- [Cas15] F. Castella, *p -adic heights of Heegner points and Beilinson–Flach classes*. J. London Math. Soc. 96(2017), no.1, 156–180. <http://dx.doi.org/10.1112/jlms.12058>
- [CE98] R. F. Coleman and B. Edixhoven, *On the semi-simplicity of the U_p -operator on modular forms*. Math. Ann. 310(1998), no. 1, 119–127. <http://dx.doi.org/10.1007/s002080050140>
- [Das16] S. Dasgupta, *Factorization of p -adic Rankin L -series*. Invent. Math. 205(2016), no. 1, 221–268. <http://dx.doi.org/10.1007/s00222-015-0634-4>
- [Hid85] H. Hida, *A p -adic measure attached to the zeta functions associated with two elliptic modular forms. I*. Invent. Math. 79(1985), no. 1, 159–195. <http://dx.doi.org/10.1007/BF01388661>
- [Hid88] ———, *A p -adic measure attached to the zeta functions associated with two elliptic modular forms. II*. Ann. Inst. Fourier (Grenoble) 38(1988), no. 3, 1–83. <http://dx.doi.org/10.5802/aif.1141>
- [JSW15] D. Jetchev, C. Skinner, and X. Wan, *The Birch–Swinnerton–Dyer formula for elliptic curves of analytic rank one*. [arxiv:1512.06894](https://arxiv.org/abs/1512.06894)
- [Kat04] K. Kato, *P -adic Hodge theory and values of zeta functions of modular forms*. Cohomologies p -adiques et applications arithmétiques. III. Astérisque 295(2004), ix, 117–290.
- [KLZ17] G. Kings, D. Loeffler, and S. L. Zerbes, *Rankin–Eisenstein classes and explicit reciprocity laws*. Cambridge J. Math. 5(2017), no. 1, 1–122. <http://dx.doi.org/10.4310/CJM.2017.v5.n1.a1>
- [LLZ14] A. Lei, D. Loeffler, and S. L. Zerbes, *Euler systems for Rankin–Selberg convolutions of modular forms*. Ann. of Math. (2) 180(2014), no. 2, 653–771. <http://dx.doi.org/10.4007/annals.2014.180.2.6>
- [LZ16] D. Loeffler and S. L. Zerbes, *Rankin–Eisenstein classes in Coleman families*. Res. Math. Sci. 3(2016), no. 29, 53. <http://dx.doi.org/10.1186/s40687-016-0077-6>
- [My91] V. K. My, *Rankin non-Archimedean convolutions of unbounded growth*. Mat. Sb. (N.S.) 182(1991), no. 2, 164–174; translation in Math. USSR-Sb. 72(1992), no. 1, 151–161.
- [Oht00] M. Ohta, *Ordinary p -adic étale cohomology groups attached to towers of elliptic modular curves. II*. Math. Ann. 318(2000), no. 3, 557–583. <http://dx.doi.org/10.1007/s002080000119>
- [Pan82] A. Panchishkin, *Le prolongement p -adique analytique des fonctions L de Rankin. I*. C. R. Acad. Sci. Paris Sér. I Math. 295(1982), no. 2, 51–53.
- [PR88] B. Perrin-Riou, *Fonctions L p -adiques associées à une forme modulaire et à un corps quadratique imaginaire*. J. London Math. Soc. 38(1988), no. 1, 1–32. <http://dx.doi.org/10.1112/jlms/s2-38.1.1>
- [Urb14] E. Urban, *Nearly overconvergent modular forms*. In: Iwasawa Theory 2012: State of the Art and Recent Advances, Springer Berlin, Heidelberg, 2014, pp. 401–441. http://dx.doi.org/10.1007/978-3-642-55245-8_14
- [Wan15] X. Wan, *Iwasawa main conjecture for supersingular elliptic curves*. [arxiv:1411.6352v2](https://arxiv.org/abs/1411.6352v2)

Mathematics Institute, Zeeman Building, University of Warwick, Coventry CV4 7AL, UK
e-mail: d.a.loeffler@warwick.ac.uk