

TARTU RADIOCARBON DATES XI

EVALD ILVES

Institute of Zoology and Botany, Academy of Sciences
Tartu, Estonian SSR

INTRODUCTION

This list includes dates of geologic samples measured using a single-channel liquid scintillation ^{14}C counter at the Geochemical and Statistical Laboratory, Tartu, Estonian SSR. Our modern standard is made of benzene enriched in ^{14}C and its activity is checked with NBS oxalic acid standard sample. Dates are given in conventional radiocarbon years, based on the Libby half-life of 5570 ± 30 yr. AD 1950 is the reference year. Errors refer only to 1σ standard deviation calculated from count rates involved.

GEOLOGIC SAMPLES

Estonian SSR

Haanja series

The Haanja upland in SE Estonian SSR belongs to an island deposit. Upper strata of basal Devonian deposits are at 100–160m asl. Glacial deposits 100–180m (40–50m on slopes) overlie the Devonian strata. The present upland was formed at the end of Valdai glaciation during the Haanja (Luuga) glacier retreat stage 13,500–13,200 BP. The Haanja upland consists of a variety of geologic and morphologic characteristics. Dates listed below are of sediments from center of upland, ca 300km².

Samples coll 1976 by H Mäemets (Mäetilga) and by E Ilves and H Mäemets 1980 (Karuniidu), 1981 (Tuuljärv), 1983 (Vaskna).

TA-1081. Mäetilga	3480 ± 70
Bryales peat from depth 370–380cm. Pollen Zone SB1.	
TA-1082. Mäetilga	5670 ± 100
Sapropel with admixture of lime pelite from depth 550–560cm.	
TA-1359. Karuniidu	870 ± 60
Fen peat with wood remains from depth 50–60cm. Pollen Zone SA3.	
TA-1396. Karuniidu	1100 ± 60
Fen peat with wood remains from depth 100–110cm. Pollen Zone SA2.	
TA-1397. Karuniidu	1920 ± 60
Fen peat from depth 150–160cm. Pollen Zone SA1	
TA-1398. Karuniidu	2180 ± 60
Fen peat from depth 200–210cm. Pollen Zone SA1.	
TA-1399. Karuniidu	2080 ± 60
Fen peat from depth 250–260cm. Pollen Zone SA1.	
TA-1400. Karuniidu	2500 ± 60
Fen peat from depth 300–310cm. Beginning of climatic period SA.	
TA-1401. Karuniidu	3000 ± 60
Fen peat from depth 350–360cm. Pollen Zone SB2.	

TA-1402. Karuniidu	3510 ± 60
Fen peat from depth 400–410cm. Pollen Zone SB1.	
TA-1403. Karuniidu	3560 ± 60
Bryales peat from depth 450–460cm. Pollen Zone SB1.	
TA-1404. Karuniidu	4120 ± 70
Forest peat from depth 500–510cm. Beginning of climatic period SB.	
TA-1405. Karuniidu	4300 ± 70
Forest peat from depth 520–530cm. Boundary of Atlantic and Subboreal (AT and SB) climatic periods.	
TA-1406. Karuniidu	6000 ± 80
Basal layer of forest peat on sand from depth 530–550cm. Climatic period AT.	
TA-1277. Karuniidu	1830 ± 60
Buried oak trunk found overlying sediments (while digging pond). Date is from 10 outer tree rings. Sample subm 1979 by U Riispere, Inst Zoology and Botany, Tartu, Estonian SSR.	
TA-1598. Tuuljärv	380 ± 100
<i>Carex-Sphagnum</i> peat from depth 100–105cm. Pollen Zone SA3.	
TA-1599. Tuuljärv	1670 ± 70
Forest- <i>Phragmites</i> peat from depth 140–150cm. Pollen Zone SA2.	
TA-1610. Tuuljärv	2600 ± 80
<i>Equisetum</i> -Bryales peat with remains of trees from depth 240–250cm. Beginning of Pollen Zone SA1.	
TA-1613. Tuuljärv	3040 ± 100
Forest peat from depth 320–330cm. Pollen Zone SB2.	
TA-1614. Tuuljärv	3560 ± 80
Forest peat from depth 360–370cm. Pollen Zone SB2.	
TA-1617. Tuuljärv	4070 ± 70
Forest- <i>Phragmites</i> peat from depth 450–460cm. Pollen Zone SB1.	
TA-1615. Tuuljärv	6560 ± 90
Sapropel with remains <i>Equisetum</i> from depth 480–490cm. Pollen Zone AT2.	
TA-1616. Tuuljärv	7940 ± 80
<i>Phragmites</i> -Bryales peat from depth 490–500cm. End of climatic period BO.	
TA-1515. Tuuljärv	8790 ± 80
Partially decomposed Bryales peat from depth 560–570cm. Beginning of climatic period BO.	
TA-1694. Vaskna	830 ± 60
<i>Carex</i> peat with remains of trees from depth 25–30cm. Pollen Zone SA3.	

TA-1693. Vaskna	1510 ± 70
Forest- <i>Carex</i> peat from depth 30–35cm. Boundary of Pollen Zones SA2 and SA3.	
TA-1692. Vaskna	2120 ± 60
Forest- <i>Carex</i> peat from depth 45–50cm. Pollen Zone SA2.	
TA-1691. Vaskna	2300 ± 60
Forest- <i>Phragmites</i> peat from depth 50–55cm. Boundary of Pollen Zones SA2 and SA1.	
TA-1690. Vaskna	2590 ± 60
Forest- <i>Carex</i> peat from depth 55–60cm. Boundary of climatic periods SB and SA.	
TA-1698. Vaskna	4210 ± 60
Forest- <i>Carex</i> peat from depth 70–75cm. Pollen Zone SB2.	
TA-1688. Vaskna	5030 ± 60
Forest- <i>Carex</i> peat from depth 80–85cm. Pollen Zone SB1.	
TA-1687. Vaskna	6850 ± 60
Forest-Bryales peat from depth 95–100cm. Climatic period AT.	
TA-1686. Vaskna	7250 ± 60
<i>Equisetum-Eriophorum</i> peat from depth 110–115cm. End of climatic period BO2.	
TA-1727. Vaskna	8050 ± 60
Bryales peat from depth 115–120cm. Pollen Zone BO2.	
TA-1685. Vaskna	8280 ± 60
Bryales- <i>Phragmites</i> peat from depth 125–130cm. Pollen Zone BO2.	
TA-1737. Vaskna	8730 ± 60
Bryales peat from depth 145–150cm. End of climatic period PB.	
TA-1684. Vaskna	9680 ± 70
Bryales- <i>Phragmites</i> peat from depth 160–165cm. Upper layer of climatic period PB.	
TA-1683. Vaskna	9870 ± 70
Dy from depth 170–175cm. Climatic period PB.	
TA-1682. Vaskna	9970 ± 70
Sapropel from depth 185–190cm. Climatic period PB.	
TA-1600. Vaskna	9930 ± 70
Sapropel from depth 200–205cm. Beginning of climatic period PB.	

Järvesoo series

Järvesoo bog, 90ha, lies in Saaremaa Island, 18km W of Kuressaare and is formed of fen and mesotrophic peat. Samples coll 1977 by E Ilves and A Sarv, Inst Zoology and Botany, Inst Geology, Tartu.

TA-771. Järvesoo	1420 ± 70
Peat from depth 20–30cm.	

TA-772. Järvesoo	1460 ± 70
Peat from depth 60–70cm.	
TA-773. Järvesoo	1450 ± 70
Peat from depth 70–80cm.	
TA-774. Järvesoo	1750 ± 70
Peat from depth 80–90cm.	
TA-775. Järvesoo	4090 ± 80
Peat from depth 100–110cm.	
TA-776. Järvesoo	4570 ± 80
Peat from depth 110–120cm.	
TA-778. Järvesoo	5440 ± 70
Peat from depth 130–140cm.	
TA-779A. Järvesoo	6280 ± 90
Peat from depth 160–170cm.	
TA-779B. Järvesoo	6510 ± 70
Lake lime from depth 160–170cm.	
TA-780. Järvesoo	6770 ± 90
Lake lime from depth 220–230cm.	
TA-777. Järvesoo	7610 ± 80
Lake lime from depth 260–270cm.	
TA-781. Järvesoo	8380 ± 80
Lake lime from depth 320–330cm.	
TA-782. Järvesoo	8800 ± 90
Lake lime from depth 340–350cm.	

Ukrainian SSR

Stojanov series

Stojanov bog lies in Malye Polesje near Stojanov village, Lvov oblast, Ukrainian SSR. Bog is drained and peat extracted. Peat stratum reaches thickness of 5.25m. Samples coll and subm 1978 by A T Artjushenko and L G Bezus'ko, N G Holodnyi Inst Botany, Kiev, Ukrainian SSR (Bezus'ko, Ilves & Kayutina 1980).

TA-1214. Stojanov	600 ± 70
<i>Carex</i> peat with remains of <i>Phragmites</i> from depth 35–40cm. Pollen Zone SA2.	
TA-1215. Stojanov	1130 ± 60
<i>Carex</i> peat with remains of <i>Phragmites</i> from depth 85–90cm. Boundary of Pollen Zones SA2 and SA1.	
TA-1216. Stojanov	1900 ± 70
<i>Carex</i> peat with remains of <i>Phragmites</i> from depth 140–145cm. Pollen Zone SA1.	

TA-1321. Stojanov	2680 ± 70
<i>Carex-Phragmites</i> peat from depth 202–203cm. Climatic period SB.	
TA-1322. Stojanov	3390 ± 70
<i>Carex-Phragmites</i> peat from depth 298–303cm. Pollen Zone AT2.	
TA-1323. Stojanov	4850 ± 70
<i>Carex-Phragmites</i> peat from depth 348–353cm. Pollen Zone AT2.	
TA-1217. Stojanov	6280 ± 70
<i>Carex-Phragmites</i> peat from depth 423–428cm. Pollen Zone AT1.	
TA-1218. Stojanov	7360 ± 70
<i>Phragmites</i> peat from depth 498–503cm. Pollen Zone SB.	

*Byelorussian SSR***Sudoble series**

Samples from basal sediments, 300m from W shore of Sudoble Lake, 5km SE of Zhodino, Smolevitschki dist, Minsk Obl. Samples coll and subm 1979 by I I Bogdel, V I Lenin Byelorussian State Univ, Minsk (Bogdel *et al* 1983).

TA-1219. Sudoble	2360 ± 80
Dy from depth 140–160cm. End of climatic period SB.	
TA-1220. Sudoble	3930 ± 80
Dy from depth 280–300cm. Climatic period SB.	
TA-1221. Sudoble	4960 ± 70
Dy from depth 380–400cm. Beginning of climatic period SB.	
TA-1222. Sudoble	5950 ± 80
Dy from depth 485–500cm. Climatic period AT.	
TA-1223. Sudoble	8510 ± 70
Dy from depth 580–600cm. Boundary of climatic periods BO and AT.	
TA-1224. Sudoble	9080 ± 90
Sapropel from depth 680–700cm. Basal layers of climatic period BO.	
TA-1225. Sudoble	11,160 ± 100
Sapropel with slight admixture of lake lime from depth 770–790cm. Climatic period AL.	
TA-1226. Sudoble	11,550 ± 100
Peat with admixtures of lake lime and macroparticles of wood from depth 875–885cm. Climatic period AL.	

Kobuzi series

Kobuzi bog lies in Vilei dist, 400 m N of Kobuzi village. Samples coll and subm 1979 by I Bogdel.

TA-1369. Kobuzi	730 ± 60
Peat with undecayed vegetal remains from depth 22–28cm.	

TA-1368. Kobuzi	1440 ± 70
Same type, from depth 28–35cm.	
TA-1367. Kobuzi	2750 ± 70
Same type, from depth 50–55cm.	
TA-1366. Kobuzi	4320 ± 70
Same type, from depth 75–80cm.	
TA-1365. Kobuzi	6050 ± 60
Same type, from depth 100–105cm.	
TA-1364. Kobuzi	7650 ± 70
Peat from depth 125–130cm. Color of peat turns into light brown.	
TA-1331. Kobuzi	8760 ± 70
Peat from depth 150–155cm. Color of peat turns into yellowish brown.	
TA-1330. Kobuzi	8730 ± 70
Peat from depth 175–180cm.	
TA-1329. Kobuzi	9640 ± 90
Peat with slightly decayed woody remains from depth 225–230cm.	
TA-1326. Kobuzi	9550 ± 80
Wood “A” from depth 240cm.	
TA-1327. Kobuzi	10,430 ± 90
Wood “B” from depth 240cm, from layer where partly decayed wood from trunks, roots and branches were collected.	
TA-1328. Kobuzi	9530 ± 80
Forest peat from depth 250–255cm.	
TA-1325. Kobuzi	10,430 ± 90
Forest peat from depth 275–280cm.	
TA-1324. Kobuzi	11,200 ± 90
Peat with sand from depth 295–300cm.	

*Georgian SSR***Anaklja series**

Dates of lake and bog sediments reported below refer to complex study of Inguri River basin, W Georgia. Samples coll and subm 1981 by L R Serebryannyi, Inst Geography, Tartu (Serebryannyi *et al* 1984).

TA-1303. Anaklja	3090 ± 100
Peat from depth 217–220cm. Base of Pollen Zone SB2.	
TA-1301. Anaklja	4090 ± 90
Peat from depth 315–320cm. Pollen Zone SB1.	

TA-1300. Anaklija	4530 ± 70
Peat from depth 380–385cm. Pollen Zone SB1.	
TA-1299. Anaklija	4570 ± 70
Peat from depth 530–535cm. Pollen Zone AT2.	
TA-1298. Anaklija	5640 ± 100
Peat from depth 625–632cm. Pollen Zone AT2.	

*Hungarian PR***Lovaszberen**

Sample from buried soil from Lovaszberen loess, Hungary, site with coordinates (47°18'N, 18°33'E). In 1977 while examining the section, numerous large charcoal fragments were observed (Ilves, Pecsi & Serebryannyi 1980). Sample subm 1977 by M Pecsi, Hungarian Acad Sciences.

TA-1196. Lovaszberen	20,220 ± 300
-----------------------------	---------------------

REFERENCES

- Bezus'ko, LG, Ilves, EO and Kayutina, TI 1980 New data on vegetation of Maloye Polesje in the Holocene. *Ukr Bot Zh* 37(5):35–39 (in Russian).
- Bogdol, II, Vlassov, BP, Ilves, EO and Klimanov, VA 1983 Sudoble section—A stratotype reconstruction of paleogeographic conditions of the Holocene in central Byelorussia, in *History of lakes in the USSR*. Abs, All-Union Mtg, 6th, Tallinn:30–33 (in Russian).
- Ilves, E and Mäemets, H 1987 Results of radiocarbon and palynological analyses of coastal deposits of lakes Tuuljärv and Vaskna,—Palaeohydrology of the temperate zone. *Mires and lakes, Tallinn* III:108–130.
- Ilves, E, Pecsi, M and Serebryannyi, L 1980 Absolute age of buried soil from Lovaszberen loess section. *Izv Akad Nauk Est SSR Khim Geol* 29(3):118–120 (in Russian).
- Mäemets, H 1983 Palynological and radiocarbon data on postglacial vegetational history of Haanja Heights (ESSR). *Man, vegetation and soil, Tallinn*:98–111.
- Serebryannyi, LR, Margalitadze, NA, Janelidze, Ch P, Gey, NA, Ilves, EO, Skobeva, EI, Vishnevskaja, EM, Orlov, AV 1984 Late- and postglacial history of phytolandscapes of Colchis. *Proc Acad Sci Georgian SSR Biol Ser* 10(5):306–312 (in Russian).