

## CORRIGENDUM

to the paper

### WEAK BEHAVIOUR OF FOURIER-NEUMANN SERIES

ÓSCAR CIAURRI

*Departamento de Matemáticas y Computación, Universidad de La Rioja, Edificio J. L. Vives,  
Calle Luis de Ulloa s/n, 26004 Logroño, Spain  
e-mail: oscar.ciaurri@dmc.unirioja.es*

MARIO PÉREZ

*Departamento de Matemáticas, Edificio de Matemáticas, Universidad de Zaragoza, 50009 Zaragoza, Spain  
e-mail: mperez@posta.unizar.es*

and JUAN L. VARONA

*Departamento de Matemáticas y Computación, Universidad de La Rioja, Edificio J. L. Vives,  
Calle Luis de Ulloa s/n, 26004 Logroño, Spain  
e-mail: jvarona@dmc.unirioja.es  
URL: <http://www.unirioja.es/dptos/dmc/jvarona/welcome.html>*

(Received 28 May, 2003; accepted 1 June, 2003)

A very important misprint occurs in the statement of the main theorem of the paper entitled ‘Weak behaviour of Fourier-Neumann series’ *Glasgow Math. J.* **45** (2003), 97–104. The statement in lines 3–7 of p. 99 should be replaced by the following statement.

**THEOREM.** *Let  $\alpha \geq 0$ ,  $p_1 = 4(\alpha + 1)/(2\alpha + 3)$ ,  $p_2 = 4(\alpha + 1)/(2\alpha + 1)$ . Then the partial sum operators  $S_n$  ( $n = 0, 1, 2, \dots$ ) are not uniformly bounded as operators from  $L^{p_i}(x^{-2\alpha+1})$  into  $L^{p_i, \infty}(x^{-2\alpha+1})$  but are uniformly bounded as operators from  $L^{p_i, 1}(x^{-2\alpha+1})$  into  $L^{p_i, \infty}(x^{-2\alpha+1})$  ( $i = 1, 2$ ). In the case  $-1 < \alpha < 0$ , the second statement holds with  $p_i = 4/3$  and  $p_2 = 4$ .*

The proof presented corresponds to the revised statement.