

## TOTALLY REAL ORBITS IN AFFINE QUOTIENTS OF REDUCTIVE GROUPS

H. AZAD, J. J. LOEB AND M. N. QURESHI

Let  $K$  be a compact connected Lie group and  $L$  a closed subgroup of  $K$ . In [8] M. Lassalle proves that if  $K$  is semisimple and  $L$  is a symmetric subgroup of  $K$  then the holomorphy hull of any  $K$ -invariant domain in  $K^{\mathbb{C}}/L^{\mathbb{C}}$  contains  $K/L$ . In [1] there is a similar result if  $L$  contains a maximal torus of  $K$ . The main group theoretic ingredient there was the characterization of  $K/L$  as the unique totally real  $K$ -orbit in  $K^{\mathbb{C}}/L^{\mathbb{C}}$ . On the other hand, Patrizio and Wong construct in [9] special exhaustion functions on the complexification of symmetric spaces  $K/L$  of rank 1 and find that the minimum value of their exhaustions is always achieved on  $K/L$ . By a lemma of Harvey and Wells [6] one knows that the set where a strictly plurisubharmonic (briefly s.p.s.h) function achieves its minimum is totally real. There is a related result in [2, Lemma 1.3] which states that if  $\phi$  is any differentiable function on a complex manifold  $M$  then the form  $dd^{\mathbb{C}}\phi$  vanishes identically on any real submanifold  $N$  contained in the critical set of  $\phi$ ; in particular if  $\phi$  is s.p.s.h then  $N$  must be totally real. In view of these results we give in this note a description of all totally real  $K$ -orbits in the affine quotients  $K^{\mathbb{C}}/L^{\mathbb{C}}$  of  $K/L$ . Our main result is as follows:

PROPOSITION. *Let  $G = K^{\mathbb{C}}$ ,  $H = L^{\mathbb{C}}$ . The group  $L$  has finitely many totally real orbits in  $G/H$  if and only if  $N(H^{\circ})/H^{\circ}$  is finite,  $H^{\circ}$  being the connected component of  $H$  and  $N(H^{\circ})$  its normalizer in  $G$ , and in this case there is a unique totally real  $K$ -orbit in  $G/H$ .*

This proposition has the following consequence.

COROLLARY. *If  $N(H^{\circ})/H^{\circ}$  is finite then any  $K$ -invariant s.p.s.h. function on  $G/H$  is proper and achieves its minimum value on  $K/L$ . Moreover, the holomorphy hull of any  $K$ -invariant domain in  $G/H$  meets  $K/L$ .*

---

Received July 30, 1992.

This Corollary follows from the Proposition, using the main results of [2] and repeating the arguments of Corollary 2 of [1].

The appendix verifies that the assumptions of the Corollary hold if  $K$  is semi-simple and  $K/L$  is a symmetric space, thereby recovering results of Lassalle and Patrizio–Wong mentioned above.

From the Corollary we see that if  $N(H^\circ)/H^\circ$  is finite then any  $K$ -invariant s.p.s.h. function on  $G/H$  is a canonical exhaustion function in the sense of [9]. In Eliashberg–Gromov [5, Th. 1.4 A] it is stated that any two exhausting strictly plurisubharmonic functions on a Stein manifold  $M$  give the same symplectic structure. This is not quite true as e.g. the function  $\|z\|^2$  and  $\log(1 + \|z\|^2)$  ( $z \in \mathbf{C}^n$ ) are exhausting strictly plurisubharmonic functions, but their associated forms  $dd^c\|z\|^2$  and  $dd^c\log(1 + \|z\|^2)$  are not symplectically equivalent, as they give, infinite and finite volume to  $\mathbf{C}^n$ . However, by imposing an extra geometric condition, namely that the metric be complete and Ricci flat, it is in some cases possible to pin down a unique s.p.s.h. exhaustion function. This happens in the case of complex symmetric varieties of rank 1, where the existence of a complete Ricci-flat metric is assured by a theorem of Bando–Kobayashi [3]. Their conditions can be verified by using the description of invariant Chern forms given in Borel–Hirzebruch [4]. It would be interesting to understand, in the spirit of [9], the geometric significance of these distinguished metrics.

Our notation is standard. In particular, if  $H$  is a subgroup of  $G$ , then  $N_G(H)$  and  $Z_G(H)$  denote the normalizer and centralizer of  $H$  in  $G$ . Also,  $H^\circ$  denotes the connected component of identity of  $H$  and  $x_y$  the conjugate  $xyx^{-1}$ .

*Proof of the Proposition.* Let  $G$  and  $K$  be as in the statement of the Proposition. We note that totally real  $K$ -orbits in  $G/H$  are precisely those of half the dimension of  $G/H$ . Moreover, if  $\pi : G/H^\circ \rightarrow G/H$  is the natural map and  $\mathcal{Q}$  is a totally real  $K$ -orbit in  $G/H$  then  $\pi^{-1}(\mathcal{Q})$  is a union of totally real  $K$ -orbits which are permuted by the right action of the finite group  $H/H^\circ$  on  $G/H^\circ$ . Therefore, we may assume that  $H$  is connected. Let  $G = KP$  be the Cartan decomposition of  $G$  [7]. We give the proof in several steps, some of which are of independent interest.

*Step 1.* If  $p = e^x \in P$  centralizes  $y \in \text{Lie}(G)$  then the 1-parameter subgroup  $\{e^{rx} : r \in \mathbf{R}\}$  also centralizes  $y$ .

*Proof.* There is a faithful representation of  $G$  in  $GL(n, \mathbf{C})$  in which  $K$  is represented by unitary matrices and  $P$  by Hermitian matrices. Now if  $x$  is a Hermitian matrix and  $e^{nx}$  ( $n \in \mathbf{Z}, n > 0$ ) commutes with a matrix  $y$  then, taking into account that  $e^x$  has positive eigenvalues, one sees readily that  $e^x$  also commutes with  $y$  and therefore so does  $e^{rx}$  ( $r \in \mathbf{R}$ ).

*Step 2.* If  $L$  is connected and  $n \in N_G(H)$  then  $n$  factorizes as  $n = kpx$ , where  $k \in K \cap N(H)$ ,  $p \in P \cap Z_G(H)$  and  $x \in H$  (recall that  $H = L^{\mathbf{C}}$  and  $G = K^{\mathbf{C}}$ ).

*Proof.* Let  $n \in N(H)$ . Now  $L$  is a maximal compact subgroup of  $H$  so, by conjugacy of maximal compact subgroups, we have  ${}^x n L = L$  for some  $x \in H$ . Let  $xn = kp$  be the Cartan decomposition of  $xn$  with  $k \in K$  and  $p \in P$ . Taking into account that  $K$  normalizes  $P$  we see that if  $k, k_1 \in K$  and  $p \in P$  with  $pkp^{-1} = k_1$ , then  $k = k_1$ . Therefore  ${}^p L = {}^{k^{-1}} L \subset K$  shows that  $p$  centralizes  $L$  and therefore  $H$  too, and  $k$  normalizes  $H$ . Hence  $n = x^{-1}kp = k(k^{-1}x^{-1}k)p = kp(k^{-1}x^{-1}k) = kpx'$  where  $x' = k^{-1}x^{-1}k \in H$ ,  $k \in K \cap N(H)$  and  $p \in P \cap Z_G(H)$ .

*Step 3.* If  $L$  is connected and  $N = N(H)/H$  is finite, then  $N$  has representatives in  $K$ .

*Proof.* Let  $n \in N(H)$  and  $n = kpx$  be the factorization of  $n$  given by step 2. Let  $p = e^x$ . The 1-parameter subgroup  $Z = \{e^{rx} : r \in \mathbf{R}\}$  is, by step 1, in  $Z_G(H)$ . Since  $ZH/H$  is in the finite group  $N(H)/H$ , we must have  $Z \subset H$ . Therefore  $N(H)/H$  has representatives in  $K$ .

*Step 4.* For a connected group  $L$ , the orbits of  $N_K(H)H/H$  on  $N(H)/H$  parametrize the totally real  $K$ -orbits in  $G/H$ .

*Proof.* A  $K$ -orbit  $\Omega$  in  $G/H$  is totally real if and only if  $\dim(\Omega) = \dim(K/L)$ . Let  $\xi_0 = eH$  and let  $Kx\xi_0$  be totally real in  $G/H$ . So  $\dim(K \cap xHx^{-1}) = \dim(L)$  and therefore  $\dim({}^{x^{-1}} K \cap H) = \dim(L)$ . By conjugacy of maximal compact subgroups, the group  $({}^{x^{-1}} K \cap H)^\circ$  is conjugate in  $H = L^{\mathbf{C}}$  to  $L$  and therefore  $(K \cap xHx^{-1})^\circ$  is conjugate in  $G = K^{\mathbf{C}}$  to  $L$ , say by an element  $kp$ , where  $k \in K$  and  $p \in P$ . As in step 2, the element  $p$  centralizes  $L$  and so  $(K \cap xHx^{-1})^\circ = kLk^{-1}$ . Hence  $kLk^{-1} \subset xHx^{-1}$  and  $k^{-1}x \in N(H)$ , so  $x = kn$  for some  $n \in N(H)$ . Conversely if  $x = kn$  with  $n \in N(H)$ , then  $K \cap xHx^{-1} = K \cap kHk^{-1} \cong K \cap H = L$ . Therefore, a  $K$ -orbit in  $G/H$  is totally real if and only if a representative of the  $K$ -orbit can be chosen in  $N(H)$ . Finally, if  $n_1, n_2 \in N(H)$  and

$kn_1(H) = n_2(H)$  with  $k \in K$  then  $k \in N_K(H)$ . Therefore the orbits of the compact group  $N_K(H)H/H \cong N_K(H)/L$  on  $N(H)/H$  parametrize the totally real  $K$ -orbits in  $G/H$ .

*Step 5. Conclusion of the proof.* Suppose  $K$  has finitely many totally real orbits in  $G/H$ . Then it also has finitely many such orbits in  $G/H^\circ$ , so we may assume that  $H$  is connected. By step 4, the compact group  $N_K(H)H/H$  has finitely many orbits on the Stein manifold  $N(H)/H$ . Therefore,  $N(H)/H$  must be finite and by step 3 it must have representatives in  $K$ . Hence there is a unique totally real  $K$ -orbit in  $G/H$  if and only if  $N(H^\circ)/H^\circ$  is finite. This completes the proof of the proposition.

APPENDIX. In this appendix we verify that the assumptions of the proposition hold for the complexification  $K^{\mathbb{C}}/L^{\mathbb{C}}$  of a symmetric space  $K/L$ , with  $K$  being compact.

PROPOSITION. *If  $K \supset L$  is a symmetric pair then  $K^{\mathbb{C}}/L^{\mathbb{C}}$  has exactly one totally real  $K$ -orbit ( $K$ -semisimple).*

*Proof.* We have  $\dot{K}^{\mathbb{C}} = \dot{L}^{\mathbb{C}} \oplus \mathfrak{m}$  where  $\mathfrak{m}$  is  $\text{ad}(L^{\mathbb{C}})$  invariant so  $\text{Lie}(N(L^{\mathbb{C}})) = \dot{L}^{\mathbb{C}} \oplus Z_{L^{\mathbb{C}}}(\mathfrak{m})$  as vector spaces.

Hence  $N(L^{\mathbb{C}})/L^{\mathbb{C}}$  is finite if and only if  $Z_{L^{\mathbb{C}}}(\mathfrak{m}) = 0$ . The proposition follows from the following lemma.

LEMMA. *If  $G \supset H$  is a symmetric pair with  $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$ , where  $[\mathfrak{h}, \mathfrak{m}] \subset \mathfrak{m}$ ,  $[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{h}$  then adjoint representation of  $H$  on  $\mathfrak{m}$  is irreducible, provided  $G$  is simple.*

*Proof.* The Killing form is a non-degenerate form. The function  $\Theta(x + y) = x - y$  is an automorphism of  $G$ . Since the Killing form is invariant under all automorphisms, we see that  $(\mathfrak{h}, \mathfrak{m}) = 0$ . Hence the Killing form is nondegenerate on both  $\mathfrak{h}$  and  $\mathfrak{m}$ .

Let  $\mathfrak{m}_1 \subset \mathfrak{m}$  be an  $H$ -invariant subspace and  $\mathfrak{m}_2 (= \mathfrak{m}_1^\perp)$  its orthocomplement. Now  $([\mathfrak{m}_1, \mathfrak{m}_2], \mathfrak{m}) = 0$  as  $(\mathfrak{m}, \mathfrak{m}) \subset \mathfrak{h}$ . Moreover as  $([x, y], z) = (x, [y, z])$  we see that  $([\mathfrak{m}_1, \mathfrak{m}_2], \mathfrak{h}) = 0$ . Hence  $[\mathfrak{m}_1, \mathfrak{m}_2] = 0$  by non-degeneracy of the Killing form.

Now  $\mathfrak{m}_1 + [\mathfrak{m}_1, \mathfrak{m}_2]$  is an  $H$ -submodule. Moreover

$$\begin{aligned} [\mathfrak{m}_1 + [\mathfrak{m}_1, \mathfrak{m}_1], \mathfrak{m}] &= [\mathfrak{m}_1 + [\mathfrak{m}_1, \mathfrak{m}_1], \mathfrak{m}_1 + \mathfrak{m}_2] \\ &\subset [\mathfrak{m}_1, \mathfrak{m}_1] + \mathfrak{m}_1 + [\mathfrak{m}_1, \mathfrak{m}_2] + [[\mathfrak{m}_1, \mathfrak{m}_1], \mathfrak{m}_2]. \end{aligned}$$

Since  $[\mathfrak{m}_1, \mathfrak{m}_2] = 0$  we have by Jacobi identity  $[[\mathfrak{m}_1, \mathfrak{m}_1], \mathfrak{m}_2] = 0$ . Hence  $[\mathfrak{m}_1 + [\mathfrak{m}_1, \mathfrak{m}_1], \mathfrak{m}] \subset \mathfrak{m}_1 + [\mathfrak{m}_1, \mathfrak{m}_1]$ . Thus if  $G$  is semisimple then  $\mathfrak{m}_1 + [\mathfrak{m}_1, \mathfrak{m}_1]$  is an ideal. In particular if  $G$  is simple then  $\mathfrak{m}_1 + [\mathfrak{m}_1, \mathfrak{m}_1] \subset \mathfrak{m} \oplus \mathfrak{h}$  shows that  $\mathfrak{m}_1 = 0$  or  $\mathfrak{m}_1 = \mathfrak{m}$ . Hence in this case  $\mathfrak{m}$  is an irreducible submodule.

In particular if  $K \supset L$  is a symmetric pair and  $K$  is simple then in the decomposition of  $K^{\mathbb{C}} = L^{\mathbb{C}} \oplus \mathfrak{m}$  we must have  $Z_{L^{\mathbb{C}}}(\mathfrak{m}) = 0$ . So in this case there is exactly one totally real  $K$ -orbit.

GENERAL CASE. Let  $G = G_1 \oplus \cdots \oplus G_r$  be semisimple. If  $\theta$  is an involutory automorphism then either  $\theta(G_i) = G_i$  or else the orbit of  $\theta$  on  $G_i$  is of length 2.

Hence  $G = \bigoplus_{i=1}^{r_0} G_i \oplus \bigoplus_{j=1}^{r_1} (G_j + \theta G_j)$ . In  $G_i \oplus \theta(G_i)$  the fixed point set is  $(\xi \oplus \theta(\xi))$  and  $(-1)$  eigen-space is  $(\xi - \theta(\xi))$ .

For a symmetric pair  $L^{\mathbb{C}} = G_{\theta}$  and  $\mathfrak{m} = G_{-1}$ . Hence  $Z_{L^{\mathbb{C}}}(\mathfrak{m})$  is a sum of  $Z_{L^{\mathbb{C}}}(\mathfrak{m}_i)$ . Since each sum is zero we see that for a symmetric pair  $K^{\mathbb{C}} \subset L^{\mathbb{C}}$  there is exactly one totally real  $K$ -orbit.

REFERENCES

[ 1 ] H. Azad and J. J. Loeb, On a theorem of Kempf and Ness, *Indiana University Math. J.*, **39**, No. 1 (1990), 61–65.  
 [ 2 ] —, Plurisubharmonic functions and the Kempf-Ness theorem, *Bull. London Math. Soc.*, **25** (1993), 162–168.  
 [ 3 ] S. Bando and R. Kobayashi, Ricci flat Kähler metrics on affine algebraic manifolds. II, *Math. Ann.*, **287** (1990), 175–180.  
 [ 4 ] A. Borel and F. Hirzebruch, Characteristic classes and homogeneous spaces, *Amer. J. Math.*, **80** (1958), 459–538.  
 [ 5 ] Y. Eliashberg and M. Gromov, Convex symplectic manifolds, *Proc. Symp. Pure Math.*, Vol. 52, Part 2, 135–162, AMS, 1991.  
 [ 6 ] F. R. Harvey and R. O. Wells, Zero sets of non-negative strictly plurisubharmonic functions, *Math. Ann.*, **201** (1973), 165–170.  
 [ 7 ] G. Helgason, *Differential Geometry and Symmetric Spaces*, New York, Academic Press, 1978.  
 [ 8 ] M. Lassalle, Série de Laurent des fonction holomorphes dans la complexification d'un espace symétrique compact, *Ann. Scient. Ec. Norm Sup.*, t. 11 (1978), 167–210.

- [9] G. Patrizio and P. M. Wong, Stein manifolds with compact symmetric center, *Math. Ann.*, **289** (1991), 355–382.

H. Azad  
*Department of Mathematics*  
*Quaid-i-Azam University*  
*Islamabad, Pakistan*

J. J. Loeb  
*Université D'Angers*  
*Faculté des Sciences*  
*Department de Mathématiques*  
*49045 Angers, France*

M. N. Qureshi  
*Department of Mathematics*  
*Quaid-i-Azam University*  
*Islamabad, Pakistan*