

## CORRIGENDUM: LIMITING CASES OF BOARDMAN'S FIVE HALVES THEOREM

MICHAEL C. CRABB<sup>1</sup> AND PEDRO L. Q. PERGHER<sup>2</sup>

<sup>1</sup>*Department of Mathematics, University of Aberdeen,  
Aberdeen AB24 3UE, UK (m.crabb@abdn.ac.uk)*

<sup>2</sup>*Departamento de Matemática, Universidade Federal de São Carlos, Caixa Postal 676,  
São Carlos, SP 13565-905, Brazil (pergher@dm.ufscar.br)*

(Received 15 April 2015)

Published in *Proceedings of the Edinburgh Mathematical Society* **56**(3) (2013), 723–732.

*Abstract* We rectify two omissions in the list of generators and include a brief discussion of the localization theorem of Kosniowski and Stong.

*Keywords:* localization theorem; five halves theorem; involution; fixed-point data; equivariant cobordism class; Stiefel–Whitney class

2010 *Mathematics subject classification:* Primary 57R85  
Secondary 57R75

### 1. Corrections

We are grateful to J. M. Boardman (private communication, published as [2]) for pointing out two omissions in the list of generators given in [3].

The cases in which  $c = 2$  in the main Theorems 1.2 and 3.4 should be corrected as follows.

Theorem 1.2 (p. 724) should read

$$c = 2: \quad 2 \text{ if } k = 1, \quad 9 \text{ if } k = 2, \quad 13 \text{ if } k = 3, \quad 14 \text{ if } k \geq 4.$$

Theorem 3.4 (p. 730) should read

$$\begin{aligned} c = 2: \quad & \text{if } k \geq 1, \quad b^{k-1} \cdot x_3^{(2)}, \quad b^{k-1} \cdot \gamma(x_2^{(1)}), \\ & \text{and, if } k \geq 2, \quad b^{k-2} \cdot (x_4^{(2)})^2, \quad b^{k-2} \cdot (y_4^{(2)})^2, \quad b^{k-2} \cdot \gamma(x_3^{(2)}) \cdot x_4^{(2)}, \\ & \quad b^{k-2} \cdot x_6^{(3)} \cdot x_2^{(1)}, \quad b^{k-2} \cdot \gamma^3(x_5^{(4)}), \quad b^{k-2} \cdot x_8^{(4)}, \quad b^{k-2} \cdot \gamma(x_7^{(4)}), \\ & \text{and, if } k \geq 3, \quad b^{k-3} \cdot \gamma^2(x_7^{(4)}) \cdot x_4^{(2)}, \quad b^{k-3} \cdot \gamma^2(x_7^{(4)}) \cdot y_4^{(2)}, \quad b^{k-3} \cdot \gamma^2(x_{11}^{(6)}), \\ & \quad b^{k-3} \cdot x_2^{(1)} \cdot z_{11}^{(5)}, \\ & \text{and, if } k \geq 4, \quad b^{k-4} \cdot (\gamma^2(x_7^{(4)}))^2. \end{aligned}$$

These require the following corrections to the text on p. 729. To the list of exclusions when  $c = 2$  must be added, if  $k \geq 3$ ,  $((6, 2_{k-3}), \omega_0)$ . Also, in the paragraph below the list, the dimension of the group  $(\mathcal{N}_3^{\mathbb{Z}_2})^{(2)}$  should be corrected to  $\dim(\mathcal{N}_3^{\mathbb{Z}_2})^{(2)} = 2$ .

We note too that the exceptional case in which  $c = 3$  and  $n = 2k - 1$  should read  $\omega = (2_{k-1}), \omega' = (1)$ .

**2. The localization theorem**

We take this opportunity to place the result of Kosniowski and Stong [4] that provided the basic input into [3] in the context of what is now standard localization theory.

Cohomology with  $\mathbb{Z}_2$ -coefficients will be denoted by  $H^*$ . For a  $\mathbb{Z}_2$ -space  $M$  we write  $H_{\mathbb{Z}_2}^*(M) = H^*(E\mathbb{Z}_2 \times_{\mathbb{Z}_2} M)$  for the equivariant Borel cohomology and let  $t \in H_{\mathbb{Z}_2}^1(*)$  be the generator, that is, the Euler class of the universal real line bundle over  $B\mathbb{Z}_2$ . We have a restriction map  $i^* : H_{\mathbb{Z}_2}^*(M) \rightarrow H^*(M)$ . If  $\mathbb{Z}_2$  acts trivially on  $M$ , then  $H_{\mathbb{Z}_2}^*(M) = H^*(M) \otimes \mathbb{Z}_2[t]$ .

Using the notation and terminology of [3], we can state the localization theorem for  $\mathbb{Z}_2$ -Borel cohomology as follows.

**Lemma 2.1.** *Consider an  $m$ -dimensional  $\mathbb{Z}_2$ -manifold  $M$  with fixed-point data  $(F^j, \eta_j), j = 0, \dots, m$ . Suppose that  $u \in H_{\mathbb{Z}_2}^m(M)$ . Then*

$$i^*(u)[M] = \sum_{j=0}^m (e(\eta_j)^{-1}u^{(j)})[F^j] \in \mathbb{Z}_2,$$

where  $u^{(j)} \in H_{\mathbb{Z}_2}^m(F^j)$  is the restriction of  $u$  to  $F^j$  and  $e(\eta_j) \in H_{\mathbb{Z}_2}^{m-j}(F^j)$  is the equivariant Euler class of  $\eta_j$ .

More explicitly, the equivariant Euler class  $e(\eta_j)$  and its inverse can be written as

$$\begin{aligned} e(\eta_j) &= t^{m-j} + w_1(\eta_j)t^{m-j-1} + \dots + w_{m-j}(\eta_j) \in H^*(F^j) \otimes \mathbb{Z}_2[t], \\ e(\eta_j)^{-1} &= t^{j-m}(1 + w_1(-\eta_j)t^{-1} + \dots + w_j(-\eta_j)t^{-j}) \in H^*(F^j) \otimes \mathbb{Z}_2[t, t^{-1}]. \end{aligned}$$

The class  $u^{(j)}$  may be expanded as  $u_m^{(j)} + u_{m-1}^{(j)}t + \dots + u_0^{(j)}t^m$ , where  $u_i^{(j)} \in H^i(F^j)$ , so that

$$(e(\eta_j)^{-1}u^{(j)})[F^j] = \sum_{i=0}^j (w_{j-i}(-\eta_j)u_i^{(j)})[F^j] \in \mathbb{Z}_2.$$

The result of Kosniowski and Stong [3, Proposition 2.5] is proved, when  $f(X_1, \dots, X_m)$  is homogeneous of degree  $d \leq m$ , by taking  $u = t^{m-d}v$ , where  $v$  is obtained by substituting in  $f(X_1, \dots, X_m)$  the  $r$ th Stiefel–Whitney class of  $E\mathbb{Z}_2 \times_{\mathbb{Z}_2} TM$  for the  $r$ th elementary symmetric function in the  $X_i$ .

**Proof.** This may be proved by following the argument given by Atiyah and Segal in [1, Theorem 2.12] to establish the corresponding result for  $K$ -theory. □

**References**

1. M. F. ATIYAH AND G. B. SEGAL, The index of elliptic operators: II, *Annals Math.* **87** (1968), 531–545.
2. J. M. BOARDMAN, *Math. Rev.* (2014), MR3109755.
3. M. C. CRABB AND P. L. Q. PERGHER, Limiting cases of Boardman's five halves theorem, *Proc. Edinb. Math. Soc.* **56**(3) (2013), 723–732.
4. C. KOSNIOWSKI AND R. E. STONG, Involutions and characteristic numbers, *Topology* **17** (1978), 309–330.