

LOWER BOUNDS FOR FUNDAMENTAL UNITS OF REAL QUADRATIC FIELDS

KOSHI TOMITA AND KOUJI YAMAMURO

Abstract. Let d be a square-free positive integer and $l(d)$ be the period length of the simple continued fraction expansion of ω_d , where ω_d is integral basis of $\mathbb{Z}[\sqrt{d}]$. Let $\varepsilon_d = (t_d + u_d\sqrt{d})/2$ (> 1) be the fundamental unit of the real quadratic field $\mathbb{Q}(\sqrt{d})$. In this paper new lower bounds for ε_d , t_d , and u_d are described in terms of $l(d)$. The lower bounds of ε_d are sharper than the known bounds and those of t_d and u_d have been yet unknown. In order to show the strength of the method of the proof, some interesting examples of d are given for which ε_d and Yokoi's d -invariants are determined explicitly in relation to continued fractions of the form $[a_0, \overline{1, \dots, 1, a_{l(d)}}]$.

Introduction

For a positive square-free integer d , let D be the discriminant of the real quadratic field $\mathbb{Q}(\sqrt{d})$ and $l(d)$ be the period length in the simple continued fraction expansion of the algebraic integer $\omega_d = (\sigma_d - 1 + \sqrt{d})/\sigma_d$, where $\sigma_d = 1$ (resp. 2) for $d \not\equiv 1 \pmod{4}$ (resp. $d \equiv 1 \pmod{4}$). It is well-known that the fundamental unit $\varepsilon_d = (t_d + u_d\sqrt{d})/2$ (> 1) of $\mathbb{Q}(\sqrt{d})$ has lower bounds that increase with $l(d)$. For example, there are $((1 + \sqrt{5})/2)^{l(d)}$ (see, for example [1, p. 240]) and $\sqrt{D}(3/2)^{l(d)-2}$ (see [2, p. 98]). For d with sufficiently large $l(d)$, ε_d is much larger than these lower bounds. So we will calculate a sharper lower bound in terms of $l(d)$ to study the sufficiently large ε_d for d . Furthermore, we will calculate the lower bounds for t_d and u_d in terms of $l(d)$ that have been yet unknown. In order to study Yokoi's d -invariants, which are concerned with the class number one problem for real quadratic fields, we need to investigate t_d and u_d . We have obtained the following theorem.

Received October 24, 2000.

2000 Mathematics Subject Classification: Primary 11R11; Secondary 11R27.

THEOREM 1. Let d be a positive square-free integer with $l(d) \geq 2$ and D be the discriminant of $\mathbb{Q}(\sqrt{d})$. Then we have

$$\varepsilon_d > \begin{cases} \frac{\sqrt{D}}{\sqrt{5}} \left(\frac{1 + \sqrt{5}}{2} \right)^{l(d)-1} & \text{if } l(d) \text{ is even,} \\ \frac{1}{\sqrt{5}} \left(\sqrt{D} - \frac{\sqrt{5}-1}{2} \right) \cdot \left(\frac{1 + \sqrt{5}}{2} \right)^{l(d)} & \text{if } l(d) \text{ is odd.} \end{cases}$$

Moreover, for t_d and u_d in $\varepsilon_d = (t_d + u_d\sqrt{d})/2$ (> 1), we have

$$t_d > \begin{cases} \frac{\sqrt{D}}{\sqrt{5}} \left(\frac{1 + \sqrt{5}}{2} \right)^{l(d)-1} & \text{if } l(d) \text{ is even,} \\ \frac{1}{\sqrt{5}} (\sqrt{D} - \sqrt{5} + 1) \cdot \left(\frac{1 + \sqrt{5}}{2} \right)^{l(d)} & \text{if } l(d) \text{ is odd} \end{cases}$$

and

$$u_d > \begin{cases} \frac{2}{\sigma_d \sqrt{5}} \left(\frac{1 + \sqrt{5}}{2} \right)^{l(d)-1} & \text{if } l(d) \text{ is even,} \\ \frac{2}{\sigma_d \sqrt{5}} \left(\frac{1 + \sqrt{5}}{2} \right)^{l(d)-2} & \text{if } l(d) \text{ is odd.} \end{cases}$$

In this paper, the simple continued fraction with period l is generally denoted by $[a_0, \overline{a_1, \dots, a_l}]$, and $[x]$ means the greatest integer less than or equal to x . Let $\{F_i\}$ be the Fibonacci numbers determined by $F_0 = 0$, $F_1 = 1$, $F_{i+1} = F_i + F_{i-1}$ ($i \geq 1$). The Fibonacci numbers play an important role in showing Theorem 1, because we use the inequality $\varepsilon_d \geq ([\omega_d] + \omega_d)F_{l(d)} + F_{l(d)-1}$ (resp. $([\omega_d] - 1 + \omega_d)F_{l(d)} + F_{l(d)-1}$) in the case that $d \not\equiv 1 \pmod{4}$ (resp. $d \equiv 1 \pmod{4}$). We are interested in whether the equality is possible. We got an affirmative answer and we have discovered real quadratic fields with $m_d \neq 0$. Here m_d is one of Yokoi's d -invariant, and it is defined by $m_d = [u_d^2/t_d]$. Another is defined by $n_d = [t_d/u_d^2]$. We know that there exist only finitely many d satisfying both class number one and $n_d \neq 0$ (i.e. $m_d = 0$) (see [7, p. 188]). So it is very important to investigate the case that $m_d \neq 0$ (i.e. $n_d = 0$). Now these results are stated as follows:

THEOREM 2. *Let d be a positive square-free integer and l be a positive integer satisfying $l \geq 2$. Assume that*

$$d = (2F_l + 1)^2 + 8F_{l-1} + 4.$$

Then $d \equiv 1 \pmod{4}$, $\omega_d = [F_l + 1, \overline{1, \dots, 1, 2F_l + 1}]$, and $l(d) = l$ hold.

Moreover, we have

$$\varepsilon_d = \frac{1}{2}(2F_l^2 + F_l + 2F_{l-1} + F_l\sqrt{d}) \quad (= ([\omega_d] - 1 + \omega_d)F_l + F_{l-1}),$$

$$\begin{cases} t_d = 2F_l^2 + F_l + 2F_{l-1}, \\ u_d = F_l, \end{cases}$$

and

$$n_d = \begin{cases} 5 & \text{if } l = 2, \\ 3 & \text{if } l = 3, \\ 2 & \text{if } l \geq 5. \end{cases}$$

THEOREM 3. *Let d be a positive square-free integer and l be a positive integer satisfying that $l \geq 2$ and $l \equiv 1, 2, \text{ or } 4 \pmod{6}$. Assume that*

$$d = ((F_l + 1)/2)^2 + F_{l-1} + 1.$$

Then $d \not\equiv 1 \pmod{4}$, $\omega_d = [(F_l + 1)/2, \overline{1, \dots, 1, F_l + 1}]$, and $l(d) = l$ hold.

Moreover, we have

$$\varepsilon_d = \frac{1}{2}(F_l^2 + F_l + 2F_{l-1} + 2F_l\sqrt{d}) \quad (= ([\omega_d] + \omega_d)F_l + F_{l-1}),$$

$$\begin{cases} t_d = F_l^2 + F_l + 2F_{l-1}, \\ u_d = 2F_l, \end{cases}$$

and

$$m_d = \begin{cases} 1 & \text{if } l = 2, \\ 2 & \text{if } l = 4, \\ 3 & \text{if } l \geq 7. \end{cases}$$

Remark 1. Our aim is to show that, for any l , there exists d satisfying $l(d) = l$ and $\omega_d = [a_0, \overline{1, \dots, 1, a_{l(d)}}]$. The case that $l = 6n + 1$ was treated in [4], but our proof is simpler than theirs. Their aim is to consider Eisenstein's problem.

Preliminary

In order to prove our theorems, we need several lemmas.

LEMMA 1. For a square-free positive integer d , we suppose $\omega_d = [a_0, \overline{a_1, \dots, a_l}]$. Moreover let q_i be the integers determined by $q_0 = 0, q_1 = 1, q_{i+1} = a_i q_i + q_{i-1}$ ($i \geq 1$). Then the fundamental unit $\varepsilon_d = (t_d + u_d \sqrt{d})/2$ (> 1) of $\mathbb{Q}(\sqrt{d})$ is given by the following formula:

If $d \not\equiv 1 \pmod{4}$, then

$$\varepsilon_d = (a_0 + \sqrt{d})q_{l(d)} + q_{l(d)-1}, \quad \begin{cases} t_d = 2(a_0 q_{l(d)} + q_{l(d)-1}), \\ u_d = 2q_{l(d)}. \end{cases}$$

If $d \equiv 1 \pmod{4}$, then

$$\varepsilon_d = \left(\frac{2a_0 - 1 + \sqrt{d}}{2} \right) q_{l(d)} + q_{l(d)-1}, \quad \begin{cases} t_d = (2a_0 - 1)q_{l(d)} + 2q_{l(d)-1}, \\ u_d = q_{l(d)}. \end{cases}$$

Proof is omitted (see proof of Lemma 1 in [3]).

LEMMA 2. For a positive square-free integer d , denote by D the discriminant of $\mathbb{Q}(\sqrt{d})$. Then we have

$$\varepsilon_d > (\sqrt{D} - 1)F_{l(d)} + F_{l(d)-1}.$$

Moreover, for t_d and u_d in $\varepsilon_d = (t_d + u_d \sqrt{d})/2$ (> 1), we have

$$t_d > (\sqrt{D} - 2)F_{l(d)} + 2F_{l(d)-1} \quad \text{and} \quad u_d \geq \left(\frac{2}{\sigma_d} \right) F_{l(d)}.$$

Proof. In the case that $d \not\equiv 1 \pmod{4}$, since $a_0 > \sqrt{d} - 1$ and $q_i \geq F_i$ for any integer i (≥ 1), from Lemma 1 we get

$$\begin{aligned} \varepsilon_d &\geq (a_0 + \sqrt{d})F_{l(d)} + F_{l(d)-1} \\ &> (2\sqrt{d} - 1)F_{l(d)} + F_{l(d)-1} \\ &= (\sqrt{D} - 1)F_{l(d)} + F_{l(d)-1}. \end{aligned}$$

For t_d and u_d , in the case that $d \not\equiv 1 \pmod{4}$, we have

$$t_d > 2\{(\sqrt{d} - 1)F_{l(d)} + F_{l(d)-1}\} = (\sqrt{D} - 2)F_{l(d)} + 2F_{l(d)-1}$$

and

$$u_d \geq 2F_{l(d)}.$$

In the case that $d \equiv 1 \pmod{4}$, we get the results in the same way. □

We get the following lemma by straightforward calculations.

LEMMA 3. For $i \geq 1$,

$$F_i > \begin{cases} \frac{1}{\sqrt{5}} \left(\frac{1 + \sqrt{5}}{2} \right)^{i-1} & \text{if } i \text{ is even,} \\ \frac{1}{\sqrt{5}} \left(\frac{1 + \sqrt{5}}{2} \right)^i & \text{if } i \text{ is odd.} \end{cases}$$

Proof of Theorem 1

Proof of Theorem 1. We put $\alpha = (1 + \sqrt{5})/2$. First we shall show the lower bound of ε_d . From Lemma 2, we know $\varepsilon_d > (\sqrt{D} - 1)F_{l(d)} + F_{l(d)-1}$.

In the case that $l(d)$ (≥ 3) is odd, we have, from Lemma 3,

$$\begin{aligned} (\sqrt{D} - 1)F_{l(d)} + F_{l(d)-1} &> (\sqrt{D} - 1)\frac{\alpha^{l(d)}}{\sqrt{5}} + \frac{1}{\sqrt{5}}\alpha^{l(d)-2} \\ &= \frac{\alpha^{l(d)}}{\sqrt{5}} \left\{ \sqrt{D} - \frac{\sqrt{5} - 1}{2} \right\}. \end{aligned}$$

This proves the odd case.

In the case that $l(d)$ (≥ 2) is even, we have

$$\begin{aligned} (\sqrt{D} - 1)F_{l(d)} + F_{l(d)-1} &> (\sqrt{D} - 1)\frac{\alpha^{l(d)-1}}{\sqrt{5}} + \frac{\alpha^{l(d)-1}}{\sqrt{5}} \\ &= \frac{\sqrt{D}}{\sqrt{5}}\alpha^{l(d)-1}. \end{aligned}$$

Next we shall show the lower bounds of t_d and u_d . From Lemma 2, we know

$$t_d > (\sqrt{D} - 2)F_{l(d)} + 2F_{l(d)-1}.$$

In the case that $l(d)$ (≥ 3) is odd, we have

$$\begin{aligned} (\sqrt{D} - 2)F_{l(d)} + 2F_{l(d)-1} &> (\sqrt{D} - 2)\frac{\alpha^{l(d)}}{\sqrt{5}} + 2\frac{\alpha^{l(d)-2}}{\sqrt{5}} \\ &= \frac{\alpha^{l(d)}}{\sqrt{5}}(\sqrt{D} - \sqrt{5} + 1). \end{aligned}$$

In the case that $l(d) (\geq 2)$ is even, we have

$$\begin{aligned} (\sqrt{D} - 2)F_{l(d)} + 2F_{l(d)-1} &> (\sqrt{D} - 2)\frac{\alpha^{l(d)-1}}{\sqrt{5}} + 2\frac{\alpha^{l(d)-1}}{\sqrt{5}} \\ &= \frac{\sqrt{D}}{\sqrt{5}}\alpha^{l(d)-1}. \end{aligned}$$

From Lemma 2 and Lemma 3, we can get the lower bound of u_d in a similar way as in the proof of t_d . Theorem has been completely proved. □

From Theorem 1, we get the following corollary for the period $l(d)$ and Yokoi’s d -invariant m_d :

COROLLARY. *If there exist a positive integer M and a positive square-free integer d such that $d > 13$ and*

$$l(d) \geq \frac{\log(M + 1) + \log(\sqrt{5d}) - \log 2}{\log\left(\frac{1 + \sqrt{5}}{2}\right)} + 1,$$

then $m_d > M$.

Proof. From the assumption, we have

$$\frac{\sqrt{D}}{\sqrt{5}}\left(\frac{1 + \sqrt{5}}{2}\right)^{l(d)-1} \geq (M + 1)d.$$

Since it holds that

$$\frac{1}{\sqrt{5}}\left(\sqrt{D} - \frac{\sqrt{5} - 1}{2}\right) \cdot \left(\frac{1 + \sqrt{5}}{2}\right)^i > \frac{\sqrt{D}}{\sqrt{5}}\left(\frac{1 + \sqrt{5}}{2}\right)^{i-1}$$

for any positive integer i , we get, from Theorem 1,

$$\varepsilon_d > (M + 1)d.$$

Moreover, $m_d = [\varepsilon_d/d]$ if $d > 13$ from Theorem 1.1 in [8], therefore we have

$$m_d > \frac{\varepsilon_d}{d} - 1 > M.$$

□

Remark 2. We describe the comparison between the lower bound in Theorem 1 and the two well-known lower bounds given in the introduction of this paper. From the proof of Corollary, our lower bound for d with odd $l(d)$ is greater than $(\sqrt{D}/\sqrt{5}) \cdot ((1 + \sqrt{5})/2)^{l(d)-1}$. Moreover, if $D > 13$, then we have the following:

$$\frac{\sqrt{D}}{\sqrt{5}} \left(\frac{1 + \sqrt{5}}{2}\right)^{l(d)-1} - \left(\frac{1 + \sqrt{5}}{2}\right)^{l(d)} > 0.22 \left(\frac{1 + \sqrt{5}}{2}\right)^{l(d)-1}.$$

Furthermore, if $l(d) \geq 7$, then

$$\frac{\sqrt{D}}{\sqrt{5}} \left(\frac{1 + \sqrt{5}}{2}\right)^{l(d)-1} - \sqrt{D} \left(\frac{3}{2}\right)^{l(d)-2} > 0.038\sqrt{D} \left(\frac{1 + \sqrt{5}}{2}\right)^{l(d)-2}.$$

Hence, our lower bounds are sharper than theirs.

Proof of Theorems 2 and 3

Proof of Theorems 2 and 3. First we shall prove the first half of Theorem 3. Suppose that $l \equiv 1, 2$, or $4 \pmod{6}$. Since it holds that

$$F_{l+6} \equiv F_l \pmod{4} \quad (l \geq 0),$$

we have $d \not\equiv 1 \pmod{4}$. We put

$$\omega_R = (F_l + 1)/2 + [(F_l + 1)/2, \underbrace{1, \dots, 1}_{l-1}, F_l + 1].$$

Then we have

$$\omega_R = F_l + 1 + \frac{1}{1 + \dots + 1} + \frac{1}{\omega_R}.$$

By a straightforward induction argument, we obtain that

$$\omega_R = F_l + 1 + \frac{F_{l-1}\omega_R + F_{l-2}}{F_l\omega_R + F_{l-1}}.$$

Here, using $F_l = F_{l-1} + F_{l-2}$ ($l \geq 2$), we have

$$\omega_R^2 - (F_l + 1)\omega_R - (F_{l-1} + 1) = 0.$$

Since $\omega_R > 0$, it holds that

$$\omega_R = \frac{F_l + 1}{2} + \sqrt{d}.$$

Hence, we obtain that $\omega_d = [(F_l + 1)/2, \overline{1, \dots, 1, F_l + 1}]$ and $l(d) = l$, and we can determine ε_d, t_d and u_d from Lemma 1. In the case of Theorem 2, we put

$$\omega_R = F_l + [F_l + 1, \underbrace{\overline{1, \dots, 1}}_{l-1}, 2F_l + 1].$$

Since we have $\omega_R = F_l + \omega_d$ in the same way, we can determine ε_d, t_d and u_d .

Next we shall show the remaining part of Theorem 3. Since $1/F_l$ is monotone decreasing in l , it holds that

$$3.4 < 4 \left(1 + \frac{3}{F_l}\right)^{-1} \leq 4 \left(1 + \frac{1}{F_l} + \frac{2F_{l-1}}{F_l^2}\right)^{-1} < 4$$

for $l \geq 7$. Hence, we have $m_d = 3$ for $l \geq 7$ from definition of m_d . And we can get m_d in the case that $l = 2$ or 4 from straightforward calculations.

Lastly we shall show the remaining part of Theorem 2. We have $n_d = 2$ for $l \geq 5$ from the definition of n_d , because

$$2 \leq n_d \leq 2 + \frac{1}{F_l} + \frac{2F_{l-1}}{F_l^2} < 2 + \frac{3}{F_l} < 3$$

for $l \geq 5$. And we can get n_d in the case that $l = 2$ or 3 from straightforward calculations. Theorems have been proved. □

Table: Square-free positive integers d with $2 < l(d) \leq 15$ represented by the Fibonacci numbers:

d	$\frac{l(d)}{(\text{mod } 6)}$	$l(d)$	h_d	$F_{l(d)}$	ω_d
3	2	2	1	1	$[1, \overline{1}, 2]$
21	2	2	1	1	$[2, \overline{1}, 3]$
37	3	3	1	2	$[3, \overline{1, 1}, 5]$
7	4	4	1	3	$[2, \overline{1, 1, 1}, 4]$
69	4	4	1	3	$[4, \overline{1, 1, 1}, 7]$
149	5	5	1	5	$[6, \overline{1, 1, 1, 1}, 11]$
58	1	7	2	13	$[7, \overline{1, 1, 1, 1, 1, 1}, 14]$
797	1	7	1	13	$[14, \overline{1, 1, 1, 1, 1, 1}, 27]$
4933	3	9	3	34	$[35, \overline{1, 1, 1, 1, 1, 1, 1, 1}, 69]$
32485	5	11	8	89	$[90, \overline{1, 1, 1, 1, 1, 1, 1, 1, 1, 1}, 179]$
84237	0	12	6	144	$[145, \overline{1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1}, 289]$

Table: Square-free positive integers d with $2 < l(d) \leq 15$ represented by the Fibonacci numbers:

d	$\frac{l(d)}{\pmod{6}}$	$l(d)$	h_d	$F_{l(d)}$	ω_d
13834	1	13	22	233	[117, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 234]
219245	1	13	12	233	[234, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 467]
1493861	3	15	20	610	[611, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1221]

Finally, we have the above table for d concerning Theorem 2 and Theorem 3. Here h_d is the class number of $\mathbb{Q}(\sqrt{d})$.

REFERENCES

[1] R. A. Mollin, *Quadratics*, CRC Press, Boca Rato, FL., 1996.
 [2] R. Sasaki, *A characterization of certain real quadratic fields*, Proc. Japan Acad., **62**, Ser. A (1986), no. 3, 97–100.
 [3] K. Tomita, *Explicit representation of fundamental units of some quadratic fields*, Proc. Japan Acad., **71**, Ser. A (1995), no. 2, 41–43.
 [4] K. S. Williams and N. Buck, *Comparison of the lengths of the continued fractions of \sqrt{D} and $\frac{1}{2}(1 + \sqrt{D})$* , Proc. Amer. Math. Soc., **120** (1994), no. 4, 995–1002.
 [5] H. Yokoi, *The fundamental unit and class number one problem of real quadratic fields with prime discriminant*, Nagoya Math. J., **120** (1990), 51–59.
 [6] H. Yokoi, *a note on class number one problem for real quadratic fields*, Proc. Japan Acad., **69**, Ser. A (1993), 22–26.
 [7] H. Yokoi, *The fundamental unit and bounds for class numbers of real quadratic fields*, Nagoya Math. J., **124** (1991), 181–197.
 [8] H. Yokoi, *New invariants and class number problem in real quadratic fields*, Nagoya Math. J., **132** (1993), 175–197.

Koshi Tomita
 Department of Mathematics
 Meijo University
 Shiogamaguchi 1-501, Tenpaku-ku, Nagoya, 468-8502
 Japan
 tomita@ccmfs.meijo-u.ac.jp

Kouji Yamamuro
 Department of Liberal Arts
 Aichi Konan College
 Omatsubara 172, Takaya-cho, Konan-shi, Aichi, 483-8086
 Japan
 yamamuro@math.nagoya-u.ac.jp