

Review

Cite this article: Rzepka-Marot I, Gebhardt N, Nowak J, Bruns B, Friederich H-C and Nikendei C (2025). Stabilization interventions in the treatment of traumatized refugees: A scoping review. *Cambridge Prisms: Global Mental Health*, **12**, e73, 1–16
<https://doi.org/10.1017/gmh.2025.10028>

Received: 16 August 2024

Revised: 04 May 2025

Accepted: 29 May 2025

Keywords:


refugee; posttraumatic stress; cross-cultural; forced displacement; stabilization

Corresponding author:

Irja Rzepka-Marot;

Email: irja.rzepka-marot@med.uni-heidelberg.de

Stabilization interventions in the treatment of traumatized refugees: A scoping review

Irja Rzepka-Marot^{1,2} , Nadja Gebhardt^{1,2}, Jonathan Nowak^{1,2}, Bastian Bruns^{1,2}, Hans-Christoph Friederich^{1,2} and Christoph Nikendei^{1,2}

¹Department of General Internal Medicine and Psychosomatics, University Hospital Heidelberg, Germany and ²Deutsches Zentrum für Psychische Gesundheit, DZPG (German Centre for Mental Health – Partner Site Heidelberg/Mannheim/Ulm), Germany

Abstract

Refugees and forced migrants are particularly susceptible to trauma-related disorders, due exposure to traumatic events before, during or after displacement. In trauma therapy, the concept of psychological stabilization refers to the improvement of a patient's capacity to manage symptoms and emotions associated with traumatic experiences. While exposure-based therapies are widely recommended for treating posttraumatic stress disorder (PTSD), stabilizing interventions may offer a valuable alternative, particularly given the unique challenges in refugee care. This scoping review aims to provide a comprehensive overview of stabilizing, non exposure-based interventions for traumatized refugees. A systematic search identified 31 relevant studies featuring diverse interventions, settings, and outcomes. Most studies showed a significant reduction in PTSD symptoms compared to waitlist (six studies), treatment as usual (three studies) and pre-post analyses (nine studies), though nine studies found no difference between intervention and comparison group. Notably, two studies found the stabilizing approach less effective than the comparison group, and two reported no symptom reduction in pre-post analysis. Heterogeneity among the examined interventions as well as living conditions was high and limited the generalizability of the results. Further studies should take these environmental factors into consideration.

Impact statement

This scoping review investigates the potential of stabilizing interventions as an alternative approach to exposure-based therapies for treating trauma-related disorders in refugee populations, a group particularly vulnerable to trauma. By offering a comprehensive review of nonexposure-based interventions, this study provides valuable insights into the current state of research on stabilizing interventions for refugees under different living conditions. The 31 studies included in the review were comprised of over 15 different interventions implemented through various formats. They showed mixed results, but most reported a significant reduction in PTSD symptoms compared to waitlist, treatment as usual or in pre-post analyses. These findings highlight the importance of future research that considers the different living conditions of refugees.

Introduction

By the end of 2023, approximately 117 million individuals worldwide – around 1 % of the global population – were forced to leave their places of residence. This displacement occurred within their country of origin, neighboring countries or countries further away (UNHCR, 2024). As refugees are frequently exposed to traumatic events (Abu Suhaiban et al., 2019; Nesterko et al., 2019; Acarturk et al., 2021), they are disproportionately affected by trauma-related disorders compared to the general population, with most studies reporting a prevalence rate of posttraumatic stress disorder (PTSD) exceeding 30% (Kaltenbach et al., 2018; Blackmore et al., 2020). PTSD is characterized by re-experiencing through intrusions, flashbacks or nightmares, hyperarousal and avoidance of trauma-related stimuli (World Health Organisation, 2020), which can significantly diminish the quality of life (Monson et al., 2017; Lefebvre et al., 2021). Moreover, in refugee populations, mental health impairments are closely associated with integration difficulties (Schick et al., 2016). These reasons call for efficient and timely treatment.

Recent research has demonstrated the efficacy of exposure-based therapy interventions for the treatment of PTSD (McLean et al., 2022). As a result, it is now the standard treatment recommendation in several treatment guidelines (International Society for Traumatic Stress Studies Guidelines Committee, 2018; Hamblen et al., 2019; Schäfer et al., 2019). A defining

© The Author(s), 2025. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

 Cambridge
Prisms

 CAMBRIDGE
UNIVERSITY PRESS

characteristic of exposure-based interventions is that patients are confronted with traumatic memories or trauma-related stimuli in a therapeutic manner, with the objective of processing the associated emotions (Rothbaum and Schwartz, 2002). However, exposure-based interventions are also associated with a higher dropout rate than other psychotherapeutic interventions, possibly related to the distress resulting from the confrontation with intense negative emotions (Lewis *et al.*, 2020). Outside of controlled clinical study conditions, various factors on the clinician side, such as concerns about worsening symptoms, or on the patient side, such as comorbidities, also contribute to the infrequent implementation of exposure-based therapies. These factors can lead to the adaptation of manualized interventions to better suit individual cases, rather than being strictly implemented as originally designed (Najavits, 2015). As exposure-based interventions might lead to considerable treatment-associated distress through the reprocessing of traumatic experiences and the accompanying emotions (Foa *et al.*, 2002), an adequate level of psychological stability is warranted. Moreover, the stability of the therapeutic relationship – encompassing both continuity of treatment, particularly during exposure to traumatic memories and the reliability of the therapist–patient bond to prevent therapy discontinuation – is essential in order to successfully conduct exposure-based interventions (Gjerstad *et al.*, 2024).

The prerequisites for exposure-based interventions are particularly hard to fulfill when treating PTSD in refugee populations, as there are many legal and structural obstacles which must be overcome to provide refugees with healthcare (Giacco *et al.*, 2014). In general, the development of a stable therapeutic relationship can be impeded by a high degree of mistrust often displayed by individuals who have lived through interpersonal trauma (Hembree *et al.*, 2003; Olatunji *et al.*, 2009). The language barrier is often identified as a significant challenge in clinical healthcare settings, rendering the establishment of a stable therapeutic relationship more challenging. The lack of interpreters (Bell and Zech, 2009) or inadequate reimbursement policies often impede patients from accessing necessary and adequate treatment of any kind (Helmboldt *et al.*, 2019). Access to adequate mental health care often depends on a person's asylum status (Bell and Zech, 2009). Moreover, the uncertainty regarding legal status and housing, commonly faced by refugees, along with cultural challenges, can threaten psychological stability. This may deter mental health professionals from providing exposure-based interventions, especially when treatment continuity cannot be guaranteed (Bell and Zech, 2009). The systemic challenges of mental health care for refugees apply to all mental disorders and therapeutic approaches. They also have an impact on the implementation and execution of studies, making it challenging to collect data of high quality (Panter-Brick *et al.*, 2020; Hinchey *et al.*, 2023b). Nevertheless, attempts have been made to assess the efficacy of the exposure-based treatments for refugees and asylum seekers, with promising results (Kaltenbach *et al.*, 2020), including narrative exposure therapy (NET) therapy in a refugee camp in Uganda (Neuner *et al.*, 2008) and eye movement desensitization and reprocessing (EMDR) in a refugee camp in Turkey (Acarturk *et al.*, 2015, 2016; Yurtsever *et al.*, 2018). When examining individual studies, the following aspects should be considered: it is important to note that the improvement in PTSD symptoms among participants of NET therapy (Neuner *et al.*, 2008) was reported at the 1-year follow-up, during which most participants in this group no longer resided in the camp, raising questions about whether the symptom improvement was attributable to the intervention or improved living conditions (Mundt *et al.*, 2014). During the follow-up of a group EMDR therapy, no differences were observed between

the intervention and control groups, despite an initial reduction in symptoms in the intervention group. The authors primarily attributed this outcome to the persistently stressful living conditions in the refugee camp (Yurtsever *et al.*, 2018). In a meta-analysis by Turrini *et al.*, no significant effectiveness was found for NET and EMDR in refugee populations (Turrini *et al.*, 2019). This finding contrasts with the results of a meta-analysis conducted by Nosè *et al.*, which specifically investigated psychological interventions for refugees in high-income countries and demonstrated the effectiveness of NET in this setting (Nosè *et al.*, 2017). Nonetheless, it should be noted at this point that a meta-analysis by Turrini *et al.* also identified trauma-focused cognitive behavioral therapy (CBT) as an effective intervention, with sustained effects at follow-up despite the presence of postmigratory stressors (Turrini *et al.*, 2019). While trauma exposure-based interventions have shown effectiveness, albeit with variable levels of evidence, alternative treatment approaches for situations in which the prerequisites for exposure-based interventions are not met could represent a valuable addition to improving the mental health of refugee populations.

Stabilizing interventions, which are commonly well-established in clinical settings (Rosner *et al.*, 2015; Equit *et al.*, 2018), represent an alternative treatment approach. They can reduce trauma-related symptoms but can also serve as a preparation for exposure-based interventions as part of a phase-based approach (Willis *et al.*, 2023). They are designed to assist trauma survivors in managing trauma-related symptoms without using maladaptive regulation strategies. In psycho-traumatology, the term “stable” is commonly used to describe a person who is capable of coping with trauma-related stimuli, emotions and memories, without risk of serious deterioration in their general physical and mental well-being. This encompasses the absence of behaviors such as self-harm, suicidal ideation, substance abuse and dissociative episodes (Reddemann, 2011). In this context, stability mainly refers to a trauma survivor's inner stability (e.g., coping with symptoms) but also includes external aspects and risk assessments, such as social stability (social support system and network), physical and psychological safety (perpetrator contact, living conditions) and the nature of the patient–therapist relationship (Sack and Gromes, 2013; Seidler *et al.*, 2015). While many definitions of stabilization interventions exist, our review uses the definition proposed by Luise Reddemann, as it aligns with the current clinical understanding of stabilization in treating trauma-related disorders. According to Reddemann, stabilization interventions aim to enhance symptom management, emotion regulation and the acquisition of new competencies (Reddemann, 2011; Reddemann and Piedfort-Marin, 2017). This is achieved through regaining a sense of control (Herman, 1992), interpersonal safety (Willis *et al.*, 2023) and strengthening socio-psychological skills (Ter Heide *et al.*, 2016). Stabilization techniques do not make use of traumatic memories. Thus, psychological stability, treatment continuity and the development of a stable therapeutic relationship are of lesser importance than exposure-based therapy.

The necessity of stabilization interventions is a topic of critical debate. Concerns have been raised that stabilization interventions might delay the implementation of evidence-based exposure-based interventions (Neuner *et al.*, 2008; De Jongh *et al.*, 2016). While many treatment manuals for exposure-based interventions, such as EMDR, incorporate stabilization elements (Foa *et al.*, 2008; Shapiro, 2017), usually evaluations assess the therapy manual as a whole (Rosner *et al.*, 2015). Therefore, the specific impact of the stabilization elements cannot be determined independently from the overall treatment effect and systematic evidence for the efficacy of stabilizing interventions is lacking (National Institute for Health

and Care Excellence, 2018; Berliner et al., 2019). Thus, the aim of this scoping review is to provide an overview of the current research on stabilizing, nonexposure-based interventions for refugees with trauma-related disorders through a systematic search of the literature. In light of the expansive scope of this definition, the review will initially present the interventions conducted in the included studies, followed by an analysis of the study designs, including participants and outcomes.

Methods

Search strategy

As this was the first review on stabilizing interventions in refugee populations and considering the large variety of implementations of stabilizing interventions, we decided to conduct a scoping review of the literature. Herein, we adhered to the guidelines laid out by the PRISMA-ScR checklist (Tricco et al., 2018). We conducted a systematic search to identify studies that examine stabilizing interventions and evaluate their effect on the symptom burden of PTSD in adult refugees. With regard to the criterion of whether a study examines a stabilizing approach, we have used the above-mentioned definition by Luise Reddemann (2011) as a guideline. The final decision as to whether an intervention was “stabilizing” was the subject of discussion among the reviewers. The screening process was conducted by three independent reviewers (IR, NG, JN). The search was conducted on PubMed, Embase, Web of Science, PsycInfo and CINAHL.

Inclusion criteria

Studies were included if the sample was comprised of refugees or forced migrants worldwide at all stages of flight, i.e. internally displaced persons, people in refugee camps or the postmigration phase at various stages of the asylum process. We included intervention studies, such as randomized controlled trials, as well as single interventions from all regions of the world. The respective interventions could be performed by professionals or lay providers. Individual and group interventions were included. Further, we included studies that evaluated the effect of stabilizing interventions on reducing PTSD symptom load, with PTSD as either the primary or secondary outcome. Initially, we planned to include only studies focusing on adult refugees ≥18 years. However, some projects targeted communities, families or “youth,” with some individuals being under 18 years. Therefore, we adjusted our criteria to include studies in which the majority of participants were adult refugees. Exclusion criteria were other types of publications such as abstracts, conference papers or dissertations, single case studies, systematic reviews and meta-analyses, studies conducted among only underage refugees or qualitative studies. In addition, we excluded studies that did not provide sufficient details about stabilizing interventions, multimodal interventions and interventions that included trauma exposure elements. No date restriction was placed on this search. Only papers in English or German language were included. Additionally, we conducted a forward reference search to identify further publications on the topic. Search terms related to the population were: refugees OR asylum seekers OR forced migration OR displaced people. Search terms related to the outcome were PTSD or posttraumatic stress disorder OR trauma OR traumatized. Search terms related to therapy were therapy OR intervention OR treatment OR psychotherapy OR stabilization. The full search

Table 1. Key search terms for EMBASE

Population	refugee*, “asylum seeker*”, “displaced people”, “displaced person*” “displaced population”, “forced migra*”, refugee/exp., “forced migrant”/exp OR/1–8:ti,ab AND ([english]/lim OR [german]/lim)
Intervention	“therap*”, intervention*, treat*, stabiliz*, stabilis*, psychother*, psychosoc*, therapy/exp., intervention/exp OR/1–9:ti,ab AND ([english]/lim OR [german]/lim)
Outcome	PTSD, “Posttraumatic stress disorder,” “Post-traumatic stress disorder,” traumatized, trauma, “trauma-affected,” Trauma/exp., Psychotrauma/exp., Posttraumatic stress disorder/exp OR/1–9:ti,ab

Note: Date of search: 10.11.2023.

strategy for EMBASE can be found in Table 1. The protocol of this scoping review can be assessed at OSF (<https://osf.io/z3dcy>).

Data extraction and management

For each included study, information on author and year of publication, study design, country of study conduction, study population, sample size and gender distribution, inclusion criteria, information on the stabilizing intervention, PTSD outcome measure and PTSD symptom outcome was collected by the reviewers on a data collection form. The full screening process is displayed in Figure 1.

Risk of bias assessment

For each included study, we also conducted a risk of bias evaluation. For this purpose, we utilized the Cochrane evaluation tools, specifically the revised RoB 2 tool for randomized controlled trials (Sterne et al., 2019) and the ROBINS-I tool for nonrandomized studies (Sterne et al., 2016).

Results

Study selection and procedure

We indentified a total of 5,115 studies after automated removal of duplicats (see Figure 1). Following the screening of titles and abstracts, 5,048 results were excluded. The full text was assessed for eligibility for 67 studies. A total of seven studies were excluded because the intervention included exposure-based elements. Another 11 studies were excluded because the specifications in the study design did not meet the definition of stabilization (Kruse et al., 2009; Renner et al., 2011; Jespersen and Vuust, 2012; Ter Heide and Smid, 2015; Stammel et al., 2017; Yurtsever et al., 2018; Shultz et al., 2019; Trilesnik et al., 2019; Park et al., 2020; Gever et al., 2023; Graef-Calliess et al., 2023). Three studies were excluded because the available information about the examined intervention was insufficient (Neuner et al., 2010; Rees et al., 2013, 2014). With regard to the intervention, six studies lacked any measure of PTSD symptomatology (Renner et al., 2008; Sonne et al., 2016, 2021; Acarturk et al., 2022; Aizik-Reebs et al., 2022; Orang et al., 2022). Three studies could not be found (Renner and Peltzer, 2008; Kayal et al., 2013; Sonne et al., 2019), and six were conference




Figure 1. PRISMA 2020 flow diagram.

abstracts or nonpeer-reviewed dissertations (Mojica-Castillo, 2003; Stenmark *et al.*, 2008; Ekstrøm *et al.*, 2016; Bryant, 2022; De Graaff *et al.*, 2022; Stöckli *et al.*, 2023). A total of 31 studies were included in our final analysis. An overview of the included studies can be found in Table 2. The 31 studies included a total of 2,759 participants (1,192 male, 1,269 female, 298 not known) receiving a stabilizing intervention. An overview of the included interventions can be found in Table 2.

Examined stabilizing interventions

The 31 studies analyzed presented a variety of stabilizing treatment approaches. A total of 15 interventions were conducted in a group

format (Yeomans *et al.*, 2010; Stanford *et al.*, 2014; Im *et al.*, 2018; Zehetmair *et al.*, 2018; Alsheikh Ali, 2020; Koch *et al.*, 2020; Lancaster and Gaede, 2020; Tol *et al.*, 2020; Aizik-Reeb *et al.*, 2021; Akhtar *et al.*, 2021; Acarturk *et al.*, 2022; Bryant *et al.* 2022a,b; Griggs *et al.*, 2022; Hasha *et al.*, 2022), 14 in individual sessions (Neuner *et al.*, 2004; Ter Heide *et al.*, 2011; Hensel-Dittmann *et al.*, 2011; Stenmark *et al.*, 2013; Meffert *et al.*, 2014; Ter Heide *et al.*, 2016; Brakemeier *et al.*, 2017; Altawil *et al.*, 2018; Carlsson *et al.*, 2018; De Graaff *et al.*, 2020; Barhoma *et al.*, 2021; Knefel *et al.*, 2022; Orang *et al.*, 2022; De *et al.*, 2023) and two as smartphone-based interventions (Mazzulla *et al.*, 2021; Röhr *et al.*, 2021).

Seven studies explored the effectiveness of Problem Management Plus (PM+), a transdiagnostic intervention developed by the World

Table 2. Overview of included studies

Author & year of publication	Study design	Country	Participants	Inclusion criteria	Stabilizing intervention	PTSD outcome measure	Outcome for PTSD symptomatology
Acarturk et al. (2022)	Pilot RCT with enhanced care as usual	Turkey	Syrian refugees	K10 ¹ > 15 WHODAS ² 2.0 > 16	Group Problem Management Plus (gPM+)	PCL-5 ³	Significant effect of time of both interventions ($F(2,88) = 10.01$, $p = .000$, $d = .66$.)
Aizik-Reebs et al. (2021)	RCT with waitlist	Israel	Eritrean refugees	Not specified	Mindfulness-based Trauma Recovery for Refugees (MBTR-R)	HTQ ⁴	Significant reduction of PTSD symptoms relative to waitlist group $F(1,74) = 12.44$, $p = .001$, $\eta^2 = .17$
Akhtar et al. (2021)	Feasibility RCT with enhanced care as usual	Jordan	Syrian refugees	K10 > 16 WHODAS ² 2.0 > 17	Group Problem Management Plus (gPM+)	PCL-5 ³	Descriptive reduction of mean for PTSD symptoms in intervention group (pre: 32.00 (19.21), post: 22.86 (17.00))
Alsheikh Ali et al. (2020)	Quasi experimental RCT with control group	Jordan	Syrian refugee women	PCL-5 ³ , Ryff's psychological well-being scales (PWB),	Counseling with focus on alienation, loss and grief, hope and hopelessness, and psychological problems	PCL-5 ³	Significant reduction of PTSD symptoms relative to waitlist control group $F(1,35) = 6.69$, $p = .01$
Altawil et al. (2018)	Comparison of 3 interventions, non-randomized	Gaza	Palestinian refugees (12–75 yrs)	Participants with symptoms of PTSD	Community Wellness Focussing (CWP)	PTSD-SRII scale ⁵	Reduction of PTSD symptoms in CWF group ($t(113) = 23.12$, $p < 0.01$)
Barhoma et al. (2021)	RCT with Stress Management vs. Cognitive Restructuring – follow-up	Denmark	Refugees in Denmark	PTSD according to ICD 10	Stress Management (SM)	HTQ ⁴	Nonsignificant interaction effect on PTSD symptoms six ($p = .441$, $d = 0.19$) and 18-month posttreatments ($p = .724$, $d = 0.14$)
Brakemeier et al. (2017)	Feasibility study	Germany	Syrian refugees	Affective/ anxiety/ somatization/ eating disorder, PTSD, substance abuse according to DSM-IV	Interpersonal therapy	PDS ⁶	Nonsignificant reduction of mean for PTSD symptoms in pre–post analysis (pre: 30.57 ± 13.29 ; post: 20.24 ± 12.59)
Bryant (2022)	RCT with enhanced usual care	Jordan	Syrian refugees	K10 ¹ ≥ 16 WHODAS 2.0 ² ≥ 17	Group Problem Management Plus (gPM+)	PCL-5 ³	No significant differences between conditions for changes in PTSD after 3 months (adjusted mean difference – 0.68, 95% CI –4.01 to 2.66; $p = 0.69$)
Bryant (2022)	RCT with enhanced usual care – follow-up	Jordan	Syrian refugees	K10 ¹ ≥ 16 WHODAS 2.0 ² ≥ 17	Group Problem Management Plus (gPM+)	PCL-5 ³	No significant differences between conditions for changes in PTSD after 12 months (adjusted mean difference: –0.06 (95% CI –3.48 to 3.36) $p = 0.97$)
Carlsson et al. (2018)	RCT with Stress Management vs. Cognitive Restructuring	Denmark	Refugees in Denmark	PTSD according to ICD 10	Stress Management (SM)	HTQ ⁴	No significant difference of change for PTSD symptom between groups (difference mean pretreatment: –0.03 (0.08), posttreatment = 0.04 (0.11), $p = 0.07$ (0.09), $p = 0.45$, $d = 0.16$)
De Graaf et al. (2020)	Pilot RCT with care as usual (CAU)	Netherlands	Syrian refugees	K10 ¹ > 15 WHODAS 2.0 ² > 16	Problem Management Plus (PM+) by peer providers	PCL-5 ³	Significant interaction effect in favor of PM+/CAU between time and condition for symptoms of PTSD ($\chi^2(2) = 9.07$; $p = 0.010$)
De Graaf et al. (2023)	RCT with care as usual (CAU)	Netherlands	Syrian refugees	K10 ¹ > 15 WHODAS 2.0 ² > 16	Problem Management Plus (PM+) by peer providers	PCL-5 ³	At 3-month FU, PM+/CAU had greater reductions on PTSD symptoms ($p = 0.0005$, Cohen's $d = 0.39$)

(Continued)

Table 2. (Continued)

Author & year of publication	Study design	Country	Participants	Inclusion criteria	Stabilizing intervention	PTSD outcome measure	Outcome for PTSD symptomatology
Griggs et al. (2022)	Pilot pre-post evaluation	Great Britain	refugees from 22 different countries	Not specified	Manualised stabilization based on CBT (Moving on after trauma, MOAT)	IES-R ⁷	Significant reduction of mean for PTSD symptoms in pre-post analysis ($p = .001$, $\eta^2 = .393$)
Hasha et al. (2022)	RCT with waitlist	Norway	Syrian refugees	IES-R ⁷ > 24	Teaching Recovery Techniques (TRT)	IES-R ⁷	No significant reduction of mean for PTSD relative to waitlist (-1.3 ($-8.7, 6.2$))
Hensel-Dittmann et al. (2011)	RCT Narrative Exposure Therapy vs. Stress Inoculation Training	Germany	Not specified	"history of experiencing organized violence, current PTSD diagnosis"	Stress Inoculation Training (SIT)	CAPS ⁸	Significant time-treatment interaction [$F(3, 52) = 3.08$; $p = 0.05$, $d = 1.42$] in favor for NET
Im et al. (2018)	Pre-Post evaluation	Kenya	Somali refugees	Not specified	Trauma-Informed Psychoeducation (TIPE) delivered by lay counselors	PCL-C ⁹	Significant reduction of mean for PTSD symptoms in pre-post analysis for no/low PTSD (pre: 27.42 (6.67), post: 34.48 (12.83), $t(95) = -.476$, $p < 0.001$) or high PTSD (pre: 50.09 (7.52), post: 31.93 (13.86), $t(44) = 8.188$, $p < 0.001$)
Knefel et al. (2022)	RCT with treatment as usual (TAU)	Austria	Afghan refugees	RHS-15 ¹⁰ > 12 or RHS-15-stress scale >5	Problem Management Plus, adapted version (aPM+)	ITQ ¹¹	Significant reduction of mean for PTSD symptoms in pre-post analysis for aPM+/TAU group (pre: 13.67 (4.22), post: 10.96 (5.50), $p = .005$, $d_z = 0.61$)
Koch et al. (2020)	RCT with waitlist	Germany	Afghan refugees	Male (15–21 yrs), reporting exposure to traumatic events and difficulties in emotion regulation	Skills-Training of Affect Regulation (STARC)	PCL-5 ³	Significant Condition \times Time interactions for PTSD symptoms (Δ dSTARC-Waitlist = 1.19)
Lancaster and Gaede et al. (2020)	Pre-post evaluation	Iraq	IDP ¹²	Not specified	Resilience-based intervention (GROW) provided by paraprofessionals	SPTSS ¹³	Significant reduction of mean for PTSD symptoms in pre-post analysis ($t(765) = 32.22$, $p < .001$, $d = 1.16$)
Mazzulla et al. (2021)	Pre-post evaluation	USA	Somali-Bantu and Nepali-Bhutanese refugees	Not specified	Language-free NESTT app based on cognitive behavioral and acceptance therapy techniques	RHS-15 ¹⁰	Significant reduction of mean for PTSD symptoms in pre-post analysis ($t(17) = 12.23$, $p < .001$, $d = 2.83$)
Meffert et al. (2014)	Pilot RCT with waitlist	Egypt	Sudanese refugees	HTQ ⁴ > 2.3	Interpersonal therapy	HTQ ⁴	Significant reduction of mean for PTSD symptoms in pre-post analysis (pre: 2.92 (0.44), post: 1.76 (0.49), difference: -1.16 (0.46); effect size for group assignment: 2.52)
Neuner et al. (2004)	RCT with NET ¹⁴ vs. supportive counseling	Uganda	Sudanese refugees	Not specified	Supportive counseling (SC)	PDS ⁶	significant Time \times Treatment interaction for PTSD symptoms in favor of NET (Wilks's λ .78), $F(6, 54) = 4.30$, $p = .01$, $\eta^2 = .31$
Orang et al. (2022)	RCT with waitlist	Germany	Refugees from Africa, Middle East, Arabic countries	Not specified	Value based counseling (VBC)	PCL-5 ³	Significant reduction of mean for PTSD symptoms in intervention group relative to waitlist (adjusted difference 17.15, 95% CI [10.49, 23.81], effect size 0.76, $p < .001$)
Röhr et al. (2021)	RCT with waitlist	Germany	Syrian refugees	PDS ⁶ -Score ≥ 11	Sanadak App, based on cognitive behavioral therapy	PDS ⁶	Nonsignificant reduction of PTSD symptoms in both groups after 4 weeks (Diff -0.90 , 95% CI -0.24 to 0.47 ; $p = .52$)

(Continued)

Table 2. (Continued)

Author & year of publication	Study design	Country	Participants	Inclusion criteria	Stabilizing intervention	PTSD outcome measure	Outcome for PTSD symptomatology
Stanford et al. (2014)	Pre-post evaluation	Libya	IDP ¹²	Not specified	HOPE-curriculum	PCL-5 ³	Significant reduction of PTSD symptoms in pre-post analysis (t(26) = 3.45, p < 0.01)
Stenmark et al. (2013)	RCT multicenter study with NET ¹⁴ vs. treatment as usual (TAU)	Norway	Refugees from different countries	PTSD diagnosis according to the DSM IV criteria	According to reports: help with such as sleep problems, depressive symptoms, problems related to asylum status, and other practical matters	CAPS ⁸	Significant reduction of PTSD symptoms in both groups (significant main effect of time (F(2, 121.3) 30.11, p < .0001), significant main effect of treatment (F(1,71.1) 4.44, p < .05), significant Time x Treatment interaction (F(2, 122.5) 7.55, p < .001)
Ter Heide et al. (2011)	Pilot RCT with EMDR vs. Stabilization	Netherlands	refugees from different countries	SCID ¹⁵ Modul PTSD and parts of the MINI ¹⁶	Stabilization with the focus on the “here and now”	HTQ ⁴	No significant of PTSD symptom reduction in either condition, but nonsignificant deterioration of PTSD symptoms for stabilization in pre-post analysis (pre:2.74 (0.27), post: 3.04 (0.25))
Ter Heide et al. (2016)	RCT with EMDR vs. Stabilization as usual	Netherlands	Refugees from different countries	PTSD diagnosis according to the DSM-IV	Stabilizing interventions according to patient's needs	CAPS ⁸ , HTQ ⁴	No significant differences between the two groups were found in either linear or quadratic slopes and effect sizes between the groups were small
Tol et al. (2020)	Cluster RCT with enhanced usual care	Uganda	Sudanese female refugees	Kessler 6 ¹⁷ ≥ 5	Self-Help Plus, based on ACT	PCL-6 ¹⁸	Significant reduction of PTSD symptoms posttreatment in intervention group compared to control group (mixed model analysis: −3.53 (−4.67 to −2.38), p < 0.0001 effect size = −0.68)
Yeomans et al. (2010)	RCT with waitlist control	Burundi	IDP ⁶ in Burundi	Not specified	“Healing and Reconciling Our Communities” with and without psychoeducation workshop	HTQ ⁴	Significant reduction of PTSD symptoms in both intervention groups (F(2, 117) = 6.87, p < .01, partial η ² = .11), with participants in the condition without psychoeducation showing a trend for having less severe PTSD symptoms
Zehetmair et al. (2018)	Pre-post evaluation	Germany	English speaking refugees	≥3 symptoms in PC-PTSD-5 ¹⁹ questionnaire	Mindfulness based and imaginative stabilization techniques	PC-PTSD-5 ¹⁹	No significant reduction of PTSD symptoms in pre-post analysis (z = −1.93, p = 0.06, r = −0.47)

Note: ALTAWIL, M., NEL, P., ASKER, A., SAMARA, M. & HARROLD, D. 2008. The effects of chronic war trauma among Palestinian children. *Children: The invisible victims of war-An interdisciplinary study: Peterborough: DSM Technical Publications Ltd.* ¹K10: Kessler Distress Scale, 10 items, ²WHODAS=World Health Organization Disability Assessment Schedule, ³PCL-5: Posttraumatic Stress Disorder Checklist, ⁴HTQ: Harvard Trauma Questionnaire, ⁵PTSD-SR11 Scale: see Altawil et al. (2008), ⁶PDS: Posttraumatic Diagnostic Scale, ⁷IES-R: Impact of Event Scale-Revised, ⁸CAPS: Clinician Administered Scale, ⁹PCL-C: PTSD Checklist – civilian version, ¹⁰RHS-15: Refugee Health Screener-15, ¹¹ITQ: International Trauma Questionnaire, ¹²IDP: Internally displaced person, ¹³SPTSS: Screen for Posttraumatic Stress Symptoms, ¹⁴NET: Narrative Exposure Therapy, ¹⁵SCID: structured clinical interview, for DSM-IV Axis I Disorders (SCID-I), Modul PTSD, ¹⁶MINI: Mini International Neuropsychiatric Interview, ¹⁷Kessler 6: Kessler Distress Scale, 6 items, ¹⁸PCL-6: PTSD Checklist – 6 items version, ¹⁹PC-PTSD-5: Primary-Care PTSD-5.

Health Organization (WHO) that addresses common mental health issues. PM+ offers strategies to manage stress and addresses problems through different techniques such as relaxation, problem-solving, behavioral activation and enhancing social support (Dawson et al., 2015). This intervention modality has been conducted by both professional (Kantor et al., 2017) and peer providers (De Graaff et al., 2020, 2023), in individual and group formats (Bryant et al. 2022a,b) and across diverse living conditions (Akhtar et al., 2021; Knefel et al., 2022).

Three studies explored the efficacy of mindfulness-based interventions. These included, for instance, mindfulness-based stress reduction for refugees (MBSR-R) (Aizik-Reebs et al., 2021) and

mindfulness-related and imaginative stabilization techniques that incorporated exercises such as guided imagery. The effectiveness of Acceptance and Commitment Therapy (ACT) (Tol et al., 2020), which integrates elements of mindfulness strategies with CBT, was also evaluated (Self-Help Plus) (Hayes and Pierson, 2005).

Furthermore, three studies examined stress management (SM) as an intervention approach (Carlsson et al., 2018; Barhoma et al., 2021). SM or stress inoculation training (SIT) (Hensel-Dittmann et al., 2011), both encompass relaxation techniques, attention diversion and behavioral activation. The underlying assumption is that

inadequate coping strategies may precipitate pathological stress (Lazarus and Folkman, 1984).

A similar approach was followed by three CBT-based interventions, which specifically addressed the symptom clusters of PTSD rather than general distress (Moving on after trauma, MOAT; teaching recovery techniques (TRT); HOPE-curriculum) (Stanford et al., 2014; Griggs et al., 2022; Hasha et al., 2022).

Two studies investigated interpersonal therapy as an intervention (Meffert et al., 2014; Brakemeier et al., 2017). Interpersonal therapy addresses interpersonal problems as significant contributing factors to the development and progression of mental health impairments, including grief, role transitions, role disputes and interpersonal deficiencies such as social isolation (Lipsitz and Markowitz, 2013). The authors of the respective publications examined this therapeutic approach in more detail, arguing that both traumatic experiences, especially those of an interpersonal nature and forced displacement adversely affect interpersonal relationships and can perpetuate negative cycles (Meffert et al., 2014).

Additionally, we included two smartphone-based interventions (Sanadak app; NESTT app) (Mazzulla et al., 2021; Röhr et al., 2021). Both were based on a CBT approach, with one being language-free and incorporating ACT techniques (Mazzulla et al., 2021).

One study explored a transdiagnostic approach to improve emotion regulation skills (Skills-Training of Affect Regulation (STARC)) (Koch et al., 2020). The training has been culturally adapted to the needs of Afghan refugees based on the model of Dialectic Behavioral Therapy (DBT) (Linehan, 2014) and Skills Training in Affective and Interpersonal Regulation (STAIR) (Cloitre et al., 2010).

The remaining 11 studies examined additional approaches to stabilization, including the evaluation of different kinds of counseling. These included value-based counseling (VBC) (Orang et al., 2022) and counseling that emphasizes grieving and loss, hope and hopelessness and alienation (Alsheikh Ali, 2020). Other interventions focused on increasing resilience and dealing with feelings of helplessness (Community Wellness Focussing (CWP)) (Altawil et al., 2018), or on fostering an individual's religiousness, thankfulness, kindness, hope and courage (GROW) (Lancaster and Gaede, 2020).

Two studies investigated a manual that, in addition to psychoeducation and regulation strategies, also addresses the topics of stigma, migration stress, collective trauma (Trauma-Informed Psychoeducation, TIPE) (Im et al., 2018) and, respectively, the healing of interpersonal relationships (Healing and Reconciling Our Communities) (Yeomans et al., 2010).

Finally, three studies employed stabilization (Ter Heide et al., 2011, 2016), counseling (Neuner et al., 2004) or treatment as usual as a control condition, focusing on "sleep problems, depressive symptoms, problems related to asylum status and other practical matters" (Stenmark et al., 2008). One of the studies focused on enhancing physical safety and well-being as well as the implementation of body-oriented interventions to ease PTSD-related symptoms (Ter Heide et al., 2011). The second study focused on the enhancement of emotional regulation and the development of relational skills (Ter Heide et al., 2016). The third study employed a nonstructured counseling intervention, tailored to the patients needs with the aim of controlling for non-specific treatment effects (Neuner et al., 2004).

Study designs

Different study designs were used to assess the effects of the interventions reviewed. Of the 31 studies, 23 were RCTs (Neuner

et al., 2004; Yeomans et al., 2010; Hensel-Dittmann et al., 2011; Ter Heide et al., 2011, 2016; Stenmark et al., 2013; Meffert et al., 2014; Carlsson et al., 2018; Alsheikh Ali, 2020; De Graaff et al., 2020, 2023; Koch et al., 2020; Tol et al., 2020; Aizik-Reebs et al., 2021; Akhtar et al., 2021; Barhoma et al., 2021; Röhr et al., 2021; Acarturk et al., 2022; Bryant et al., 2022a, b; Knefel et al., 2022; Hasha et al., 2022; Orang et al., 2022), of which four were pilot evaluations (Meffert et al., 2014; De Graaff et al., 2020; Akhtar et al., 2021; Acarturk et al., 2022). Eight studies compared the intervention with a wait-list control (Yeomans et al., 2010; Meffert et al., 2014; Alsheikh Ali, 2020; Koch et al., 2020; Aizik-Reebs et al., 2021; Röhr et al., 2021; Hasha et al., 2022; Orang et al., 2022), eight further studies had an active control group, mostly compared with treatment as usual (De Graaff et al., 2020, 2023; Tol et al., 2020; Akhtar et al., 2021; Knefel et al., 2022; Acarturk et al., 2022; Bryant et al., 2022a, b). Four studies used a stabilizing intervention as an active control group for NET (Neuner et al., 2004; Stenmark et al., 2013) or EMDR (Ter Heide et al., 2011, 2016). Three studies compared two different interventions, one of which can be classified as stabilizing (Hensel-Dittmann et al., 2011; Carlsson et al., 2018; Barhoma et al., 2021). The remaining eight studies were conducted as a pre-post analysis (Stanford et al., 2014; Brakemeier et al., 2017; Altawil et al., 2018; Im et al., 2018; Zehetmair et al., 2018; Lancaster and Gaede, 2020; Mazzulla et al., 2021; Griggs et al., 2022), with one study comparing three interventions and assigning participants according to need rather than randomly (Altawil et al., 2018).

Participant mental health burden

Inclusion criteria differed between studies in terms of the mental health burden of the study populations: 12 of the studies included refugees with a formal diagnosis of PTSD, assessed with questionnaires such as the HTQ, PDS, IES-R, PC-PTSD-5 (Meffert et al., 2014; Zehetmair et al., 2018; Alsheikh Ali, 2020; Röhr et al., 2021; Hasha et al., 2022) or described as according to ICD-10 or DSM-IV/V criteria (Ter Heide et al., 2011, 2016; Stenmark et al., 2013; Carlsson et al., 2018; Barhoma et al., 2021), or only "with PTSD" without displaying the diagnostic process or tools (Hensel-Dittmann et al., 2011; Altawil et al., 2018), or reporting exposure to traumatic events (Koch et al., 2020). One study included refugees if they met DSM-IV criteria for PTSD, affective, anxiety, somatization, eating disorders or substance abuse (Brakemeier et al., 2017). In seven other studies, refugees were included if they showed a general level of psychological distress (De Graaff et al., 2020, 2023; Akhtar et al., 2021; Acarturk et al., 2022; Bryant et al., 2022a, b; Knefel et al., 2022), as measured with the Kessler Distress Scale, Kessler 6, WHO Disability Assessment Schedule 2.0 (WHODAS 2.0) or Refugee Health Screener 15 (RHS-15). Nine studies did not specify their inclusion criteria (Neuner et al., 2004; Yeomans et al., 2010; Stanford et al., 2014; Im et al., 2018; Lancaster and Gaede, 2020; Aizik-Reebs et al., 2021; Mazzulla et al., 2021; Griggs et al., 2022; Orang et al., 2022). One study (Altawil et al., 2018) compared three interventions aimed at individuals and communities who had been exposed to severely traumatizing experiences.

Outcome

The studies reported varying outcomes for the investigated treatment approaches (see Table 3). Nine studies showed significant reductions in PTSD symptoms for participants in the stabilizing intervention group: compared with treatment as usual, Self-Help Plus, PM+ and Mindfulness-based Trauma Recovery for Refugees

(De Graaff et al., 2020, 2023; Tol et al., 2020) showed positive outcomes. Compared with a waitlist control group, counseling that emphasizes grieving and loss, hope and hopelessness and alienation, TRTs, Skills-Training of Affect Regulation, Healing and Reconciling Our Communities and VBC (Yeomans et al., 2010; Alsheikh Ali, 2020; Koch et al., 2020; Aizik-Reebs et al., 2021; Hasha et al., 2022; Orang et al., 2022), showed favorable outcomes in six interventions. Eight studies found significant reductions in PTSD symptoms in pre-post analyses of CBT-based stabilization (MOAT), trauma-informed psychoeducation (TIPE), resilience-based intervention (GROW), peer-lead recovery group, community wellness focusing, an adapted version of PM+, interpersonal therapy or a CBT- and ACT-based (NESTT) (Meffert et al., 2014; Stanford et al., 2014; Altawil et al., 2018; Im et al., 2018; Lancaster and Gaede, 2020; Mazzulla et al., 2021; Griggs et al., 2022; Knefel et al., 2022). One study using gPM+ reported a reduction in mean PTSD symptom scores without testing for statistical significance (Akhtar et al., 2021).

Nine studies found no difference in the efficacy of a stabilizing approach in reducing PTSD symptoms: compared with treatment as usual, PM+, gPM+ did not show a better outcome (C Acarturk et al., 2022; Bryant et al. 2022a, b). When a stabilizing approach was the control condition (TAU: support with accompanying problems and practical matters), there was a significant symptom reduction in both groups compared with NET, with NET showing a greater reduction (Stenmark et al., 2013). The use of a CBT-based self-help app (Sanadak) compared with waitlist did not show significant differences in PTSD symptom reduction (Röhr et al., 2021). SM compared with cognitive restructuring also showed no significant differences in outcome, with both approaches showing nonsignificant symptom reductions (Carlsson et al., 2018; Barhoma et al., 2021). When EMDR was compared with stabilization with the focus on the here and now or “according to the patients needs,” no significant differences between the two groups were described (Ter Heide et al., 2011, 2016), with one study reporting a nonsignificant deterioration of PTSD symptomatology for the stabilization group (Ter Heide et al., 2011). The stabilizing approach was inferior to another intervention (NET) in two studies (Neuner et al., 2004; Hensel-Dittmann et al., 2011). Two further studies assessing interpersonal therapy or the use of guided imagery reported no reduction in symptoms (Brakemeier et al., 2017; Zehetmair et al., 2018). A brief summary of the results can be found in Table 3.

Risk of bias assessment

The results of the risk of bias assessment are presented in two separate figures: Figure 2 for randomized studies and Figure 3 for

nonrandomized studies. It is evident that the majority of studies exhibit a high overall risk of bias.

Discussion

The aim of this scoping review was to give an overview of the current literature on stabilizing, nonexposure-based interventions for refugees with trauma-related disorders. With a total number of 31 studies examined, six trials reported a significant reduction of PTSD symptom burden after conduction of a stabilizing intervention when compared to waitlist, three when compared to treatment as usual, nine in a pre-post analysis. Nine studies found no difference in the effectiveness of PTSD symptom reduction when compared to another intervention.

Implemented stabilization interventions and underlying definitions of stabilization

The question of how to define stabilization played a significant role throughout the review. Over 15 different interventions were implemented, often in varying formats (e.g., group or individual sessions, professional or lay counselors, app-based approaches) and in some

Table 3. Summary of the results

Stabilizing compared against TAU	• Three studies showing significant symptom reduction of PTSD
Stabilizing intervention against waitlist	• Six studies showing significant symptom reduction of PTSD
Stabilizing intervention in pre-post analysis	• Nine studies showing significant symptom reduction of PTSD • Two studies showing no significant symptom reduction of PTSD
Stabilizing intervention in comparison with another intervention	• Nine studies no showing any difference on PTSD symptom reduction • Two studies showing inferiority of stabilizing intervention in terms of symptom reduction of PTSD




Figure 2. Risk assessment for randomized-controlled trials (RoB2).




Figure 3. Risk assessment for non-randomized trials (ROBINS-I).

cases with limited details about the interventions. This lack of specificity made precise differentiation between the interventions challenging. It is noteworthy that, in most publications, the interventions were not explicitly described as stabilizing, though the reviewed interventions encompassed various dimensions of stabilization. Several approaches targeted emotional stabilization, employing methods such as skills training and SM to enhance individuals' capacity to manage anxiety and overwhelming emotions (Carlsson et al., 2018; Koch et al., 2020). Mindfulness-based interventions were also prominent, fostering decentering, self-compassion and reduced emotional reactivity (Aizik-Reeb et al., 2021; Aizik-Reeb et al., 2022). These can counteract typical symptoms such as hyperarousal, emotional numbness or negative mood and cognition or feelings of guilt and shame. Other interventions addressed interpersonal difficulties, which are often central to trauma-related disorders, particularly in the context of interpersonal trauma (Meffert et al., 2014; Alsheikh Ali, 2020). One study focused explicitly on rebuilding community relationships in postcivil war Burundi, emphasizing that restoring trust was pivotal in alleviating PTSD symptoms (Yeomans et al., 2010). Efforts to address everyday challenges, contributing to social stabilization, were exemplified by the Problem Management Plus intervention (Knefel et al., 2022), alongside general counseling approaches and "treatment as usual," which also provided support in navigating asylum processes (Stenmark et al., 2013).

It is possible that with a different underlying referential definition, further or different studies would have been included. For example, one study comparing social media-based drama, music and art therapy was excluded because the interventions described did not align with the definitional criteria used as a reference framework (Gever et al., 2023). On the other hand, studies were included in which psychological counseling was conducted, with one study addressing "problems related to asylum status, and other practical matters," among other aspects (Stenmark et al., 2013). This raises the question of where the boundaries between psychotherapeutic stabilization, dealing with everyday struggles and social work support lie. Existing literature highlights that prolonged asylum procedures, temporary housing and language barriers exacerbate PTSD symptoms among

refugees (Li et al., 2016; Kartal et al., 2019). Accordingly, interventions that help to deal with social difficulties could possibly have a stabilizing effect and contribute to the reduction of PTSD symptomatology. Therefore, it may be important to emphasize the aspect of social stabilization more prominently in therapeutic interventions and further investigate its impact.

Contextual factors influencing the effectiveness of stabilization interventions

The overall analysis indicates that the effectiveness of the stabilizing interventions is closely associated with the participants' living conditions. Some studies found no significant effects of the stabilizing intervention on PTSD symptoms. The reasons given by the authors included the intervention's focus on addressing daily struggles, while the absence of exposure-based therapy was suggested as a factor in preventing changes in PTSD symptoms (Hasha et al., 2022). In another study, challenging environmental conditions such as poverty, separation from family, concern for their safety in the country of origin, and loneliness were identified as limiting factors for therapeutic success (Bryant, 2022; Bryant et al., 2022b), as the same intervention was successful under different living conditions (De Graaff et al. 2020, 2023). Similarly, another study hypothesized that living conditions in a reception center, combined with an uncertain residency status, contributed to the lack of improvement in PTSD symptoms (Zehetmair et al., 2018). It appears that the environmental conditions serve both as an explanatory model for why an intervention does not lead to symptom reduction but are also directly addressed as part of the therapy for stabilization. An uncontrolled study demonstrated that the number of postmigratory stressors, as well as ongoing conflicts in the country of origin, were associated with reduced symptom improvement through therapy. Additionally, an insecure residency status was linked to an increased likelihood of therapy dropout (Djelantik et al., 2020). However, a longitudinal study indicated that psychosocial interventions addressing postmigratory stressors primarily led to symptom reduction in depression and anxiety but not in PTSD (Schick et al.,

2018). Although no definitive conclusions can be drawn in this regard, the significant relevance of postmigratory stressors as an influencing factor can be acknowledged. These stressors repeatedly emerge as a critical issue in therapeutic settings (Bruhn et al., 2018).

The aspect of individual living conditions and the question of whether interventions are adapted to these conditions or even specifically address coping with them has been highlighted by many authors. The need to address daily stressors and postmigration difficulties (Knefel et al. 2022; De Graaff et al. 2023), which are known to contribute to a higher symptom burden (Gleeson et al., 2020), was emphasized. For example, Lancaster and Gaede (2020) investigated a resilience-based approach that aims to foster a person's religiousness, gratitude, kindness, hope and courage in order to help them become more resilient. This program relies on non-professional providers in a group format and posits that resilience is a necessary skill for living and surviving in a refugee camp. Moreover, the intervention problem management plus focuses on problem-solving strategies, SM, behavioral activation and strengthening social support networks. This aims to address psychosocial challenges (De Graaff et al., 2020; Bryant, 2022; Bryant et al., 2022b; De et al., 2023). It is important to note, however, that peri- and postmigratory stressors vary significantly in every context. This scoping review included studies on refugees worldwide, whose living conditions are difficult to compare. For instance, individuals living in refugee camps face different challenges than those who have arrived in a destination country and are seeking asylum. Even within the latter group, significant differences exist between those with secure residency status and those without it regarding their mental health burden (Laban et al., 2008) and effectiveness of therapeutic interventions (Ter Heide and Smid, 2015). A systematic analysis of the WHO intervention PM+ reveals that the same intervention produces highly heterogeneous outcomes across different settings (Schäfer et al., 2023). A comparable large-scale comparison is not available for other stabilization-focused studies with refugees or asylum seekers. To minimize the variability introduced by external factors, different interventions could be compared under similar circumstances rather than comparing the same intervention under different circumstances.

Limited resources and resulting adaptations of stabilization interventions

The challenge of limited resources is a recurring issue in refugee treatment settings and frequently necessitates context-specific adaptations of interventions. Some of the respective examined interventions are easily learnable and implementable by laypersons or peers (Meffert et al., 2014; Stanford et al., 2014; Lancaster and Gaede, 2020; De Graaff et al., 2023). This highlights the challenge of limited resources, as interventions delivered by laypersons can reach a larger population compared to those requiring trained mental health professionals. It should be noted that there is also a study on NET conducted by laypersons (Neuner et al., 2008), though this is not the case for other forms of exposure therapy, such as EMDR. Additionally, other forms of resource constraints can impact the execution of studies. For example, a study conducted in a camp for internally displaced persons reported shortages of paper and printing facilities for therapy materials (Stanford et al., 2014). Group interventions also appear to be a better format for effective resource utilization. This is also reflected in the studies presented, as 15 of the 31 studies were conducted in a group format. Resource considerations also underpinned the two studies examining interventions via apps, as these are flexible and, once established,

resource-efficient. Mazulla et al. addressed another aspect by creating a language-free app to reach a larger number of people (Mazulla et al., 2021). The current state of knowledge about smartphone-based mental health interventions for refugees was evaluated in a systematic review. The authors summarized that, up to now, none of the apps examined sufficiently met the needs of the target group (El-Haj-Mohamad et al., 2023). These factors should not only be accounted for in research studies but also considered when designing interventions for implementation outside of a research context. Not only due to limited resources but also to address shared mechanisms underlying common mental health issues, some authors have implemented transdiagnostic interventions that target PTSD alongside other conditions (Koch et al., 2020; Knefel et al., 2022). This approach was motivated by the frequent comorbidities associated with PTSD and the recognition that psychological distress among refugees extends beyond PTSD (Fazel et al., 2005; Hinchey et al., 2023a), despite the latter often being the primary focus of research (Akhtar et al., 2021; Acarturk et al., 2022; Bryant et al. 2022a, b; Knefel et al., 2022). A broader focus also allows for reaching more individuals experiencing psychological distress, which is particularly relevant in resource-limited settings, such as refugee camps or mass accommodations. Moreover, such approaches do not rely on formal diagnoses or corresponding specific interventions, making them more resource-efficient. Additionally, avoiding formal diagnostic procedures can help mitigate potential stigma, which might otherwise hinder access to effective treatment (Lancaster and Gaede, 2020). A systematic analysis of the barriers to mental health care among refugee populations also showed that it was primarily self-stigma and the fear of social consequences that prevented those affected from seeking professional help (Byrow et al., 2020). Another transdiagnostic intervention employed in refugee populations is the common elements treatment approach, which shows a significant reduction in PTSD symptom burden (Bolton et al., 2014; Bogdanov et al., 2021). However, these studies were excluded from this review because the manual for participants in low- and middle-income countries includes gradual exposure and in-vivo exposure (Murray et al., 2014).

Needs and limitations of cultural adaptation of stabilization interventions

The studies included also repeatedly highlighted the limited applicability of Western concepts, which may not meet the needs of the target population (Altawil et al., 2018; Im et al., 2018). For SIT, the lack of cultural adaptation of the intervention for non-Western patients was also identified as a possible explanation as to why the intervention, contrary to the hypothesis, did not lead to significant symptom improvement (Hensel-Dittmann et al., 2011). The use of peers in delivering interventions could better address this aspect during the implementation of the intervention and overcome the language and cultural barriers, thus representing a lower-threshold access to psychosocial care. In some of the interventions, adaptations to the culture of the sample had already taken place within the intervention. For example, the "Community Wellness Focussing" intervention, which was carried out in Gaza, included a session on "Proverbs and Quran exercises" (Altawil et al., 2018). In a study conducted in Burundi in a camp for internally displaced persons on the other hand, the focus was on restoring social relationships within the community (Yeomans et al., 2010). Collaborative work in the community was also a major part of the study with trauma-informed psychoeducation for Somali refugees in Kenya, with sessions on "Stigma, Collective trauma, Collective healing" (Im et al., 2018). In another study, efforts were made to adapt

the intervention to the sample by consulting an advisor from the same country. In this case, gender issues and the timing of the intervention were adapted. However, no further details were disclosed (Hasha *et al.*, 2022).

Implications and summary

It becomes apparent that the authors of the presented studies provided varied responses to the challenge of addressing the complex treatment conditions of traumatized refugees, which is reflected in the heterogeneity of the study designs, interventions and results. The living conditions in which these studies were examined differ considerably. The systemic difficulties in providing psychosocial support for refugees also have an impact on the conduct of studies. The studies presented here, which were conducted under the living conditions in humanitarian settings, are subject to a variety of factors that influence the results, which must be considered in detail (Panter-Brick *et al.*, 2020; Hinchey *et al.*, 2023b). In 18 out of the 31 studies presented, a significant reduction in PTSD symptoms was observed as a result of the stabilizing intervention. However, a meta-analysis would be necessary to validly assess the effectiveness of stabilizing interventions for refugees, as the methods used in this review cannot provide definitive conclusions on this matter. The focus of the evaluation should consider the living conditions of refugees, as the outcomes of the same intervention may not readily translate from refugee camps to life circumstances in the country of resettlement (Acarturk *et al.*, 2022; Bryant *et al.*, 2022a), even though many factors, such as insecure residency status (Laban *et al.*, 2008), family separation (Fogden *et al.*, 2020), temporary housing (Ziersch *et al.*, 2017; Leiler *et al.*, 2019), unemployment (Lai *et al.*, 2022) and language acquisition (Kartal *et al.*, 2019), are already known to contribute to psychological distress. Furthermore, studies published in non-English languages or found in the gray literature could also be included to provide a more comprehensive and globally representative overview of stabilization approaches. Given the contextual diversity among refugee populations, qualitative studies could furthermore deepen the understanding of how refugees interpret and experience stabilization interventions in relation to their cultural, political and social realities, taking host-country-specific context into account.

Limitations

Some limitations of this scoping review should be acknowledged: both during the screening process and the final selection of studies, stabilization was understood according to the definition provided by Luise Reddemann (Reddemann, 2011; Reddemann and Piedfort-Marin, 2017). Consequently, studies that did not align with this definition or did not offer sufficient information to ascertain their fit were excluded. With regard to the results of the bias assessment, it can be seen that limitations arise primarily due to missing data, which appear to be understandable under the living circumstances mentioned with high fluctuation and social and economic deprivation. In addition, it should be mentioned that there is a certain risk of bias in all studies, as the interventions cannot be blinded and the measurements were always patient-reported outcomes. Another important limitation should be acknowledged: among the included studies, nine did not clearly specify their inclusion criteria. In eight studies, participants were included based on a general psychological

burden without further specification, and one study also included participants with other diagnoses. Seven studies reported the presence of a PTSD diagnosis but did not provide details about the diagnostic procedures. Only six studies explicitly defined a PTSD diagnosis as an inclusion criterion and described the respective diagnostic instruments. However, the majority of the studies identified the reduction of trauma-related symptoms as their objective, even if PTSD was not defined as an inclusion criterion. A similar issue applies to the age inclusion criterion: some studies also targeted participants under 18 years old, and, in a few studies, the age of the participants was not entirely clear, leading to ambiguities in defining the target population. We chose not to exclude certain studies based on the following rationale: Refugees are frequently exposed to traumatic events, which is associated with a significantly higher prevalence of PTSD compared to the general population. Although current clinical guidelines recommend exposure-based interventions, such approaches cannot always be implemented for the reasons outlined in our introduction. A strict, linear logic – where only studies are included that demonstrate a direct, simple link between a traumatic event, a formally diagnosed PTSD using a standardized instrument, and an intervention specifically tailored to PTSD symptoms – would have led to an overly narrow selection. Such an approach would not sufficiently reflect the complexity of this research question and research environment, as elaborated in various aspects throughout the discussion. To account for this complexity and to offer a more comprehensive picture of the existing evidence, we deliberately applied broader inclusion criteria in our review.

Conclusion

The findings of this scoping review indicate that a range of stabilizing interventions for refugees have been explored. The 31 studies that were included yielded heterogeneous results, with most showing significant PTSD symptom reduction compared to waitlist, treatment as usual or in pre-post analyses, though some found no differences between interventions, and a few reported the stabilizing approach as less effective or ineffective. Both the heterogeneity of the interventions and the environmental conditions under which the studies were conducted limit the generalizability of the results. Future studies or publications should place greater emphasis on incorporating the specific living conditions under which their results are obtained. Additionally, qualitative studies could provide valuable insights by engaging affected individuals to identify factors they perceive as contributing to symptom improvement.

Open peer review. To view the open peer review materials for this article, please visit <http://doi.org/10.1017/gmh.2025.10028>.

Data availability statement. Data sharing is not applicable – no new data is generated.

Author contribution. IR: conceptualization, data curation, investigation, formal analysis, writing – original draft, writing – reviewing & editing, NG: investigation, formal analysis, writing – reviewing & editing, JN: investigation, formal analysis, writing – reviewing & editing, BB: investigation, formal analysis, writing – rewriting and editing, HCF: resources, CN: supervision, resources, writing – reviewing & editing.

Financial support. This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Competing interests. The authors declare none.

References

- Abu Suhaiban H, Grasser LR and Javanbakht A (2019) Mental health of refugees and torture survivors: A critical review of prevalence, predictors, and integrated care. *International Journal of Environmental Research and Public Health* **16**(13), 2309.
- Acarturk C, Konuk E, Cetinkaya M, Senay I, Sijbrandij M, Cuijpers P and Aker T (2015) EMDR for Syrian refugees with posttraumatic stress disorder symptoms: Results of a pilot randomized controlled trial. *European Journal of Psychotraumatology* **6**(1), 27414.
- Acarturk C, Konuk E, Cetinkaya M, Senay I, Sijbrandij M, Gulen B and Cuijpers P (2016) The efficacy of eye movement desensitization and reprocessing for post-traumatic stress disorder and depression among Syrian refugees: Results of a randomized controlled trial. *Psychological Medicine* **46**(12), 2583–2593.
- Acarturk C, McGrath M, Roberts B, Ilkursorun Z, Cuijpers P, Sijbrandij M, Sondorp E, Ventevogel P, McKee M, Fuhr DC and On behalf of the Sc (2021) Prevalence and predictors of common mental disorders among Syrian refugees in Istanbul, Turkey: A cross-sectional study. *Social Psychiatry and Psychiatric Epidemiology* **56**(3), 475–484. <https://doi.org/10.1007/s00127-020-01941-6>.
- Acarturk C, Uygun E, Ilkursorun Z, Yurtbakan T, Kurt G, Adam-Troian J, Senay I, Bryant R, Cuijpers P and Kiselev N (2022) Group problem management plus (PM+) to decrease psychological distress among Syrian refugees in Turkey: A pilot randomised controlled trial. *BMC Psychiatry* **22**, 1–11.
- Azizik-Reebs A, Amir I, Yuval K, Hadash Y and Bernstein A (2022) Candidate mechanisms of action of mindfulness-based trauma recovery for refugees (MBTR-R): Self-compassion and self-criticism. *Journal of Consulting and Clinical Psychology* **90**(2), 107–122. <https://doi.org/10.1037/ccp0000716>.
- Azizik-Reebs A, Yuval K, Hadash Y, Gebreyohans Gebremariam S and Bernstein A (2021) Mindfulness-based trauma recovery for refugees (MBTR-R): Randomized waitlist-control evidence of efficacy and safety. *Clinical Psychological Science* **9**(6), 1164–1184. <https://doi.org/10.1177/2167702621998641>.
- Akhtar A, Giardinelli L, Bawaneh A, Awwad M, Al-Hayek H, Whitney C, Jordans MJ, Sijbrandij M, Cuijpers P and Dawson K (2021) Feasibility trial of a scalable transdiagnostic group psychological intervention for Syrians residing in a refugee camp. *European Journal of Psychotraumatology* **12**(1), 1932295.
- Alsheikh Ali ASS (2020). Efficiency of intervention counseling program on the enhanced psychological well-being and reduced post-traumatic stress disorder symptoms among Syrian women refugee survivors. *Clinical Practice and Epidemiology in Mental Health: CP & EMH* **16** (Suppl-1), 134.
- Altawil MA, El Asam A and Khadaroo A (2018) The effectiveness of therapeutic and psychological intervention programs in PTC-GAZA. *Journal of Child & Adolescent Trauma* **11**, 473–486.
- Barhoma M, Sonne C, Lommen MJ, Mortensen EL and Carlsson J (2021) Stress management versus cognitive restructuring in trauma-affected refugees – A follow-up study on a pragmatic randomised trial. *Journal of Affective Disorders* **294**, 628–637.
- Bell P and Zech E (2009) Access to mental health for asylum seekers in the European Union. An analysis of disparities between legal rights and reality. *Archives of Public Health* **67**(1), 1–15. <https://doi.org/10.1186/0778-7367-67-1-30>.
- Berliner L, Bisson J, Cloitre M, Forbes D, Jensen TK, Lewis C, Monson CM, Olff M, Pilling S and Riggs DS (2019) The international society for traumatic stress studies new guidelines for the prevention and treatment of posttraumatic stress disorder: Methodology and development process. *Journal of Traumatic Stress* **32**(4), 475–483.
- Blackmore R, Boyle JA, Fazel M, Ranasinha S, Gray KM, Fitzgerald G, Misso M and Gibson-Helm M (2020) The prevalence of mental illness in refugees and asylum seekers: A systematic review and meta-analysis. *PLoS Medicine* **17**(9), e1003337. <https://doi.org/10.1371/journal.pmed.1003337>.
- Bogdanov S, Augustinavicius J, Bass JK, Metz K, Skavenski S, Singh NS, Moore Q, Haroz EE, Kane J, Doty B, Murray L and Bolton P (2021) A randomized-controlled trial of community-based transdiagnostic psychotherapy for veterans and internally displaced persons in Ukraine. *Global Mental Health* **8**, e32. <https://doi.org/10.1017/gmh.2021.27>.
- Bolton P, Bass JK, Zangana GAS, Kamal T, Murray SMI, Kaysen D, Lejuez CW, Lindgren K, Pagoto S, Murray LK, Van Wyk SS, Ahmed AMA, Mohammad Amin NM and Rosenblum M (2014) A randomized controlled trial of mental health interventions for survivors of systematic violence in Kurdistan, northern Iraq. *BMC Psychiatry* **14**(1), 360. <https://doi.org/10.1186/s12888-014-0360-2>.
- Brakemeier E-L, Zimmermann J, Erz E, Bollmann S, Rump S, von Kempfki V, Grossmüller T, Mitelman A, Gehrisch J and Spies J (2017) Interpersonelles Integratives Modellprojekt für Geflüchtete mit psychischen Störungen. *Psychotherapeut* **62**, 322–332.
- Bruhn M, Rees S, Mohsin M, Silove D and Carlsson J (2018) The range and impact of postmigration stressors during treatment of trauma-affected refugees. *The Journal of Nervous and Mental Disease* **206**(1), 61–68.
- Bryant R (2022) A randomised controlled trial of a lay-provider psychological intervention for common mental disorders for Syrian refugees in Jordan. *Australian and New Zealand Journal of Psychiatry* **56**(SUPPL 1), 120–121. <https://doi.org/10.1177/00048674221088686>.
- Bryant RA, Bawaneh A, Awwad M, Al-Hayek H, Giardinelli L, Whitney C, Jordans MJ, Cuijpers P, Sijbrandij M and Ventevogel P (2022a) Effectiveness of a brief group behavioral intervention for common mental disorders in Syrian refugees in Jordan: A randomized controlled trial. *PLoS Medicine* **19**(3), e1003949.
- Bryant RA, Bawaneh A, Awwad M, Al-Hayek H, Giardinelli L, Whitney C, Jordans MJ, Cuijpers P, Sijbrandij M and Ventevogel P (2022b) Twelve-month follow-up of a randomised clinical trial of a brief group psychological intervention for common mental disorders in Syrian refugees in Jordan. *Epidemiology and Psychiatric Sciences* **31**, e81.
- Byrow Y, Pajak R, Specker P and Nickerson A (2020) Perceptions of mental health and perceived barriers to mental health help-seeking amongst refugees: A systematic review. *Clinical Psychology Review* **75**, 101812. <https://doi.org/10.1016/j.cpr.2019.101812>.
- Carlsson J, Sonne C, Vindbjerg E and Mortensen EL (2018) Stress management versus cognitive restructuring in trauma-affected refugees – A pragmatic randomised study. *Psychiatry Research* **266**, 116–123.
- Cloitre M, Stovall-McClough KC, Noonan K, Zorbas P, Cherry S, Jackson CL, Gan W and Petkova E (2010) Treatment for PTSD related to childhood abuse: A randomized controlled trial. *American Journal of Psychiatry* **167**(8), 915–924.
- Dawson KS, Bryant RA, Harper M, Kuowei Tay A, Rahman A, Schafer A and van Ommeren M (2015) Problem management plus (PM+): A WHO transdiagnostic psychological intervention for common mental health problems. *World Psychiatry* **14**(3), 354–357. <https://doi.org/10.1002/wps.20255>.
- De Graaff A, Cuijpers P, McDaid D, Park A, Woodward A, Bryant R, Fuhr D, Kieft B, Minkenberg E and Sijbrandij M (2020) Peer-provided problem management plus (PM+) for adult Syrian refugees: A pilot randomised controlled trial on effectiveness and cost-effectiveness. *Epidemiology and Psychiatric Sciences* **29**, e162.
- De Graaff A, Sijbrandij M and Cuijpers P (2022) Scalable psychological interventions for Syrian refugees: Preliminary results of a randomized controlled trial on the peer-refugee delivered problem management plus (PM+) intervention in the Netherlands. *European Psychiatry* **65**, S636. <https://doi.org/10.1192/j.eurpsy.2022.1632>.
- De Graaff AM, Cuijpers P, Twisk JW, Kieft B, Hunaidy S, Elsayy M, Gorgis N, Bouman TK, Lommen MJ and Acarturk C (2023) Peer-provided psychological intervention for Syrian refugees: Results of a randomised controlled trial on the effectiveness of problem management plus. *BMJ Mental Health* **26**(1), e300637.
- De Jongh A, Resick PA, Zoellner LA, Van Minnen A, Lee CW, Monson CM, Foa EB, Wheeler K, Broeke E and Feeny N (2016) Critical analysis of the current treatment guidelines for complex PTSD in adults. *Depression and Anxiety* **33**(5), 359–369.
- Djelantik AAAMJ, de A, Kuiper D, Kleber RJ, Boelen PA and Smid GE (2020) Post-migration stressors and their association with symptom reduction and non-completion during treatment for traumatic grief in refugees. *Frontiers in Psychiatry* **11**, 407. <https://doi.org/10.3389/fpsy.2020.00407>.
- Ekström M, Carlsson J, Sonne C and Mortensen EL (2016) Stress management versus cognitive restructuring: A randomized clinical study on traumatized refugees. *European Psychiatry* **33**, S399–S400. <https://doi.org/10.1016/j.eurpsy.2016.01.1437>.

- El-Haj-Mohamad R, Nohr L, Niemeyer H, Böttche M and Knaevelsrud C (2023) Smartphone-delivered mental health care interventions for refugees: A systematic review of the literature. *Glob Ment Health (Camb)* 10, e6. <https://doi.org/10.1017/gmh.2022.61>.
- Equit M, Maurer S, Michael T and Köllner V (2018) Konfrontation oder Stabilisierung: Wie planen Verhaltenstherapeuten die Behandlung bei Post-traumatischer Belastungsstörung? *Verhaltenstherapie* 28(1), 7–14.
- Fazel M, Wheeler J and Danesh J (2005) Prevalence of serious mental disorder in 7000 refugees resettled in western countries: A systematic review. *The Lancet* 365(9467), 1309–1314. [https://doi.org/10.1016/S0140-6736\(05\)61027-6](https://doi.org/10.1016/S0140-6736(05)61027-6).
- Foa EB, Chrestman KR and Gilboa-Schechtman E (2008) *Prolonged Exposure Therapy for Adolescents with PTSD Emotional Processing of Traumatic Experiences, Therapist Guide*. New York: Oxford University Press.
- Foa EB, Zoellner LA, Feeny NC, Hembree EA and Alvarez-Conrad J (2002) Does imaginal exposure exacerbate PTSD symptoms? *Journal of Consulting and Clinical Psychology* 70(4), 1022. <https://doi.org/10.1037//0022-006x.70.4.1022>.
- Fogden G, Berle D and Steel Z (2020) The impact of family separation and worry about family on psychological adjustment in refugees resettled in Australia. *Journal of Traumatic Stress* 33(6), 894–907. <https://doi.org/10.1002/jts.22568>.
- Gever VC, Iyendo TO, Obiugo-Muoh UO, Okunade JK, Agujiobi-Odoh N, Udengwu N, Talabi FO and Nwokolo PN (2023) Comparing the effect of social media-based drama, music and art therapies on reduction in post-traumatic symptoms among Nigerian refugees of Russia's invasion of Ukraine. *Journal of Pediatric Nursing* 68, e96–e102. <https://doi.org/10.1016/j.pedn.2022.11.018>.
- Giacco D, Matanov A and Priebe S (2014) Providing mental healthcare to immigrants: Current challenges and new strategies. *Current Opinion in Psychiatry* 27(4), 282–288. <https://doi.org/10.1097/ycp.0000000000000065>.
- Gjerstad SF, Nordin L, Poulsen S, Spadaro EFA and Palic S (2024) How is trauma-focused therapy experienced by adults with PTSD? A systematic review of qualitative studies. *BMC psychology* 12(1), 135.
- Gleeson C, Frost R, Sherwood L, Shevlin M, Hyland P, Halpin R, Murphy J and Silove D (2020) Post-migration factors and mental health outcomes in asylum-seeking and refugee populations: A systematic review. *European Journal of Psychotraumatology* 11(1), 1793567. <https://doi.org/10.1080/20008198.2020.1793567>.
- Graef-Calliess IT, Erdmann L, Mohwinkel V, Özkan I, Finkelstein D, Loos K, Penteker G and Trilesnik B (2023) Post-migration living difficulties, discrimination, and mental health of traumatized refugees in Germany: Data from the refuKey project for timely and need-adapted treatment in a stepped-care setting. *International Review of Psychiatry* 35(3–4), 339–351. <https://doi.org/10.1080/09540261.2022.2164181>.
- Griggs M, Liu C and Cooper K (2022) Pilot evaluation of a group stabilisation intervention for refugees and asylum seekers with PTSD. *Behavioural and Cognitive Psychotherapy* 50(1), 111–116. <https://doi.org/10.1017/S135246582100028X>.
- Hamblen JL, Norman SB, Sonis JH, Phelps AJ, Bisson JI, Nunes VD, Megnin-Viggars O, Forbes D, Riggs DS and Schnurr PP (2019) A guide to guidelines for the treatment of posttraumatic stress disorder in adults: An update. *Psychotherapy* 56(3), 359.
- Hasha W, Igland J, Fadnes LT, Kumar BN, Heltne UM and Diaz E (2022) Effect of a self-help group intervention using teaching recovery techniques to improve mental health among Syrian refugees in Norway: A randomized controlled trial. *International Journal of Mental Health Systems* 16(1), 47.
- Hayes SC and Pierson H (2005) Acceptance and commitment therapy. In: Freeman, A., Felgoise, S.H., Nezu, C.M., Nezu, A.M., Reinecke, M.A. (eds), *Encyclopedia of Cognitive Behavior Therapy*. Boston, MA: Springer. https://doi.org/10.1007/0-306-48581-8_1.
- Helmboldt L, Nikendei C and Kindermann D (2019) Sprachmittlung – Einsatz von DolmetscherInnen in der Traumatherapie von Geflüchteten. *Psychother. Dialog (Print)* 20(02), 95–99.
- Hembree EA, Rauch SAM and Foa EB (2003) Beyond the manual: The insider's guide to prolonged exposure therapy for PTSD. *Cognitive and Behavioral Practice* 10(1), 22–30. [https://doi.org/10.1016/S1077-7229\(03\)80005-6](https://doi.org/10.1016/S1077-7229(03)80005-6).
- Hensel-Dittmann D, Schauer M, Ruf M, Catani C, Odenwald M, Elbert T and Neuner F (2011) Treatment of traumatized victims of war and torture: A randomized controlled comparison of narrative exposure therapy and stress inoculation training. *Psychotherapy and Psychosomatics* 80(6), 345–352. <https://doi.org/10.1159/000327253>.
- Herman JL (1992) *Trauma and Recovery: The Aftermath of Violence-From Domestic Abuse to Political Terror*. New York: Hachette UK.
- Hinchey L, Nashef R, Bazzi C, Gorski K and Javanbakht A (2023a) The longitudinal impact of war exposure on psychopathology in Syrian and Iraqi refugee youth. *International Journal of Social Psychiatry* 69(7), 1833–1836.
- Hinchey LME, Khalil D and Javanbakht A (2023b) Practical approaches to conducting biopsychosocial research with refugee and internally displaced communities. *Comprehensive Psychoneuroendocrinology* 16, 100217. <https://doi.org/10.1016/j.cpnec.2023.100217>.
- Im H, Jettner JF, Warsame AH, Isse MM, Khoury D and Ross AI (2018) Trauma-informed psychoeducation for Somali refugee youth in urban Kenya: Effects on PTSD and psychosocial outcomes. *Journal of Child & Adolescent Trauma* 11, 431–441.
- International Society for Traumatic Stress Studies Guidelines Committee (2018) *Posttraumatic Stress Disorder Prevention and Treatment Guidelines Methodology and Recommendations*. Oakbrook Terrace, IL: Author.
- Jespersen KV and Vuust P (2012) The effect of relaxation music listening on sleep quality in traumatized refugees: A pilot study. *Journal of Music Therapy* 49(2), 205–229. <https://doi.org/10.1093/jmt/49.2.205>.
- Kaltenbach E, Hermenau K, Schauer M, Dohrmann K, Elbert T and Schallinski I (2020) Trajectories of posttraumatic stress symptoms during and after narrative exposure therapy (NET) in refugees. *BMC Psychiatry* 20, 1–14.
- Kaltenbach E, Schauer M, Hermenau K, Elbert T and Schallinski I (2018) Course of mental health in refugees – A one year panel survey. *Frontiers in Psychiatry* 9, 352. <https://doi.org/10.3389/fpsy.2018.00352>.
- Kantor V, Knefel M and Lueger-Schuster B (2017) Perceived barriers and facilitators of mental health service utilization in adult trauma survivors: A systematic review. *Clinical Psychology Review* 52, 52–68. <https://doi.org/10.1016/j.cpr.2016.12.001>.
- Kartal D, Alkemade N and Kiropoulos L (2019) Trauma and mental health in resettled refugees: Mediating effect of host language acquisition on posttraumatic stress disorder, depressive and anxiety symptoms. *Transcultural Psychiatry* 56(1), 3–23.
- Kayal H, Gratton J, Blumberg J and Walsh E (2013) XIII ESTSS Conference: “Trauma and its clinical pathways: PTSD and beyond”, Bologna, June 2013: Pre Conference Workshops and Opening Keynote Address, June 6. *European Journal of Psychotraumatology* 4. <https://doi.org/10.3402/ejpt.v4i0.21270>.
- Knefel M, Kantor V, Weindl D, Schiess-Jokanovic J, Nicholson AA, Verginer I, Schäfer I and Lueger-Schuster B (2022) A brief transdiagnostic psychological intervention for afghan asylum seekers and refugees in Austria: A randomized controlled trial. *European Journal of Psychotraumatology* 13(1), 2068911.
- Koch T, Ehring T and Liedl A (2020) Effectiveness of a transdiagnostic group intervention to enhance emotion regulation in young afghan refugees: A pilot randomized controlled study. *Behaviour Research and Therapy* 132, 103689.
- Kruse J, Joksimovic L, Cavka M, Wöller W and Schmitz N (2009) Effects of trauma-focused psychotherapy upon war refugees. *Journal of Traumatic Stress* 22(6), 585–592. <https://doi.org/10.1002/jts.20477>.
- Laban CJ, Komproe IH, Gernaat HB and de Jong JT (2008) The impact of a long asylum procedure on quality of life, disability and physical health in Iraqi asylum seekers in the Netherlands. *Social Psychiatry and Psychiatric Epidemiology* 43(7), 507–515. <https://doi.org/10.1007/s00127-008-0333-1>.
- Lai H, Due C and Ziersch A (2022) The relationship between employment and health for people from refugee and asylum-seeking backgrounds: A systematic review of quantitative studies. *SSM - Population Health* 18, 101075. <https://doi.org/10.1016/j.ssmph.2022.101075>.
- Lancaster SL and Gaede C (2020) A test of a resilience based intervention for mental health problems in Iraqi internally displaced person camps. *Anxiety, Stress, & Coping* 33(6), 698–705.
- Lazarus RS and Folkman S (1984) *Stress, Appraisal, and Coping*. New York: Springer publishing company.
- Lefebvre C, Fortin C and Guay S (2021) Quality of life after violent crime: The impact of acute stress disorder, posttraumatic stress disorder, and other consequences. *Journal of Traumatic Stress* 34(3), 526–537. <https://doi.org/10.1002/jts.22623>.

- Leiler A, Bjärtå A, Ekdahl J and Wasteson E (2019) Mental health and quality of life among asylum seekers and refugees living in refugee housing facilities in Sweden. *Social Psychiatry and Psychiatric Epidemiology* 54(5), 543–551.
- Lewis C, Roberts NP, Andrew M, Starling E and Bisson JI (2020) Psychological therapies for post-traumatic stress disorder in adults: Systematic review and meta-analysis. *European Journal of Psychotraumatology* 11(1), 1729633. <https://doi.org/10.1080/20008198.2020.1729633>.
- Li SS, Liddell BJ and Nickerson A (2016) The relationship between post-migration stress and psychological disorders in refugees and asylum seekers. *Current Psychiatry Reports* 18(9), 1–9.
- Linehan M (2014) *DBT Skills Training Manual*. New York: Guilford Publications.
- Lipsitz JD and Markowitz JC (2013) Mechanisms of change in interpersonal therapy (IPT). *Clinical Psychology Review* 33(8), 1134–1147. <https://doi.org/10.1016/j.cpr.2013.09.002>.
- Mazzulla EC, Fondacaro KM, Weldon H, Dibble M and Price M (2021) Addressing the disparity in refugee mental health services: A pilot study of a traumatic stress intervention utilizing a language-free mhealth application. *Journal of Technology in Behavioral Science* 6(4), 599–608.
- McLean CP, Levy HC, Miller ML and Tolin DF (2022) Exposure therapy for PTSD: A meta-analysis. *Clinical Psychology Review* 91, 102115. <https://doi.org/10.1016/j.cpr.2021.102115>.
- Meffert SM, Abdo AO, Alla OAA, Elmakki YOM, Omer AA, Yousif S, Metzler TJ and Marmar CR (2014) A pilot randomized controlled trial of interpersonal psychotherapy for Sudanese refugees in Cairo. *Egypt. Psychological Trauma: Theory, Research, Practice, and Policy* 6(3), 240.
- Mojica-Castillo S (2003) *The Effectiveness of a Psychosocial Group Intervention on Older Bosnian Female Refugees in Diminishing Loneliness*. 64, Ann Arbor: ProQuest Information & Learning.
- Monson E, Caron J, McCloskey K and Brunet A (2017) Longitudinal analysis of quality of life across the trauma spectrum. *Psychological Trauma* 9(5), 605–612. <https://doi.org/10.1037/tra0000254>.
- Mundt AP, Wünsche P, Heinz A and Pross C (2014) Evaluating interventions for posttraumatic stress disorder in low and middle income countries: Narrative exposure therapy. *Interventions* 12(2), 250–266.
- Murray LK, Dorsey S, Haroz E, Lee C, Alsiahy MM, Haydary A, Weiss WM and Bolton P (2014) A common elements treatment approach for adult mental health problems in low-and middle-income countries. *Cognitive and Behavioral Practice* 21(2), 111–123.
- Najavits LM (2015) The problem of dropout from “gold standard” PTSD therapies. *F1000prime reports* 7, 43. <https://doi.org/10.12703/P7-43>
- National Institute for Health and Care Excellence (2018) Post-traumatic Stress Disorder. Available at www.nice.org.uk/guidance/ng116 (accessed 18 October 2024).
- Nesterko Y, Jäckle D, Friedrich M, Holzapfel L and Glaesmer H (2019) Prevalence of post-traumatic stress disorder, depression and somatisation in recently arrived refugees in Germany: An epidemiological study. *Epidemiology and Psychiatric Sciences* 29, e40. <https://doi.org/10.1017/S2045796019000325>.
- Neuner F (2008) Stabilisierung vor Konfrontation in der Traumatherapie – Grundregel oder mythos? *Verhaltenstherapie* 18(2), 109–118.
- Neuner F, Kurreck S, Ruf M, Odenwald M, Elbert T and Schauer M (2010) Can asylum-seekers with posttraumatic stress disorder be successfully treated? A randomized controlled pilot study. *Cognitive Behaviour Therapy* 39(2), 81–91. <https://doi.org/10.1080/16506070903121042>.
- Neuner F, Onyut PL, Ertl V, Odenwald M, Schauer E and Elbert T (2008) Treatment of posttraumatic stress disorder by trained lay counselors in an African refugee settlement: A randomized controlled trial. *Journal of Consulting and Clinical Psychology* 76(4), 686.
- Neuner F, Schauer M, Klaschik C, Karunakara U and Elbert T (2004) A comparison of narrative exposure therapy, supportive counseling, and psychoeducation for treating posttraumatic stress disorder in an African refugee settlement. *Journal of Consulting and Clinical Psychology* 72(4), 579–587.
- Nosè M, Ballette F, Bighelli I, Turrini G, Purgato M, Tol W, Priebe S and Barbui C (2017) Psychosocial interventions for post-traumatic stress disorder in refugees and asylum seekers resettled in high-income countries: Systematic review and meta-analysis. *PLoS One* 12(2), e0171030.
- Olatunji BO, Deacon BJ and Abramowitz JS (2009) The cruelest cure? Ethical issues in the implementation of exposure-based treatments. *Cognitive and Behavioral Practice* 16(2), 172–180. <https://doi.org/10.1016/j.cbpra.2008.07.003>.
- Orang TM, Missmahl I, Thoele A-M, Valensise L, Brenner A, Gardisi M, Peter H and Kluge U (2022) New directions in the mental health care of migrants, including refugees - a randomized controlled trial investigating the efficacy of value-based counselling. *Clin Psychol Psychother* 29(4), 1433–1446. <https://doi.org/10.1002/cpp.2728>.
- Panther-Brick C, Eggerman M, Ager A, Hadfield K and Dajani R (2020) Measuring the psychosocial, biological, and cognitive signatures of profound stress in humanitarian settings: Impacts, challenges, and strategies in the field. *Conflict and Health* 14, 1–7.
- Park JK, Park J, Elbert T and Kim SJ (2020) Effects of narrative exposure therapy on posttraumatic stress disorder, depression, and insomnia in traumatized north Korean refugee youth. *Journal of Traumatic Stress* 33(3), 353–359. <https://doi.org/10.1002/jts.22492>.
- Reddemann L (2011) Stabilisierung in der Traumatherapie: Eine Standortbestimmung. *Trauma und Gewalt* 5(3), 256–263.
- Reddemann L and Piedfort-Marin O (2017) Stabilization in the treatment of complex post-traumatic stress disorders: Concepts and principles. *European Journal of Trauma & Dissociation* 1(1), 11–17.
- Rees B, Travis F, Shapiro D and Chant R (2013) Reduction in posttraumatic stress symptoms in Congolese refugees practicing transcendental meditation. *Journal of Traumatic Stress* 26(2), 295–298. <https://doi.org/10.1002/jts.21790>.
- Rees B, Travis F, Shapiro D and Chant R (2014) Significant reductions in posttraumatic stress symptoms in Congolese refugees within 10 days of transcendental meditation practice. *Journal of Traumatic Stress* 27(1), 112–115. <https://doi.org/10.1002/jts.21883>.
- Renner W, Bänninger-Huber E and Peltzer K (2011) Culture-sensitive and resource oriented peer (CROP)-groups as a community based intervention for trauma survivors: A randomized controlled pilot study with refugees and asylum seekers from Chechnya. *Australasian Journal of Disaster and Trauma Studies* 2011(1), 1–13.
- Renner W, Lind M and Ottomeyer K (2008) Psychodramatische Gruppentherapie bei traumatisierten Migrantinnen - neue Ergebnisse einer Evaluationsstudie. *Zeitschrift für Psychotraumatologie, Psychotherapiewissenschaft und Psychologische Medizin (Roland Asanger Verlag GmbH)* 6(1), 89–95.
- Renner W and Peltzer K (2008) Culture-sensitive and resource oriented peer-groups (CROP-G) as a community based intervention for trauma survivors: A pilot randomized trial with asylum seekers and refugees from Chechnya. *International Journal of Psychology* 43(3–4), 258–258.
- Röhr S, Jung FU, Pabst A, Grochtdreis T, Dams J, Nagl M, Renner A, Hoffmann R, König H-H and Kersting A (2021) A self-help app for Syrian refugees with posttraumatic stress (Sanadak): Randomized controlled trial. *JMIR mHealth and uHealth* 9(1), e24807. <https://doi.org/10.2196/24807>.
- Rosner R, Henkel C, Ginkel K and Mestel R (2015) Was passiert nach der stationären Stabilisierung mit komplex traumatisierten PTB-Patientinnen? *Zeitschrift für Psychiatrie, Psychologie und Psychotherapie* 58(2), 127–135. <https://doi.org/10.1024/1661-4747/a000017>
- Rothbaum BO and Schwartz AC (2002) Exposure therapy for posttraumatic stress disorder. *American Journal of Psychotherapy* 56(1), 59–75.
- Sack M and Gromes B (2013) Ressourcenorientierte Behandlungsstrategien in der Traumatherapie. *PiD - Psychotherapie im Dialog* 14(01), 30–35.
- Schäfer I, Gast U, Hofmann A, Knaevelsrud C, Lampe A, Liebermann P, Lotzin A, Maercker A, Rosner R and Wöller W (2019) *S3-Leitlinie Post-traumatische Belastungsstörung*. Berlin: Springer.
- Schäfer SK, Thomas LM, Lindner S and Lieb K (2023) World Health Organization’s low-intensity psychosocial interventions: A systematic review and meta-analysis of the effects of problem management plus and step-by-step. *World Psychiatry* 22(3), 449–462. <https://doi.org/10.1002/wps.21129>.
- Schick M, Morina N, Mistridis P, Schnyder U, Bryant RA and Nickerson A (2018) Changes in post-migration living difficulties predict treatment outcome in traumatized refugees. *Frontiers in Psychiatry* 9, 476. <https://doi.org/10.3389/fpsy.2018.00476>.
- Schick M, Zumwald A, Knöpfli B, Nickerson A, Bryant RA, Schnyder U, Müller J and Morina N (2016) Challenging future, challenging past: The relationship of social integration and psychological impairment in traumatized refugees. *European Journal of Psychotraumatology* 7, 28057. <https://doi.org/10.3402/ejpt.v7.28057>.
- Seidler GH, Freyberger HJ and Maercker A (2015) *Handbuch der Psychotraumatologie*. Stuttgart: Klett-Cotta.

- Shapiro F (2017) *Eye Movement Desensitization and Reprocessing (EMDR) Therapy: Basic Principles, Protocols, and Procedures*. New York: Guilford Publications.
- Shultz JM, Verdelli H, Ceballos AG, Hernandez LJ, Espinel Z, Helpman L, Neria Y and Araya R (2019) A pilot study of a stepped-care brief intervention to help psychologically-distressed women displaced by conflict in Bogota, Colombia. *Global Mental Health*, 6, e28. <https://doi.org/10.1017/gmh.2019.26>.
- Sonne C, Carlsson J, Bech P, Vindbjerg E, Mortensen EL and Elklit A (2016) Psychosocial predictors of treatment outcome for trauma-affected refugees. *European Journal of Psychotraumatology* 7, 30907. <https://doi.org/10.3402/ejpt.v7.30907>.
- Sonne C, Mortensen EL, Palic S and Carlsson J (2019) Predictors of treatment outcomes for refugees with posttraumatic stress disorder. *European Neuropsychopharmacology* 29, S147–S148. <https://doi.org/10.1016/j.euroneuro.2019.09.236>.
- Sonne C, Mortensen EL, Silove D, Palic S and Carlsson J (2021) Predictors of treatment outcomes for trauma-affected refugees – Results from two randomised trials. *Journal of Affective Disorders* 282, 194–202. <https://doi.org/10.1016/j.jad.2020.12.095>.
- Stammel N, Knaevelsrud C, Schock K, Walther LCS, Wenk-Ansohn M and Böttche M (2017) Multidisciplinary treatment for traumatized refugees in a naturalistic setting: Symptom courses and predictors. *European Journal of Psychotraumatology* 8(sup2), 1377552. <https://doi.org/10.1080/2008198.2017.1377552>.
- Stanford MS, Elverson TM, Padilla JI and Rogers EB (2014) Feasibility and efficacy of a peer-led recovery group program for war-related trauma in Libya. *South Africa Journal of Psychology* 44(1), 97–105.
- Stenmark H, Catani C, Elbert T and Gotestam KG (2008) Narrative exposure therapy compared to treatment as usual for refugees with PTSD - preliminary results from a randomized controlled trial. *European Psychiatry* 23, S90–S90. <https://doi.org/10.1016/j.eurpsy.2008.01.700>.
- Stenmark H, Catani C, Neuner F, Elbert T and Holen A (2013) Treating PTSD in refugees and asylum seekers within the general health care system. A randomized controlled multicenter study. *Behaviour Research and Therapy* 51(10), 641–647. <https://doi.org/10.1016/j.brat.2013.07.002>.
- Sterne JA, Hernán MA, Reeves BC, Savović J, Berkman ND, Viswanathan M, Henry D, Altman DG, Ansari MT and Boutron I (2016) ROBINS-I: A tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 355, i4919.
- Sterne JA, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, Cates CJ, Cheng H-Y, Corbett MS and Eldridge SM (2019) RoB 2: A revised tool for assessing risk of bias in randomised trials. *BMJ* 366, i4898.
- Stöckli RT, Zöllner V, Burchert S, Haji F, Hosmann M, Aebbersold M, Wabiszczewicz J, Heim E, Knaevelsrud C and Berger T (2023) A psychosocial intervention (sui app) to enhance quality of life in Arabic-speaking refugees in Switzerland: Cultural adaptation, recruitment and preliminary results of an RCT. *Early Intervention in Psychiatry* 17, 303. <https://doi.org/10.1111/eip.13409>.
- Ter Heide FJ and Smid GE (2015) Difficult to treat? A comparison of the effectiveness of treatment as usual in refugees and non-refugees. *BJPsych Bull* 39(4), 182–186. <https://doi.org/10.1192/pb.bp.114.047928>.
- Ter Heide FJJ, Mooren T, Kleijn W, de A and Kleber R (2011) EMDR versus stabilisation in traumatised asylum seekers and refugees: Results of a pilot study. *European Journal of Psychotraumatology* 2(1), 5881.
- Ter Heide FJJ, Mooren TM, van de R, de A and Kleber RJ (2016) Eye movement desensitisation and reprocessing therapy v. stabilisation as usual for refugees: Randomised controlled trial. *British Journal of Psychiatry* 209(4), 311–318. <https://doi.org/10.1192/bjp.bp.115.167775>.
- Tol WA, Leku MR, Lakin DP, Carswell K, Augustinavicius J, Adaku A, Au TM, Brown FL, Bryant RA and Garcia-Moreno C (2020) Guided self-help to reduce psychological distress in south Sudanese female refugees in Uganda: A cluster randomised trial. *The Lancet Global Health* 8(2), e254–e263.
- Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, Moher D, Peters MD, Horsley T and Weeks L (2018) PRISMA extension for scoping reviews (PRISMA-ScR): Checklist and explanation. *Annals of Internal Medicine* 169(7), 467–473.
- Trilesnik B, Altunoz U, Wesolowski J, Eckhoff L, Ozkan I, Loos K, Penteker G and Graef-Calliess IT (2019) Implementing a need-adapted stepped-care model for mental health of refugees: Preliminary data of the state-funded project "Refukey". *Frontiers in Psychiatry* 10(SEP). <https://doi.org/10.3389/fpsy.2019.00688>.
- Turrini G, Purgato M, Acarturk C, Anttila M, Au T, Ballette F, Bird M, Carswell K, Churchill R and Cuijpers P (2019) Efficacy and acceptability of psychosocial interventions in asylum seekers and refugees: Systematic review and meta-analysis. *Epidemiology and Psychiatric Sciences* 28(4), 376–388.
- UNHCR (2024) Refugee Data Finder. Available at <https://www.unhcr.org/refugee-statistics/> (accessed 23 July 2024).
- Willis DN, Dowling APC and O'Reilly PG (2023) Stabilisation and phase-orientated psychological treatment for posttraumatic stress disorder: A systematic review and meta-analysis. *European Journal of Trauma & Dissociation* 7(1), 100311. <https://doi.org/10.1016/j.ejtd.2022.100311>.
- World Health Organisation (2020) ICD-11 for Mortality and Morbidity Statistics. Available at <https://icd.who.int/browse11/l-m/en#/http%3a%2f%2fid.who.int%2fid%2fentity%2f2070699808> (accessed 29 May 2024).
- Yeomans PD, Forman EM, Herbert JD and Yuen E (2010) A randomized trial of a reconciliation workshop with and without PTSD psychoeducation in Burundian sample. *Journal of Traumatic Stress* 23(3), 305–312.
- Yurtsever, Konuk E, Akyüz T, Zat Z, Tükel F, Çetinkaya M, Savran C and Shapiro E (2018) An eye movement desensitization and reprocessing (EMDR) group intervention for Syrian refugees with post-traumatic stress symptoms: Results of a randomized controlled trial. *Frontiers in Psychology* 9, 493. <https://doi.org/10.3389/fpsyg.2018.00493>.
- Zehetmair C, Kaufmann C, Tegeler I, Kindermann D, Junne F, Zipfel S, Herpertz SC, Herzog W and Nikendei C (2018) Psychotherapeutic group intervention for traumatized male refugees using imaginative stabilization techniques – A pilot study in a German reception center. *Frontiers in Psychiatry* 9, 533. <https://doi.org/10.3389/fpsy.2018.00533>.
- Ziersch A, Walsh M, Due C and Duivesteyn E (2017) Exploring the relationship between housing and health for refugees and asylum seekers in South Australia: A qualitative study. *International Journal of Environmental Research and Public Health* 14(9), 1036.