

RESTRICTED PRINCIPAL CLUSTER SETS OF A CERTAIN HOLOMORPHIC FUNCTION

JOHN T. GRESSER

Let D be the open unit disk and let K be the unit circle. We say that α is an arc at $\zeta \in K$ if α is contained in D and is the image of a continuous function $z = z(t)$ ($0 \leq t < 1$) such that $z(t) \rightarrow \zeta$ as $t \rightarrow 1$. We call α a segment at ζ if the function $z = z(t)$ is linear in t . If P is a property which is meaningful for each point of K , we say that nearly every point of K has property P if the exceptional set is a set of first Baire category in K . We assume that the reader is familiar with the rudiments of cluster set theory, and in particular with the terms ambiguous point, Meier point, and Plessner point of a function (cf. [4] or [7]). If f is a function which maps D into the Riemann sphere, if $\zeta \in K$ and α is an arc at ζ , then $C(f, \zeta, \alpha)$ will denote the arc-cluster set of f at ζ along α . The principal cluster set of f at ζ is defined to be the set

$$H(f, \zeta) = \bigcap_{\alpha} C(f, \zeta, \alpha),$$

where α ranges over all arcs of ζ . The nontangential principal cluster set of f at ζ is defined to be the set

$$H^*(f, \zeta) = \bigcap_{\alpha} C(f, \zeta, \alpha),$$

where α ranges over all arcs at ζ for which there is a Stolz angle at ζ containing α . Finally, if Δ is a Stolz angle at ζ , we define the set

$$H_{\Delta}(f, \zeta) = \bigcap_{\alpha} C(f, \zeta, \alpha),$$

where α ranges over all arcs at ζ such that $\alpha \subseteq \Delta$.

Let Δ be a Stolz angle at $\zeta_0 = 1$, and for each $\zeta \in K$ let $\Delta(\zeta)$ be the Stolz angle at ζ which is obtained by rotating Δ about the origin. We

mention as background for this paper two well-known results by E. F. Collingwood. Let f be an arbitrary complex-valued function defined in D . Then

$$C_{A(\zeta)}(f, \zeta) = C(f, \zeta)$$

for nearly every point $\zeta \in K$, and

$$C_B(f, \zeta) = C(f, \zeta)$$

for all but a countable number of points $\zeta \in K$. Here $C_B(f, \zeta)$ denotes the boundary cluster set of f at ζ (cf. [4, pp. 80, 82]). More recently, using analogous definitions and techniques, the boundary principal cluster set of f at ζ , denoted by $B\Pi(f, \zeta)$, has been investigated [5]. This set describes the behavior of $\Pi(f, \xi)$ at points ξ which are near and distinct from ζ . It has been shown that if f is continuous in D , then

$$B\Pi(f, \zeta) = \Pi(f, \zeta)$$

for nearly every point $\zeta \in K$ [5, Theorem 9]. This naturally suggests the question of whether for a continuous function f in D it follows that

$$\Pi_{A(\zeta)}(f, \zeta) = \Pi(f, \zeta)$$

for nearly every point $\zeta \in K$. The purpose of this paper is to answer this question in the negative, even for a holomorphic function; we show, in fact, that an even stronger inequality is possible. We use the symbol \subset in the sequel to denote proper set inclusion.

THEOREM. *Let Δ be a Stolz angle at $\zeta_0 = 1$. Then there exists a holomorphic function f in D such that*

$$\Pi^*(f, \zeta) \subset \Pi_{\Delta(\zeta)}(f, \zeta)$$

for nearly every point $\zeta \in K$.

Proof. Our construction is a modification of a method presented by F. Bagemihl in [2]. If ρ is a segment at ζ , we let $\theta(\rho)$ denote the angle ($0 < \theta(\rho) < \pi$) between ρ and the forward tangent to the unit circle at ζ , and we let $\ell(\rho)$ denote the length of ρ .

For each ternary fraction

$$t = 0 \cdot t_1 t_2 t_3 \dots$$

where each t_j is zero or two, we denote by

$$b(t) = 0 \cdot b_1 b_2 b_3 \dots$$

the binary fraction such that for each $j = 1, 2, 3, \dots$

$$b_j = \begin{cases} 0 & \text{if } t_j = 0 \\ 1 & \text{if } t_j = 2. \end{cases}$$

The set T of all such ternary fractions is the Cantor “middle thirds” set, and the set of all corresponding binary fractions is the closed unit interval. We set $T^* = T - \{1\}$, and for each $t \in T^*$ we let

$$\zeta_t = e^{2\pi i b(t)}.$$

We let θ_0 be any fixed number ($0 < \theta_0 < \pi$) such that $\theta(\rho) > \theta_0$ for every segment ρ at $\zeta_0 = 1$ which is contained in Δ . Then for each $t \in T^*$ we let ρ_t be the segment at ζ_t defined by

$$\theta(\rho_t) = (1 - t)\theta_0/4, \quad \text{and} \quad \ell(\rho_t) = (1 - \sin(\pi t/2)) \sin(\theta_0/4).$$

Since $\ell(\rho_t) < \sin(\theta(\rho_t))$ for $t \in T^*$, it follows that the line perpendicular to ρ_t and passing through the origin does not intersect ρ_t . In addition, $\theta(\rho_t)$ is a decreasing function of t which is always less than $\pi/4$. These observations easily imply that for each $t \in T^*$ the segments in the collection

$$\{\rho_s : s \in T^*, s \geq t, \text{ and } \arg \zeta_t \leq \arg \zeta_s \leq \arg \zeta_t + \pi/2\}$$

are mutually nonintersecting. Thus in order to establish that the segments in the collection

$$P = \{\rho_t : t \in T^*\}$$

are mutually nonintersecting, it suffices to show that for each $t \in T^*$ such that ζ_t lies in the fourth quadrant, the corresponding segment ρ_t lies in the lower half of D . This follows from the observation that for each $t \in T^*$ with $t > 7/9$ we have $t > b(t) > 3/4$ and hence

$$\ell(\rho_t) < 1 - \sin(\pi b(t)/2) < 1 - \cos(2\pi b(t)) < |\sin(2\pi b(t))|.$$

There is a countably dense subset E of K such that for each $\zeta \in E$ there are two segments, say α_ζ and β_ζ , at ζ belonging to P , while for each $\zeta \in K - E$ there is exactly one segment at ζ belonging to P . For each $\zeta \in E$ we let τ_ζ be the segment at ζ defined by

$$\theta(\tau_\zeta) = (1/2)(\theta(\alpha_\zeta) + \theta(\beta_\zeta)) \quad \text{and} \quad \ell(\tau_\zeta) = (1/2)(\ell(\alpha_\zeta) + \ell(\beta_\zeta)),$$

and we let

$$Q = \{\tau_\zeta : \zeta \in E\}.$$

It follows easily from [3, Theorem 1, p. 187–8] that there is a holomorphic function f defined in D such that for every $\rho \in P$

$$f(z) \longrightarrow 0 \text{ as } |z| \longrightarrow 1 \text{ along } \rho,$$

and for every $\tau \in Q$

$$f(z) \longrightarrow \infty \text{ as } |z| \longrightarrow 1 \text{ along } \tau.$$

Because E , and $K - E$ are both dense in K , we have that $0, \infty \in C(f, \zeta)$ for every $\zeta \in K$, and hence no point of K is a Meier point of f . Therefore, nearly every point of K is a Plessner point of f [6, Theorem 6, p. 330]. We will show that

$$(1) \quad \infty \in \Pi_{\mathcal{A}(\zeta)}(f, \zeta)$$

for every Plessner point ζ which is not an ambiguous point of f . Since there are at most countably many ambiguous points of f [1, Theorem 2, p. 380], and since $\Pi^*(f, \zeta) \subseteq \{0\}$ for every $\zeta \in K$, the verification of (1) will complete the proof.

Let ζ be a Plessner point of f which is not an ambiguous point of f , and suppose, to the contrary, that γ is an arc at ζ with $\gamma \subseteq \mathcal{A}(\zeta)$ such that $\infty \notin C(f, \zeta, \gamma)$. If ρ is the segment at ζ belonging to P , then $\infty \in C(f, \zeta, \rho)$. Since the region bounded by γ , ρ , and any third convenient arc in D contains a Stolz angle, it follows that the full cluster set of f at ζ restricted to this subregion is total. By the Gross-Iversen theorem [4, Theorem 5.8, p. 101] it follows that ∞ is an asymptotic value of f at ζ , and this contradicts our assumption that ζ is not an ambiguous point of f . Thus (1) is established and the proof is complete.

REFERENCES

- [1] F. Bagemihl, Curvilinear cluster sets of arbitrary functions, Proc. Nat. Acad. Sci., **41** (1955), 379–382.
- [2] F. Bagemihl, The chordal and horocyclic principal cluster sets of a certain holomorphic function, Yokohama Math. J., **16** (1968), 11–14.
- [3] F. Bagemihl and W. Seidel, Some boundary properties of analytic functions, Math. Zeitschr., **61** (1954), 186–199.
- [4] E. F. Collingwood and A. J. Lohwater, The theory of cluster sets, Cambridge, 1966.

- [5] J. T. Gresser, The local behavior of principal and chordal principal cluster sets, *Trans. Amer. Math. Soc.*, **165** (1972), 323–332.
- [6] K. Meier, Über die Randwerte der meromorphen Funktionen, *Math. Ann.*, **142** (1961), 328–344.
- [7] K. Noshiro, *Cluster sets*, Berlin, 1960.

Bowling Green State University