INSTRUCTIONS FOR CONTRIBUTORS

SUBMISSION OF MANUSCRIPTS

Manuscripts should be submitted online via our manuscript submission and tracking site, http://www.editorialmanager.com/psm/. Full instructions for electronic submission are available directly from this site. To facilitate rapid reviewing, communications for peer review will be electronic and authors will need to supply a current e-mail address when registering to use the system.

Papers for publication from Europe (except those on genetic topics, irrespective of country), and all papers on imaging topics, should be submitted to the UK Office.

Papers from the Americas, Asia, Africa, Australasia and the Middle East (except those dealing with imaging topics), and all papers dealing with genetic topics, irrespective of country, should be sent to US Office.

Generally papers should not have text more than 4500 words in length (excluding these sections) and should not have more than a combined total of 5 tables and/or figures. Papers shorter than these limits are encouraged. For papers of unusual importance the editors may waive these requirements. A structured abstract of no more than 250 words should be given at the beginning of the article using the headings: Background; Methods; Results; Conclusions. The name of an author to whom correspondence should be sent must be indicated and a full postal address given in the footnote. Any acknowledgements should be placed at the end of the text (before the References section)

Declaration of Interest: A statement must be provided in the acknowledgements listing all financial support received for the work and, for all authors, any financial involvement (including employment, fees, share ownership) or affiliation with any organization whose financial interests may be affected by material in the manuscript, or which might potentially bias it. This applies to all papers including editorials and letters to the editor.

Contributors should also note the following:

- 1. S.I. units should be used throughout in text, figures and tables.
- 2. Authors should spell out in full any abbreviations used in their manuscripts.
- 3. Foreign quotations and phrases should be followed by a translation.
- 4. If necessary, guidelines for statistical presentation may be found in: **Altman DG, Gore SM, Gardner MJ & Pocock SJ** (1983). Statistical guidelines for contributors to medical journals. *British Medical Journal* 286, 1489–1493.

REFERENCES (1) The Harvard (author-date) system should be used in the text and a complete list of References cited given at the end of the article. In a text citation of a work by more than two authors cite the first author's name followed by et al. (but the names of all of the authors should be given in the References section). Where several references are cited together they should be listed in rising date order. (2) The References section should be supplied in alphabetical order (authors' names in **bold**, journal titles in full), following the text. Some examples follow:

Miller PM, Byrne M, Hodges A, Lawrie SM, Johnstone EC (2002). Childhood behaviour, psychotic symptoms and psychosis onset in young people at high risk of schizophrenia: early findings from the Edinburgh high risk study. *Psychological Medicine* 32, 173–179.

Cleckley HJ (1941). The Mask of Sanity, 2nd edn. Mosby: St. Louis, MO.

Brewer WJ, Wood SJ, DeLuca C, Pantelis C (2006). Models of olfaction for exploring neurodevelopment. In *Olfaction and the Brain* (ed. W. J. Brewer, D. Castle and C. Pantelis), pp. 97–121. Cambridge University Press: Cambridge.

(3) Online citations

doi (when published online prior to printed issue)

Lauritsen MB, Pedersen CB, Mortensen CB (2004). The incidence and prevalence of pervasive developmental disorders: a Danish population-based study. *Psychological Medicine*. Published online: 21 October 2004. doi:10.1017/S0033291704002387.

URL

World Bank (2003). Quantitative techniques for health equity analysis – Technical Notes (http://siteresources.worldbank. org/INTPAH/Resources/Publications/Quantitative-Techniques/health.eq tn07.pdf). Accessed 15 February 2006.

[Authors are requested to print-out and keep a copy of any online-only material, in case the URL changes or is no longer maintained.]

FIGURES AND TABLES Only essential figures and tables should be included. Further tables, figures, photographs and appendices, may be included with the online version on the journal website. To ensure that your figures are reproduced to the highest possible standards, Cambridge Journals recommends the following formats and resolutions for supplying electronic figures. Please ensure that your figures are saved at final publication size and are in our recommended file formats. Following these guidelines will result in high quality images being reproduced in both the print and the online versions of the journal. Line artwork: Format: tif or eps, Colour mode: black and white (also known as 1-bit), Resolution: 1200 dpi; Combination artwork (line/tone): Format: tif or eps, Colour mode: grayscale (also known as 8-bit), Resolution: 800 dpi; Black and white halftone artwork: Format: tif, Colour mode: grayscale (also known as 8-bit), Resolution: 300 dpi; Colour halftone artwork: Format: tif, Colour mode: CMYK colour, Resolution: 300 dpi. All photographs, graphs, and diagrams should be referred to as figures and should be numbered consecutively in Arabic numerals. Captions for figures should be typed double-spaced on separate sheets. Tables Tables should be typed above the table.

PROOFS AND OFFPRINTS Page proofs will be sent to the author designated to receive correspondence. corrections other than to printer's errors may be charged to the author. The corresponding author of each paper will receive a PDF file of their article and hard copy offprints may be purchased if they are ordered on the form supplied when the proof is returned.

PSYCHOLOGICAL MEDICINE

CONTENTS

REVIEW ARTICLES Depressive psychopathology in first-episode schizophrenia spectrum disorders: a systematic review, meta-analysis and meta-regression		Are children with unrecognised psychiatric disorders being excluded from school? A secondary analysis of the British Child and Adolescent Mental Health Surveys 2004 and 2007 Parker C, Tejerina-Arreal M, Henley W, Goodman R, Logan S &	0504
Herniman SE, Allott K, Phillips LJ, Wood SJ, Uren J, Mallawaarachchi SR & Cotton SM	2463	Ford T Change in IQ in schizophrenia patients and their siblings:	2561
Impairments of large-scale functional networks in attention-deficit/hyperactivity disorder: a meta-analysis of resting-state functional connectivity		a controlled longitudinal study Van Haren NEM, Van Dam DS, Stellato RK & Genetic Risk and Outcome of Psychosis (GROUP) investigators	2573
Gao Y, Shuai D, Bu X, Hu X, Tang S, Zhang L, Li H, Hu X, Lu L, Gong Q & Huang X	2475	Continuity of genetic and environmental influences on clinically assessed major depression from ages 18 to 45	
The relationship between depressive symptoms and negative symptoms in people with non-affective psychosis: a meta-analysis		Torvik FA, Gustavson K, Ystrom E, Rosenström TH, Gillespie N, Reichborn-Kjennerud T & Kendler KS	2582
Edwards CJ, Garety P & Hardy A	2486	Negative self-evaluation and the genesis of internal threat: beyond a continuum of suicidal thought and behaviour	
ORIGINAL ARTICLES		Butter S, Shevlin M & Murphy J	2591
Polygenic risk for schizophrenia and season of birth within the UK Biobank cohort		Assessing cross-national invariance of the Community Assessment of Psychic Experiences (CAPE)	
Escott-Price V, Smith DJ, Kendall K, Ward J, Kirov G, Owen MJ, Walters J & O'Donovan MC	2499	Pignon B, Peyre H, Ferchiou A, van Os J, Rutten BPF, Murray RM, Morgan C, Leboyer M, Schürhoff F, Szöke A &	
Weight change increases the odds of psychological distress in middle age: bidirectional analyses from the Whitehall II Study		EU-GEI WP2 Group Author Effect of selective serotonin reuptake inhibitor on prefrontal-striatal connectivity is dependent on the level of	2600
Knüppel A, Shipley MJ, Llewellyn CH & Brunner EJ	2505	TNF- α in patients with major depressive disorder	
No sex differences in the origins of covariation between social and physical aggression		Liu K, Zhao X, Lu X, Zhu X, Chen H, Wang M, Yan W, Jing L, Deng Y, Yu L, Wu H, Wen G, Sun X & Lv Z	2608
Slawinski BL, Klump KL & Alexandra Burt S	2515	Visual attention in adults with attention-deficit/hyperactivity	
Mediators of increased self-harm and suicidal ideation in sexual minority youth: a longitudinal study		disorder before and after stimulant treatment Low A-M, Vangkilde S, le Sommer J, Fagerlund B, Glenthøj B,	
Oginni OA, Robinson EJ, Jones A, Rahman Q & Rimes KA	2524	Jepsen JRM, Bundesen C, Petersen A & Habekost T	2617
Frontostriatal functional connectivity and striatal dopamine synthesis capacity in schizophrenia in terms of antipsychotic responsiveness: an [18F]DOPA PET and fMRI study		Cognitive change in cognitive-behavioural therapy v. pharmacotherapy for adult depression: a longitudinal mediation analysis	
Kim S, Jung WH, Howes OD, Veronese M, Turkheimer FE, Lee Y-S, Lee JS, Kim E & Kwon JS	2533	Quigley L, Dozois DJA, Bagby RM, Lobo DSS, Ravindran L & Quilty LC	2626
Risk of non-affective psychotic disorder or bipolar disorder in autism spectrum disorder: a longitudinal register-based study in the Netherlands		CORRESPONDENCE Design choices when estimating the treated incidence of schizophrenia	
Schalbroeck R, Termorshuizen F, Visser E, van Amelsvoort T & Selten JP	2543	Hogerzeil SJ & van Hemert AM	2635
		INVITED LETTER REJOINDER	
Posttraumatic stress disorder and incidence of thyroid dysfunction in women		Framing the incidence of psychotic disorders: the case for context	
Jung SJ, Kang JH, Roberts AL, Nishimi K, Chen Q, Sumner JA, Kubzansky L & Koenen KC	2551	Edwards J, Rodrigues R & Anderson KK	2637



