

# ON RANDOM INTERPOLATION

P. ERDÖS

(received 23 March 1959)

In a recent paper Salem and Zygmund [1] proved the following result:  
Put

$$\alpha_\nu = \alpha_\nu^{(n)} = \frac{2\pi\nu}{2n+1} \quad (\nu = 0, 1, \dots, 2n)$$

and denote the  $\varphi_\nu(t)$  the  $\nu$ -th Rademacher function. Denote by  $L_n(t, \theta)$  the unique trigonometric polynomial (in  $\theta$ ) of degree not exceeding  $n$  for which

$$L_n(t, \alpha_\nu) = \varphi_\nu(t) \quad (\nu = 0, 1, \dots, 2n).$$

Denote  $M_n(t) = \max_{0 \leq \theta < 2\pi} |L_n(t, \theta)|$ . Then for almost all  $t$

$$\overline{\lim}_{n \rightarrow \infty} \frac{M_n(t)}{(\log n)^{\frac{1}{2}}} \leq 2.$$

I am going to prove the following sharper

**THEOREM 1.** For almost all  $t$

$$\lim_{n \rightarrow \infty} \frac{M_n(t)}{\log \log n} = \overline{\lim}_{n \rightarrow \infty} \frac{M_n(t)}{\log \log n} = \frac{2}{\pi}.$$

Instead of Theorem 1 we shall prove the following stronger (throughout this paper  $c_1, c_2, \dots$  will denote suitable positive constants)

**THEOREM 2.** To every  $c_1$  there exists a constant  $c_2 = c_2(c_1)$  so that for  $n > n_0(c_1, c_2)$  the measure of the set in  $t$  for which

$$\frac{2}{\pi} \log \log n - c_2 < M_n(t) < \frac{2}{\pi} \log \log n + c_2$$

is not satisfied, is less than  $1/n^{c_1}$ .

Theorem 1 follows immediately from Theorem 2 by the Borel-Cantelli Lemma. Thus we only have to prove Theorem 2.

First we need two simple combinatorial lemmas. Let  $m$  be a sufficiently large integer, we define for  $1 \leq i < m$  (for the purpose of these lemmas)

$$\varphi_{m+i}(t) = \varphi_i(t), \quad \varphi_{-i}(t) = \varphi_{m-i}(t).$$

LEMMA 1. Let  $m > m_0(c_1)$ . Then neglecting a set in  $t$  of measure less than  $1/2m^{c_1}$  there exists for every  $t$  a  $k$ ,  $0 \leq k \leq m$  satisfying

$$(1) \quad \varphi_{k+i}(t) = \varphi_{k-1-i}(t) = (-1)^i \text{ for all } 0 \leq i < \frac{1}{2} \log m.$$

The measure of the set in  $t$  for which  $k = [r \log m]$  satisfies (1) is clearly equal to

$$2^{-2\left[\left(\frac{\log m}{2}\right)+1\right]} < 2^{-\log m}.$$

But there are clearly  $[m/\log m] + 1$  possible choices of  $r$  (i.e.  $r$  can take all the values  $0 \leq r < m/\log m$ ). Thus by an obvious independence argument the measure of the set in  $t$  for which none of the possible choices of  $r$  satisfies (1) is less than

$$(1 - 2^{-\log m})^{m/\log m} < \frac{1}{2} m^{-c_1}$$

for every  $c_1$  if  $m > m_0(c_1)$ , which proves Lemma 1.

LEMMA 2. To every  $c_1$  there exists a  $c_3$  so that for  $m > m_0(c_1, c_3)$  neglecting a set (in  $t$ ) of measure less than  $\frac{1}{2}m - c_1$  we have for every  $t, r, (0 \leq r \leq m)$  and  $\nu, (-m/2 < \nu < m/2)$

$$(2) \quad s_{\nu,k}(t) = \left| \sum_{i=0}^{\nu} (-1)^i \varphi_{k+i}(t) \right| < c_3 \nu^{\frac{1}{2}} (\log m)^{\frac{1}{2}}.$$

It is well known that the measure of the set in  $t$  for which

$$\left| \sum_{i=0}^{\nu} (-1)^i \varphi_{k+i}(t) \right| \geq c_3 \nu^{\frac{1}{2}} (\log m)^{\frac{1}{2}}$$

holds is less than

$$(3) \quad c_4 c^{-\frac{1}{2}c_3^2 \log m} < \frac{1}{2} m^{-c_1-2}$$

for sufficiently large  $c_3$ . In (3) there are fewer than  $m^2$  possible choices for  $r$  and  $\nu$ , thus Lemma 2 clearly follows from (3).

Now we are ready to prove our Theorem. (Define for  $0 < \nu \leq n$   $\alpha_{-\nu} = \alpha_{2n-\nu}, \alpha_{2n+\nu} = \alpha_{\nu}$ ). It is well known that

$$(4) \quad L_n(t, \theta) = \frac{1}{2n+1} \sum_{\nu=0}^{2n} \varphi_{\nu}(t) D_n(\theta - \alpha_{\nu})$$

where  $D_n(\theta) = \sin(n + \frac{1}{2})\theta / \sin \frac{1}{2}\theta$  is the Dirichlet kernel. Let  $\alpha_k \leq \theta < \alpha_{k+1}$ . We have

$$(5) \quad L_n(t, \theta) = \frac{1}{2n+1} \left( \sum_{\nu=0}^n \varphi_{\nu}(t) D_n(\theta - \alpha_{k+\nu}) + \sum_{\nu=1}^n \varphi_{\nu}(t) D_n(\theta - \alpha_{k-\nu}) \right) = \Sigma_1 + \Sigma_2.$$

Now we consider only the  $t$  which satisfy Lemmas 1 and 2, (put  $m = 2n$ ), by our Lemmas we thus neglect a set in  $t$  of measure less than  $n^{-c_1}$ . Put

$$(6) \quad \Sigma_1 = \Sigma'_1 + \Sigma''_1$$

where in  $\Sigma'_1$   $0 \leq \nu \leq [\log n]$  and in  $\Sigma''_1$   $[\log n] \leq \nu \leq n$ . We evidently have by  $|D_n(\theta)| \leq 2n + 1$  and a simple computation

$$(7) \quad \begin{aligned} \Sigma'_1 &\leq \frac{1}{2n + 1} \sum_{0 \leq \nu \leq [\log n]} |D_n(\theta - \alpha_{k+\nu})| \\ &\leq 1 + \frac{1}{2n + 1} \sum_{1 \leq r \leq \log n} \frac{1}{\sin \frac{r\pi}{2n + 1}} < \frac{1}{\pi} \log \log n + c_4. \end{aligned}$$

Further by partial summation and Lemma 2

$$(8) \quad \begin{aligned} \Sigma''_1 &= \frac{1}{2n + 1} \sum_{[\log n] < \nu \leq n} (s_{\nu,k}(t) - s_{\nu-1,k}(t)) (-1)^\nu D_n(\theta - \alpha_{k+\nu}) \\ &= \frac{1}{2n + 1} \sum_{\nu=[\log n]+1}^{n-1} s_{\nu,k}(t) ((-1)^\nu D_n(\theta - \alpha_{k+\nu}) - (-1)^{\nu+1} D_n(\theta - \alpha_{k+\nu+1})) \\ &\quad - \frac{1}{2n + 1} s_{[\log n],k}(t) (-1)^{[\log n]+1} D_n(\theta - \alpha_{k+[\log n]+1}) \\ &\quad + \frac{1}{2n + 1} s_{n,k}(t) (-1)^n D_n(\theta - \alpha_{k+n}) \\ &\leq \frac{1}{2n + 1} c_5 (\log n)^{\frac{1}{2}} \sum_{\nu > \log n} \nu^{-\frac{1}{2}} + c_6 < c_7 \end{aligned}$$

since a simple computation shows that for  $\alpha_k \leq \theta < \alpha_{k+1}$

$$|D_n(\theta - \alpha_{k+\nu})| < c_8 \frac{n}{\nu}$$

and

$$|D_n(\theta - \alpha_{k+\nu}) + D_n(\theta - \alpha_{k+\nu+1})| < c_8 \frac{n}{\nu^2}.$$

(6), (7) and (8) implies

$$(9) \quad \Sigma_1 < \frac{\log \log n}{\pi} + c_4 + c_7.$$

Similarly we can show

$$(10) \quad \Sigma_2 < \frac{\log \log n}{\pi} + c_9$$

(9), (10) and (5) implies that for our  $t$  (i.e. for all  $t$  neglecting a set in  $t$  of measure  $< n^{-c_1}$ ).

$$(11) \quad |L_n(t, \theta)| < \frac{2}{\pi} \log \log n + c_{10}.$$

Let now  $k$  satisfy Lemma 1 and put  $\theta_0 = \pi(2k + 1)/(2n + 1)$ . Then we have by (4) and the definition of  $k$

$$\begin{aligned}
 L_n(t, \theta_0) &= \frac{1}{2n+1} \sum_{\nu=0}^n \varphi_\nu(t) D_n(\theta_0 - \alpha_\nu) \\
 (12) \quad &= \frac{1}{2n+1} \sum_{|\nu-k| < \frac{1}{2} \log n} |D_n(\theta_0 - \alpha_\nu)| + \frac{1}{2n+1} \sum_{|\nu-k| \geq \frac{1}{2} \log n} \varphi_\nu(t) D_n(\theta_0 - \alpha_\nu) \\
 &= \Sigma_1 + \Sigma_2
 \end{aligned}$$

Further clearly

$$\begin{aligned}
 (13) \quad \Sigma_1 &= \frac{1}{2n+1} \sum_{|r| < \frac{1}{2} \log n} \frac{2}{\sin \frac{(2r+1)\pi}{2(2n+1)}} \\
 &= \frac{1}{2n+1} \sum_{|r| < \frac{1}{2} \log n} \frac{2}{\frac{(2r+1)\pi}{2(2n+1)} + o\left(\frac{r^3}{n^3}\right)} > \frac{2}{\pi} \log \log n - c_{11}.
 \end{aligned}$$

As in (8) we can show that

$$(14) \quad |\Sigma_2| < c_{12}.$$

(12), (13) and (14) implies

$$(15) \quad |L_n(t, \theta_0)| > \frac{2}{\pi} \log \log n - c_{11} - c_{12}.$$

(11) and (15) complete the proof of Theorem 2.

By more complicated arguments we could prove the following sharper,

**THEOREM 3.** There exists an absolute constant  $C$  so that neglecting a set in  $t$  whose measure tends to 0 as  $n$  tends to infinity we have

$$M_n(\theta) = \frac{2}{\pi} \log \log n + c + o(1).$$

(The exceptional set whose measure goes to 0 depends on  $n$ ).

Using the methods of another paper by Salem and Zygmund [2] we can prove the following

**THEOREM 4.** There exists a distribution function  $(\psi(\alpha))$  (i.e.  $\psi(\alpha)$ ,  $-\infty < \alpha < \infty$  is non decreasing,  $\psi(-\infty) = 0$ ,  $\psi(+\infty) = 1$ ), so that, neglecting a set in  $t$  whose measure tends to 0 as  $n$  tends to infinity, we have

$$m(\theta: L_n(t, \theta) < \alpha) \rightarrow \psi(\alpha).$$

In other words: If we neglect a set in  $t$  of measure tending to 0 (the exceptional set may depend on  $n$ ) we have for a  $t$  not belonging to this exceptional set the following situation: The measure of the set in  $\theta$  for which  $L_n(t, \theta) < \alpha$  holds, equals  $\psi(\alpha) + o(1)$ .

We do not discuss in this paper the proofs of Theorem 3 and 4.

**References**

- [1] Salem and Zygmund, Third Berkeley Symposium for probability and statistics, Vol. 2, 243–246.
- [2] Salem and Zygmund, Acta Math. 91 (1954).

Technion, Haifa.