

hallucination, which in the case of our patient was eradicated, so the contribution of this case could enrich the current bibliography.

Conclusions: This is unfrequently presentation of Charles Bonnet syndrom.

Disclosure of Interest: None Declared

EPV1540

Complexities of the Prodromal Phase of First-Episode Psychosis: A Longitudinal and Phenomenological Diagnostic Approach

A. Guiomar^{1*}, C. P. Osório¹, P. M. Martins¹, P. Macedo¹ and M. de Barros¹

¹Psychiatry, Unidade Local de Saúde do Tâmega e Sousa, Penafiel, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2025.2049

Introduction: A 26-year-old man presented with his first-episode psychosis (FEP) following a 15-month period marked by a progressive sense of estrangement from his surroundings, ideas of reference, persistent anxiety, difficulty focusing, and social withdrawal. Two years prior, he began stimulant treatment for suspected attention-deficit/hyperactivity disorder (ADHD), though he discontinued the medication shortly after, as he perceived no improvement. Over the past year, he became increasingly distant from friends and eventually resigned from his job. About three months before hospitalization, he began experiencing first-rank symptoms of schizophrenia. This case will serve as a starting point to discuss the complexities of diagnosing the prodromal phase of FEP.

Objectives: This clinical review aims to examine the phenomenology of the prodromal phase of FEP and address the diagnostic challenges posed by symptom similarities between this phase and neurodevelopmental conditions like ADHD.

Methods: A literature review was conducted using the PubMed database, covering studies from the past 20 years. Studies were selected if they included phenomenological descriptions of the prodromal phase in FEP and/ or examined the impact of neurodevelopmental conditions on the emergence of psychosis.

Results: The review identified several key phenomenological markers characterizing the prodromal phase of FEP, which can aid in distinguishing it from other psychiatric conditions. The prodromal phase of FEP is frequently marked by subtle but progressive alterations in cognition, perception, and affect, including experiences such as derealization-depersonalization, ideas of reference, paranoid ideation, and social withdrawal. Evidence suggests that prodromal symptoms intensify over time, evolving from vague unease to specific disruptions in reality testing. Although ADHD and the prodromal phase of a FEP may share some overlapping characteristics - particularly when symptoms are assessed in a cross-sectional manner - ADHD symptoms are generally regarded as stable traits that persist consistently into adulthood.

Conclusions: This case underscores the need for careful differential diagnosis, especially when evaluating individuals in high-risk age groups for psychosis who present with subtle symptoms that do not clearly fit a single diagnostic category. In such cases, clinicians should avoid premature conclusions and instead adopt a longitudinal and comprehensive approach, considering whether genetic, neurodevelopmental, or social risk factors may be contributing to the presentation. A phenomenological perspective can help

clinicians detect subtle yet significant shifts in perception, cognition, and affect, enhancing diagnostic accuracy and enabling timely intervention.

Disclosure of Interest: None Declared

EPV1541

A literature review of first-episode psychosis, a perspective on the future

M. C. Herrero Rodriguez^{1*}, Z. Martin García¹, C. Vall García¹ and J. M. Martínez Sánchez¹

¹Psychiatry, Río Hortega University Hospital, Valladolid, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2025.2050

Introduction: First-episode psychosis is one of the major challenges of mental health research worldwide because it is a traumatic experience for patients and their families. Patients who experience these episodes may experience fear, distress, and isolation.

Objectives: The early phase of psychosis is a critical period when long-term outcome is predictable and biological, psychological and psychosocial influences are developing and display maximal plasticity. This phase presents important opportunities for secondary prevention and delaying treatment may affect the chance of recovery. The main goal is to reduce the duration of untreated psychosis and ensure that, in addition to symptom remission, there is also psychosocial recovery. Currently, the clinical and research focus in psychotic disorders has shifted toward first episode psychosis, early detection of the prodromal phase of psychosis, and an effective integrated treatment model known as "Early Intervention."

Methods: Selective review of the literature on first episode psychosis.

Results: The studies strongly support the efficacy of antipsychotic medication as both acute and maintenance treatment for patients with a first episode of psychosis.

Conclusions: Early intervention may improve outcomes in first episode psychosis. The use of new antipsychotics with greater efficacy and fewer side effects may improve medication adherence and reduce morbidity associated with repeated relapses. However, the optimal duration of maintenance treatment has not been determined and a long duration of untreated psychosis may be associated with a poorer treatment response. Finally, services for Early Intervention should be easily accessible, non-threatening and non-stigmatising.

Disclosure of Interest: None Declared

EPV1542

Does negative voice content enhance strong priors and conditioned hallucinations?

C. Jacquet^{1*}, C. Dondé^{2,3} and C. Bortolon^{1,4,5}

¹38000, Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, LIP/PC2S;

²38000, Centre Hospitalier Universitaire Grenoble Alpes, GIN, Grenoble;

³38120, Centre Hospitalier Alpes Isère, Saint-Egrève; ⁴Institut

Universitaire de France, Paris and ⁵38000, C3R – Réhabilitation psychosociale et remédiation cognitive, Centre Hospitalier Alpes Isère, Grenoble, France