

On the profinite rigidity of lattices in higher rank Lie groups

BY HOLGER KAMMEYER

*Mathematical Institute, Building 25.22, Heinrich Heine University Düsseldorf,
Universitätsstraße 1, 40225 Düsseldorf, Germany.*

e-mail: holger.kammeyer@hhu.de

AND STEFFEN KIONKE

*FernUniversität in Hagen, Fakultät für Mathematik und Informatik, Universitätsstraße
11, 58097 Hagen, Germany.*

e-mail: steffen.kionke@fernuni-hagen.de

(Received 13 April 2021; revised 27 July 2022; accepted 21 July 2022)

Abstract

We investigate which higher rank simple Lie groups admit profinitely but not abstractly commensurable lattices. We show that no such examples exist for the complex forms of type E_8 , F_4 , and G_2 . In contrast, there are arbitrarily many such examples in all other higher rank Lie groups, except possibly $\mathrm{SL}_{2n+1}(\mathbb{R})$, $\mathrm{SL}_{2n+1}(\mathbb{C})$, $\mathrm{SL}_n(\mathbb{H})$, or groups of type E_6 .

2020 Mathematics Subject Classification: 22E40, 20E18 (Primary)

1. Introduction

If two residually finite groups are commensurable, so are the profinite completions. Thus for any class \mathcal{C} of residually finite groups, the converse *rigidity* question arises: if two groups from \mathcal{C} are profinitely commensurable, are they abstractly commensurable? It is known that arithmetic groups with the congruence subgroup property can be used to construct lattices in higher rank Lie groups which are profinitely isomorphic but not abstractly isomorphic; see [1, 2, 11, 17]. Aka showed [2] that the profinite isomorphism class of a higher rank lattice contains only finitely many abstract commensurability types of lattices. Here we address the question whether every simple Lie group G of higher rank actually admits profinitely isomorphic but non-commensurable lattices. The flexibility of the familiar construction might be taken as evidence, that the answer should be affirmative. However, it turns out that the question is more delicate, and the answer depends on G .

THEOREM 1.1. *Let G be a connected simple higher rank Lie group with finite center which is:*

- (i) neither a complex form of type E_8 , F_4 , or G_2 ;
- (ii) nor a real or complex form of type E_6 ;
- (iii) nor locally isomorphic to $\mathrm{SL}_{2m+1}(\mathbb{R})$, $\mathrm{SL}_{2m+1}(\mathbb{C})$, or $\mathrm{SL}_m(\mathbb{H})$.

Then for each $n \geq 2$, there exist cocompact lattices $\Gamma_1, \dots, \Gamma_n \subset G$ which are pairwise profinitely isomorphic but pairwise not commensurable.

So in most cases rigidity fails in a strong sense but the first three groups form a surprising exception.

THEOREM 1.2. *Let G be a connected simple complex Lie group of type E_8 , F_4 , or G_2 and let $\Gamma_1, \Gamma_2 \subset G$ be lattices such that Γ_1 is profinitely commensurable with Γ_2 . Then Γ_1 is commensurable with Γ_2 .*

Going through the census, the complete list of higher rank simple Lie groups currently not covered by either Theorem 1.1 or Theorem 1.2 is given by $\mathrm{SL}_{2m-1}(\mathbb{R})$ and $\mathrm{SL}_{2m-1}(\mathbb{C})$, and $\mathrm{SL}_m(\mathbb{H})$ for $m \geq 2$, as well as all the E_6 -forms: $E_{6(6)}$, $E_{6(2)}$, $E_{6(-14)}$, $E_{6(-26)}$, and $E_6(\mathbb{C})$. If the congruence subgroup property was known to hold true for higher rank groups of type A_n and E_6 , all these groups would likewise fall under the conclusion of Theorem 1.1. For instance, the non-isomorphic but locally isomorphic algebraic groups of types A_n , D_{2n+1} and E_6 in [25, theorem 9.12] would give rise to examples. Let us stress however that our result is unconditional. Though the congruence subgroup problem is generally open in type A_n and D_4 , we were able to exploit the partial progress made in the literature to the extent that the type A_n groups $\mathrm{SL}_{2m}(\mathbb{R})$ and $\mathrm{SL}_{2m}(\mathbb{C})$, as well as $\mathrm{SU}(p, q)$ for $p, q \geq 2$, and all the type D_4 groups $\mathrm{SO}^0(6, 2)$, $\mathrm{SO}^0(5, 3)$, $\mathrm{SO}^0(4, 4)$, and $\mathrm{SO}^*(8)$ are included in Theorem 1.1.

The lattices constructed here are intrinsically cocompact, since our approach is based on arithmetic groups defined over number fields of large degree. The construction of profinitely isomorphic *non-cocompact* lattices requires different methods; e.g., similar to the construction in [25, theorem 9.12]. In Proposition 3.1 we illustrate this by showing that $\mathrm{SL}_m(\mathbb{R})$, $\mathrm{SL}_m(\mathbb{C})$ and $\mathrm{SL}_m(\mathbb{H})$, where $m \geq 6$ is composite, admit profinitely isomorphic, non-commensurable, non-cocompact lattices.

For the various types of G , we employ varying methods to construct the families $\Gamma_1, \dots, \Gamma_n \subset G$ of profinitely isomorphic but non-commensurable lattices in Theorem 1.1. There does not seem to be a uniform approach that would work in all cases. This is however different for the question that is more commonly addressed under the term *profinite rigidity* in the literature: if two groups have isomorphic profinite completions, are they isomorphic? Here the congruence subgroup property, whenever it is known to hold for G , can be used to construct profinitely isomorphic but non-isomorphic congruence subgroups in a uniform way.

THEOREM 1.3. *Let G be a higher rank connected simple Lie group with trivial center and not isomorphic to $\mathrm{PSL}_m(\mathbb{H})$ or to a complex or real form of type E_6 . Then there exist arbitrarily many cocompact lattices in G which are pairwise profinitely isomorphic but pairwise not isomorphic.*

Again, the exceptions can likely be omitted. They only owe to the incomplete status of the congruence subgroup problem. We see that profinite rigidity in the usual sense also fails for $E_8(\mathbb{C})$, $F_4(\mathbb{C})$, and $G_2(\mathbb{C})$, and most likely fails for lattices in all higher rank simple

Lie groups. However, the lattices we construct in the proof of Theorem 1.3 are arithmetic subgroups of different congruence levels. While they are not isomorphic, they are commensurable which is why they should not really be considered as distinct lattices. This is why we find it more on point to ask for profinitely commensurable lattices in G of higher rank which are not commensurable.

Let us remark that profinite rigidity questions are typically only posed for residually finite groups to avoid trivial examples like $\widehat{\Gamma \times \Lambda} \cong \widehat{\Gamma} \times \widehat{\Lambda} \cong \widehat{\Gamma}$ for any group Γ and any infinite simple group Λ . Most simple Lie groups with finite center are linear and hence lattices are residually finite. However, there are simple Lie groups with finite center which admit lattices that are not residually finite. An example is given by the four-fold covering of $\mathrm{Sp}_n(\mathbb{R})$ [7]. In these cases it is nevertheless reasonable to consider the profinite completion because the kernel of the completion homomorphism $\Gamma \rightarrow \widehat{\Gamma}$ is merely a finite central subgroup.

Profinite rigidity of groups and related problems receive considerable attention in current research activities; see [26] for a survey. Fundamental groups of 3-manifolds, see [5] and references therein, and lattices in Lie groups [12, 30] are the main objects of interest.

We briefly sketch the proofs of Theorems 1.1 and 1.2. As we just commented, the proof of Theorem 1.1 splits up into various cases. At this point, we shall only present the most common line of arguments that works for most real forms G . We may assume $G = \mathbf{G}(\mathbb{R})$ for an absolutely simple, simply connected algebraic \mathbb{R} -group \mathbf{G} . We construct linear algebraic groups $\mathbf{G}_1, \dots, \mathbf{G}_n$ over a suitable totally real number field k such that \mathbf{G}_i is isomorphic to \mathbf{G} at exactly one real place of k and is compact at all other real places. The lattices arise as arithmetic subgroups $\Gamma_i \subseteq \mathbf{G}_i(k)$, ensuring that the congruence subgroup property holds for the groups \mathbf{G}_i (special attention is needed in type A_m and D_4). The core of the argument is a local-global principle which *almost* allows us to achieve that the groups $\mathbf{G}_1, \dots, \mathbf{G}_n$ are isomorphic at all finite places of k . If the groups \mathbf{G}_i were isomorphic at all finite places of k , then the congruence subgroup property implies that the groups are profinitely commensurable (by Theorem 2.5 and Lemma 2.6). If the field k has no non-trivial automorphisms, then Margulis superrigidity can be used to deduce that the arithmetic lattices are not commensurable (Theorem 2.7). There is however a caveat: The local-global principle only allows to control the isomorphism type except for one finite place. But since there exists only a finite number of possible p -adic types for the groups \mathbf{G}_i , we can infer from Dirichlet’s box principle that for arbitrary large n , arbitrarily many of the groups \mathbf{G}_i are isomorphic at every finite place.

To prove Theorem 1.2, we apply Margulis arithmeticity to conclude that for $i = 1, 2$, Γ_i is commensurable with an arithmetic subgroup in a k_i -group \mathbf{G}_i over some number field k_i with precisely one complex place such that \mathbf{G}_i is anisotropic at all real places of k_i . The congruence subgroup property, which is known in the exceptional types under consideration, effects that $\widehat{\Gamma}_i$ is commensurable with the finite adèle points $\mathbf{G}_i(\mathbb{A}_{k_i}^f)$ of \mathbf{G}_i . Hence $\mathbf{G}_1(\mathbb{A}_{k_1}^f)$ is commensurable with $\mathbf{G}_2(\mathbb{A}_{k_2}^f)$ from which we conclude that the number fields k_1 and k_2 are *arithmetically equivalent*, meaning they have equal Dedekind zeta function. There exists a myriad of arithmetically equivalent number fields, also among totally real ones, which are not isomorphic. However, a theorem due to Chinburg–Hamilton–Long–Reid says that arithmetically equivalent number fields with precisely one complex place are isomorphic [6, corollary 1.4]. We conclude that \mathbf{G}_2 is a k_1 -twist of \mathbf{G}_1 . These are classified by the noncommutative Galois cohomology set $H^1(k_1, \mathbf{G}_1)$ because \mathbf{G}_1 has trivial center and trivial outer automorphism group. The *Hasse principle* for simply connected groups in combination with

M. Kneser's vanishing result for Galois cohomology over p -adic fields therefore implies that \mathbf{G}_1 is actually k_1 -isomorphic to \mathbf{G}_2 , hence Γ_1 is commensurable with Γ_2 .

Structure of the article

We discuss some preliminaries in Section 2. Sections 3, 4 and 5 are dedicated to the proofs of Theorems 1.1, 1.2, and 1.3, respectively.

2. Preliminaries

2.1. Notation

Let k be an algebraic number field. The set of places of k is denoted by $V(k) = V_\infty(k) \cup V_f(k)$; it is the union of the set of archimedean places $V_\infty(k)$ and the set of finite places $V_f(k)$. The completion of k at $v \in V(k)$ is denoted by k_v . The ring of adeles (respectively of finite adeles) of k is \mathbb{A}_k (resp. \mathbb{A}_k^f).

2.2. Number fields without automorphisms

The following result is well-known; e.g., [19, proposition 2.3].

LEMMA 2.1. *There are totally real number fields of arbitrarily large degree over \mathbb{Q} with trivial automorphism group.*

Proof. For (arbitrarily large) $n \geq 3$, let K/\mathbb{Q} be a totally real Galois extension with Galois group $\text{Gal}(K/\mathbb{Q}) \cong S_n$. Let $H \leq \text{Gal}(K/\mathbb{Q})$ be a subgroup isomorphic to S_{n-1} . The fixed field $k = K^H$ is totally real, $[k:\mathbb{Q}] = n$ and k has a trivial automorphism group. Indeed, an automorphism of k extended to K/\mathbb{Q} normalizes H but S_{n-1} is self-normalising in S_n .

Similarly, there is the following result for ‘‘almost’’ totally real fields.

LEMMA 2.2. *There are number fields of arbitrarily large degree over \mathbb{Q} with precisely one complex place and with trivial automorphism group.*

Proof. Fix a prime number $p > 2$ and an irreducible rational polynomial P of degree p with exactly two non-real roots. Let K be the splitting field of P with Galois group $G \subseteq S_p$. Given $a \in K$ with $P(a) = 0$, the subgroup $H \leq G$ corresponding to $k = \mathbb{Q}(a)$ has index p . So the group $G \subseteq S_p$ contains an element of order p which must be a full p -cycle because p is prime. Moreover, the non-real roots of P are complex conjugates of one another, hence complex conjugation exhibits a nontrivial transposition in G . A symmetric group of prime order is generated by any full cycle and any transposition, so $G = S_p$. We observe (as in Lemma 2.1) that the stabilizer H of a is actually self-normalising. Hence $k = \mathbb{Q}(a)$ has trivial automorphism group.

2.3. Finite coverings of Lie groups

The following lemma will be applied to reduce the proof to the case where the Lie group G is given by the \mathbb{R} - or \mathbb{C} -points of a linear algebraic group.

LEMMA 2.3. *Let $f: G_1 \rightarrow G$ and $g: G \rightarrow G_0$ be homomorphisms of Lie groups with finite kernels. Assume that G is linear. If G possesses n cocompact lattices which are pairwise profinitely commensurable but pairwise not commensurable, then the same holds true for G_1 and G_0 .*

Proof. The linearity will only be used to ensure that all lattices in G are residually finite. It suffices to treat the case $n = 2$. Let $\Gamma_1, \Gamma_2 \subseteq G$ be two non-commensurable, profinitely commensurable lattices. Then $\Delta_i = f^{-1}(\Gamma_i)$ is a cocompact lattice in G_1 . Let $K_i \subseteq \Delta_i$ be the kernel of the completion $\Delta_i \rightarrow \widehat{\Delta}_i$. Since Γ_i is residually finite, $K_i \subseteq \ker(f)$. Since $\ker(f)$ is finite, there is a finite index normal subgroup $\Delta'_i \subseteq \Delta_i$ which intersects $\ker(f)$ exactly in K_i and thus the profinite completion of Δ'_i is isomorphic to the profinite completion of $f(\Delta'_i)$ which is a finite index normal subgroup of Γ_i . Therefore Δ'_1, Δ'_2 are profinitely commensurable. However, these groups are not commensurable, since every isomorphism between finite index subgroups maps the completion kernel K_1 to the kernel K_2 , i.e., it induces an isomorphism of finite index subgroups of Γ_1 and Γ_2 .

Since Γ_1, Γ_2 are residually finite and $\ker(g)$ is finite, there are finite index subgroups Γ'_1, Γ'_2 which do not intersect $\ker(g)$ so that they are isomorphic to lattices in G_0 . As finite index subgroups of Γ_1, Γ_2 they are still profinitely commensurable but not commensurable.

2.4. *The congruence subgroup property and its consequences*

A key ingredient in the proof is the *congruence subgroup property* (CSP): the statement that the kernel $C(k, \mathbf{G})$ of the canonical homomorphism $\widehat{\mathbf{G}}(k) \rightarrow \overline{\mathbf{G}}(k)$ from the arithmetic completion to the congruence completion of the k -rational points of certain k -groups \mathbf{G} is finite. We will apply various special cases in which the congruence subgroup property is known to hold true.

THEOREM 2.4. *Let \mathbf{G} be a simply connected absolutely almost simple linear algebraic k -group which either is k -isotropic or has type:*

- (i) B_l ($l \geq 2$);
- (ii) C_l ($l \geq 2$)
- (iii) D_l ($l \geq 5$)
- (iv) E_7, E_8, F_4, G_2 ,

or let \mathbf{G} be the type ${}^2A_{m-1}$ group $\mathbf{G} = \mathbf{SU}_m(K, h)$ where h is a nondegenerate m -dimensional Hermitian form over a quadratic extension K/k with $m \geq 3$. Assume moreover that $\sum_{v \in V_\infty(k)} \text{rank}_{k_v} \mathbf{G} \geq 2$ and that k is not totally imaginary. Then the congruence kernel $C(k, \mathbf{G})$ has order at most two. If \mathbf{G} is not topologically simply connected at some real place of k , then $C(k, \mathbf{G})$ is trivial.

The theorem is the essence of decades of research on the congruence subgroup problem. References are [8; 21, theorem 9.1, p. 512, theorem 9.5, p. 513, corollary 9.7, p. 515, theorems 9.23 and 9.24; 22, main theorem]. A survey article providing extensive information on CSP can be found in [23]. The following result is another main tool for us.

THEOREM 2.5. *Let k be an algebraic number field and let \mathbf{G} be a simply connected simple linear algebraic group over k . Let $\Gamma \subseteq \mathbf{G}(k)$ be an arithmetic subgroup. Assume that $G_\infty = \prod_{v \in V_\infty(k)} \mathbf{G}(k_v)$ is not compact.*

- (a) $\Gamma \subseteq G_\infty$ is a lattice. If $\mathbf{G}(k_v)$ is compact for some $v \in V_\infty(k)$, then Γ is cocompact.
- (b) Assume that \mathbf{G} has CSP. Then the profinite completion $\widehat{\Gamma}$ is commensurable with the open compact subgroups of $\mathbf{G}\left(\mathbb{A}_k^f\right)$.

Proof. Part (a) is a famous result of Borel and Harish-Chandra [3]. To prove part (b), note that by CSP, the kernel of the map $\varphi: \widehat{\Gamma} \rightarrow \mathbf{G}(\mathbb{A}_k^f)$ is finite. Passing to a finite index subgroup of Γ , we can assume that φ is injective. The strong approximation theorem holds since \mathbf{G} is simply connected [21, theorem 7.12] and implies that the image of φ is open (and compact since $\widehat{\Gamma}$ is compact).

Finally, we will need to know that a collection of local isomorphisms of algebraic groups assembles to an adelic isomorphism.

LEMMA 2.6. *Let k be an algebraic number field and let \mathbf{G}, \mathbf{H} be two semi-simple linear algebraic groups over k . If \mathbf{G}, \mathbf{H} are isomorphic at all finite places, i.e., $\mathbf{G} \times_k k_v \cong \mathbf{H} \times_k k_v$ for all $v \in V_f(k)$, then*

$$\mathbf{G}(\mathbb{A}_k^f) \cong \mathbf{H}(\mathbb{A}_k^f)$$

as topological groups.

Proof. By assumption the topological groups $\mathbf{G}(k_v)$ and $\mathbf{H}(k_v)$ are isomorphic at all finite places v of k . We pick models of \mathbf{G}, \mathbf{H} over the ring of integers \mathcal{O}_k of k . Then for all but finitely many places $v \in V_f(k)$, the compact subgroups $\mathbf{G}(\mathcal{O}_{k,v}), \mathbf{H}(\mathcal{O}_{k,v})$ are hyperspecial [32, 3.9.1] and hence isomorphic [32, 2.5]. We deduce that

$$\begin{aligned} \mathbf{G}(\mathbb{A}_k^f) &= \varinjlim_S \prod_{v \in S} \mathbf{G}(k_v) \times \prod_{v \notin S} \mathbf{G}(\mathcal{O}_{k,v}) \\ &\cong \varinjlim_S \prod_{v \in S} \mathbf{H}(k_v) \times \prod_{v \notin S} \mathbf{H}(\mathcal{O}_{k,v}) \cong \mathbf{H}(\mathbb{A}_k^f), \end{aligned}$$

where the direct limit runs over all finite sets S of finite places of k .

2.5. Margulis superrigidity

Margulis superrigidity will be used to show that certain arithmetic lattices are not abstractly commensurable.

THEOREM 2.7 (Margulis). *Let k_1, k_2 be number fields and let $\mathbf{G}_1, \mathbf{G}_2$ be simply connected, absolutely almost simple linear algebraic groups over k_1 and k_2 respectively. Assume that $\sum_{v \in V_\infty(k_j)} \text{rk}_{k_j}(\mathbf{G}_j(k_{j,v})) \geq 2$ for all $j \in \{1, 2\}$. The arithmetic subgroups $\Gamma_1 \subset \mathbf{G}_1(k_1)$ and $\Gamma_2 \subset \mathbf{G}_2(k_2)$ are commensurable if and only if there is an isomorphism of fields $\sigma: k_1 \rightarrow k_2$ and a k_2 -isomorphism of algebraic groups $\eta: {}^\sigma \mathbf{G}_1 \rightarrow \mathbf{G}_2$.*

Proof. After passing to finite index subgroups, we can assume that there exists an isomorphism $\delta: \Gamma_1 \xrightarrow{\cong} \Gamma_2$. Then by Margulis superrigidity [18, theorem (C), p. 259], there exists $\sigma: k_1 \rightarrow k_2$ and an epimorphism $\eta: {}^\sigma \mathbf{G}_1 \rightarrow \mathbf{G}_2$ such that δ agrees with η on a finite index subgroup of Γ_1 . Without loss of generality $\delta = \eta|_{\Gamma_1}$. The same argument applied to δ^{-1} implies that σ is an isomorphism and further (since \mathbf{G}_2 is simply connected) that η is an isomorphism.

Conversely, if σ and η exist, then Γ_1 and Γ_2 are (isomorphic to) arithmetic subgroups of the same algebraic group and are thus commensurable.

3. Proof of Theorem 1.1

In this section, k denotes a number field with $n := [k : \mathbb{Q}] > 2$ and with trivial automorphism group. We will say k has type I if it is totally real; see Lemma 2.1. In this case k_1, \dots, k_n denote the completions of k at the archimedean places. We will say k has type II if it has exactly one complex place; see Lemma 2.2. In this case k_1, \dots, k_{n-1} denote the completions at the archimedean places of k with $k_1 \cong \mathbb{C}$.

3.1. Reduction to $G = \mathbf{G}(\mathbb{R})$ or $G = \mathbf{G}(\mathbb{C})$

As a first step of the proof of Theorem 1.1, we observe that we may assume that the Lie group G is the group of real or complex points of a simply connected simple algebraic group. To this end, let G be a connected simple Lie group with finite center. Let \mathfrak{g} be the Lie algebra of G . Since the center $Z(G)$ of G is finite and the adjoint group $G/Z(G)$, being a subgroup of $\text{Aut}(\mathfrak{g})$, is linear, we may assume by Lemma 2.3 that G has trivial center.

If $\mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}$ is simple, then the linear algebraic \mathbb{R} -group $\mathbf{G}_{\text{ad}} = \text{Aut}_{\mathbb{R}}(\mathfrak{g})$ is absolutely simple and satisfies $\mathbf{G}_{\text{ad}}(\mathbb{R})^0 = G$. Let \mathbf{G} be the simply connected covering of \mathbf{G}_{ad} . By Lemma 2.3 it is sufficient to show for every n that $\mathbf{G}(\mathbb{R})$ has n profinitely isomorphic lattices which are not commensurable.

If $\mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C}$ is not simple, then \mathfrak{g} possesses a complex structure which turns it into a simple Lie algebra over \mathbb{C} . For any such structure, the \mathbb{C} -group $\mathbf{G}_{\text{ad}} = \text{Aut}_{\mathbb{C}}(\mathfrak{g})$ is (absolutely) simple and satisfies $\mathbf{G}_{\text{ad}}(\mathbb{C}) = G$. Again \mathbf{G} will denote the simply connected covering group of \mathbf{G}_{ad} and it is enough to find n non-commensurable profinitely isomorphic lattices in $\mathbf{G}(\mathbb{C})$.

With these remarks, we associated to each local isomorphism class of Lie groups G as in Theorem 1.1 a connected simply connected absolutely almost simple linear algebraic \mathbb{K} -group \mathbf{G} with $\mathbb{K} = \mathbb{R}$ or \mathbb{C} which is unique up to \mathbb{K} -isomorphism. Our task now is to find n profinitely commensurable lattices in $\mathbf{G}(\mathbb{K})$ which are not commensurable. The common intersection of the profinite completions then corresponds to non-commensurable lattices which are profinitely isomorphic, see [27, proposition 3.2.2, p. 80] and, for instance, [10, proposition 6.39, p. 159].

3.2. Overview of the proof

Let us first assume that $\mathbb{K} = \mathbb{R}$, in which case we take k of type I. The general idea of the proof in this case was already outlined in the introduction. At this point we have to point out, however, that the argument only goes through provided the following two requirements are met.

- (i) We need to assume that \mathbf{G} and the \mathbb{R} -anisotropic real form \mathbf{G}^u with $\mathbf{G} \times_{\mathbb{R}} \mathbb{C} \cong_{\mathbb{C}} \mathbf{G}^u \times_{\mathbb{R}} \mathbb{C}$ are inner forms of each other. This is automatic unless the Dynkin diagram has symmetries, meaning \mathbf{G} has type A_m, D_m , or E_6 . In these cases, one can read off from the Tits indices [31, table II], whether the condition is satisfied: In type A_m , the condition fails for $\text{SL}_{m+1}(\mathbb{R})$ and $\text{SL}_{m+1}(\mathbb{H})$, while the groups $\text{SU}(r, s)$ with $r + s = m + 1$ satisfy this requirement. In type D_m , the groups $\text{SO}^*(2m) = \text{SO}(m, \mathbb{H})$ are inner twists of the compact form $\text{SO}(2m)$. For the groups $\text{SO}^0(r, s)$ with $r + s = 2m$, the condition is satisfied if r and s are even and fails if r and s are odd. Finally, in type E_6 , the condition is satisfied for $E_{6(2)}$ and $E_{6(-14)}$ and fails for $E_{6(6)}$ and $E_{6(-26)}$.
- (ii) We need that k -anisotropic forms of \mathbf{G} defined over k satisfy the congruence subgroup property. This is still generally open in type A_m, D_4 and E_6 .

Property (ii) forces us to exclude type E_6 altogether. In type A_m , however, CSP is known for special unitary groups $SU(K/k, h)$ of hermitian forms h over a quadratic field extension K/k as we stated in Theorem 2.4. This allows us to prove Theorem 1.1 for the groups $SU(r, s)$ in Section 3.3. There we also present a workaround that allows us to include the groups $SL_{2m}(\mathbb{R})$ in spite of the failure of property (i).

Similarly, Kneser [14] has shown that CSP holds for spinor groups. Using this, we can construct the required non-commensurable but profinitely isomorphic lattices in all the real forms $SO^0(6, 2)$, $SO^0(5, 3)$, $SO^0(4, 4)$, and $SO^*(8)$ of type D_4 as arithmetic spinor groups. So it is no issue for us that CSP is still open for anisotropic type ${}^{3,6}D_4$ forms (whose unique inner quasi-split twist has a splitting field extension of degree 3 or 6, so called *triality forms*). Note also that property (i) fails in some D_m cases with $m \geq 4$. Therefore, we will sort out the type D_m groups with $m \geq 4$ separately in Section 3.4.

With the special cases taken care of, we treat the remaining \mathbb{R} -groups of type B_m, C_m, E_7, E_8, F_4 , and G_2 in Section 3.5. For all these, the general strategy applies because (i) and (ii) are satisfied.

Finally, we give the proof of Theorem 1.1 for $\mathbb{K} = \mathbb{C}$ and \mathbf{G} of type B_m, C_m, D_m with $m \geq 5$, and E_7 in Section 3.6. Also in the complex case, type A_m and D_4 need special attention because of the incomplete status of CSP. The type D_4 group $SO_8(\mathbb{C})$ can again be covered by Kneser’s result so that it was more convenient to include it in Section 3.4. A similar trick as for $SL_{2m}(\mathbb{R})$ also allows us to cover the type A_{2m-1} group $SL_{2m}(\mathbb{C})$. This argument is included in Section 3.3.

3.3. Type A_\bullet : $SU(r, s)$, $SL_{2m}(\mathbb{R})$ and $SL_{2m}(\mathbb{C})$

As a first instance we consider the case when G is either of the groups $SU(r, s)$ with $r, s \geq 2$, $SL_{2m}(\mathbb{R})$ and $SL_{2m}(\mathbb{C})$ with $m \geq 2$. In these concrete examples it is instructive how local-global principles are key to our investigation.

It is wellknown that hermitian forms for the extension \mathbb{C}/\mathbb{R} are classified by dimension and signature. We also recall that hermitian forms for a quadratic extension E/F of p -adic fields are classified by dimension and discriminant $d \in \{\pm 1\}$ (i.e., $d = 1$ exactly if the determinant lies in the image of the norm $N_{E/F}: E^\times \rightarrow F^\times$); see e.g. [9, theorem 3.1].

Fix a dimension $m \geq 2$. Suppose that $K = k(\sqrt{a})$ is a quadratic extension of k . Let $V_\infty^K(k)$ denote the set of non-split archimedean places of k , i.e., the set of real places v where a is negative with respect to the embedding $k \rightarrow k_v$. We will use the following result of Landherr [16] on the existence of hermitian forms with prescribed local properties: Choosing a pair of nonnegative integers (r_i, s_i) with $r_i + s_i = m$ for each $v_i \in V_\infty^K(k)$ and choosing $d_v \in \{\pm 1\}$ for each $v \in V_f(k)$, there exists a K/k -hermitian form of dimension m with signature (r_i, s_i) at $v_i \in V_\infty^K(k)$ and discriminant $d_v \in \{\pm 1\}$ at $v \in V_f(k)$ if and only if:

- (i) $d_v = 1$ for almost all $v \in V_f(k)$;
- (ii) $d_v = 1$ whenever v splits in K ; and
- (iii) $\prod_{v_i \in V_\infty^K(k)} (-1)^{s_i} \prod_{v \in V_f(k)} d_v = 1$.

The hermitian form is uniquely determined by this data.

3.3.1. $SU(r, s)$ with $r, s \geq 2$

We take k of type I and fix some $w^0 \in V_f(k)$. By weak approximation, there is an element $a \in k^\times$ which is negative at all real places and is a non-square in k_{v^0} . Define $K = k(\sqrt{a})$

and observe that by construction all archimedean places are non-split. In addition, the finite place w^0 does not split in K . Using the result of Landherr, there is a unique hermitian form h_j such that:

- (i) the signature at the j th real place is (r, s) ;
- (ii) the signature at all other real places is $(r + s, 0)$;
- (iii) the discriminant at w^0 is $d_{w^0} = (-1)^s$; and
- (iv) the discriminant is $d_v = 1$ for every $v \in V_f(k) \setminus \{w^0\}$.

These hermitian forms define linear algebraic groups $\mathbf{SU}(h_j)$ over k and by Theorem 2.5 (a) arithmetic subgroups $\Gamma_j \subset \mathbf{SU}(h_j)(k)$ which are cocompact lattices in $\mathbf{SU}(h_j)(k_j) = \mathbf{SU}(r, s)$. These lattices have a congruence kernel of order at most two by Theorem 2.4 (Since the compact Lie group $\mathbf{SU}(r + s)$ is topologically simply connected, the congruence kernel might however be nontrivial.) In any case, hermitian forms over p -adic fields are classified by dimension and discriminant [9, theorem 3.1], so the forms h_j are isometric at each finite place. By Lemma 2.6 the groups $\mathbf{SU}(h_j)(\mathbb{A}_k^f)$ are all isomorphic. It follows immediately from Theorem 2.5 (b) that Γ_i and Γ_j are pairwise profinitely commensurable. Finally, we observe that the groups $\mathbf{SU}(h_j)$ are not isomorphic. By construction k has no non-trivial automorphisms and therefore superrigidity via Theorem 2.7 (we recall that $r, s \geq 2$) implies that Γ_i and Γ_j are not commensurable for $i \neq j$.

3.3.2. $\mathrm{SL}_{2m}(\mathbb{R})$ and $\mathrm{SL}_{2m}(\mathbb{C})$ for $m \geq 2$

Let \mathbb{K} denote either \mathbb{R} or \mathbb{C} and let $G = \mathrm{SL}_{2m}(\mathbb{K})$. If $\mathbb{K} = \mathbb{R}$ we take k of type I and otherwise of type II. Pick a rational prime number p which splits completely in k and let w_1, \dots, w_n denote the places of k dividing p , i.e., $k_{w_i} \cong \mathbb{Q}_p$. We fix an additional finite place $w^0 \in V_f(k)$ which is distinct from w_1, \dots, w_n . By weak approximation, there is an element $a \in k^\times$ which satisfies: a is negative in k_i for all $i \geq 2$, represents a prescribed non-square element $x \in \mathbb{Q}_p^\times$ modulo squares at the places w_1, \dots, w_n and is a non-square at w^0 . If $\mathbb{K} = \mathbb{R}$ we can arrange that, in addition, a is positive in k_1 . Define $K = k(\sqrt{a})$. Then $V_\infty^K(k)$ contains all but the first archimedean places. By construction, the places w_1, \dots, w_n, w^0 are not split in K and the quadratic extensions K_{w_i}/k_{w_i} are isomorphic to $\mathbb{Q}_p(\sqrt{x})/\mathbb{Q}_p$. Using the result of Landherr, there is a unique hermitian form h_j such that:

- (i) the signature at k_2, \dots, k_n is $(2m, 0)$;
- (ii) the discriminant at w^0 and w_j equals -1 ; and
- (iii) the discriminant is $d_v = 1$ for every $v \in V_f(k) \setminus \{w_j, w^0\}$.

By construction, $\mathbf{SU}(h_j)(k_i) \cong \mathbf{SU}(2m)$ for all $i \geq 2$. We observe that $\mathbf{SU}(h_j)(k_1) \cong \mathrm{SL}_{2m}(\mathbb{K})$ due to the choices of k and a . The groups $\mathbf{SU}(h_j)$ are pairwise non-isomorphic, since they are non-isomorphic at one of the finite places w_1, \dots, w_n . Here it is essential that $2m$ is even; only under this assumption the classification [32, 4.4] entails that non-isomorphic hermitian forms have non-isomorphic special unitary groups. We claim that the topological groups $\mathbf{SU}(h_j)(\mathbb{A}_k^f) \cong \mathbf{SU}(h_i)(\mathbb{A}_k^f)$ are isomorphic for all i, j . Indeed, the groups $\mathbf{SU}(h_j)$ are isomorphic at all finite places except for w_1, \dots, w_n ; here a permutation of the places w_1, \dots, w_n yields an isomorphism

$$\prod_{\ell=1}^n \mathbf{SU}(h_j)(k_{w_\ell}) \cong \prod_{\ell=1}^n \mathbf{SU}(h_i)(k_{w_\ell}).$$

The argument used in the proof of Lemma 2.6 implies $\mathbf{SU}(h_j) \left(\mathbb{A}_k^f \right) \cong \mathbf{SU}(h_i) \left(\mathbb{A}_k^f \right)$ and we can proceed as above to obtain the lattices $\Gamma_1, \dots, \Gamma_n$ as arithmetic subgroups of $\Gamma_j \subseteq \mathbf{SU}(h_j)(k)$.

3.4. Type D_\bullet : $\mathbf{SO}^0(r, s)$ with $r + s$ even and $r + s \geq 8$ and $\mathbf{SO}_8(\mathbb{C})$

Fix an odd prime number p which completely splits in k and let w_1, \dots, w_n denote the finite places of k dividing p . We fix an additional finite place $w^0 \in V_f(k)$ which is distinct from w_1, \dots, w_n .

3.4.1. $\mathbf{SO}^0(r, s)$ with $r + s$ even and $r + s \geq 8$

The argument here is similar to the argument above for $\mathbf{SU}(r, s)$, now using the corresponding local-global principle for quadratic forms. Recall that quadratic forms over \mathbb{R} are classified by their dimension and signature; quadratic forms over p -adic fields are classified by dimension, determinant (modulo squares) and the Hasse invariant (see [28, section 6.4]).

Let k be of type I. We will use [28, chapter 6, theorem 6.10] to construct quadratic forms q_1, \dots, q_n of dimension $r + s$ over k such that:

- (i) q_j has signature (r, s) at the first real place but is positive definite over all other real places;
- (ii) q_1, \dots, q_n are isometric at every finite place outside w_1, \dots, w_n ;
- (iii) q_i is non-split at the place w_i , but split at w_j for all $j \neq i$.

In addition, we can achieve that $\mathbf{Spin}(q_i)(k_{w_i}) \cong \mathbf{Spin}(q_j)(k_{w_j})$ as topological groups for all i, j , which entails

$$\mathbf{Spin}(q_i) \left(\mathbb{A}_k^f \right) \cong \mathbf{Spin}(q_j) \left(\mathbb{A}_k^f \right)$$

using the argument of Lemma 2.6. By a result of Kneser, the groups $\mathbf{Spin}(q_j)$ have CSP; see [14, 11.1]. As before, it follows from Theorems 2.5 and 2.7 that arithmetic subgroups of the algebraic groups $\mathbf{Spin}(q_j)$ give rise to profinitely commensurable cocompact lattices in $\mathbf{SO}^0(r, s)$ which are not commensurable.

By weak approximation, there is an element $a \in k^\times$ such that $(-1)^s a$ is positive at the first real place, a is positive at all other real places and such that $(-1)^{(r+s)/2} a$ is a square in k_{w_i} for all $i \in \{1, 2, \dots, n\}$. We define q_j to be the unique form of determinant a which has signature (r, s) at the first real place, is positive definite at all other real places, has Hasse invariant -1 at w_j , and has Hasse invariant 1 at the finite places not equal to w^0, w_j . The Hasse invariant at w^0 is then determined by the product formula. It depends on r, s , and n , but not on j .

We observe that q_j is split at w_i for all $i \neq j$, since q_j has $r + s$ variables, and the determinant $(-1)^{(r+s)/2}$ (i.e., a modulo squares) and the Hasse invariant 1 are equal to the determinant and Hasse invariant of the $(r + s)/2$ -fold orthogonal sum of hyperbolic planes $\langle 1, -1 \rangle$. On the other hand, q_j is non-split at w_j , because the Hasse invariant of q_j and the split form differ. We observe that modulo the canonical isomorphism $k_{w_i} \cong \mathbb{Q}_p \cong k_{w_j}$ the forms are isometric and hence the groups $\mathbf{Spin}(q_i)(k_{w_i})$ and $\mathbf{Spin}(q_j)(k_{w_j})$ are isomorphic as topological groups. However, for $i \neq j$ the Witt indices of q_i and q_j at w_j differ, hence the k_{w_j} -rank of the groups $\mathbf{Spin}(q_i)$ and $\mathbf{Spin}(q_j)$ are different. So these algebraic groups are not isomorphic over k . Note that our discussion also covers the group $G = \mathbf{SO}^*(8)$ which happens to be locally isomorphic to $\mathbf{SO}^0(6, 2)$.

3.4.2. $SO_8(\mathbb{C})$

Let k be of type II. Recall that w_1, \dots, w_n denote the finite places of k dividing p and w^0 denotes a finite place away from p . For $i = 1, \dots, n$, there exists a non-degenerate quadratic k -form q_i of rank eight, with trivial determinant, such that q_i is anisotropic at all real places of k , such that it has Hasse invariant -1 at w_i and w^0 , and such that it has Hasse invariant 1 elsewhere. Over \mathbb{Q}_p , every nondegenerate quadratic form in at least four variables represents 1 . For $j = 1, \dots, n$, we thus have a decomposition $q_i \otimes_k k_{w_j} \cong \langle 1, 1, 1, 1 \rangle \oplus h_{ij}$ for some rank four quadratic \mathbb{Q}_p -form h_{ij} with trivial discriminant and the same Hasse invariant as $q_i \otimes_k k_{w_j}$. By our choice of Hasse invariants, h_{ij} is \mathbb{Q}_p -anisotropic if and only if $i = j$ [29, theorem 6 (iii), p. 36]. Since the form $\langle 1, 1, 1, 1 \rangle$ is metabolic over \mathbb{Q}_p (we assumed p is odd), the form q_i has Witt index two at w_i and Witt index four at w_j for $j \neq i$. This shows that the group $\mathbf{G}_i = \mathbf{Spin}(q_i)$ has \mathbb{Q}_p -rank two at w_i and \mathbb{Q}_p -rank four at w_j for $j \neq i$. The groups \mathbf{G}_i have moreover CSP by [14, 11.1], so that as above we can conclude that $\mathbf{G}_i(\mathbb{C}) \cong \mathbf{Spin}_8(\mathbb{C})$ has n profinitely commensurable cocompact lattices which are not commensurable.

3.5. The remaining real forms

Suppose G is locally isomorphic to $\mathbf{G}(\mathbb{R})$, where \mathbf{G} is a connected, simply connected absolutely almost simple algebraic \mathbb{R} -group. Assume that \mathbf{G} is neither of type A_m, E_6 , nor isomorphic to $\mathbf{Spin}(r, s)$. Let \mathbf{G}^u be the compact real form of \mathbf{G} . Under these assumptions \mathbf{G}^u is an inner form of \mathbf{G} as we saw in Section 3.2. Let \mathbf{G}_{qs} denote the unique quasi-split inner form of \mathbf{G} and \mathbf{G}^u .

Let k be of type I. By [4, proposition 1.10], we can choose a quasi-split, absolutely simple, simply connected algebraic group \mathbf{G}_{qs}^0 over k such that $\mathbf{G}_{qs}^0 \times_k k_i \cong \mathbf{G}_{qs}$ for every $i \in \{1, 2, \dots, n\}$. Fix a nonarchimedean place $w^0 \in V_f(k)$. Then by [24, theorem 1], for $i = 1, \dots, n$, there exists an inner k -twist \mathbf{G}_i of \mathbf{G}_{qs}^0 such that

- (i) $\mathbf{G}_i \times_k k_i$ is isomorphic to \mathbf{G} while
- (ii) $\mathbf{G}_i \times_k k_j$ is isomorphic to \mathbf{G}^u for $j \neq i$ and
- (iii) \mathbf{G}_i is isomorphic to \mathbf{G}_{qs}^0 at every finite place $w \neq w^0$.

Here, it is essential that \mathbf{G} and \mathbf{G}^u are inner twists of each other. Over p -adic fields, there only exist a finite number of inner twists of a given absolutely simple group. More precisely, in our context the non-abelian Galois cohomology $H^1(k_{w^0}, \text{Ad}(\mathbf{G}_i)) \cong H^2(k_{w^0}, Z(\mathbf{G}_i))$ has cardinality at most four [13, Satz 2 and table on p. 254] because we assume \mathbf{G} , hence \mathbf{G}_i , is not of type A_m . Hence if $n > 4(n' - 1)$, the pigeon hole principle guarantees that at least n' of the groups $\mathbf{G}_1, \dots, \mathbf{G}_n$ are isomorphic at all finite places. Without loss of generality, let us assume that the first n' groups $\mathbf{G}_1, \dots, \mathbf{G}_{n'}$ have this property; in particular, by Lemma 2.6,

$$\mathbf{G}_1\left(\mathbb{A}_k^f\right) \cong \mathbf{G}_2\left(\mathbb{A}_k^f\right) \cong \dots \cong \mathbf{G}_{n'}\left(\mathbb{A}_k^f\right). \tag{3.1}$$

Pick arithmetic subgroups $\Gamma_i \subseteq \mathbf{G}_i(k)$. The k -group \mathbf{G}_i is neither of type A_m , nor E_6 , nor D_4 (in which case it would be a spinor group) and we have $\text{rank}_{k_i} \mathbf{G}_i = \text{rank} G \geq 2$. Hence the congruence kernel $C(k, \mathbf{G}_i)$ has order at most two by Theorem 2.4.

One more time, by Theorem 2.5 and Theorem 2.7, we obtain n' profinitely commensurable but non-commensurable cocompact lattices $\Gamma_1, \dots, \Gamma_{n'}$ in $\mathbf{G}(\mathbb{R})$.

3.6. *The remaining complex forms*

Finally, let \mathbf{G} be a simply connected simple \mathbb{C} -group of type B_m ($m \geq 2$), C_m ($m \geq 2$), D_m ($m \geq 5$) or E_7 . In the following, all uniqueness statements for algebraic groups are meant up to isomorphism over the field of definition of the group. Let \mathbf{G}^u be the unique \mathbb{R} -anisotropic \mathbb{R} -group with $\mathbf{G}^u \times_{\mathbb{R}} \mathbb{C} = \mathbf{G}$. Let the \mathbb{R} -group \mathbf{G}_{qs} be the unique quasi split inner twist of \mathbf{G}^u . Finally, let \mathbf{G}^0 be the unique \mathbb{Q} -split \mathbb{Q} -group with $\mathbf{G}^0 \times_{\mathbb{Q}} \mathbb{C} = \mathbf{G}$.

Let k be of type II. Fix an odd prime p that splits in k and let w_1, \dots, w_n be the places of k over p . By [4, proposition 1.10], we find a quasi-split, absolutely simple, simply connected algebraic group \mathbf{G}_{qs}^0 over k which is isomorphic to \mathbf{G}_{qs} at all real places of k and is isomorphic to $\mathbf{G}^0 \times \mathbb{Q}_p$ at w_1, \dots, w_n . According to Kneser [13, Satz 2 and table on p. 254], our assumption on the type of \mathbf{G} implies that there exists a non-trivial inner \mathbb{Q}_p -twist \mathbf{G}^p of $\mathbf{G}^0 \times_{\mathbb{Q}} \mathbb{Q}_p$. Fix another non-archimedean place w^0 which does not lie over p . By [24, theorem 1], for each $i = 1, \dots, n$, there exists an inner k -twist \mathbf{G}_i of \mathbf{G}_{qs}^0 such that:

- (i) \mathbf{G}_i is isomorphic to \mathbf{G}^u at every real place of k ;
- (ii) $\mathbf{G}_i \times_k k_{w_i}$ is isomorphic to \mathbf{G}^p ,
- (iii) $\mathbf{G}_i \times_k k_{w_j}$ is isomorphic to $\mathbf{G}^0 \times_{\mathbb{Q}} \mathbb{Q}_p$ for $j \neq i$;
- (iv) \mathbf{G}_i is isomorphic to \mathbf{G}_{qs}^0 at every finite place $w \notin \{w^0, w_1, \dots, w_n\}$.

By the same pigeon hole argument as above, we may assume that the first n' groups $\mathbf{G}_1, \dots, \mathbf{G}_{n'}$ are also isomorphic at w^0 . Since p splits in k , swapping any two places over p defines an automorphism of \mathbb{A}_k^f . It follows from the argument in Lemma 2.6 that the groups $\mathbf{G}_1(\mathbb{A}_k^f), \dots, \mathbf{G}_{n'}(\mathbb{A}_k^f)$ are pairwise isomorphic as topological groups. Since CSP is known for \mathbf{G}_i by Theorem 2.4, any arithmetic subgroups $\Gamma_1, \dots, \Gamma_{n'}$ of $\mathbf{G}_1, \dots, \mathbf{G}_{n'}$ are pairwise profinitely commensurable cocompact lattices in $\mathbf{G}(\mathbb{C})$ (Theorem 2.5) which are pairwise non-commensurable because k has no automorphism which could interchange the places w_1, \dots, w_n (Theorem 2.7).

3.7. *Non-cocompact lattices in special linear groups*

Our methods above exclusively produce cocompact lattices. To complement this, we sketch a mechanism to come up with profinitely isomorphic, non-commensurable, non-cocompact lattices in special linear groups.

PROPOSITION 3.1. *Let G be either $\text{SL}_m(\mathbb{R})$, $\text{SL}_m(\mathbb{C})$, or $\text{SL}_m(\mathbb{H})$ where $m \geq 6$ is a composite number. Then there are non-cocompact lattices $\Gamma_1, \Gamma_2 \subseteq G$ which are profinitely isomorphic but not commensurable.*

Proof. Assume that $G = \text{SL}_m(\mathbb{R})$ where $m \geq 6$ is composite. We write $m = dk$ with $d \geq 3$ and $k \geq 2$. Let C, D be two finite dimensional central division \mathbb{Q} -algebras of degree d , i.e., $\dim_{\mathbb{Q}} C = \dim_{\mathbb{Q}} D = d^2$. We assume that for every prime number p , at least one of the algebras $C_p = \mathbb{Q}_p \otimes_{\mathbb{Q}} C$ and $D_p = \mathbb{Q}_p \otimes_{\mathbb{Q}} D$ splits, this means, is isomorphic to $M_d(\mathbb{Q}_p)$. In addition, we assume that $\mathbb{R} \otimes_{\mathbb{Q}} C$ and $\mathbb{R} \otimes_{\mathbb{Q}} D$ are split. It follows from the theorem of Albert–Brauer–Hasse–Noether and the resulting description of the Brauer group $\text{Br}(\mathbb{Q})$ (see e.g. [20, section 18.5]), that there are infinitely many such pairs of division algebras (it would be interesting to have concrete examples though).

By construction, the central simple algebras $C \otimes_{\mathbb{Q}} D$ and $C \otimes_{\mathbb{Q}} D^{\text{op}}$ have index d (see [20, section 18.6, corollary]), i.e.,

$$C \otimes_{\mathbb{Q}} D \cong M_d(E_1), \quad C \otimes_{\mathbb{Q}} D^{\text{op}} \cong M_d(E_2)$$

for two division \mathbb{Q} -algebras E_1, E_2 of degree d . We define two central simple \mathbb{Q} -algebras

$$A = M_k(E_1), \quad B = M_k(E_2).$$

Since $d \geq 3$ is the order of $[C]$ and $[D]$ in $\text{Br}(\mathbb{Q})$ (see [20, section 18.6]), we deduce that the algebras C and D are not isomorphic to their opposite algebras, that is $C \not\cong C^{\text{op}}$ and $D \not\cong D^{\text{op}}$. It follows (using $[A] = [E_1] = [C][D]$ and $[B] = [E_2] = [C][D]^{-1}$ in $\text{Br}(\mathbb{Q})$) that A is neither isomorphic to B nor to B^{op} . Therefore, the associated reduced norm-one groups $\mathbf{G} = \text{SL}_1(A)$ and $\mathbf{H} = \text{SL}_1(B)$ are not isomorphic. In fact, reduced norm-one groups are isomorphic if and only if the underlying algebras are isomorphic or opposite isomorphic (see [15, (26.9) and (26.11)]). For every prime number p we have $\mathbf{G}(\mathbb{Q}_p) \cong \mathbf{H}(\mathbb{Q}_p)$. Indeed, suppose C splits at p , then $E_{1,p} \cong D_p$ and $E_{2,p} \cong D_p^{\text{op}}$, and consequently $A_p = M_k(D_p) \cong B_p^{\text{op}}$. If on the other hand D splits at p , then $A_p \cong M_k(C_p) \cong B_p$.

Since C and D split over the real numbers, we have $\mathbf{G}(\mathbb{R}) \cong \mathbf{H}(\mathbb{R}) \cong \text{SL}_m(\mathbb{R})$. The groups \mathbf{G}, \mathbf{H} are isotropic (because $k \geq 2$) and thus have the congruence subgroup property; see Theorem 2.4. As above, one can show that arithmetic subgroups of \mathbf{G} and \mathbf{H} are lattices in $\text{SL}_m(\mathbb{R})$ which are profinitely commensurable but not commensurable.

If we replace \mathbb{Q} by an imaginary quadratic number field, the same construction yields profinitely isomorphic lattices in $\text{SL}_m(\mathbb{C})$. In order to obtain lattices in $\text{SL}_m(\mathbb{H})$, we vary the argument and write $2m = dk$ as a product of $k \geq 2$ and an even number $d \geq 4$. We choose C, D of degree d as before, now assuming that C is ramified over \mathbb{R} and D splits over \mathbb{R} .

4. Proof of Theorem 1.2

In this section we show that profinitely commensurable lattices in a connected simple complex Lie group G of type E_8, F_4 , or G_2 are abstractly commensurable. Three features of G are used to conclude Theorem 1.2: G is simply connected, has trivial center, and has no Dynkin diagram symmetries. In particular, G is uniquely determined by its Lie algebra \mathfrak{g} : we have $G \cong \mathbf{G}(\mathbb{C})$ for the linear algebraic \mathbb{C} -group $\mathbf{G} = \text{Aut}_{\mathbb{C}}(\mathfrak{g})$.

So let Γ_1 and Γ_2 be two profinitely commensurable lattices in $\mathbf{G}(\mathbb{C})$. We need to show that Γ_1 is commensurable with Γ_2 . By Margulis arithmeticity [18, theorem IX.1.11 and p. 293/294], for $i = 1, 2$, there exists a dense number subfield $k_i \subset \mathbb{C}$ whose remaining infinite places are real and there exists a simply connected absolutely almost simple k_i -group \mathbf{G}_i which is anisotropic at all real places of k_i such that for any k_i -embedding $\mathbf{G}_i \subset \mathbf{GL}_r$ the group of k_i -integral points $\mathbf{G}_i(\mathcal{O}_{k_i})$ is commensurable with Γ_i . Since commensurable groups are also profinitely commensurable, it follows that $\mathbf{G}_1(\mathcal{O}_{k_1})$ is profinitely commensurable with $\mathbf{G}_2(\mathcal{O}_{k_2})$. Consequently by [11, theorem 4], k_1 is arithmetically equivalent to k_2 . From [6, corollary 1.4], we conclude that actually k_1 is abstractly isomorphic to k_2 , hence the two subfields of \mathbb{C} are either equal or complex conjugates of one another. Replacing k_2 and \mathbf{G}_2 with the complex conjugate field and group if need be, we may assume $k_1 = k_2 = k$.

Again owing to the exceptional type at hand, the groups \mathbf{G}_i have trivial center and no Dynkin diagram symmetries. We thus have $\text{Aut}(\mathbf{G}_i) \cong \mathbf{G}_i$, so the k -isomorphism type of \mathbf{G}_2 is classified by a class $\alpha \in H^1(k, \mathbf{G}_1)$ in non-commutative Galois cohomology with values in \mathbf{G}_1 . Since \mathbf{G}_1 is isomorphic to \mathbf{G}_2 at all infinite places of k , we see that α reduces to the distinguished point in $H^1(k_v, \mathbf{G}_1)$ for all $v \in V_{\infty}(k)$. But by the main theorem of Galois

cohomology of simply connected groups [21, Theorem 6.6, p. 289], we have a bijection of pointed sets

$$\theta : H^1(k, \mathbf{G}_1) \xrightarrow{\cong} \prod_{v \in V_\infty(k)} H^1(k_v, \mathbf{G}_1).$$

So α is the trivial twist, hence \mathbf{G}_1 is k -isomorphic to \mathbf{G}_2 . In particular, $\mathbf{G}_1(\mathcal{O}_k)$ is commensurable with $\mathbf{G}_2(\mathcal{O}_k)$, and so is Γ_1 with Γ_2 .

5. Proof of Theorem 1.3

Finally, in this section we give the proof of Theorem 1.3. Let \mathfrak{g} be the Lie algebra of G . The simple linear algebraic \mathbb{R} -group $\mathbf{G}_{\text{ad}} = \text{Aut}_{\mathbb{R}}(\mathfrak{g})$, satisfies $\mathbf{G}_{\text{ad}}(\mathbb{R})^0 = G$. Let $\tilde{\mathbf{G}}$ be the simply connected covering of \mathbf{G}_{ad} . We will construct profinitely isomorphic, non-isomorphic cocompact lattices $\Gamma_1, \dots, \Gamma_n \subseteq \tilde{\mathbf{G}}(\mathbb{R})$ which do not intersect the center and are thus isomorphic to lattices in G .

If $\tilde{\mathbf{G}}$ is absolutely simple, then we take k of type I and set $\mathbf{G} = \tilde{\mathbf{G}}$. If $\tilde{\mathbf{G}}$ is not absolutely simple, then $\tilde{\mathbf{G}} = \text{Res}_{\mathbb{C}/\mathbb{R}}(\mathbf{G})$ for a simply connected simple algebraic \mathbb{C} -group \mathbf{G} . In this case we choose k of type II. Let p be a rational prime number which splits completely in k with finite places w_1, \dots, w_n dividing p and $k_{w_i} \cong \mathbb{Q}_p$ for all i .

By Theorem B in [4] there is a simply connected, absolutely simple algebraic k -group \mathbf{H} such that:

- (i) $\mathbf{H} \times_k k_1 \cong \mathbf{G}$;
- (ii) $\mathbf{H}(k_j)$ is compact for all $j \geq 2$; and
- (iii) $\mathbf{H}(k_{w_i}) \cong \mathbf{H}(k_{w_j})$ for all $i, j \in \{1, \dots, n\}$.

If G is of type A_m (using $G \neq \text{PSL}_m(\mathbb{H})$) we can take \mathbf{H} to be a special unitary group. If G is of type D_4 , we can take \mathbf{H} to be a spinor group. By construction the group \mathbf{H} then has the congruence subgroup property and we can find an arithmetic subgroup $\Gamma \subseteq \mathbf{H}(k)$ which does not intersect the congruence kernel, so that $\hat{\Gamma} \cong \bar{\Gamma} \subseteq \mathbf{H}(\mathbb{A}_k^f)$; here $\bar{\Gamma}$ denotes the closure of Γ in $\mathbf{H}(\mathbb{A}_k^f)$. Passing to a finite index subgroup if need be, we may assume that Γ intersects the center of \mathbf{H} trivially, too.

We decompose the ring of finite adeles as $\mathbb{A}_k^f = \prod_{i=1}^n k_{w_i} \times \mathbb{A}_k^{f,p'}$ into the p - and p' -part. We can find open compact subgroups $K_{0,i} \subseteq \mathbf{H}(k_{w_i})$ such that $K_{0,i} \cong K_{0,j}$ for all i, j and such that

$$\prod_{i=1}^n K_{0,i} \times K_f^{p'} \subseteq \bar{\Gamma},$$

where $K_f^{p'}$ is some open compact subgroup of $\mathbf{H}(\mathbb{A}_k^{f,p'})$. We choose open compact subgroups $K_{1,i} \subseteq K_{0,i}$ again such that $K_{1,i} \cong K_{1,j}$ for all i, j of sufficiently large index satisfying $|K_{0,i} : K_{1,i}| > |\mathbf{Z}(\mathbf{H})(k)|$. Now we define the arithmetic groups

$$\Gamma_j = \Gamma \cap \prod_{i=1}^n K_{\delta_{ij},i} \times K_f^{p'},$$

where $\delta_{i,j}$ is the Kronecker–Delta. By construction $\hat{\Gamma}_j \cong \prod_{i=1}^n K_{\delta_{ij},i} \times K_f^{p'}$, hence the groups $\Gamma_1, \dots, \Gamma_n$ are profinitely isomorphic. Being subgroups of Γ , they do not intersect the center.

It remains to prove that the arithmetic groups $\Gamma_1, \dots, \Gamma_n$ are pairwise non-isomorphic. We explain this for Γ_1 and Γ_2 . Suppose for a contradiction that $\phi: \Gamma_1 \rightarrow \Gamma_2$ is an isomorphism. As Γ_2 is a subgroup of $\mathbf{H}(k)$, it follows from Margulis superrigidity [18, theorem (5), p.5] (using that k has no automorphisms) that there is an automorphism η of \mathbf{H} defined over k and a homomorphism $\nu: \Gamma_1 \rightarrow \mathbf{Z}(\mathbf{H})(k)$ such that

$$\phi(\gamma) = \nu(\gamma)\eta(\gamma)$$

for all $\gamma \in \Gamma_1$. Fix a Haar measure on $\mathbf{H}(k_{w_1})$. Using the inclusion of Γ_2 into $\mathbf{H}(k_{w_1})$, we see that the closure $\overline{\phi(\Gamma_1)}$ of $\phi(\Gamma_1)$ in $\mathbf{H}(k_{w_1})$ is $K_{0,1}$. On the other hand, $\overline{\phi(\Gamma_1)}$ is contained in $\mathbf{Z}(\mathbf{H})(k)\eta(K_{1,1})$. This can be used to derive a contradiction, since the Haar measure of the latter is strictly smaller than the Haar measure of $K_{0,1}$. Recall that $\mathbf{H}(k_1)$ is unimodular [18, I(2.2.3)] and that the inner automorphism group of \mathbf{H} has finite index in the automorphism group of \mathbf{H} . We deduce that η preserves the Haar measure and therefore

$$\begin{aligned} \text{vol}(K_{0,1}) &= \text{vol}(\overline{\phi(\Gamma_1)}) \leq |\mathbf{Z}(\mathbf{H})(k)|\text{vol}(\eta(K_{1,1})) \\ &= |\mathbf{Z}(\mathbf{H})(k)|\text{vol}(K_{0,1})|K_{0,1} : K_{1,1}|^{-1} < \text{vol}(K_{0,1}) \end{aligned}$$

which is a contradiction.

Acknowledgements. It is our pleasure to thank G. Mantilla Soler and S. Kühnlein for helpful discussions. We are grateful for financial support from the German Research Foundation [DFG 338540207 (SPP 2026/1), DFG 441848266 (SPP 2026/2), and DFG 281869850 (RTG 2229)].

REFERENCES

[1] M. AKA. Profinite completions and Kazhdan’s property (T). *Groups Geom. Dyn.* **6** (2012), no. 2, 221–229.
 [2] M. AKA. Arithmetic groups with isomorphic finite quotients. *J. Algebra* **352** (2012), 322–340.
 [3] A. BOREL and HARISH-CHANDRA. Arithmetic subgroups of algebraic groups. *Ann. of Math.* (2) **75** (1962), 485–535.
 [4] A. BOREL and G. HARDER. Existence of discrete cocompact subgroups of reductive groups over local fields. *J. Reine Angew. Math.* **298** (1978), 53–64.
 [5] M. R. BRIDSON, D. B. MCREYNOLDS, A. W. REID and R. SPITLER. Absolute profinite rigidity and hyperbolic geometry. *Ann. of Math.* (2) **192** (2020), no. 3, 679–719.
 [6] T. CHINBURG, E. HAMILTON, D. D. LONG and A. W. REID. Geodesics and commensurability classes of arithmetic hyperbolic 3-manifolds. *Duke Math. J.* **145** (2008), no. 1, 25–44.
 [7] P. DELIGNE. Extensions centrales non residuellement finies de groupes arithmetiques, C. R. Acad. Sci. Paris Sér. A-B **287** (1978), no. 4, A203–A208 (French, with English summary).
 [8] P. GILLE. Le probleme de Kneser-Tits. *Astérisque* **326** (2009), Exp. No. 983, vii, 39–81 (2010), (French, with French summary). *Seminaire Bourbaki*. vol. 2007/2008.
 [9] R. JACOBOWITZ. Hermitian forms over local fields, *Amer. J. Math.* **84** (1962), 441–465.
 [10] H. KAMMEYER. Introduction to ℓ^2 invariants. *Lecture Notes in Mathematics*, vol. 2247, (Springer, Cham, 2019).
 [11] H. KAMMEYER. Profinite commensurability of S-arithmetic groups. *Acta Arith.* **197** (2021), no. 3, 311–330.
 [12] H. KAMMEYER, S. KIONKE, J. RAIMBAULT and R. SAUER. Profinite invariants of arithmetic groups. *Forum Math. Sigma* **8** (2020), paper no. e54, 22.
 [13] M. KNESER. Galois-Kohomologie halbeinfacher algebraischer Gruppen über p-adischen Körpern. II, *Math. Z.* **89** (1965), 250–272.
 [14] M. KNESER. Normalteiler ganzzahliger Spingruppen. *J. Reine Angew. Math.* **311(312)** (1979), 191–214 (German).

- [15] M.-A. KNUS, A. MERKURJEV, M. ROST and J.-P. TIGNOL. *The book of involutions* American Mathematical Society Colloquium Publications, vol. 44 (American Mathematical Society, Providence, RI, 1998). With a preface in French by J. Tits. MR1632779
- [16] W. LANDHERR. Äquivalenz Hermitescher Formen über einem beliebigen algebraischen Zahlkörper. *Abh. Math. Sem. Univ. Hamburg* **11** (1935), no. 1, 245–248. (German).
- [17] A. LUBOTZKY. Finiteness properties and profinite completions. *Bull. Lond. Math. Soc.* **46** (2014), no. 1, 103–110.
- [18] G. A. MARGULIS. Discrete subgroups of semisimple Lie groups. *Ergeb. Math. Grenzgeb.* (3), vol. 17 (Springer-Verlag, Berlin, 1991).
- [19] J. S. MILNE and J. SUH. Nonhomeomorphic conjugates of connected Shimura varieties. *Amer. J. Math.* **132** (2010), no. 3, 731–750.
- [20] R. S. PIERCE. Associative Algebras. *Graduate Texts in Math.* vol. 88 (Springer-Verlag, New York, 1982).
- [21] V. PLATONOV and A. RAPINCHUK. *Algebraic groups and number theory. Pure and Applied Mathematics*, vol. 139 (Academic Press, Inc., Boston, MA, 1994). Translated from the 1991 Russian original by Rachel Rowen.
- [22] G. PRASAD and A. S. RAPINCHUK. Computation of the metaplectic kernel. *Inst. Hautes Études Sci. Publ. Math.* **84** (1996), 91–187 (1997).
- [23] G. PRASAD and A. S. RAPINCHUK. Developments on the congruence subgroup problem after the work of Bass, Milnor and Serre (2008). e-print arXiv:0809.1622.
- [24] G. PRASAD and A. S. RAPINCHUK. On the existence of isotropic forms of semi-simple algebraic groups over number fields with prescribed local behavior. *Adv. Math.* **207** (2006), no. 2, 646–660.
- [25] G. PRASAD and A. S. RAPINCHUK. Weakly commensurable arithmetic groups and isospectral locally symmetric spaces. *Publ. Math. Inst. Hautes Études Sci.* **109** (2009), 113–184.
- [26] A. W. REID. Profinite rigidity. Proceedings of the International Congress of Mathematicians— Rio de Janeiro 2018. Vol. II. Invited lectures (World Sci. Publ., Hackensack, NJ, 2018), pp. 1193–1216.
- [27] L. RIBES and P. ZALESKII. Profinite groups. 2nd ed., *Ergeb. Math. Grenzgeb.* (3). Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 40 (Springer-Verlag, Berlin, 2010).
- [28] W. SCHARLAU. *Quadratic and Hermitian forms Grundlehren Math. Wiss.* [Fundamental Principles of Mathematical Sciences], vol. 270 (Springer-Verlag, Berlin, 1985).
- [29] J.-P. SERRE. *A Course in Arithmetic* (Springer-Verlag, New York-Heidelberg, 1973). Translated from the French; *Graduate Texts in Math.*, No. 7.
- [30] M. STOVER. Lattices in $PU(n, 1)$ that are not profinitely rigid. *Proc. Amer. Math. Soc.* **147** (2019), no. 12, 5055–5062.
- [31] J. TITS. Classification of algebraic semisimple groups. *Algebraic Groups and Discontinuous Subgroups* (Proc. Sympos. Pure Math., Boulder, Colorado, 1965), (Amer. Math. Soc., Providence, R.I., 1966, 1966), pp. 33–62.
- [32] J. TITS. Reductive groups over local fields. Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Proc. Sympos. Pure Math., XXXIII. (Amer. Math. Soc., Providence, R.I., 1979), pp. 29–69