

On Reducibility and Uнитарizability for Classical p -Adic Groups, Some General Results

Marko Tadić

Abstract. The aim of this paper is to prove two general results on parabolic induction of classical p -adic groups (actually, one of them holds also in the archimedean case), and to obtain from them some consequences about irreducible unitarizable representations. One of these consequences is a reduction of the unitarizability problem for these groups. This reduction is similar to the reduction of the unitarizability problem to the case of real infinitesimal character for real reductive groups.

1 Introduction

Let F be a p -adic field. The normalized absolute value on F will be denoted by $|\cdot|_F$. Denote by $\nu: GL(n, F) \rightarrow \mathbb{R}^\times$ the character $\nu(g) = |\det(g)|_F$. For two smooth representations π_1 and π_2 of $GL(n_1, F)$ and $GL(n_2, F)$ respectively, we can consider $\pi_1 \otimes \pi_2$ as a representation of the Levi factor of the maximal parabolic subgroup which contains upper triangular matrices and whose Levi factor is isomorphic to $GL(n_1, F) \times GL(n_2, F)$. Following Bernstein and Zelevinsky, we denote by $\pi_1 \times \pi_2$ the representation of $GL(n_1 + n_2, F)$ parabolically induced by $\pi_1 \otimes \pi_2$ (see Section 2 for more details regarding notation). For an irreducible essentially square integrable representation δ of $GL(n, F)$ there exists a unique $e(\delta) \in \mathbb{R}$ and a unique unitarizable (irreducible square integrable) representation δ^u , such that $\delta = \nu^{e(\delta)} \delta^u$.

We shall fix a tower of symplectic or odd-orthogonal spaces over F . We shall denote by S_n the group with split rank n of isometries of determinant one of a space from the tower. Since Levi factors of parabolic subgroups are direct products of general linear groups and smaller groups S_m for a smooth representations π and σ of $GL(p, F)$ and S_q respectively, we can parabolically induce $\pi \otimes \sigma$ and get a representation of S_{p+q} (similarly as in the case of general linear groups). We shall denote the parabolically induced representation by $\pi \rtimes \sigma$.

The aim of this paper is to prove two general results regarding parabolic induction of classical p -adic groups (actually, one of them holds also in the archimedean case), and to obtain from them some consequences about irreducible unitarizable representations, like a reduction of the problem of unitarizability for these groups.

The first of the two main results of the paper gives a sufficient condition that $\pi \rtimes \sigma$ is irreducible, for irreducible representations π and σ (see Proposition 3.2).

Received by the editors May 27, 2006; revised January 12, 2007.

The author was partly supported by Croatian Ministry of Science, Education and Sports grant 037-0372794-2804.

AMS subject classification: Primary: 22E50; secondary: 22E35.

©Canadian Mathematical Society 2009.

This result could be also obtained by the (powerful) methods of [7], after necessary modifications. Instead of explaining this technical result here, we shall recall of the most important consequence. An irreducible representation σ of S_q will be called weakly real if, whenever we have an embedding

$$(1.1) \quad \sigma \hookrightarrow \rho_1 \times \cdots \times \rho_l \rtimes \sigma',$$

where ρ_1, \dots, ρ_l are irreducible cuspidal representations of general linear groups and σ' is an irreducible cuspidal representation of some $S_{q'}$, then $\rho_i^u \cong \tilde{\rho}_i^u$ for all $i \in \{1, \dots, l\}$ ($\tilde{\rho}$ denotes the contragredient representation of ρ).

Theorem 1.1 *If σ is an irreducible unitarizable representation of some S_q , then there exist an irreducible unitarizable representation π of a general linear group and a weakly real irreducible unitarizable representation σ' of some $S_{q'}$ such that $\sigma \cong \pi \rtimes \sigma'$.*

Since we have a classification of irreducible unitarizable representations of general linear groups [14], the above theorem reduces the problem of classification of irreducible unitarizable representations of classical groups to the problem of classification of weakly real irreducible unitarizable representations of classical groups. (Such a reduction in [21] for irreducible square integrable representations of classical p -adic groups was an important initial step in the process of classifying irreducible square integrable representations of classical p -adic groups, which ended with such a classification modulo cuspidal data in [8, 9].)

Suppose that an irreducible representation σ of S_m is a subquotient of a principal series representation $\text{Ind}_{P_\emptyset}^{S_m}(\chi)$. If σ is weakly real, then χ is real valued. In particular, if σ is a weakly real irreducible representation with Iwahori fixed vector, then its infinitesimal character is determined by (A_\emptyset, χ) , and χ is a real valued unramified character. Therefore, the above theorem is similar to the reduction of the unitarizability problem to the case of real infinitesimal character for real reductive groups. The above theorem is also related to the reduction obtained in [3].

In this paper we also get an upper bound for the places where weakly real irreducible unitarizable representations of classical p -adic groups can show up. We explain this bound now. Let ρ be an irreducible self dual cuspidal representation of a general linear group and let σ' be an irreducible cuspidal representation of some S_q . Then there exists a unique $\alpha_{\rho, \sigma'} \geq 0$ such that $\nu^{\alpha_{\rho, \sigma'}} \rho \rtimes \sigma'$ reduces.

Proposition 1.2 *Suppose that σ is a weakly real irreducible unitarizable representations of a classical group. Fix any embedding as in (1.1) and fix $i_0 \in \{1, \dots, l\}$. Denote by π_1, \dots, π_n all the representations from $\{\rho_1, \dots, \rho_l\}$ for which $\rho_i^u \cong \rho_{i_0}^u$. Write*

$$\{ |e(\pi_i)| ; |e(\pi_i)| > \alpha_{\rho_{i_0}^u, \sigma'} \text{ and } 1 \leq i \leq n \} = \{ \alpha_1, \dots, \alpha_\ell \},$$

where $\ell \geq 0$ and $\alpha_1 < \alpha_2 < \dots < \alpha_\ell$. Then

- (i) $\alpha_1 - \alpha_{\rho_{i_0}^u, \sigma'} \leq 1$ if $\ell \geq 1$, and $\alpha_i - \alpha_{i-1} \leq 1$ for each $i = 2, 3, \dots, \ell$.
- (ii) $\alpha_i \leq \alpha_{\rho_{i_0}^u, \sigma'} + i; \quad i = 1, \dots, \ell$.

One of the main results of this paper is the identification of some irreducible subquotients of parabolically induced representations. It is natural to expect such a result if we consider (conjectural) local Langlands correspondences (and functoriality in its simplest form).

Let D be the set of all equivalence classes of irreducible essentially square integrable representation of $GL(n, F)$'s, $n \geq 1$. Denote $D_+ = \{\delta \in D; e(\delta) > 0\}$. The set of finite multisets in D (resp. D_+) will be denoted by $M(D)$ (resp. $M(D_+)$). We add multisets in a natural way. The Langlands classification for general linear groups attaches to each $d \in M(D)$ an irreducible representation $L(d)$ (see Section 5 for details).

Denote by $T(S)$ the set of all equivalence classes of irreducible tempered representations of groups S_q , $q \geq 0$. Then the Langlands classification for groups S_q parametrizes irreducible representations of these groups by elements of $M(D_+) \times T(S)$. To $(d', \tau) \in M(D_+) \times T(S)$ is attached $L(d', \tau)$ (see Section 5 for details).

For $d = (\delta_1, \dots, \delta_k) \in M(D)$ denote $\lambda(d) = \delta_1 \times \dots \times \delta_k$. We shall denote by d^\dagger the element of $M(D_+)$ which we get from d by removing all unitarizable δ_i 's and changing all δ_i 's for which $e(\delta_i) < 0$, by $\tilde{\delta}_i$. Denote by d_u the multiset in $M(D)$ which we get from d removing all δ_i 's which are not unitarizable.

Proposition 1.3 *Let $d \in M(D)$ and $t = (d', \tau) \in M(D_+) \times T(S)$. Denote by $T_{d,\tau}$ the set of all (equivalence classes of) irreducible subrepresentations of $\lambda(d_u) \rtimes \tau$. Then each of the representations $L(d^\dagger + d'; \tau')$, $\tau' \in T_{d,\tau}$ is a subquotient of $L(d) \rtimes L(d'; \tau)$. If F is of characteristic 0, then the multiplicity of each of these representations in $L(d) \rtimes L(d'; \tau)$ is 1.*

From this result we can conclude reducibility of some representations that are interesting for the construction of irreducible unitary representations of groups S_q (see the Section 5). Further work on understanding such reducibilities will be related to understanding of Arthur's R -groups (see Corollary 5.5 and [1]).

The above result is a generalization of the following result of A. V. Zelevinsky from the case of general linear groups (which played an important role in the process of classifying of irreducible unitary representations of these groups):

$$L(d_1 + d_2) \text{ is a subquotient of } L(d_1) \times L(d_2) \text{ with multiplicity 1.}$$

Let us recall that the above result holds also in the archimedean case (see [15]).

Let $F \subset F'$ be a separable quadratic extension, and let θ be a non-trivial element of the Galois group. We can consider towers of unitary groups associated with this extension (see [9]). When we fix one such series, all the results of this paper, after changing contragredients $\tilde{\pi}$ to representations $g \mapsto \tilde{\pi}(\theta(g))$ in them, hold also for unitary groups with the same proofs.

We expect that the results of this paper will also hold for orthogonal groups $O(2n)$, with basically the same proofs.

The results of this paper may be considered as initial steps of the work on the unitarizability problem for classical p -adic groups in general case. Some of the initial ideas in that direction are presented in the last section of [22]. We are finishing the

sequel of this paper, in which we give evidence for the approach to the unitarizability which we are starting here (and in [22]).

We now give more information about the content of this paper, section by section. Section 2 introduces notation. Section 3 presents the sufficient condition for irreducibility of parabolically induced representations. In Section 4 we list some consequences of this result mainly related to the problem of unitarizability. The description of Langlands parameters of some distinguished subquotients of parabolically induced representations $L(d) \rtimes L(d', \tau)$ is proved in Section 5. There we present some applications of this result to the unitarizability problem. In Section 6 we collect some simple previously known results on the reducibility of parabolic induction, which are useful in studying unitarizability.

2 Notation

First we shall introduce the basic notation for the representation theory of general linear groups over a p -adic field F . We shall use mainly the notation of Bernstein and Zelevinsky from [23]. The normalized absolute value on F will be denoted by $|\cdot|_F$. The character $g \mapsto |\det|_F, GL(n, F) \rightarrow \mathbb{R}^\times$ is denoted by ν . For each irreducible essentially square integrable representation δ of $GL(n, F)$ there exists a unique, up to an equivalence, unitarizable irreducible square integrable representation δ^u , and $e(\delta) \in \mathbb{R}$, such that $\delta = \nu^{e(\delta)} \delta^u$.

There is a natural map from the category of smooth representations of a reductive p -adic group G of finite length, to the Grothendieck group of this category. It will be denoted by ss (and is called a *semi-simplification*). An irreducible representation π of G can be considered in a natural way as an element of the Grothendieck group.

Let π_i be a smooth representation of $GL(n_i, F)$ for $i = 1, 2$. Then $\pi_1 \times \pi_2$ will denote the smooth representation of $GL(n_1 + n_2, F)$ parabolically induced by $\pi_1 \otimes \pi_2$ from an appropriate maximal parabolic subgroup, which is standard with respect to the subgroup of the upper triangular matrices (the parabolic induction that we consider is normalized). The sum of Grothendieck groups R_n of the category of smooth representations of $GL(n, F)$ of finite length is denoted by R . Then \times lifts in a natural way to a mapping $R \times R \rightarrow R$ which will again be denoted by \times . This mapping naturally factors through a mapping $R \otimes R \rightarrow R$, which will be denoted by m . For an admissible representation π of $GL(n, F)$, the sum of semi simplifications of the Jacquet modules for standard parabolic subgroups which have Levi subgroups $GL(k, F) \times GL(n-k, F)$, $0 \leq k \leq n$, defines an element of $R \otimes R$ in a natural way. (See [23] for a precise definition. The Jacquet modules that we consider in this paper will be always normalized.) This can be additively extended to a mapping $m^*: R \rightarrow R \otimes R$. In this way R becomes a Hopf algebra.

Let π be an irreducible representation of $GL(n, F)$. Then there exist irreducible cuspidal representations ρ_1, \dots, ρ_k of general linear groups such that π is isomorphic to a subquotient of $\rho_1 \times \dots \times \rho_k$. The multiset of equivalence classes (ρ_1, \dots, ρ_k) is called the cuspidal support of π (it depends only on the the equivalence class of π). It is denoted by $\text{supp}(\pi)$,

Now we shall introduce basic notation for the representation theory of the classical p -adic groups. We shall follow the notation of [17] and [9]. Fix a Witt tower $V \in \mathcal{V}$

of symplectic vector spaces over F , or of orthogonal vector spaces which starts with an anisotropic space of odd dimension. We shall denote by $S(V)$ the group of isometries of $V \in \mathcal{V}$ of determinant 1 (this is automatically satisfied in the symplectic case). The group of split rank n will be denoted by S_n . Now the sum of Grothendieck groups $R_n(S)$ of categories of smooth representations of S_n of finite length, is denoted by $R(S)$. Similarly as in the case of general linear groups, using parabolic induction, one defines $\pi \rtimes \sigma$ for a smooth representation π of a general linear group over F and a smooth representation σ of S_m . Now \rtimes lifts in a natural way to a mapping $R \times R(S) \rightarrow R$ which is again denoted by \rtimes (factorization through $R \otimes R(S)$ is denoted by μ). In this way $R(S)$ becomes R -module. Recall that $\pi \rtimes \sigma = \tilde{\pi} \rtimes \sigma$ in R ($\tilde{\pi}$ denotes the contragredient of π).

The Jacquet module of a representation π of S_n for the standard maximal parabolic subgroup whose Levi factor is a direct product of $GL(k, F)$ and a classical group, is denoted by $s_{(k)}(\pi)$. Suppose that π has finite length. The sum of semi simplifications of $s_{(k)}(\pi)$, $0 \leq k \leq n$, is denoted by $\mu^*(\pi)$. We can consider $\mu^*(\pi) \in R \otimes R(S)$. Extending additively to $\mu^* : R(S) \rightarrow R \otimes R(S)$, one gets the structure of R -comodule on $R(S)$.

Observe that $R \otimes R(S)$ is in a natural way $R \otimes R$ -module (the multiplication is denoted again by \rtimes). Further, denote by $\sim : R \rightarrow R$ the contragredient map and by $\kappa : R \otimes R \rightarrow R \otimes R, \sum x_i \otimes y_i \mapsto y_i \otimes x_i$. Let

$$M^* = (m \otimes \text{id}_R) \circ (\sim \otimes m^*) \circ \kappa \circ m^*.$$

Then $\mu^*(\pi \rtimes \sigma) = M^*(\pi) \rtimes \mu^*(\sigma)$ for admissible representations π and σ of $GL(n, F)$ and S_m respectively (or, for elements of R and $R(S)$ respectively). In other words, if $m^*(\pi) = \sum_i x_i \otimes y_i$, then

$$M^*(\pi) = \sum_i (m \otimes \text{id}_R)(\tilde{y}_i \times m^*(x_i)).$$

The term of $M^*(\pi)$ in $R \otimes R_0$ will be denoted by $M_{GL}^*(\pi)$. Then

$$M_{GL}^*(\pi) = [m \circ (\text{id}_R \otimes \sim) \circ m^*(\pi)] \otimes 1$$

(since R is commutative). Thus if $m^*(\pi) = \sum_i x_i \otimes y_i$, then

$$M_{GL}^*(\pi) = \sum_i x_i \times \tilde{y}_i.$$

The term $M^*(\pi)$ in $R_0 \otimes R$ is simply $1 \otimes \pi$.

Let τ be an irreducible representation of some classical group S_ℓ . Then it is a sub-quotient of $\rho_1 \times \cdots \times \rho_k \rtimes \sigma$ for some irreducible cuspidal representations ρ_1, \dots, ρ_k of general linear groups and an irreducible cuspidal representation σ of some S_m . The representation σ is called the partial cuspidal support of τ and is denoted by τ_{cusp} . If $\rho_1 \times \cdots \times \rho_k$ is a representation of $GL(p, F)$, then the Jacquet module $s_{(p)}(\tau)$ will be denoted by $s_{GL}(\tau)$. An irreducible cuspidal representation ρ of a general linear group

is called a factor of τ if there exists an irreducible subquotient $\pi \otimes_{\tau_{\text{cusp}}} \text{of } s_{GL}(\tau)$ such that ρ is in the cuspidal support of π . Then the set of all factors of τ is contained in

$$\{\rho_1, \tilde{\rho}_1, \rho_2, \tilde{\rho}_2, \dots, \rho_k, \tilde{\rho}_k\}$$

(recall that our τ is a subquotient of $\rho_1 \times \dots \times \rho_k \rtimes \sigma$). Further, for each $1 \leq i \leq k$, at least one representation from $\{\rho_i, \tilde{\rho}_i\}$ is a factor of τ .

We have a natural ordering \leq on the Grothendieck group of the category of smooth representations of a reductive p -adic group G , of finite length. Therefore, there are natural orderings on $R, R(S), R \otimes R$ and $R \otimes R(S)$. If we write $\pi_1 \leq \pi_2$ for two admissible representations, we shall actually mean inequality between their images in the Grothendieck group (*i.e.*, between their semi simplifications). We shall quite often use two forms of Frobenius reciprocity. Let $P = MN$ be a parabolic subgroup of G . Suppose that π and σ are smooth representations of G and M respectively. The Jacquet module of π with respect to $P = MN$ will be denoted by $r_P^G(\pi)$. Then Frobenius reciprocity says that we have a canonical isomorphism

$$\text{Hom}_G(\pi, \text{Ind}_P^G(\sigma)) \cong \text{Hom}_M(r_P^G(\pi), \sigma)$$

Let A_\emptyset be a maximal split torus in a connected reductive group G over F . Suppose that P contains A_\emptyset . Then we can find Levi subgroup M which contains A_\emptyset (such M is unique). Let \bar{P} be the opposite parabolic subgroup (this is the unique parabolic subgroup which contains A_\emptyset , whose Levi subgroup is M and which satisfies $P \cap \bar{P} = M$). Then the second form of Frobenius reciprocity is

$$\text{Hom}_G(\text{Ind}_{\bar{P}}^G(\sigma), \pi) \cong \text{Hom}_M(\sigma, r_{\bar{P}}^G(\pi)).$$

While the Frobenius reciprocity is an elementary fact, the second form of it is not. If π and σ are admissible (the case which we shall use), this follows from [4]. For general smooth π and σ , this is proved by J. Bernstein.

The following case of Frobenius reciprocity is of interest here. Let τ and ω be irreducible representations of $GL(p, F)$ and S_q respectively. Let π be an admissible representation of S_{p+q} . Now Frobenius reciprocity implies

$$\text{Hom}_{S_{p+q}}(\pi, \tau \rtimes \omega) \cong \text{Hom}_{GL(p,F) \times S_q}(s_{(p)}(\pi), \tau \otimes \omega).$$

From the second form of Frobenius reciprocity we get the isomorphism

$$\text{Hom}_{S_{p+q}}(\tau \rtimes \omega, \pi) \cong \text{Hom}_{GL(p,F) \times S_q}(\tau \otimes \omega, r_{\bar{P}}^{S_{p+q}}(\pi)),$$

where \bar{P} denotes the parabolic subgroup opposite to the standard parabolic subgroup which has $GL(p, F) \times S_q$ for the Levi subgroup. After conjugation by a suitable element of the Weyl group (which carries \bar{P} to P), we get

$$\text{Hom}_{S_{p+q}}(\tau \rtimes \omega, \pi) \cong \text{Hom}_{GL(p,F) \times S_q}(\tilde{\tau} \otimes \omega, s_{(p)}(\pi)).$$

Recall that in the above formula τ must be irreducible. We could write the above formulas in a more general setting, not necessarily for maximal parabolic subgroups.

Let ρ and σ be unitarizable irreducible cuspidal representations of a general linear group and of S_n respectively. Then if $\nu^\alpha \rho \rtimes \sigma$ reduces for some $\alpha \in \mathbb{R}$, then $\rho \cong \tilde{\rho}$. Further, if $\rho \cong \tilde{\rho}$, then we always have a reduction for unique $\alpha \geq 0$. This reducibility point will be denoted by $\alpha_{\rho,\sigma}$. In all known examples the following holds:

$$(2.1) \quad \alpha_{\rho,\sigma} \in (1/2)\mathbb{Z}.$$

F. Shahidi has proved this to be the case if σ is generic (see [12] and [13]). This is expected to hold in general.

3 A Criterion for Irreducibility

Lemma 3.1 *Let π be an irreducible representation of a classical group S_q and let ρ be an irreducible cuspidal representation of a general linear group $GL(p, F)$. Suppose*

- (1) $\rho \not\cong \tilde{\rho}$.
- (2) $\rho \rtimes \pi_{\text{cusp}}$ is irreducible.
- (3) $\rho \times \rho'$ and $\tilde{\rho} \times \rho'$ are irreducible for any factor ρ' of π .
- (4) Neither ρ nor $\tilde{\rho}$ is a factor of π .

Then $\rho \rtimes \pi$ is irreducible.

Proof Observe that if ρ' is an irreducible cuspidal representation of a general linear group such that $\rho' \otimes \pi' \leq \mu^*(\pi)$ for some non-zero π , then ρ' is a factor of π .

Recall that

$$\mu^*(\rho \rtimes \pi) = (1 \otimes \rho + \rho \otimes 1 + \tilde{\rho} \otimes 1) \rtimes \mu^*(\pi).$$

Suppose that an irreducible representation $\rho \otimes \tau$ (resp. $\tilde{\rho} \otimes \tau$) is a subquotient of $\mu^*(\rho \rtimes \pi)$. Since ρ (resp. $\tilde{\rho}$) is not a factor of π , and $\rho \not\cong \tilde{\rho}$, the above formula implies that we can get $\rho \otimes \tau$ (resp. $\tilde{\rho} \otimes \tau$) only from $(\rho \otimes 1) \rtimes \mu^*(\pi)$ (resp. $(\tilde{\rho} \otimes 1) \rtimes \mu^*(\pi)$). Considering the grading, we get that $\rho \otimes \tau$ (resp. $\tilde{\rho} \otimes \tau$) can come only from $(\rho \otimes 1) \rtimes (R_0 \otimes R(S))$ (resp. $(\tilde{\rho} \otimes 1) \rtimes (R_0 \otimes R(S))$). The only term of $\mu^*(\rho \rtimes \pi)$ in $R_0 \otimes R(S)$ is $1 \otimes \pi$. Thus $\rho \otimes \tau \leq (\rho \otimes 1) \rtimes (1 \otimes \pi) = \rho \otimes \pi$ (resp. $\tilde{\rho} \otimes \tau \leq \tilde{\rho} \otimes \pi$).

The above discussion implies that the multiplicities of $\rho \otimes \pi$ and $\tilde{\rho} \otimes \pi$ in $\mu^*(\rho \rtimes \pi)$ are both one. Moreover, we have shown that if $\rho \otimes \tau$ (resp. $\tilde{\rho} \otimes \tau$) is an irreducible subquotient of $\mu^*(\rho \rtimes \pi)$, then it must be $\rho \otimes \pi$ (resp. $\tilde{\rho} \otimes \pi$).

By Frobenius reciprocity, any irreducible subrepresentation of $\rho \rtimes \pi$ has $\rho \otimes \pi$ for a quotient of its Jacquet module. Therefore, $\rho \rtimes \pi$ has a unique irreducible subrepresentation. Denote it by π_s . Obviously π_s can be characterized as an irreducible subquotient of $\rho \rtimes \pi$ which has $\rho \otimes \pi$ for a subquotient of its Jacquet module.

Take any irreducible quotient of $\rho \rtimes \pi$. Denote it by π_q . Now the second form of Frobenius reciprocity implies that $\tilde{\rho} \otimes \pi$ is a subrepresentation of the Jacquet module of π_q . Therefore, $\rho \rtimes \pi$ has a unique irreducible quotient (and it has $\tilde{\rho} \otimes \pi$ for a subquotient of its Jacquet module). Further, π_q can be characterized as a subquotient of $\rho \rtimes \pi$ which has $\tilde{\rho} \otimes \pi$ for a subquotient of its Jacquet module).

We know $\pi \hookrightarrow \tau_1 \times \cdots \times \tau_\ell \rtimes \pi_{\text{cusp}}$ for some irreducible cuspidal representations τ_i 's of general linear groups. Then

$$\pi_s \hookrightarrow \rho \rtimes \pi \hookrightarrow \rho \times \tau_1 \times \cdots \times \tau_\ell \rtimes \sigma.$$

The conditions of the lemma imply, after a series of isomorphisms, $\rho \times \tau_1 \times \cdots \times \tau_\ell \rtimes \sigma \cong \tilde{\rho} \times \tau_1 \times \cdots \times \tau_\ell \rtimes \sigma$. Thus, $\tilde{\rho} \otimes \tau_1 \otimes \cdots \otimes \tau_\ell \otimes \sigma$ is in the Jacquet module of π_s . By transitivity of Jacquet modules, it must come from some subquotient in $s_{(p)}(\pi_s)$, which must be of the form $\tilde{\rho} \otimes \alpha$. But a comment at the beginning of proof implies that $\alpha = \pi$, which implies $\pi_s = \pi_q$. Therefore $\pi_s = \pi_q = \rho \rtimes \pi$. Thus, we have proved irreducibility in this case. ■

The above lemma (and the proposition bellow) could be proved by the methods of [7]. (We expect that the more sophisticated methods of [7] will enable the proof of the lemma without assuming (1).)

For a set X of smooth representations, we denote $\tilde{X} = \{\tilde{\pi}; \pi \in X\}$.

Proposition 3.2 *Let π be an irreducible representation of a classical group S_q .*

(i) *Let X be a set of irreducible cuspidal representations of general linear groups which satisfy the following.*

- (a) $\nu^{\pm 1} \rho \notin \tilde{X}$, for any $\rho \in X$.
- (b) $X \cap \tilde{X} = \emptyset$.
- (c) *There is no element in $X \cup \tilde{X}$ which is a factor of π .*
- (d) $\rho \rtimes \pi_{\text{cusp}}$ is irreducible for any $\rho \in X$.
- (e) $\rho \times \rho'$ and $\tilde{\rho} \times \rho'$ are irreducible for any $\rho \in X$ and any factor ρ' of π .

Suppose that θ is an irreducible representation of a general linear group whose cuspidal support is contained in X . Then $\theta \rtimes \pi$ is irreducible.

(ii) *Suppose that we can find sets X and Y of (equivalence classes of) irreducible cuspidal representations of general linear groups such that $X \cup \tilde{X} \cup Y \cup \tilde{Y}$ contains all the factors of π , $X \cap (Y \cup \tilde{Y}) = \emptyset$, and that hold conditions (a), (b), and (d) from (i). Further suppose that $\rho \times \rho'$ and $\tilde{\rho} \times \rho'$ are irreducible for all $\rho \in X \cup \tilde{X}$ and $\rho' \in Y$ (i.e., that holds condition (e) from (i) for all $\rho \in X \cup \tilde{X}$ and ρ' in Y).*

Then there exists an irreducible representation θ of a general linear whose cuspidal support is contained in X (i.e., each representation of the support), and there exists an irreducible representation π' of a classical group whose factors are all contained in $Y \cup \tilde{Y}$, such that $\pi \cong \theta \rtimes \pi'$. The partial cuspidal support of π' is π_{cusp} . Further, π determines θ and π' as above up to equivalence.

If X is a subset of the set of all the factors of π , then each representation from X shows up in the cuspidal support of θ .

Proof (i) Let π, X , and θ satisfy the conditions of (i). Note that Lemma 3.1 implies that $\varphi \rtimes \pi$ is irreducible for any $\varphi \in X$.

Recall that

$$\mu^*(\theta \rtimes \pi) = M^*(\theta) \rtimes \mu^*(\pi).$$

Suppose that an irreducible representation $\theta \otimes \tau$ (resp. $\tilde{\theta} \otimes \tau$) is a subquotient of $\mu^*(\theta \rtimes \pi)$. Since there does not exist a factor ρ of π such that ρ or $\tilde{\rho}$ is in the cuspidal

support of θ (recall that no element of $X \cup \tilde{X}$ is a factor of π by (c) and θ is supported by X), the above formula implies that we can get $\theta \otimes \tau$ (resp. $\tilde{\theta} \otimes \tau$) only from $M^*(\theta) \rtimes (1 \otimes \pi)$. Considering the grading, we get that $\theta \otimes \tau$ (resp. $\tilde{\theta} \otimes \tau$) can come only from $M_{GL}^*(\theta) \rtimes (1 \otimes \pi)$. Now the formula for $M_{GL}^*(\theta)$ from the previous section, the condition that θ is supported by X , and $X \cap \tilde{X} = \emptyset$ imply that $\theta \otimes \tau$ (resp. $\tilde{\theta} \otimes \tau$) can come only from $(\theta \otimes 1) \rtimes (1 \otimes \pi) = \theta \otimes \pi$ (resp. $(\tilde{\theta} \otimes 1) \rtimes (1 \otimes \pi) = \tilde{\theta} \otimes \pi$).

Thus if $\theta \otimes \tau$ (resp. $\tilde{\theta} \otimes \tau$) is a subquotient of $\mu^*(\theta \rtimes \pi)$, then it must be $\theta \otimes \pi$ (resp. $\tilde{\theta} \otimes \pi$). Further, the multiplicity of $\theta \otimes \pi$ (resp. $\tilde{\theta} \otimes \pi$) in $\mu^*(\pi)$ is 1. This implies that $\theta \rtimes \pi$ has a unique irreducible subrepresentation, which we shall denote by π_s . We can characterize π_s as a unique irreducible subquotient of $\theta \rtimes \pi$ which has $\theta \otimes \pi$ for a subquotient of appropriate Jacquet module. Further, $\theta \rtimes \pi$ has unique irreducible quotient, which we denote by π_q . Similarly as before, π_q can be characterized as a unique irreducible subquotient of $\theta \rtimes \pi$ which has $\tilde{\theta} \otimes \pi$ for a subquotient of a Jacquet module.

Actually, by the argument from the beginning of the proof, we can get more than we stated above. We can conclude the following: suppose θ is a representation of $GL(p, F)$ and $\varphi \otimes \tau$ is an irreducible subquotient of $s_{(p)}(\theta \rtimes \pi)$ such that the cuspidal support of φ is contained in X (resp. \tilde{X}), then $\varphi \otimes \tau = \theta \otimes \pi$ (resp. $\varphi \otimes \tau = \tilde{\theta} \otimes \pi$).

Let $\varphi \otimes \tau$ be an irreducible subquotient of $s_{(p)}(\pi_s)$, such that φ is supported in $X \cup \tilde{X}$, with maximal possible number of representations in $\text{supp}(\varphi)$ (counted with multiplicities), which are in \tilde{X} (note that there is at least one element that satisfies this: $\theta \otimes \pi$). If there are no elements of $\text{supp}(\varphi)$ which are in X , then they are all in \tilde{X} , and thus $\varphi \otimes \tau = \tilde{\theta} \otimes \pi$ by the remark in the above paragraph. Thus $\pi_s = \pi_q$ by the above characterization of π_q , which implies that $\theta \rtimes \pi$ is irreducible.

Therefore it remains to consider the case where at least one element of $\text{supp}(\varphi)$ is in X . Then we can find an irreducible quotient $\varphi_1 \otimes \tau'$ of $s_{(p)}(\pi_s)$, such that the cuspidal support of φ_1 is the same as of φ (this follows easily using decomposition determined by infinitesimal characters). By Frobenius reciprocity, $\pi_s \hookrightarrow \varphi_1 \rtimes \tau'$. Write $\varphi_1 = \varphi_- \times \varphi_+$, where the cuspidal support of φ_+ (resp. φ_-) is contained in X (resp. \tilde{X}). We can do this, according to assumptions of the lemma (use assumption (a)).

We know that $\varphi_+ \hookrightarrow \tau_1 \times \cdots \times \tau_{\ell-1} \times \tau_\ell$, for some $\tau_i \in X$ and $\ell \geq 1$. This implies that we have an embedding $\pi_s \hookrightarrow \varphi_- \times \tau_1 \times \cdots \times \tau_\ell \rtimes \tau'$. Note that $\tau_\ell \rtimes \tau'$ is irreducible by Lemma 3.1 (observe that τ_ℓ and τ' satisfy the conditions of Lemma 3.1). Therefore $\tau_1 \times \cdots \times \tau_{\ell-1} \times \tau_\ell \rtimes \tau' \cong \tau_1 \times \cdots \times \tau_{\ell-1} \times \tilde{\tau}_\ell \rtimes \tau'$.

Thus $\pi_s \hookrightarrow \varphi_- \times \tau_1 \times \cdots \times \tau_{\ell-1} \times \tilde{\tau}_\ell \rtimes \tau'$, which implies (using Frobenius reciprocity) that $\varphi_- \otimes \tau_1 \otimes \cdots \otimes \tau_{\ell-1} \otimes \tilde{\tau}_\ell \otimes \tau'$ is a quotient of the appropriate Jacquet module of π_s . By the transitivity of Jacquet modules, we must be able to get it from an irreducible subquotient of $s_{(p)}(\pi_s)$, say from $\varphi_2 \otimes \tau''$. Note that the support of φ_2 is again contained in $X \cup \tilde{X}$, and further, that the cuspidal support of φ_2 has one representation more in \tilde{X} than the cuspidal support of φ_1 , and therefore than the cuspidal support of φ . This is a contradiction, which completes the proof of (i).

(ii) Suppose that π , X , and Y satisfy the conditions of (ii). The representation $s_{GL}(\pi)$ has some irreducible quotient. Denote such a quotient by $\pi_1 \otimes \pi_{\text{cusp}}$. Now the cuspidal support of π_1 is contained in $X \cup \tilde{X} \cup Y \cup \tilde{Y}$. Conditions on X and Y imply

that there exist irreducible representations π_X and π_Y such that $\pi = \pi_X \times \pi_Y$ and the cuspidal support of π_X is contained in $X \cup \tilde{X}$ and of π_Y in $Y \cup \tilde{Y}$. Now Frobenius reciprocity implies that π is a subrepresentation of $\pi_X \times \pi_Y \rtimes \pi_{\text{cusp}}$. Therefore, there exists an irreducible subquotient σ_Y of $\pi_Y \times \pi_{\text{cusp}}$ such that π is subquotient of $\pi_X \rtimes \sigma_Y$. Clearly, all the factors of σ_Y are contained in $Y \cup \tilde{Y}$.

We can write $\pi_X = \pi_X^+ \times \pi_X^-$ in a such way that cuspidal supports of π_X^+ and π_X^- are contained in X and \tilde{X} respectively (use conditions (a) and (b)). Since $\pi_X^- \rtimes \sigma_Y$ and $(\pi_X^-)^\sim \rtimes \sigma_Y$ have the same composition series, π is a subquotient of $\pi_X^+ \times (\pi_X^-)^\sim \rtimes \sigma_Y$. Now there exists an irreducible subquotient π'_X of $\pi_X^+ \times (\pi_X^-)^\sim$ such that π is a subquotient of $\pi'_X \rtimes \sigma_Y$. Clearly, the support of π'_X is contained in X . Now from (i) we get that $\pi'_X \rtimes \sigma_Y$ is irreducible. Thus $\pi \cong \pi'_X \rtimes \sigma_Y$. This proves the existence of decomposition in (ii).

Suppose that $\theta' \rtimes \pi'$ and $\theta'' \rtimes \pi''$ are two decompositions of π considered in (ii). Recall that if π is parabolically induced by two irreducible cuspidal representations, then they must be conjugated by the Weyl group. The action of the Weyl group and conditions on X and Y imply that θ' and θ'' are representations of the same general linear group. Denote it by $GL(l, F)$. Moreover, the representations θ and θ' must have the same cuspidal support. Frobenius reciprocity implies that $\theta'' \otimes \pi''$ is a subquotient $s_{(l)}(\pi) = s_{(l)}(\theta' \rtimes \pi')$. Now the first part of the proof of (i) implies that $\theta' \otimes \pi' \cong \theta'' \otimes \pi''$, which implies $\theta' \cong \theta''$ and $\pi' \cong \pi''$. ■

4 Some Consequences

Bernstein defined rigid representations of general linear groups in [2]. We shall recall his definition, and extend it to the classical groups.

Definitions 4.1 (i) Let τ be an irreducible representation of a general linear group. Then τ is called *rigid* if each ρ in the cuspidal support of τ satisfies $e(\rho) \in (1/2)\mathbb{Z}$. Further, τ is called *strongly rigid*, if it is rigid, and if it satisfies the following condition: if ρ, ρ' are in the cuspidal support of τ and $\rho^u \cong (\rho')^u$, then $e(\rho) - e(\rho') \in \mathbb{Z}$.

(ii) An irreducible representation π of S_n will be called *weakly real* if $\tilde{\rho}^u \cong \rho^u$ for any factor ρ of π . It will be called *rigid* if it is weakly real and if $e(\rho) \in (1/2)\mathbb{Z}$ for any factor ρ of π . Further, it will be called *strongly rigid* if it is weakly real and $e(\rho) + \alpha_{\rho^u, \pi_{\text{cusp}}}$ and $e(\rho) - \alpha_{\rho^u, \pi_{\text{cusp}}} \in \mathbb{Z}$ for any factor ρ of π .

Note that if π is strongly rigid, it is rigid. Further, ρ^u and π_{cusp} satisfy (2.1) for any factor ρ of a strongly rigid representation π .

The Hermitian contragredient of a representation π will be denoted by π^+ (i.e., π^+ is the complex conjugate of $\tilde{\pi}$). Representation π is called Hermitian if $\pi \cong \pi^+$. Each admissible unitarizable representation is Hermitian.

From the Langlands classification (see the following section) and the formula for the contragredient in it (see [18]) it follows that each weakly real representation is Hermitian.

Denote by \mathcal{C} the set of all the equivalence classes of irreducible cuspidal representations of general linear groups $GL(n, F)$, $n \geq 1$. The subset of the unitarizable classes

is denoted by \mathcal{C}_u .

Theorem 4.2 *Let π be an irreducible representation of S_n .*

- (i) *Suppose that π is unitarizable. Then there exists an irreducible unitarizable representation θ of a general linear group and a weakly real irreducible unitarizable representation π' of some $S_{n'}$, such that $\pi \cong \theta \rtimes \pi'$.*
- (ii) *Let \mathcal{C}'_u be a subset of \mathcal{C}_u satisfying $\mathcal{C}'_u \cap \widehat{\mathcal{C}'_u} = \emptyset$ such that $\mathcal{C}'_u \cup \widehat{\mathcal{C}'_u}$ contains all $\rho \in \mathcal{C}_u$ which are not self dual. Denote*

$$\mathcal{C}' = \{\nu^\alpha \rho; \alpha \in \mathbb{R}, \rho \in \mathcal{C}'_u\}.$$

Then there exists an irreducible representation θ of a general linear group with support contained in \mathcal{C}' , and a weakly real irreducible representation π' of some $S_{n'}$ such that

$$\pi \cong \theta \rtimes \pi'.$$

Moreover, π determines such θ and π' up to an equivalence. Further, π is unitarizable (resp. Hermitian) if and only both θ and π' are unitarizable (resp. Hermitian).

Proof Clearly, one can find a set \mathcal{C}'_u which satisfies the conditions in (ii). Obviously, (ii) implies (i). Therefore, it is enough to prove (ii).

Denote by X the set \mathcal{C}' from (ii) and let Y_u be the set of all self dual representations in \mathcal{C} . Let $Y = \{\nu^\alpha \rho; \alpha \in \mathbb{R}, \rho \in Y_u\}$. Obviously, X and Y satisfy (ii) of Proposition 3.2. Applying this proposition, we get the decomposition of π and the unicity of the decomposition.

Obviously, if θ and π' are unitarizable (resp. Hermitian), then π is unitarizable (resp. Hermitian).

Suppose that $\pi = \theta \rtimes \pi'$ is Hermitian. Then $\theta \rtimes \pi' \cong \theta^+ \rtimes (\pi')^+$. Now Frobenius reciprocity implies that $\theta^+ \otimes (\pi')^+$ is a subquotient of a Jacquet module of $\pi = \theta \rtimes \pi'$. Note that θ^+ is supported again in X . Similarly as in the proof of unicity in (ii) it follows that cuspidal supports of θ and θ^+ must be the same. Now the first part of the proof of (i) of Proposition 3.2 implies $\theta^+ \otimes (\pi')^+ \cong \theta \otimes \pi'$, which implies $\theta^+ \cong \theta$ and $(\pi')^+ \cong \pi'$, i.e., θ and π' are Hermitian.

Suppose now that π is unitarizable. Then it is Hermitian. The above part of the proof implies that $\theta \otimes \pi'$ is Hermitian. Now we know that $\theta \otimes \pi'$ is unitarizable (see for example construction (d) in the third section of [19]). This implies that θ and π' are unitarizable. The proof of (ii) is now complete. ■

Since we have a classification of irreducible unitarizable representations of general linear groups, the above theorem reduces the classification of irreducible unitarizable representations of groups S_n to the classification of weakly real irreducible unitarizable representations of groups S_n .

Let $Z \subseteq \mathcal{C}_u$. Denote by $\text{Irr}_Z^{\mathbb{R}}$ the set of all equivalence classes of irreducible representations of general linear groups supported in $\{\nu^\alpha \rho; \alpha \in \mathbb{R}\}$. For $Z \subseteq \mathcal{C}_u$ and an irreducible cuspidal representation σ of S_m , let $\mathcal{J}(Z; \sigma)$ denote the set of all equivalence classes of irreducible representations π of groups S_n 's for which we can find $\alpha_1, \dots, \alpha_k \in \mathbb{R}$ and $\rho_1, \dots, \rho_k \in Z$ such that π is isomorphic to a subquotient of $\nu^{\alpha_1} \rho_1 \times \dots \times \nu^{\alpha_k} \rho_k \rtimes \sigma$. We shall point out two interesting consequences of the above theorem.

Corollary 4.3 *Let σ be an irreducible cuspidal representation of S_m .*

- (i) *Let \mathcal{C}'_u be as in Theorem 4.2(ii). Then $\theta \mapsto \theta \rtimes \sigma$ defines a bijection of $\text{Irr}^{\mathbb{R}}_{\mathcal{C}'_u}$ onto $\mathcal{J}(\mathcal{C}'_u; \sigma)$, and θ is unitarizable if and only if $\theta \rtimes \sigma$ is unitarizable.*
- (ii) *Let $\rho \in \mathcal{C}_u$, $\rho \not\cong \tilde{\rho}$. Then $\theta \mapsto \theta \rtimes \sigma$ defines a bijection of $\text{Irr}^{\mathbb{R}}_{\rho}$ onto $\mathcal{J}(\rho; \sigma)$. Moreover, θ is unitarizable if and only if $\theta \rtimes \sigma$ is unitarizable.*

In particular, in this way we get reduction of unitarizability in $\mathcal{J}(\rho; \sigma)$ to the well known case of general linear groups.

Proof (i) Mapping $\theta \mapsto \theta \rtimes \sigma$ carries $\text{Irr}^{\mathbb{R}}_{\mathcal{C}'_u}$ to $\mathcal{J}(\mathcal{C}'_u; \sigma)$ since $\theta \rtimes \sigma$ is irreducible by (i) of Proposition 3.2. From (ii) of the same proposition follows that it is surjective. To see injectivity, suppose $\theta \rtimes \sigma \cong \theta' \rtimes \sigma$. Then $\theta \otimes \sigma$ is a subquotient of the Jacquet module of $\theta' \rtimes \sigma$. Now the first part of the proof of (i) of Proposition 3.2 implies $\theta \cong \theta'$. The unitarizability claim follows in the same way as in the proof of the above theorem (first one proves that θ is Hermitian if and only if $\theta \rtimes \sigma$ is Hermitian; note that cuspidal σ is always unitarizable and Hermitian).

One proves (ii) in the same way. ■

Remarks 4.4 Take $\rho_1, \dots, \rho_k \in \mathcal{C}_u$ such that sets $\{\rho_i, \tilde{\rho}_i\}$, $1 \leq i \leq k$, have two elements each, and that these sets are different.

(i) Using arguments from the proof of Theorem 4.2 (and basic properties of the representation theory of general linear groups), one easily gets that $(\theta_1, \dots, \theta_k) \mapsto \theta_1 \times \dots \times \theta_k \rtimes \sigma$ defines a bijection of $\text{Irr}^{\mathbb{R}}_{\rho_1} \times \dots \times \text{Irr}^{\mathbb{R}}_{\rho_k}$ onto $\mathcal{J}(\rho_1, \dots, \rho_k; \sigma)$, and that this bijection preserves the unitarizability in both directions.

(ii) By Corollary 4.3(ii), $(\theta_1, \dots, \theta_k) \mapsto (\theta_1 \rtimes \sigma, \dots, \theta_k \rtimes \sigma)$ defines a bijection of $\text{Irr}^{\mathbb{R}}_{\rho_1} \times \dots \times \text{Irr}^{\mathbb{R}}_{\rho_k}$ onto $\mathcal{J}(\rho_1; \sigma) \times \dots \times \mathcal{J}(\rho_k; \sigma)$, which preserves the unitarizability (in both directions). Combining this bijection with the bijection from (i), we get a bijection

$$(\pi_1, \dots, \pi_k) \mapsto \pi,$$

$$I(\rho_1; \sigma) \times \dots \times \mathcal{J}(\rho_k; \sigma) \rightarrow I(\rho_1, \dots, \rho_k; \sigma).$$

This bijection satisfies π is unitarizable if and only if π_i are unitarizable for all $1 \leq i \leq k$.

(iii) One sees easily that the bijection in (i) depends on the fact that ρ_i 's are not self dual. It is hard to expect that such a bijection can be established in the self dual case, in particular, in a way that the unitarizability is preserved in both directions (because if ρ is self dual, we have much more unitarizable representations in $\mathcal{J}(\rho; \sigma)$ than in $\text{Irr}^{\mathbb{R}}_{\rho}$). From the other side, the bijection in (ii) can be extended to the general case. Jantzen has defined such a bijection in the general case in [7] (without assuming that ρ_i are not self dual). We expect that this bijection in general preserves the unitarizability. This would reduce the problem of unitarizability to the same problem in "lines" $I(\rho; \sigma)$. Recall that (ii) of the above corollary solves the problem of unitarizability in the (easy) case of a line $I(\rho; \sigma)$, when $\rho \not\cong \tilde{\rho}$.

The following proposition is useful for constructing complementary series.

Lemma 4.5 *Let τ and π be irreducible representations of a general linear group and S_m respectively. Suppose that τ is strongly rigid and that (2.1) holds. Let $\rho^u \cong \tilde{\rho}^u$ for each ρ in the cuspidal support of τ . If π is rigid, then representations $\nu^\alpha \tau \rtimes \pi$, $\alpha \in \mathbb{R} \setminus (1/2)\mathbb{Z}$, are irreducible.*

Under the assumptions of the above lemma, using [9] we can improve the above result if $\tau \rtimes \pi$ contains a strongly rigid subquotient (but we need to assume that (BA) from [9] holds).

Proof Denote by X_0 the set of all representations which are in the support of τ . Fix α in $\mathbb{R} \setminus (1/2)\mathbb{Z}$. Let $X = \{\nu^\alpha \rho; \rho \in X_0\}$. Now one directly checks that the conditions (a)–(e) of Proposition 3.2(i) are satisfied (use that π is rigid, τ strongly rigid and that we assume (2.1)). Proposition 3.2 implies the irreducibility. ■

Theorem 4.2 reduces the unitarizability problem to the weakly real case. The following proposition provides an upper bound for unitarizability in the weakly real case.

Proposition 4.6 *Let π be an irreducible unitarizable representation of a classical group S_q . Let ρ be a factor of π such that $\rho^u \cong \tilde{\rho}^u$. Suppose that ρ_1, \dots, ρ_n are all the factors τ of π such that $\tau^u \cong \rho^u$. Write*

$$\{ |e(\rho_i)|; |e(\rho_i)| > \alpha_{\rho^u, \pi_{\text{cusp}}} \} = \{ \alpha_1, \dots, \alpha_\ell \},$$

where $\ell \geq 0$ and $\alpha_1 < \alpha_2 < \dots < \alpha_\ell$. Then

- (i) $\alpha_\ell - \alpha_{\rho^u, \pi_{\text{cusp}}} \leq 1$ if $\ell \geq 1$, and $\alpha_i - \alpha_{i-1} \leq 1$ for each $i = 2, 3, \dots, \ell$.
- (ii) $\alpha_i \leq \alpha_{\rho^u, \pi_{\text{cusp}}} + i$; $i = 1, \dots, \ell$.

Proof Obviously, it is enough to prove (i). After modifying enumeration, we can assume $|e(\rho_1)| \leq \dots \leq |e(\rho_n)|$. Suppose that we have a gap greater than 1. It must be before some $|e(\rho_i)|$. Fix one such i and take minimal i_0 such that $|e(\rho_{i_0})| = |e(\rho_i)|$. By Proposition 3.2, we can write $\pi = \tau \rtimes \pi'$, where the cuspidal support of τ is contained in $\{\rho_{i_0}^\uparrow, \dots, \rho_n^\uparrow\} \cup \{(\rho_{i_0}^\uparrow)^\sim, \dots, (\rho_n^\uparrow)^\sim\} = \{\rho_{i_0}, \dots, \rho_n\} \cup \{\tilde{\rho}_{i_0}, \dots, \tilde{\rho}_n\}$ and none of these representations or their contragredients is a factor of π' . Moreover, if $\nu^\alpha \rho_u$ is a factor of π' , then $|\alpha| < |e(\rho_i)| - 1$ (this follows from the existence of the gap).

Note that by the same proposition, $\nu^\alpha \tau \rtimes \pi'$ is irreducible for $\alpha \geq 0$.

From the unitarizability of π it follows that π' is Hermitian (use the formula for the contragredient in the Langlands classification from [18]). Now representations $\nu^\alpha \tau \rtimes \pi'$ form a continuous family of Hermitian representations, with one unitarizable representation in it (see [19] for the definition of such a family). Then all the representations in the family are unitarizable (see construction (b) from the third section of [19]). This is impossible since this family is not bounded (see [16]). ■

5 Langlands Parameter of a Subquotient

Denote by D the set of all classes of essentially square integrable representations of all $GL(n, F)$'s, $n \geq 1$. The set of all finite multisets in D is denoted by $M(D)$. Let $d = (\delta_1, \dots, \delta_l) \in M(D)$. Take a permutation p of $\{1, \dots, l\}$ such that

$$e(\delta_{p(1)}) \geq \dots \geq e(\delta_{p(l)}).$$

If $p = id$ satisfies this condition, then we say that $d = (\delta_1, \dots, \delta_l)$ is written in standard order. The representation $\lambda(d) = \delta_{p(1)} \times \dots \times \delta_{p(l)}$ has a unique irreducible quotient, which will be denoted by $L(d)$. This is the Langlands classification of irreducible representations of general linear groups. The multiplicity of $L(d)$ in $\delta_{p(1)} \times \dots \times \delta_{p(l)}$ (and in $\delta_1 \times \dots \times \delta_l$) is 1.

Let $\delta_1, \delta_2 \in D$. Suppose that $\delta_1 \times \delta_2$ reduces. Then it is a multiplicity 1 representation of length 2. One subquotient is $L(\delta_1, \delta_2)$ while the other is of the form

$$\delta_{1,2}^{\cup} \times \delta_{1,2}^{\cap}$$

for unique representations $\delta_{1,2}^{\cup}, \delta_{1,2}^{\cap} \in D$ of $GL(n_1)$ and $GL(n_2, F)$ respectively, such that $n_1 > n_2$ (see [23] for details). In the case of reducibility we have

$$(5.1) \quad \min\{e(\delta_1), e(\delta_2)\} < e(\delta_{1,2}^{\cup}), e(\delta_{1,2}^{\cap}) < \max\{e(\delta_1), e(\delta_2)\}.$$

We shall recall a well-known property of the Langlands classification (which follows from the factorization of so called long intertwining operator in the Langlands classification). If π is an irreducible subquotient of $\delta_{p(1)} \times \dots \times \delta_{p(l)}$ (or $\delta_1 \times \dots \times \delta_l$) different from $L(d)$, then there exists $1 \leq i < j \leq l$ such that $\delta_i \times \delta_j$ reduces and π is a subquotient of

$$(5.2) \quad \delta_1 \times \delta_2 \times \dots \times \delta_{i-1} \times \delta_{i,j}^{\cup} \times \delta_{i+1} \times \dots \times \delta_{j-1} \times \delta_{i,j}^{\cap} \times \delta_{j+1} \times \dots \times \delta_{l-1} \times \delta_l.$$

We can write

$$L(\delta_1, \dots, \delta_l) = \delta_1 \times \dots \times \delta_l + \sum_{c \in X_d} m_{d,c} L(c)$$

in the Grothendieck group, where X_d is a finite subset of $M(D)$ such that $m_{d,c} \neq 0$ for all $c \in X_d$ (these conditions uniquely determine X_d ; note that $m_{d,c}$ are negative). Clearly $d \notin X_d$. From the above property it follows that if $c \in X_d$, then $L(c)$ is a subquotient of (5.2) for some i and j as above.

Now we shall recall the Langlands classification for groups S_k . Let

$$D_+ = \{\delta \in D; e(\delta) > 0\}.$$

Denote by $T(S)$ the set of all classes of irreducible tempered representations of groups S_k , $k \geq 0$. We shall write an element $t = ((\gamma_1, \dots, \gamma_m), \tau) \in M(D_+) \times T(S)$ simply as $(\gamma_1, \dots, \gamma_m; \tau)$. Take a permutation p of $\{1, \dots, m\}$ such that $e(\gamma_{p(1)}) \geq \dots \geq$

$e(\gamma_{p(m)})$ (if $p = id$ satisfies this condition, then we say that $t = (\gamma_1, \dots, \gamma_m; \tau)$ is written in standard order). Then the representation

$$(5.3) \quad \gamma_{p(1)} \times \gamma_{p(2)} \times \dots \times \gamma_{p(m)} \rtimes \tau$$

has a unique irreducible quotient, which is denoted by $L(t)$. This is the Langlands classification for groups S_k 's. The multiplicity of $L(t)$ in (5.3) is 1.

Now we shall recall a property of irreducible subquotients of (5.3), similar to the property that we have described in the case of general linear groups. We can write

$$L(\gamma_1, \dots, \gamma_m; \tau) = \gamma_1 \times \dots \times \gamma_m \rtimes \tau + \sum_{s \in Y_t} n_{t,s} L(s)$$

in the Grothendieck group, where $n_{t,s} \neq 0$ for all s from a finite subset $Y_t \subseteq M(D_+) \times T(S)$ (actually, $n_{t,s}$ are negative). Again $t \notin Y_t$. Further, for each $L(s)$, $s \in Y_t$ at least one of the following two claims hold:

Claim 5.1 (i) There exist $1 \leq i < j \leq m$ and $\varepsilon_i \in \{\pm 1\}$ such that $(\gamma_i)^{\varepsilon_i} \times \gamma_j$ reduces and that $L(s)$ is subquotient of

$$\gamma_1 \times \dots \times \gamma_{i-1} \times (\gamma_{i,j,\varepsilon_i})^\cap \times \gamma_{i+1} \times \dots \times \gamma_{j-1} \times (\gamma_{i,j,\varepsilon_i})^\cup \times \gamma_{j+1} \times \dots \times \gamma_m \rtimes \tau,$$

where $(\gamma_i)^{\varepsilon_i}$ denotes γ_i if $\varepsilon_i = 1$ and $(\gamma_i)^{\varepsilon_i}$ denotes $\tilde{\gamma}_i$ if $\varepsilon_i = -1$, and $(\gamma_{i,j,\varepsilon_i})^\cap \times (\gamma_{i,j,\varepsilon_i})^\cup$ is an irreducible subquotient of $(\gamma_i)^{\varepsilon_i} \times \gamma_j$ different from $L((\gamma_i)^{\varepsilon_i}, \gamma_j)$.

(ii) There exists $1 \leq i \leq m$ such that $\gamma_i \rtimes \tau$ reduces and if we write $\gamma_i \rtimes \tau = L(\gamma_i; \tau) + K(\gamma_i; \tau)$ in the Grothendieck group, then $L(s)$ is subquotient of

$$\gamma_1 \times \dots \times \gamma_{i-1} \times \gamma_{i+1} \times \dots \times \gamma_m \rtimes K(\gamma_i; \tau).$$

We could describe subquotients in Claim 5.1(i) by two ε_i and ε_j from $\{\pm 1\}$, but the subquotients that we would get in the remaining cases are already covered by Claim 5.1.

We shall now recall an inequality which must satisfy $\pi = L(s)$ if it is an irreducible subquotient of (5.3) ($s \in M(D_+) \times T(S)$). Suppose that (5.3) is a representation of S_k . First we introduce a partial order on \mathbb{R}^k defined by

$$\begin{aligned} (x_1, \dots, x_k) \leq (y_1, \dots, y_k) \iff & \quad x_1 \leq y_1, \\ & \quad x_1 + x_2 \leq y_1 + y_2, \\ & \quad \vdots \\ & \quad x_1 + \dots + x_k \leq y_1 + \dots + y_k. \end{aligned}$$

Now we shall define $\mathbf{e}(t)$ (analogously one defines $\mathbf{e}(s)$). Suppose that $\gamma_{p(i)}$ are representations of $GL(k_i, F)$, and τ is a representation of S_l . For p is as in (5.3), set

$$\mathbf{e}(t) = (\underbrace{e(\gamma_{p(1)}), \dots, e(\gamma_{p(1)})}_{k_1 \text{ times}}, \dots, \underbrace{e(\gamma_{p(m)}), \dots, e(\gamma_{p(m)})}_{k_m \text{ times}}, \underbrace{0, 0, \dots, 0}_{l \text{ times}}).$$

If $L(s)$ is a subquotient of $\gamma_1 \times \cdots \times \gamma_m \rtimes \tau$ (t is $(\gamma_1, \dots, \gamma_m; \tau)$) such that $s \neq t$, then

$$(5.4) \quad \mathbf{e}(s) < \mathbf{e}(t).$$

An irreducible representation π of S_k is equivalent to $L(s)$, for some $s \in M(D_+) \times T(S)$. We define $\mathbf{e}(\pi)$ to be $\mathbf{e}(s)$.

Remark 5.2 If π is a subquotient of some of the representations in Claim 5.1, then $\mathbf{e}(\pi) < \mathbf{e}(t)$.

More details and explanation regarding the above properties can be found in [18]. For $\delta \in D$ which is not unitarizable, denote

$$\delta^\uparrow = \begin{cases} \delta & \text{if } e(\delta) > 0, \\ \tilde{\delta} & \text{if } e(\delta) < 0. \end{cases}$$

Let $d \in M(D)$. By d^\uparrow we shall denote the element of $M(D_+)$ which we get from d by removing all the unitarizable δ_i 's, and changing all δ_i 's for which $e(\delta_i) < 0$, by $\tilde{\delta}_i = \delta_i^\uparrow$. Denote by d_u the multiset in $M(D)$ which we get from d removing all δ_i 's which are not unitarizable.

We add multisets in an obvious way:

$$(x_1, \dots, x_n) + (y_1, \dots, y_m) = (x_1, \dots, x_n, y_1, \dots, y_m).$$

Proposition 5.3 Let $d \in M(D)$ and $t = (d', \tau) \in M(D_+) \times T(S)$. Denote by $T_{d,\tau}$ the set of all (equivalence classes of) irreducible subrepresentations of $\lambda(d_u) \rtimes \tau$. Then:

- (i) Each of the representations $L(d^\uparrow + d'; \tau')$ with $\tau' \in T_{d,\tau}$ is a subquotient of $L(d) \rtimes L(d'; \tau)$.
- (ii) Suppose that F is a field of characteristic 0. Then the multiplicity of each of $L(d^\uparrow + d'; \tau')$, $\tau' \in T_{d,\tau}$, in $L(d) \rtimes L(d'; \tau)$ is 1.

Proof Take $\delta_1, \dots, \delta_l \in D$, $\gamma_1, \dots, \gamma_m \in D_+$ and $\tau \in T(S)$ such that $d = (\delta_1, \dots, \delta_l)$ and $d' = (\gamma_1, \dots, \gamma_m)$. After modifying enumeration we can assume that $e(\delta_i) \neq 0$ for $1 \leq i \leq n$ and $e(\delta_i) = 0$ for $n + 1 \leq i \leq l$, for some $0 \leq n \leq l$. Write

$$\delta_{n+1} \times \cdots \times \delta_l \rtimes \tau = \bigoplus_{\tau' \in T_{d,\tau}} k_{\tau'} \tau'$$

where $k_{\tau'}$ are positive integers.

For the proof of the first claim of the proposition, we need to prove that each representation from the set

$$(5.5) \quad \{L(\delta_1^\uparrow, \dots, \delta_n^\uparrow, \gamma_1, \dots, \gamma_m; \tau'); \tau' \in T_{d,\tau}\}$$

is a subquotient of

$$(5.6) \quad L(\delta_1, \dots, \delta_l) \rtimes L(\gamma_1', \dots, \gamma_m'; \tau).$$

For the second claim of the proposition we need to prove that the multiplicity of each of the subquotients in (5.6) is 1.

First observe that we have in the Grothendieck group

$$\begin{aligned}
 L(d) \rtimes L(t) &= L(d) \rtimes L(d'; \tau) = L(\delta_1, \dots, \delta_l) \rtimes L(\gamma_1, \dots, \gamma_m; \tau) \\
 &= \left(\delta_1 \times \dots \times \delta_l + \sum_{c \in X_d} m_{d,c} L(c) \right) \rtimes \\
 &\quad \left(\gamma_1 \times \dots \times \gamma_m \rtimes \tau + \sum_{s \in Y_t} n_{t,s} L(s) \right) \\
 &= \delta_1 \times \dots \times \delta_l \times \gamma_1 \times \dots \times \gamma_m \rtimes \tau + \sum_{s \in Y_t} n_{t,s} \delta_1 \times \dots \times \delta_l \rtimes L(s) \\
 &\quad + \sum_{c \in X_d} m_{d,c} L(c) \times \gamma_1 \times \dots \times \gamma_m \rtimes \tau + \sum_{c \in X_d} \sum_{s \in Y_t} m_{d,c} n_{t,s} L(c) \rtimes L(s) \\
 (5.7) \quad &= \sum_{\tau' \in T_{d,\tau}} k_{\tau'} \delta_1^\uparrow \times \dots \times \delta_l^\uparrow \times \gamma_1 \times \dots \times \gamma_m \rtimes \tau' \\
 (5.8) \quad &\quad + \sum_{s \in Y_t} n_{t,s} \delta_1 \times \dots \times \delta_l \rtimes L(s) \\
 (5.9) \quad &\quad + \sum_{c \in X_d} m_{d,c} L(c) \times \gamma_1 \times \dots \times \gamma_m \rtimes \tau \\
 (5.10) \quad &\quad + \sum_{c \in X_d} \sum_{s \in Y_t} m_{d,c} n_{t,s} L(c) \rtimes L(s).
 \end{aligned}$$

Now the representations in (5.5) show up as subquotients of representations in (5.7).

If F is of characteristic 0, then all $k_{\tau'}$ are one (see [5] for split groups, the case of non-split groups goes as in [6]). Now the properties of the Langlands classification imply that the representations in (5.5) show up as subquotients of representations in (5.7) with multiplicity one.

Therefore, for the proof of the proposition, it is enough to show that representations from (5.5) do not show up as subquotients of representations in the lines (5.8), (5.9), and (5.10). It is obvious that each irreducible subquotient of each representation in the line (5.10) is already a subquotient of some representation in the line (5.8) or (5.10) (actually, of both). Therefore, for the proof of the proposition it is enough to show that representations from (5.5) do not show up as subquotients of the representations in the lines (5.8) and (5.9).

Observe that for each irreducible representation π from (5.5), $e(\pi)$ does not depend on $\tau' \in T_{d,\tau}$. We shall show below that for each irreducible subquotient σ of (5.8) or (5.9), $e(\sigma)$ is strictly smaller than $e(\pi)$, where π is from (5.5). This will prove the proposition. Observe that we know $e(\sigma) \leq e(\pi)$ by (5.4).

Suppose that some irreducible representation π is a subquotient of some term in

(5.9). Then it is a subquotient of some

$$(\delta_1 \times \cdots \times \delta_{i,j}^\cap \times \cdots \times \delta_{i,j}^\cup \times \cdots \times \delta_l) \times \gamma_1 \times \cdots \times \gamma_m \rtimes \tau$$

(see (5.2)). Observe $i \leq n$ since the tempered parabolic induction for general linear groups is irreducible.

Consider first the case $j \leq n$. Then the relation (5.1) and the definition of $\mathbf{e}(\pi)$ and $\mathbf{e}(\sigma)$ imply that $\mathbf{e}(\pi) < \mathbf{e}(\sigma)$ for σ from (5.5).

Consider now $n < j$. Then π is a subquotient of

$$(\delta_1 \times \cdots \times \delta_{i,j}^\cap \times \cdots \times \delta_n \times \cdots \times \delta_{i,j}^\cup \times \cdots \times \delta_l) \times \gamma_1 \times \cdots \times \gamma_m \rtimes \tau.$$

Write

$$(5.11) \quad (\delta_1^\uparrow, \dots, \delta_n^\uparrow, \gamma_1, \dots, \gamma_m; \tau') = (\beta_1, \dots, \beta_{n+m}; \tau')$$

such that the right hand side is in standard order (here τ' is any element from $T_{d,\tau}$).

Then $\delta_i^\uparrow = \beta_k$ for some k .

Let τ'' be an irreducible subrepresentation of

$$\delta_{n+1} \times \cdots \times \delta_{j-1} \times \delta_{j+1} \times \cdots \times \delta_l \rtimes \tau.$$

If $e(\beta_k) > e(\beta_{k+1})$ or $k = n + m$, writing

$$(\delta_1^\uparrow, \dots, (\delta_{i,j}^\cap)^\uparrow, \dots, \delta_n^\uparrow, (\delta_{i,j}^\cup)^\uparrow, \gamma_1, \dots, \gamma_m; \tau'')$$

in standard order, relation (5.1) will imply directly

$$(5.12) \quad \mathbf{e}(\pi) \leq \mathbf{e}(\delta_1^\uparrow, \dots, (\delta_{i,j}^\cap)^\uparrow, \dots, \delta_n^\uparrow, (\delta_{i,j}^\cup)^\uparrow, \gamma_1, \dots, \gamma_m; \tau'') \\ < \mathbf{e}(\delta_1^\uparrow, \dots, \delta_n^\uparrow, \gamma_1, \dots, \gamma_m; \tau') = \mathbf{e}(\beta_1, \dots, \beta_{n+m}; \tau') = \mathbf{e}(\sigma)$$

for any σ from (5.5).

If $e(b_k) = e(b_{k+1})$, then look at all $\ell \geq k$ such that $e(\beta_\ell) = e(\beta_k)$. Let ℓ_0 be a maximal such index.

Write $(\delta_1^\uparrow, \dots, (\delta_{i,j}^\cap)^\uparrow, \dots, \delta_n^\uparrow, (\delta_{i,j}^\cup)^\uparrow, \gamma_1, \dots, \gamma_m; \tau'')$ in standard order, where τ'' is an irreducible subrepresentation of $\delta_{n+1} \times \cdots \times \delta_{j-1} \times \delta_{j+1} \times \cdots \times \delta_l \rtimes \tau$ as before. We shall consider how from $\mathbf{e}(\beta_1, \dots, \beta_{n+m}; \tau')$ one gets

$$\mathbf{e}(\delta_1^\uparrow, \dots, (\delta_{i,j}^\cap)^\uparrow, \dots, \delta_n^\uparrow, (\delta_{i,j}^\cup)^\uparrow, \gamma_1, \dots, \gamma_m; \tau'').$$

Note that $\mathbf{e}(\delta_1^\uparrow, \dots, (\delta_{i,j}^\cap)^\uparrow, \dots, \delta_n^\uparrow, (\delta_{i,j}^\cup)^\uparrow, \gamma_1, \dots, \gamma_m; \tau'')$ will have the same β_h 's for $h < k$. Further, one needs to remove from $\mathbf{e}(\beta_1, \dots, \beta_{n+m}; \tau')$ elements $e(\beta_k)$ corresponding to β_k and shift elements $e(\beta_\ell)$ corresponding to β_ℓ , $k + 1 \leq \ell \leq \ell_0$, to the left.

After $e(\beta_{\ell_0})$ will come either $e(\beta_{\ell_0+1})$ or $e((\delta_{i,j}^{\cup})^\dagger)$ or $e((\delta_{i,j}^\cap)^\dagger)$. Now $e(\beta_{\ell_0+1}) < e(\beta_{\ell_0})$ and (5.1) imply again (5.12), what we needed to prove.

To complete the proof, it remains to consider π which is an irreducible subquotient of (5.8). Then it is a subquotient of some $\delta_1 \times \dots \times \delta_l \rtimes L(s)$, where $L(s)$ is a subquotient of a representation which can have following two types.

The first type is a representation for which there exist $1 \leq i < j \leq m$ and $\varepsilon_i \in \{\pm 1\}$ such that $(\gamma'_i)^{\varepsilon_i} \times \gamma'_j$ reduces and that $L(s)$ is subquotient of

$$\gamma_1 \times \dots \times (\gamma_{i,j,\varepsilon_i})^\cap \times \dots \times (\gamma_{i,j,\varepsilon_i})^\cup \times \dots \times \gamma_l \rtimes \tau.$$

Again write $(\delta_1^\dagger, \dots, \delta_n^\dagger, \gamma_1, \dots, \gamma_m; \tau') = (\beta_1, \dots, \beta_{n+m}; \tau')$ in standard order (similarly as in (5.11)). Observe

$$|e((\gamma_{i,j,\varepsilon_i})^\cap)|, |e((\gamma_{i,j,\varepsilon_i})^\cup)| < \max\{e(\gamma_i), e(\gamma_j)\}.$$

Now from (5.1) and the above relation we prove (5.12) in the same way as in the previous case.

The remaining type of the representation is one for which there exists i such that $\delta'_i \rtimes \tau$ reduces, and after writing $\gamma_i \rtimes \tau = L(\gamma_i; \tau) + K(\gamma_i; \tau)$ in the Grothendieck group, then $L(s)$ is subquotient of

$$\gamma_1 \times \dots \times \gamma_{i-1} \times \gamma_{i+1} \times \dots \times \gamma_m \rtimes K(\gamma_i; \tau).$$

Therefore, there exists a subquotient π' of $K(\gamma_i; \tau)$ such that $L(s)$ is a subquotient of $\gamma_1 \times \dots \times \gamma_{i-1} \times \gamma_{i+1} \times \dots \times \gamma_m \rtimes \pi'$. We know that $\pi' = L(r)$ for some $r \in M(D_+) \times T(S)$. Write $r = (\mu_1, \dots, \mu_a; \tau'')$ in standard order. Further,

$$\mathbf{e}(\pi') = \mathbf{e}(\mu_1, \dots, \mu_a; \tau'') = (x_1, \dots, x_b) < \mathbf{e}(\gamma_i, \tau) = \underbrace{(e(\gamma_i), \dots, e(\gamma_i))}_{u \text{ times}}, 0, \dots, 0,$$

where γ_i is a representation of $GL(u, F)$. This implies

$$\begin{aligned} x_1 \geq \dots \geq x_b \geq 0; \quad x_1 &\leq e(\gamma_i); \\ x_1 + \dots + x_u &< u e(\gamma_i); \quad x_1 + \dots + x_b \leq u e(\gamma_i). \end{aligned}$$

The third relation implies that the number v of indices j such that $x_j = e(\gamma_i)$ satisfies

$$(5.13) \quad v < u.$$

We can now write the following in standard order

$$(\delta_1, \dots, \delta_n, \gamma_1, \dots, \gamma_{i-1}, \gamma_{i+1}, \dots, \gamma_m, \mu_1, \dots, \mu_a, \tau'''),$$

where τ''' is an irreducible subrepresentation of $\delta_{n+1} \times \dots \times \delta_l \rtimes \tau''$. From (5.13) one sees similarly as above that $\mathbf{e}(\pi) < \mathbf{e}(\sigma)$ for σ from (5.5). The proof is now complete ■

Remarks 5.4 (i) Proposition 5.3 holds also in the archimedean case, in unchanged form. The proof there goes the same way as in the non-archimedean case, except that we need to make small modifications regarding composition series of representations $\delta_1 \times \delta_2, \delta_i \in D$.

(ii) Proposition 5.3 also holds for hermitian quaternionic groups in the case of characteristic 0.

(iii) The second claim of Proposition 5.3 also holds in positive characteristic. This follows from the work of J.-L. Waldspurger and V. Heiermann, but for this we do not have a written reference (the result that one needs is that standard intertwining operators span the intertwining algebra of the representation induced by an irreducible square integrable one).

The easiest way of constructing irreducible unitarizable representations of reductive groups is parabolic induction starting with irreducible unitarizable representations of proper Levi subgroups. For this, we need to know when such representations reduce, and what are irreducible constituents. A natural place to start construction for classical groups S_m 's are representations $\pi \times \tau$ where π is an irreducible unitarizable representation of a general linear group (such representations are classified) and τ is an irreducible tempered representation of a classical group S_m . (We would like to start with more general unitarizable representations τ 's, but we do not know many more general irreducible unitarizable representations of classical groups.)

For a unitarizable $\delta \in D$ and a positive integer n denote

$$u(\delta, n) = L(\nu^{(n-1)/2}\delta \times \nu^{(n-1)/2-1}\delta \times \dots \times \nu^{-(n-1)/2}\delta).$$

Then $u(\delta, n)$ is a unitarizable representation of a general linear group. These representations play a crucial role in the description of the unitary duals of general linear groups (see [14]). Each irreducible unitarizable representation of a general group is induced from a tensor product of such representations or complementary series starting with such representations.

Let us return to the classical groups and representations $\pi \times \tau$ where π is an irreducible unitarizable representation of a general linear group and τ is an irreducible tempered representation of a classical groups S_m . We can handle reducibility questions of representations $\pi \times \tau$ when π 's are complementary series pretty well using Lemma 4.5 and Proposition 3.2. Therefore, we shall concentrate now to the problem of reducibility of representations $u(\delta, n) \times \tau$.

The above proposition enables us easily to get a partial information in that direction. Let us note that we have a description of reducibility points of representations $\delta' \times \tau$ for $\delta \in D$ and τ irreducible square integrable representation, and also if τ is an irreducible tempered representation (see [10, 11]).

Corollary 5.5 *Suppose that τ is an irreducible representation of a classical group S_m , $\delta \in D$ is unitarizable and n is non-negative integer.*

- (i) *Let $\delta_1, \dots, \delta_k \in D$ be unitarizable and not self dual (i.e., $\delta_i \not\cong \tilde{\delta}_i, i = 1, \dots, k$). If n_1, \dots, n_k are positive integers and τ is weakly real or tempered, then $u(\delta_1, n_1) \times \dots \times u(\delta_k, n_k) \times \tau$ is irreducible.*

- (ii) Suppose that τ is tempered and that $\delta \rtimes \tau$ reduces. Then $u(\delta, 2n + 1) \rtimes \tau$ reduces.
- (iii) Suppose that τ is tempered. Then $u(\delta, 2) \rtimes \tau$ reduces if and only if $\nu^{1/2}\delta \rtimes \tau$ reduces.
- (iv) Suppose that τ is tempered and that $\delta \rtimes \tau$ and $\nu\delta \rtimes \tau$ are irreducible. Then $u(\delta, 3) \rtimes \tau$ is irreducible.

Proof (i) If τ is weakly real, we easily get (i) from Proposition 3.2 (see the proof of Theorem 4.2). If τ is tempered, then $\tau \cong \delta'_1 \times \cdots \times \delta'_k \rtimes \tau'$, where $\delta'_i \in D$ are unitary and not self dual, and τ' is weakly real. Now the first part of the proof implies irreducibility.

(ii) Proposition 5.3 implies (ii) directly.

(iii) Note that $L(\nu^{1/2}\delta, \nu^{1/2}\delta; \tau)$ is always a subquotient of $u(\delta, 2) \rtimes \tau$ by the above proposition. Suppose that $\nu^{1/2}\delta \rtimes \tau$ reduces. Then there exists $t \in M(D_+) \times T(S)$, $t \neq (\nu^{1/2}\delta, \tau)$ such that $L(t)$ is a subquotient of $\nu^{1/2}\delta \rtimes \tau$. Now $L((\nu^{1/2}\delta) + t)$ is a subquotient of $\nu^{1/2}\delta \times \nu^{1/2}\delta \rtimes \tau$ by the above proposition. This is a non-tempered representation. From the representation theory of general linear groups we know that $\nu^{1/2}\delta \times \nu^{1/2}\delta \rtimes \tau$ consists of $u(\delta, 2) \rtimes \tau$ and a tempered part. Therefore, $L((\nu^{1/2}\delta) + t)$ is a subquotient of $u(\delta, 2) \rtimes \tau$. Since $(\nu^{1/2}\delta) + t$ is different from $(\nu^{1/2}\delta, \tau\nu^{1/2}\delta; \tau)$, we get that $u(\delta, 2) \rtimes \tau$ is reducible.

Suppose that $\nu^{1/2}\delta \rtimes \tau$ is irreducible. Now we have the surjective intertwining

$$\nu^{1/2}\delta \times \nu^{1/2}\delta \rtimes \tau \cong \nu^{1/2}\delta \times \nu^{-1/2}\delta \rtimes \tau \rightarrow u(\delta, 2) \rtimes \tau.$$

Since the first representation has a unique irreducible quotient, $u(\delta, 2) \rtimes \tau$ must also have a unique irreducible quotient. Since the last representation is unitarizable, it must be irreducible.

(iv) Using Zelevinsky involution, the fact that it carries irreducible representations to the irreducible ones, we get easily that $L(\nu\delta, \delta) \times \delta$ is irreducible. Now we have surjective intertwinings

$$\begin{aligned} \nu\delta \times \nu\delta \times \delta \rtimes \tau &\rightarrow \nu\delta \times L(\nu\delta, \delta) \rtimes \tau \cong L(\nu\delta, \delta) \times \nu\delta \rtimes \tau \\ &\cong L(\nu\delta, \delta) \times \nu^{-1}\delta \rtimes \tau \rightarrow u(\delta, 3) \rtimes \sigma. \end{aligned}$$

This implies irreducibility (use the same argument as in the proof of (iii)). ■

Remarks 5.6 (i) After the above corollary, it is natural to ask: is $u(\delta, n) \rtimes \pi$ irreducible if $\delta' \rtimes \pi$ are irreducible for all $\delta' \in \{\nu^{(n-1)/2}\delta, \nu^{(n-1)/2-1}\delta, \dots, \nu^{-(n-1)/2}\delta\}$.

(ii) After understanding reducibilities $u(\delta, n) \rtimes \tau$, the next question would be to understand representations $u(\delta_1, n_1) \times \cdots \times u(\delta_k, n_k) \rtimes \tau$. Note that the theory of R groups takes care of the case $n_1 = \cdots = n_k = 1$. In this case we get a representation of length 2^l for some l (which we can describe in terms of relative rank one reducibilities). The general situation will be different. We shall give an example.

For a connected reductive group G , 1_G (resp. St_G) will denote the trivial one dimensional (resp. Steinberg) representation of G . Now we shall consider split odd-orthogonal groups. It is not hard to see that the representation $\text{St}_{GL(2,F)} \rtimes \text{St}_{SO(3,F)}$ is

irreducible, and that $1_{GL(2,F)} \rtimes St_{SO(3,F)}$ reduces into two irreducible pieces (clearly, $1_{GL(2,F)} \times St_{GL(2,F)}$ is irreducible). Ban and Jantzen proved in [1] that

$$1_{GL(2,F)} \times St_{GL(2,F)} \rtimes St_{SO(3,F)} = u(1_{GL(1,F)}, 2) \times u(St_{GL(2,F)}, 1) \rtimes St_{SO(3,F)}$$

reduce into three irreducible pieces.

(iii) If we fix a self dual δ , we shall often have $u(\delta, n) \rtimes \tau$ reducible. For example, at least one of the representations $u(\delta, 2n + 1) \rtimes 1_{SO(1,F)}$ or $u(\delta, 2n + 1) \rtimes 1_{Sp(0,F)}$ is always reducible (this follows from the duality established by Shahidi in [13]).

6 A Simple Case of the Reducibility Problem

Let us first recall a classification of irreducible essentially square integrable representations of general linear groups (see [23]). For $\rho \in \mathcal{C}$ and a positive integer n , the set $\Delta = [\rho, \nu^n \rho] = \{\rho, \nu \rho, \dots, \nu^n \rho\}$ is called a segment in \mathcal{C} . The representation $\nu^n \rho \times \nu^{n-1} \rho \times \dots \times \rho$ contains the unique irreducible subrepresentation, which will be denoted by $\delta(\Delta)$. It is essentially square integrable. Segments in \mathcal{C} parameterize D ($\Delta \mapsto \delta(\Delta)$ is a bijection).

We have seen that for understanding the unitarizability it is very useful to understand the reducibility of parabolic induction. One of the first steps in understanding reducibility for classical p -adic groups, is to understand the reducibility of

$$(6.1) \quad \delta \rtimes \pi$$

where δ and π are irreducible essentially square integrable representations (δ of a general linear group and π of a classical group). This question is in general hard and complicated (see [10, 11]). If π is cuspidal, the problem of reducibility of (6.1) has a simple answer (reduction). If (2.1) holds (and π is cuspidal), then $\delta(\Delta) \rtimes \pi$ is reducible if and only if $\rho \rtimes \pi$ reduces for some $\rho \in \Delta$.

The opposite case to π cuspidal is the case of δ cuspidal. We shall briefly describe the reducibility in this case.

Let ρ be an irreducible unitarizable cuspidal representation of $GL(p, F)$ and let π be an irreducible square integrable representation of S_q . If ρ is not self dual, then $\nu^\alpha \rho \rtimes \pi$ is irreducible for all $\alpha \in \mathbb{R}$. Therefore, we shall assume in the sequel that ρ is self dual.

In the sequel we shall assume that assumption (BA) from [9] holds.

For a positive integer a denote $\delta(\rho, a) = \delta([\nu^{-(a-1)/2} \rho, \nu^{(a-1)/2} \rho])$. Parity of ρ with respect to π is defined in [8] (see [9] also). Parity can be characterized in the following way: it is even (resp. odd) if $\delta(\rho, a) \rtimes \pi$ is irreducible for all odd (resp. even) positive integers a . Further, $Jord(\pi)$ is defined as the set of all pairs (ρ, a) where a is a positive integer of the same parity as the parity of ρ with respect to π , such that $\delta(\rho, a) \rtimes \pi$ is irreducible (see [8], and also [9]). Further, C. Mœglin has introduced a partially defined function ε_π . Then (admissible) triples $(Jord(\pi), \varepsilon_\pi, \pi_{\text{cusp}})$ parameterize irreducible square integrable representations of groups S_n 's (see [9] for details). In the following proposition we collect the facts about reducibility of representations (6.1) when δ is cuspidal.

Proposition 6.1 *Let $\rho \in \mathcal{C}$ be self dual, and let π be an irreducible square integrable representation of a group S_q . Suppose that (BA) from [9] holds. Denote $\text{Jord}_\rho(\pi) = \{k; (\rho, k) \in \text{Jord}_\rho(\pi)\}$. Let a be a positive integer. Then:*

- (i) *For $\alpha \in \mathbb{R}$, $\nu^\alpha \rho \rtimes \pi$ reduces if and only if $\nu^{-\alpha} \rho \rtimes \pi$ reduces.*
- (ii) *If $\alpha \in \mathbb{R} \setminus (1/2)\mathbb{Z}$, then $\nu^\alpha \rho \rtimes \pi$ is irreducible.*
- (iii) *$\rho \rtimes \pi$ reduces if and only if ρ has odd parity with respect to π and $1 \notin \text{Jord}_\rho(\pi)$.*
- (iv) *If $a \notin \text{Jord}_\rho(\pi)$, then $\nu^{(a+1)/2} \rho \rtimes \pi$ is irreducible.*
- (v) *If $a \in \text{Jord}_\rho(\pi)$ and $a + 2 \notin \text{Jord}_\rho(\pi)$, then $\nu^{(a+1)/2} \rho \rtimes \pi$ is reducible.*
- (vi) *Suppose that a and $a + 2$ are in $\text{Jord}_\rho(\pi)$. Then $\nu^{(a+1)/2} \rho \rtimes \pi$ is reducible if and only if $\varepsilon_\rho(a) = \varepsilon_\rho(a + 2)$.*
- (vii) *$\nu^{1/2} \rho \rtimes \pi$ is reducible if and only if $2 \notin \text{Jord}_\rho(\pi)$ or $2 \in \text{Jord}_\rho(\pi)$ and $\varepsilon_\rho(2) = 1$. In other words, $\nu^{1/2} \rho \rtimes \pi$ is irreducible if and only if $2 \in \text{Jord}_\rho(\pi)$ and $\varepsilon_\rho(2) = -1$.*

The first claim is a consequence of the structure of R module $R(S)$ (see Section 2). The second claim is well known (it follows also from Lemma 4.5). The third claim is just a definition of $\text{Jord}(\pi)$. The fourth claim is [8, Lemma 5.4.1(iv)], while the fifth claim is [9, Lemma 5.3]. A sketch of proof of sixth claim is in the remark after the proof of the [9, Lemma 5.3]. That proof is based on intertwining operators. We have also a proof based on the Jacquet module techniques of [20]. The seventh claim also follows applying these techniques.

Muić has described reducibility points of representations (6.1) completely in [10]. (In [11] he settles the case of more general case of standard modules.) The above proposition is a very special (starting) case of his results. The above proposition is elementary in comparison with his results. The same comment is valid for the following simple lemma, which is sometimes useful. It can be proved in a rather simple way.

Lemma 6.2 *Assume that (BA) holds. Let Δ be a segment in irreducible cuspidal representations of general linear groups and let π be an irreducible square integrable representation of a group S_q . Suppose that $\rho \rtimes \pi$ is irreducible for all $\rho \in \Delta$. Then $\delta(\Delta) \rtimes \pi$ is irreducible.*

Acknowledgment We are thankful to G. Muić for discussions on the topics studied in this paper. These discussions have helped to clarify some parts of the paper.

References

- [1] D. Ban and C. Jantzen, *R-groups and the action of intertwining operators in the nontempered case*. Int. Math. Res. Not. IMRN **2007**, no. 20, Art. ID rnm059.
- [2] J. Bernstein, *P-invariant distributions on $GL(N)$ and the classification of unitary representations of $GL(N)$ (non-Archimedean case)*. In: Lie group representations II, Lecture Notes in Mathematics 1041, Springer-Verlag, Berlin, 1984, pp. 50–102.
- [3] D. Barbasch and A. Moy, *Reduction to real infinitesimal character in affine Hecke algebras*. J. Amer. Math. Soc. **6**(1993), no. 3, 611–635.
- [4] W. Casselman, *Introduction to the theory of admissible representations of p -adic reductive groups*. preprint. <http://www.math.ubc.ca/~cass/research/pdf/p-adic-book.pdf>
- [5] D. Goldberg, *Reducibility of induced representations for $Sp(2n)$ and $SO(n)$* . Amer. J. Math. **116**(1994), no. 5, 1101–1151.

- [6] M. Hanzer, *R groups for quaternionic Hermitian groups*. Glas. Mat. **39(59)**(2004), no. 1, 31–48.
- [7] C. Jantzen, *On supports of induced representations for symplectic and odd-orthogonal groups*. Amer. J. Math. **119**(1997), no. 6, 1213–1262.
- [8] C. Mœglin, *Sur la classification des séries discrètes des groupes classiques p -adiques: paramètres de Langlands et exhaustivité*. J. Eur. Math. Soc. **4**(2002), no. 2, 143–200.
- [9] C. Mœglin and M. Tadić, *Construction of discrete series for classical p -adic groups*. J. Amer. Math. Soc. **15**(2002), no. 3, 715–786.
- [10] G. Muic, *Reducibility of generalized principal series*. Canad. J. Math. **57**(2005), no. 3, 616–647.
- [11] ———, *Reducibility of standard representations*. Pacific. J. Math. **222**(2005), no. 1, 133–168.
- [12] F. Shahidi, *A proof of Langlands' conjecture on Plancherel measures; complementary series for p -adic groups*. Ann. of Math. **132**(1990), no. 2, 273–330.
- [13] ———, *Twisted endoscopy and reducibility of induced representations for p -adic groups*. Duke Math. J. **66**(1992), no. 1, 1–41.
- [14] M. Tadić, *Classification of unitary representations in irreducible representations of general linear group (non-Archimedean case)*. Ann. Sci. École Norm. Sup. **19**(1986), no. 3, 335–382.
- [15] ———, *$GL(n, \mathbb{C})$ and $GL(n, \mathbb{R})$* . To appear in a workshop in honor of S. Gelbart on Automorphic Forms and L-functions. <http://www.hazu/~tadic/42-GL-arch.pdf>.
- [16] ———, *Geometry of dual spaces of reductive groups (non-Archimedean case)*. J. Analyse Math. **51**(1988), 139–181.
- [17] ———, *Structure arising from induction and Jacquet modules of representations of classical p -adic groups*. J. Algebra **177**(1995), no. 1, 1–33.
- [18] ———, *Representations of p -adic symplectic groups*. Compositio Math. **90**(1994), no. 2, 123–181.
- [19] ———, *An external approach to unitary representations*. Bull. Amer. Math. Soc. **28**(1993), no. 2, 212–252.
- [20] ———, *On reducibility of parabolic induction*. Israel J. Math. **107**(1998), 29–91.
- [21] ———, *On regular square integrable representations of p -adic groups*. Amer. J. Math. **120**(1998), no. 1, 159–210.
- [22] ———, *On classification of some classes of irreducible representations of classical groups*. In: Representations of real and p -adic groups, Lect. Notes Ser. Inst. Math. Sci. Natl. Singap. 2, Singapore University Press and World Scientific, Singapore, 1994, pp. 95–162.
- [23] A. V. Zelevinsky, *Induced representations of reductive p -adic groups II. On irreducible representations of $GL(n)$* . Ann. Sci. École Norm. Sup. **13**(1980), no. 2, 165–210.

Department of Mathematics, University of Zagreb, Bijenička 30, 10000 Zagreb, Croatia
 e-mail: tadic@math.hr