
INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, Journal of Helminthology,
London School of Hygiene and Tropical Medicine,
Keppel Street, London, WC1E 7HT,
England.

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief introduction (unheaded); Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration may be printed is 13.5 × 18 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space, should be indicated. Some indication of scale should normally be given on the figure. Photocopies of illustrations should also be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints of the same size as they are to appear in the Journal (maximum size 13.5 × 18 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In *Ecology and physiology of parasites* (ed A. M. Fallis) pp. 213-222. Adam Hilger Ltd.: London.

JAMES, C. and WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49-59.

50 offprints are provided free of charge; additional copies may be ordered at the proof stage.

Contents

DENNIS, E. A., SHARP, M. and DOUGLASS, R. Carbohydrate reserves and phosphatase activity in the mollusc-trematode relationship of <i>Mytilus edulis</i> L. and <i>Protoeces maculatus</i> (Looss, 1901) Odhner, 1911	1
OPUNI, E. K. and MULLER, R. L. Studies on <i>Spirometra theileri</i> (Baer, 1925) n.comb. 1. Identification and biology in the laboratory	15
COPLAND, J. W. <i>Metastrongylus</i> spp. infections of village pigs in Papua New Guinea	25
HINKS, M. I. and THOMAS, R. J. A new record of the occurrence of <i>Trichuris skrjabini</i> Baskakov, 1924 in sheep in Britain	33
WALKER, J. C. and McMILLAN, B. <i>Breinlia mackerrasae</i> n.sp. (Nematoda: Filarioidea) from the peramelid marsupial, <i>Isodon macrourus</i> , with a note on <i>Breinlia</i> sp. from the 'mumut' (Peramelidae)	39
PEREGRINE, P. C. The effects of host diet on <i>Thelastoma attenuatum</i> (Nematoda: Thelastomatidae) populations in cockroaches	47
KRITSKY, D. C. and THATCHER, V. E. Monogenetic trematodes (Monopisthocotylea: Dactylogyridae) from freshwater fishes of Colombia, South America	53
SAOUD, M. F. A., ABDEL-HAMID, M. E. and IBRAHIM, A. M. On <i>Allocreadium sudanensis</i> sp.nov. (Trematoda: Digenea) from a freshwater fish in the Sudan	67
WOOTEN, R. Observations on strigeid metacercariae in the eyes of fish from Hanningfield Reservoir, Essex, England	73
ASANJI, M. F. and WILLIAMS, M. O. Studies on the excystment of trematode metacercariae <i>in vivo</i>	85
