

FARTHEST POINTS IN W^* -COMPACT SETS

R. DEVILLE AND V.E. ZIZLER

We show that while farthest points always exist in w^* -compact sets in duals to Radon-Nikodym spaces, this is generally not the case in dual Radon-Nikodym spaces. We also show how to characterise weak compactness in terms of farthest points.

The purpose of this note is to find under what conditions the Edelstein-Asplund-Lau results on the existence of farthest points in weakly compact sets can be extended to w^* -compact sets.

To fix our notation, let C be a norm closed bounded subset of a real Banach space X and x be an element of X . We define

$$r(x) = r(x, C) = \sup\{\|x - z\| \mid z \in C\}$$

and call $r(x)$ the farthest distance from x to C . Equivalently, $r(x)$ is the radius of the smallest ball of centre x , containing C . The function r is convex as supremum of such functions, and continuous since $|r(x) - r(y)| \leq \|x - y\|$, for all $x, y \in X$. A point $z \in C$ is called a farthest point of C if there exists $x \in X$ such that $\|x - z\| = r(x)$. The existence of a farthest point of C is equivalent to the fact that the set

$$D = \{x \in X \mid (\exists z \in C)(\|x - z\| = r(x))\}$$

is non-empty. Since it follows that any farthest point of a convex set C in a locally uniformly rotund space is a strongly exposed point of C , the notion of a farthest point is widely used in the study of the extreme structure of sets. In fact, it was the first discovered method of obtaining exposed points in sets [10]. We refer the reader to [3] and [5] for unexplained notions.

Extending the results of [6] and [1], Lau showed [9] that if C is a weakly compact set in a Banach space X , then the set D defined above is dense in X and thus, in particular, C has farthest points. Other extensions of the results of [6] and [1] may be found in [11]. The connection between farthest points and strongly exposed points in

Received 19 February, 1987

Research completed while Dr Deville was a visitor at the University of Alberta. Research partially supported by NSERC (Canada)

Copyright Clearance Centre, Inc. Serial-fee code: 0004-9729/88 \$A2.00+0.00.

mentioned above led Edelstein and Lewis [7] to ask the following question: Does the existence of (strongly) exposed points in C imply the existence of farthest points in C ?

Proposition 1 gives a negative answer to this question. Moreover, it shows that Lau’s result on weakly compact sets does not always extend to w^* -compact sets in a dual space X^* . However, our Proposition 3 shows that if X has the Radon-Nikodym property then any w^* -compact set in X^* contains a farthest point, thus extending Proposition 1 of [11]. Finally, we point out in Proposition 4 that Lau’s result actually characterises weakly compact sets.

PROPOSITION 1. Let $\ell^1(\mathbb{N})$ be the Banach space of all real summable sequences $x = (x_n)$, equipped with its usual norm $\|x\| = \sum_{n=1}^\infty |x_n|$. Let

$$C = \{(a_n) \in \ell^1(\mathbb{N}) : \sum_{n=1}^\infty (|a_n| + |a_n|^2) \leq 1\}.$$

Then C is a weak*-compact convex of $\ell^1(\mathbb{N})$ and C has no farthest points.

PROOF OF PROPOSITION 1: To prove that C has no farthest points, it is enough to show that, given $x \in \ell^1(\mathbb{N})$:

- i) $r(x) = 1 + \|x\|$ and
- ii) $\|x - z\| < 1 + \|x\|$ for all $z \in C$.

To do so, first notice that if $z \in C$ then $\|z\| < 1$. Therefore, $\|x - z\| < \|x\| + \|z\| < 1 + \|x\|$ and ii) is proved. This also shows that $r(x) \leq 1 + \|x\|$. Hence, to prove i), it is enough to construct a sequence (u_n) of elements of C such that

$$\lim_{n \rightarrow \infty} \|x - u_n\| = 1 + \|x\|.$$

For each $n \in \mathbb{N}$, let $\delta_n > 0$ be such that $n(\delta_n + \delta_n^2) = 1$. Note that $0 < \delta_n < \frac{1}{n}$ and $\lim_{n \rightarrow \infty} n\delta_n = 1$. Let $u_n = (\delta_n, \dots, \delta_n, 0, 0, \dots)$ where δ_n is repeated n times. By our choice of δ_n , $u_n \in C$.

Now given $x \in \ell^1(\mathbb{N})$ and $\varepsilon > 0$, let $p \in \mathbb{N}$ be such that $\sum_{n=p+1}^\infty |x_i| < \frac{\varepsilon}{3}$. We have for $n > p$

$$\begin{aligned} \|x - u_n\| &= \sum_{i=1}^p |x_i - \delta_n| + \sum_{i=p+1}^n |x_i - \delta_n| + \sum_{i=n+1}^\infty |x_i| \\ &\geq \sum_{i=1}^p |x_i| - p\delta_n - \sum_{i=p+1}^n |x_i| + (n - p)\delta_n \\ &\geq \|x\| - \frac{\varepsilon}{3} - p\delta_n - \frac{\varepsilon}{3} + (n - p)\delta_n. \end{aligned}$$

Choose $n_0 > p$ big enough so that $(n - 2p)\delta_n > 1 - \frac{\epsilon}{3}$ for $n \geq n_0$. We have that for all $n \geq n_0$:

$$\|x - u_n\| \geq 1 + \|x\| - \epsilon.$$

Therefore,

$$\lim_{n \rightarrow \infty} \|x - u_n\| = 1 + \|x\|.$$

■

Remarks. (a) Since $\ell^1(\mathbb{N})$ has the Radon-Nikodym property, C is the norm closed convex hull of its strongly exposed points [3]. Hence, Proposition 1 gives a negative answer to the question of Edelstein and Lewis mentioned above.

(b) The proof of Proposition 1 shows that the intersection of all the balls containing C is the unit ball of $\ell^1(\mathbb{N})$. Note also that the sequence $\{u_n\}$ in the proof does not depend on x .

Before proceeding, let us recall that the subdifferential of a convex function f defined on a Banach space X is defined by

$$\partial f(x) = \{x^* \in X^* | (\forall y \in X)(\langle x^*, y - x \rangle \leq f(y) - f(x))\}.$$

It was shown in [9] that if C is a closed bounded subset of a Banach space X and $r(x)$ is the farthest distance function for C , then for any $x \in X$ and $x^* \in \partial r(x)$, we have that $\|x^*\| \leq 1$ and thus $\sup\{\langle x^*, x - z \rangle : z \in C\} \leq \|x^*\|r(x) \leq r(x)$. Moreover, Lau showed that the set

$$G = \{x \in X | (\forall x^* \in \partial r(x))(\sup\{\langle x^*, x - z \rangle : z \in C\} = r(x))\},$$

is dense G_δ in X .

We shall need the following.

LEMMA 2. *Let C be a closed bounded subset of a Banach space X and r the farthest distance function on X associated with C . Assume that r is Fréchet differentiable at $x_0 \in X$. Then*

$$r(x_0) = \sup\{\langle x_0^*, x_0 - z \rangle | z \in C\}, \quad \text{where } \partial r(x_0) = \{x_0^*\}.$$

PROOF: According to Lau's result mentioned before the statement of Lemma 2 it is enough to show that, under our assumptions, the following implication holds:

If $r(x_0) - \sup\{\langle x_0^*, x_0 - z \rangle | z \in C\} = \alpha > 0$, then there is a neighbourhood U of x_0 such that

$$r(y) - \sup\{\langle y^*, y - z \rangle | z \in C\} > 0,$$

whenever $y \in U$ and $y^* \in \partial r(y)$.

To prove that the implication holds, take $y \in X$, $y^* \in \partial r(y)$ and $z \in C$. We have:

$$\begin{aligned} |\langle y^*, y - z \rangle - \langle x_0^*, x_0 - z \rangle| &\leq |\langle y^*, y - z \rangle - \langle y^*, x_0 - z \rangle| + |\langle y^*, x_0 - z \rangle - \langle x_0^*, x_0 - z \rangle| \\ &\leq \|y^*\| \|y - x_0\| + \|y^* - x_0^*\| \|x_0 - z\|. \end{aligned}$$

Therefore,

$$|\sup\{\langle y^*, y - z \rangle | z \in C\} - \sup\{\langle x_0^*, x_0 - z \rangle | z \in C\}| \leq \|y - x_0\| + r(x_0) \|y^* - x_0^*\|.$$

Now choose a neighbourhood U of x_0 such that

$$2\|y - x_0\| + r(x_0) \|y^* - x_0^*\| < \alpha$$

whenever $y \in U$ and $y^* \in \partial r(y)$. This is possible due to the Fréchet differentiability of r at x_0 [2, Lemma 5]. If $y \in U$ and $y^* \in \partial r(y)$, then

$$\begin{aligned} (r(x_0) - \sup\{\langle x_0^*, x_0 - z \rangle | z \in C\}) - (r(y) - \sup\{\langle y^*, y - z \rangle | z \in C\}) \\ \leq |r(x_0) - r(y)| + |\sup\{\langle y^*, y - z \rangle | z \in C\} - \sup\{\langle x_0^*, x_0 - z \rangle | z \in C\}| \\ \leq \|x_0 - y\| + \|y - x_0\| + r(x_0) \|y^* - x_0^*\| < \alpha \end{aligned}$$

and this implies, by the definition of α , that

$$r(y) - \sup\{\langle y^*, y - z \rangle | z \in C\} > 0$$

whenever $y \in U$ and $y^* \in \partial r(y)$. Our implication is proven and the proof of Lemma 2 is finished. ■

PROPOSITION 3. *Let X be a Banach space with the Radon-Nikodym property, let X^* be its dual space in its usual dual norm and C be a w^* -compact subset of X^* . Then the set D of all points in X^* which have farthest points in C contains a subset D_1 dense and G_δ in X^* .*

PROOF: First notice that the farthest distance function r associated with C is w^* -lower semicontinuous as supremum of such functions. Since X has the Radon-Nikodym property it follows from [4] that r is Fréchet differentiable on a dense G_δ subset D_1 in X^* . So, to finish the proof of Proposition 3 we show that if r is Fréchet-differentiable at x , then x admits a farthest point in C . If $x \in D_1$, denote by x^* the only element of $\partial r(x)$. Since r is w^* -lower semicontinuous and Fréchet differentiable at x , $x^* \in X$ [2, Corollary 1]. Since C is w^* -compact, there is a $z_0 \in C$ such that $\langle x^*, x - z_0 \rangle = \sup\{\langle x^*, x - z \rangle | z \in C\}$. Using Lemma 2 we have $r(x) = \langle x^*, x - z_0 \rangle \leq \|x^*\| \cdot \|x - z_0\| \leq \|x - z_0\| \leq r(x)$. Thus $\|x - z_0\| = r(x)$ and the proof is finished. ■

PROPOSITION 4. Let X be a Banach space and C be a closed convex bounded subset of X . The following are equivalent:

- (i) C is weakly compact;
- (ii) for every equivalent norm $\|\cdot\|_1$ on X , $D = \{x \in X | r(x) = \|x - z\|_1 \text{ for some } z \in C\}$ is dense in X , where $r(x) = \sup\{\|x - z\|_1 | z \in C\}$.

PROOF: (i) \Rightarrow (ii) was proved by Lau in [9]. Conversely, let $\|\cdot\|$ be the norm of X and let $B(x, r)$ denote the $\|\cdot\|$ -ball centred at x and radius r . Assume without loss of generality that $C \subset B(0, 1)$. By James' theorem [5] there is a functional $f \in X^*$, $\|f\| = 1$ which does not attain its supremum on C . Let $B_1 = B(0, 6) \cap \{x \in X | -1 \leq f(x) \leq 1\}$ and $\|\cdot\|_1$ be the Minkowski functional of B_1 . Note that if $B_1(x, r)$ is the ball centred at x with radius δ with respect to the norm $\|\cdot\|_1$, then

$$B_1(x, \delta) = 6B(x, \delta) \cap \{y \in X | -\delta \leq f(x) - f(y) \leq \delta\}.$$

Pick $x_0 \in B(0, 4) \cap \{x \in X | f(x) < -3\}$. We claim that if $x \in B(x_0, 1)$, then x has no farthest point in C when X is equipped with the new norm $\|\cdot\|_1$, hence D is not dense in X .

To see it, we first show that $r(x) = \alpha$, where $\alpha = \sup\{f(z) | z \in C\} - f(x)$. Indeed, let $y \in C$; we have $|f(y) - f(x)| \leq \sup\{f(z) | z \in C\} - f(x)$ and $\|y - x\| \leq \|y\| + \|x_0\| + \|x_0 - x\| \leq 6 \leq 6\alpha$ (note that for any $y \in C$ and $x \in B(x_0, 1)$ we have $f(y) \geq -1$ and $f(x) \leq f(x_0) + f(x - x_0) < -2$ and thus $\alpha > 1$). Thus $y \in B(x, \alpha)$. Therefore $C \subset B_1(x, \alpha)$ and this shows that $r(x) \leq \alpha$. Conversely if $\delta < \alpha$, there exists $y \in C$ such that $f(y) - f(x) > \delta$ and so $y \notin B_1(x, \delta)$. This shows that $r(x) \geq \alpha$.

Now for $y \in C$, we have $\|y - x\| \leq 6 < 6\alpha$ and $|f(y) - f(x)| < \sup\{f(z) | z \in C\} - f(x) = \alpha$, hence $\|x - y\|_1 < \alpha = r(x)$ and so y is not a farthest point from x in C . ■

Remark. We would like to point out that, while differentiability properties can be used to show that the set D of points which have at least one farthest point in a w^* -compact C of a dual Banach space X is dense in X , convexity properties can be used to show the uniqueness of farthest points. More precisely, using standard rotundity arguments, we have

PROPOSITION 5. Let X be a strictly convex Banach space, and C be a norm closed bounded subset of X such that the corresponding set D is dense in X .

Then the set

$$U = \{y \in X | y \text{ has a unique farthest point in } C\}$$

is also dense in X .

PROOF: First the strict convexity of X implies that for all $x \in X$ and $\lambda > 1$,

$$B(x, \|x\|) \cap S(\lambda x, \|\lambda x\|) = \{0\}$$

where $B(x, r)$ (respectively, $S(x, r)$) denotes the ball (respectively, the sphere) of centre x and radius r .

Now let $x \in D$ and let z be a farthest point from x in C . By translation, there is no loss of generality in assuming $z = 0$. It is enough to show that for any $\lambda > 1$, $y = \lambda x$ has a unique farthest point in C , namely 0. We have

$$B(\lambda x, \|\lambda x\|) \supseteq B(x, \|x\|) \supseteq C$$

and hence $r(\lambda x) \leq \|\lambda x\|$. On the other hand,

$$\{0\} \subset C \cap S(\lambda x, \|\lambda x\|) \subset B(x, \|x\|) \cap S(\lambda x, \|\lambda x\|) = \{0\}.$$

Therefore $r(\lambda x) = \|\lambda x\|$ and the only farthest point from y in C is 0. ■

Note that if X is not strictly convex, the set U of points which admit unique farthest points can be empty. Indeed, let $X = c_0(\mathbb{N})$ and

$$C = \{(z_n) \in c_0(\mathbb{N}) \mid 0 \leq z_n \leq \frac{1}{n} \text{ for all } n\}.$$

C is a norm compact convex subset of $c_0(\mathbb{N})$, and therefore $D = c_0(\mathbb{N})$. On the other hand, if $x = (x_n) \in c_0(\mathbb{N})$, we have

$$r(x) = \max\{\max\{|x_n|, |x_n - \frac{1}{n}|\} \mid n \in \mathbb{N}\}$$

so $r(x) = |x_{n_0}|$ or $r(x) = |x_{n_0} - \frac{1}{n_0}|$, for some $n_0 \in \mathbb{N}$. Define:

$$C_1 = \{(z_n) \in C \mid z_{n_0} = 0\}$$

$$C_2 = \{(z_n) \in C \mid z_{n_0} = \frac{1}{n_0}\}.$$

If $r(x) = |x_{n_0}|$ (respectively, $r(x) = |x_{n_0} - \frac{1}{n_0}|$), C_1 (respectively, C_2) is included in the set of farthest points from x in C . In both cases $x \notin U$.

REFERENCES

- [1] E. Asplund, 'Farthest points in reflexive locally uniformly rotund Banach spaces', *Israel J. Math.* **4** (1966), 213–216.
- [2] E. Asplund, 'Fréchet differentiability of convex functions', *Acta Math.* **121** (1968), 31–47.

- [3] J. Bourgin, 'Geometric aspects of convex sets with the Radon-Nikodym property', in *Lecture Notes in Mathematics: 993* (Springer-Verlag, New York, 1983).
- [4] J.B. Collier, 'The dual of a space with the Radon-Nikodym property', *Pacific J. Math.* **64** (1976), 103–106.
- [5] J. Diestel, 'Geometry of Banach spaces—selected topics', in *Lecture Notes in Mathematics 485* (Springer-Verlag, New York, 1975).
- [6] M. Edelstein, 'Farthest points of sets in uniformly convex Banach spaces', *Israel J. Mat.* **4** (1966), 171–176.
- [7] M. Edelstein and J.E. Lewis, 'On exposed and farthest points in normed linear spaces', *J. Austral. Math. Soc.* **12** (1971), 301–308.
- [8] G. Godefroy and N. Kalton, 'The ball topology' (to appear).
- [9] K.S. Lau, 'Farthest points in weakly compact sets', *Israel J. Math.* **22** (1975), 168–174.
- [10] S. Straszewicz, 'Über exponierte Punkte abgeschlossener Punktmengen', *Fund. Math.* **24** (1935), 139–143.
- [11] V. Zizler, 'On some extremal problems in Banach spaces', *Math. Scand.* **32** (1973), 214–224.

Professor V.E. Zizler,
Department of Mathematics,
Faculty of Science,
University of Alberta,
Edmonton,
Canada. T6G 2G1

Dr R. Deville,
Equipe d'Analyse Fonctionnelle,
Université Paris VI,
Paris, France.