

Partnerships between private landowners and conservationists to protect one of the most evolutionarily distinct amphibians

ANDRÉS VALENZUELA-SÁNCHEZ^{*1,2} , SEBASTIÁN MIRANDA³
RICARDO MORENO-GONZALEZ^{4,5,6}, JULIO GERDING⁴, RAYEN CATRILEO⁴
JULES GUILLEMOT⁴, SOLEDAD DELGADO-OYARZÚN², MARICELA NÚÑEZ²
ANDREW A. CUNNINGHAM¹  and MARÍA BELÉN ZAPARARTE³

Abstract Wildlife conservation on private land is an important approach that is increasingly utilized to protect biodiversity and can help contribute to the 30 by 30 target of the Global Biodiversity Framework. In 2018, a Chilean non-profit conservation organization launched a land conservation programme aiming to build long-term partnerships with private landowners to protect critical habitat for threatened amphibians in Chilean Patagonia. Here we describe a new locality record of the micro-endemic, Endangered Barrio's frog *Insuetophrynus acarpicus* found at a site that joined the programme in 2020. Barrio's frog is ranked 11th in the list of Evolutionarily Distinct and Globally Endangered amphibians. Our systematic literature search showed that most aspects of the natural history and ecology of this species are unknown, limiting our ability to provide actionable science to inform its conservation. The newly described Barrio's frog population is the eighth known locality of this species and one of only three occurring within a protected area. Habitat quality assessments indicated optimal conditions for most of the measured habitat parameters in the high-gradient stream where the species occurs. This case study illustrates that long-term partnerships between private landowners and conservationists can be used as an effective tool to protect the habitat of highly threatened amphibians.

Keywords Amphibian conservation, Barrio's frog, Chile, EDGE species, *Insuetophrynus acarpicus*, land conservation, Patagonia, Rhinodermatidae

The supplementary material for this article is available at doi.org/10.1017/S0030605324000115

Private land conservation is an important approach that is increasingly utilized to protect biodiversity and can help contribute to the 30 by 30 target of the Global Biodiversity Framework (Cortés Capano et al., 2019; Dinerstein et al., 2019). In Chile, private initiatives are protecting at least 16,692 km² of land (Moya Ramírez et al., 2016), covering key terrestrial ecosystems that are not protected by the national system of public protected areas (Plischoff & Fuentes-Castillo, 2011). Aiming to further increase the conservation value of private lands, in 2018 the NGO Ranita de Darwin designed and launched a land conservation programme to build long-term partnerships with private landowners and protect critical habitat for threatened amphibians in Chilean Patagonia (Azat et al., 2021). Here we describe a new locality record of the micro-endemic, Endangered Barrio's frog *Insuetophrynus acarpicus* found in the Refugio de Ranitas Aldea del Viento protected area, a 7-ha property that joined the land conservation programme in 2020 (Plate 1a). This is the eighth known locality of this species and one of only three occurring within a protected area (Fig. 1).

Barrio's frog, along with the northern and southern Darwin's frogs *Rhinoderma rufum* and *Rhinoderma darwini*, are the only known members of the family Rhinodermatidae, a 70 million year-old group of neotropical frogs considered to be the oldest known hyloids, a species-rich clade encompassing 54% of extant anuran species (Feng et al., 2017). Darwin's frogs are highly threatened, particularly the northern Darwin's frog, a species last seen over 40 years ago and categorized as Critically Endangered (Possibly Extinct) on the IUCN Red List (IUCN SSC Amphibian Specialist Group, 2015). This situation highlights the importance of Barrio's frog as one of the last representatives of the Rhinodermatidae. Barrio's frog is one of the top 100 most evolutionarily distinct amphibian species (Jetz & Pyron, 2018) and is ranked 11th in the Evolutionarily Distinct and Globally Endangered (EDGE) list of amphibians (Safi et al., 2013). This Chilean micro-endemic species is categorized as Endangered because of its limited geographical distribution (1,323 km²; Fig. 1), its highly isolated populations and the continuing decline in the extent and quality of its habitat (IUCN SSC Amphibian Specialist Group, 2018). Most aspects of the natural history and ecology of the species remain

*Corresponding author, andres.valenzuela-sanchez@ioz.ac.uk

¹Institute of Zoology, Zoological Society of London, London, UK

²ONG Ranita de Darwin, Valdivia, Chile

³Fundación Escuela del Viento, Valdivia, Chile

⁴Cooperativa Calahuala, Valdivia, Chile

⁵Department of Palynology and Climate Dynamics, University of Göttingen, Göttingen, Germany

⁶Instituto de Conservación, Biodiversidad y Territorio, Universidad Austral de Chile, Valdivia, Chile

Received 5 September 2023. Revision requested 15 November 2023.

Accepted 17 January 2024. First published online 11 March 2024.



PLATE 1 (a) Signing of a voluntary land conservation agreement between the landowner (co-author SM) and the NGO Ranita de Darwin for the creation of the protected area Refugio de Ranitas Aldea del Viento, Los Pellines, Valdivia, Chile. (b) An adult Barrio's frog *Insuetophrynus acarpicus* found in this protected area. (c) The high-gradient stream where the species occurs.

unknown (Fig. 2, Supplementary Material 1), limiting our capability to implement evidence-based conservation actions.

In April 2022, we identified a Barrio's frog population in the Refugio de Ranitas Aldea del Viento protected area, Los Pellines, Chile. This population, which is 18 km from the nearest known population in Chanchan, fills a gap in the known distribution of Barrio's frogs along the coast of Valdivia (Fig. 1). We found five Barrio's frog individuals along a 200-m section at the source of one of two high-gradient streams in the private protected area (Plate 1b–c). We found most individuals under small boulders within the stream. Habitat quality assessments following the protocol by Barbour et al. (1999) indicated optimal conditions for most of the measured habitat parameters at both streams in the area, although habitat quality was slightly higher in the stream where Barrio's frog was found (Supplementary Table 1). These streams are characterized by abundant epifaunal substrate, low embeddedness and sediment deposition, minimal amounts of channel substrate exposition, stable banks, absence of channelization and frequent riffles and pools, and the riparian zone is covered by native vegetation. Similarly, in a locality near Mehuín there was a positive relationship between the occurrence of Barrio's frog and stream habitat quality measured using the same habitat assessment protocol (I. Vásquez & A. Valenzuela-Sánchez, unpubl. data, 2020). Our findings are consistent with previous field observations that this species is a stream-breeding amphibian, with individuals generally found near or in water (Méndez et al., 2006; Contreras et al., 2020; see also references in Supplementary Material 1). Motivated by the discovery of this population, the landowner (co-author SM) is considering new ways to increase the level of protection given to the species in the area (e.g. upgrading the

voluntary land conservation agreement with Ranita de Darwin by signing a conservation easement under the framework of the Derecho Real de Conservación law).

This case study illustrates that long-term partnerships between private landowners and conservationists can be used as an effective tool for protecting the habitat of highly threatened amphibians. This is particularly relevant in the case of evolutionarily distinct amphibian species, which are disproportionately susceptible to local extinction as a result of habitat loss (Greenberg et al., 2018). The voluntary land conservation agreement between a landowner and Ranita de Darwin defines clear targets (e.g. conservation of native amphibians, including Barrio's frog) and conservation goals (e.g. protection of the habitat of native amphibians and control of threats to ensure self-sustaining amphibian populations). It also sets a list of strategies to achieve these goals, including a broad array of conservation actions, such as habitat protection and restoration, management of invasive species and promotion of conservation-focused research, and it provides a monitoring plan to evaluate success and enable adaptation. An important priority in the Refugio de Ranitas Aldea del Viento is to determine the presence and impact of potential threats to Barrio's frog, especially those that could originate from surrounding lands because, to our knowledge, these areas are not managed for conservation. This will facilitate the development and prioritization of effective strategies to prevent or reduce threats. Potential direct and indirect threats to Barrio's frog in this protected area include: (1) free-roaming cattle, (2) the fungus *Batrachochytrium dendrobatidis*, (3) pollution runoff from a road 200 m away, (4) domestic sewage from nearby residential buildings, (5) freshwater abstraction, (6) illegal logging, (7) increased land subdivision



FIG. 1 Previously known localities (circles) and new locality record (square) of Barrio's frog *Insuetophrynus acarpicus* in southern Chile. The shaded area represents the extent of occurrence of Barrio's frogs from the species' IUCN Red List assessment (IUCN SSC Amphibian Specialist Group, 2018). We extracted geographical coordinates of previously known localities from Contreras et al. (2020), except for Queule, which we obtained from Méndez et al. (2006).

for residential development in nearby native forests, (8) neighbouring exotic tree plantations, and (9) climate change (Rodríguez et al., 2022).

Collaborative efforts between private landowners and conservationists have enabled the protection of critical habitat for amphibians in the USA (Milmoie, 2008; Symonds, 2008; Kuyper, 2011) and the UK (Pond Conservation, 2012). To succeed, these types of conservation initiatives need to consider the local context, including case-specific factors driving landowner participation and permanence within the programme (Selinske et al., 2015; Valenzuela-Sánchez et al., 2022). A study conducted in the northern portion of the distribution of Barrio's frog found that landowners (mainly subsistence farmers) would be willing to join a land conservation programme only if this resulted in direct government-related economic incentives (Vásquez




FIG. 2 Research publications on Barrio's frog published since the description of the species in 1970. We searched for publications using Web of Science and Google Scholar with the terms '*Insuetophrynus*' OR 'Barrio's frog'. We excluded species checklists or reviews that mentioned the species without providing novel information or analyses. The search was performed on 7 August 2023. The full list of articles is in Supplementary Material 1.

& Marchant, 2021), a mechanism not yet available in Chile. In contrast, semi-structured interviews with people participating in the Ranita de Darwin land conservation programme showed that most landowners joined the scheme primarily because of their conservation values, and increasing their knowledge regarding amphibians was one of the most prevalent landowner expectations of signing up to the programme (Hernández, 2022). Our medium-term goal is to continue targeting this type of audience in Chilean Patagonia, to incorporate additional habitat suitable for Barrio's frog in this land conservation network.

Amphibian-focused land conservation programmes can be used not only to engage people in amphibian conservation but also to communicate the importance of amphibians for ecosystem functioning and human well-being (Valenzuela-Sánchez et al., 2022). We recommend increasing public and stakeholder knowledge about amphibians as a priority in amphibian conservation, as evidence demonstrates that even a modest increase in knowledge can enhance the positive attitudes and behaviours of people towards these animals (Valenzuela-Sánchez et al., 2022). However, public knowledge regarding amphibians is generally poor (Valenzuela-Sánchez et al., 2022). For instance, prior to their inclusion in the land conservation programme led by Ranita de Darwin, most landowners lacked knowledge regarding the level of diversity and endemism of native amphibians (Hernández, 2022). Partnerships between landowners and conservationists can also be used to boost the co-production of knowledge, potentially resulting in scientific insights that are actionable for amphibian conservation (Valenzuela-Sánchez et al., 2022). As part of the Refugio de Ranitas Aldea del Viento monitoring plan we will implement a long-term capture-recapture study of the Barrio's frog population in this protected area. This study should contribute new knowledge regarding the demography and spatial ecology of this

understudied species, but most importantly it will help to inform species-specific conservation actions within this private protected area. Similar ecological studies are required to evaluate the population status of and prioritize conservation efforts for other Barrio's frog populations across the species' range.

Author contributions Study design, fieldwork: AV-S, SM, RM-G, JGe, RC, JGu, SD-O, MN, MBZ; data analysis, writing: AV-S, with contributions from RM-G, AAC, MBZ.

Acknowledgements This amphibian-focused land conservation programme is led by Ranita de Darwin in collaboration with the Zoological Society of London and the Centro de Investigación para la Sustentabilidad, Universidad Andrés Bello. This programme has been funded by Zoo Leipzig, the Weeden Foundation, The Rufford Foundation, the Stiftung Artenschutz Amphibian Conservation Fund, The Mohamed bin Zayed Species Conservation Fund and the National Geographic Society.

Conflicts of interest SM is the owner of the Refugio de Ranitas Aldea del Viento protected area.

Ethical standards This research abided by the Oryx guidelines on ethical standards, and was conducted in accordance with Chilean law under permit number 226/2021 of the Servicio Agrícola y Ganadero de Chile.

Data availability All data used in this study are available in the main text and Supplementary Material.

References

- AZAT, C., VALENZUELA-SÁNCHEZ, A., DELGADO, S., CUNNINGHAM, A.A., ALVARADO-RYBAK, M., BOURKE, J. et al. (2021) A flagship for austral temperate forest conservation: an action plan for Darwin's frogs brings key stakeholders together. *Oryx*, 55, 356–363.
- BARBOUR, M., GERRITSEN, J., SNYDER, B. & STRIBBLING, J. (1999) *Rapid Bioassessment Protocols For Use in Streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates and Fish*, 2nd edition. Office of Water, US Environmental Protection Agency, Washington, DC, USA.
- CONTRERAS, P., BUSTOS, F. & RODRÍGUEZ-JORQUERA, I. (2020) New record of Barrio's frog, *Insuetophrynus acarpicus* Barrio, 1970 (Anura, Rhinodermatidae). Los Ríos region, Chile. *Check List (Luis Felipe Toledo)*, 237, 191–199.
- CORTÉS CAPANO, G., TOIVONEN, T., SOUTULLO, A. & DI MININ, E. (2019) The emergence of private land conservation in scientific literature: a review. *Biological Conservation*, 237, 191–199.
- DINERSTEIN, E., VYNNE, C., SALA, E., JOSHI, A.R., FERNANDO, S., LOVEJOY, T.E. et al. (2019) A global deal for nature: guiding principles, milestones, and targets. *Science Advances*, 5, eaaw2869.
- FENG, Y.-J., BLACKBURN, D.C., LIANG, D., HILLIS, D.M., WAKE, D.B., CANNATELLA, D.C. & ZHANG, P. (2017) Phylogenomics reveals rapid, simultaneous diversification of three major clades of Gondwanan frogs at the Cretaceous–Paleogene boundary. *Proceedings of the National Academy of Sciences of the United States of America*, 114, E5864–E5870.
- GREENBERG, D.A., PALEN, W.J., CHAN, K.C., JETZ, W. & MOOERS, A.Ø. (2018) Evolutionarily distinct amphibians are disproportionately lost from human-modified ecosystems. *Ecology Letters*, 21, 1530–1540.
- HERNÁNDEZ, V. (2022) *Caracterización de participantes de un programa de conservación de tierras privadas enfocado en anfibios en el sur de Chile*. BSc thesis. Universidad Austral de Chile, Valdivia, Chile.
- IUCN SSC AMPHIBIAN SPECIALIST GROUP (2015) *Rhinoderma rufum*. In *The IUCN Red List of Threatened Species* 2015. dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T19514A79809567.en.
- IUCN SSC AMPHIBIAN SPECIALIST GROUP (2018) *Insuetophrynus acarpicus*. In *The IUCN Red List of Threatened Species* 2018. dx.doi.org/10.2305/IUCN.UK.2018-1.RLTS.T10832A79809262.en.
- JETZ, W. & PYRON, R.A. (2018) The interplay of past diversification and evolutionary isolation with present imperilment across the amphibian tree of life. *Nature Ecology & Evolution*, 2, 850–858.
- KUYPER, R. (2011) The role of safe harbor agreements in the recovery of listed species in California. *Endangered Species Bulletin*, 36, 10–13.
- MÉNDEZ, A.M., TORRES-PÉREZ, F., CORREA, C., SOTO, E.R., NUÑEZ, J.J., VELOSO, A. & ARMESTO, J. (2006) Genetic divergence in the endangered frog *Insuetophrynus acarpicus* (Anura: Leptodactylidae). *Herpetological Journal*, 16, 93–96.
- MILMOE, J. (2008) Partnerships to conserve amphibian habitat. *Endangered Species Bulletin*, 33, 36–37.
- MOYA RAMÍREZ, D., PALMA TORRES, M. & FERNÁNDEZ GROSSI, P. (2016) *Diagnóstico y caracterización de las iniciativas de conservación privada en Chile*. Programa de las Naciones Unidas para el Desarrollo, Santiago, Chile.
- PLISCOFF, P. & FUENTES-CASTILLO, T. (2011) Representativeness of terrestrial ecosystems in Chile's protected area system. *Environmental Conservation*, 38, 303–311.
- POND CONSERVATION (2012) *Bringing Back Clean Water to the Countryside: Million Ponds Project Year 4 Report*. Pond Conservation (now Freshwater Habitats Trust), Oxford, UK. content.freshwaterhabitats.org.uk/2013/09/MPP-Y4-REPORT-lowres.pdf [accessed January 2024].
- RODRÍGUEZ, L.J., BARBOSA, O.A., AZAT, C., ALVARADO-RYBAK, M., CORREA, C., MÉNDEZ, M.A. et al. (2022) Amphibian phylogenetic diversity in the face of future climate change: not so good news for the Chilean biodiversity hotspot. *Biodiversity and Conservation*, 31, 2587–2603.
- SAFI, K., ARMOUR-MARSHALL, K., BAILLIE, J.E.M. & ISAAC, N.J.B. (2013) Global patterns of evolutionary distinct and globally endangered amphibians and mammals. *PLOS One*, 8, e63582.
- SELINSKE, M.J., COETZEE, J., PURNELL, K. & KNIGHT, A.T. (2015) Understanding the motivations, satisfaction, and retention of landowners in private land conservation programs: landowner commitment to conservation. *Conservation Letters*, 8, 282–289.
- SYMONDS, K. (2008) Ranchers restore amphibian-friendly ponds. *Endangered Species Bulletin*, 33, 30–31.
- VALENZUELA-SÁNCHEZ, A., AGOSTINI, G., MAYNARD, L., CISTERNAS, J., HANSEN-HENDRIKX, C. & MÁRQUEZ-GARCÍA, M. (2022) Communications and education for amphibian conservation. In *The Amphibian Conservation Action Plan (ACAP): A status review and roadmap for global amphibian conservation* (eds S. Wren, A. Borzée, R. Marcec-Greaves & A. Angulo), pp. 463–502. Draft for Open Consultation. IUCN Species Survival Commission Amphibian Specialist Group. doi.org/10.32942/osf.io/brfas.
- VÁSQUEZ, Y. & MARCHANT, C. (2021) Percepciones de pequeños propietarios y su disposición a la conservación de la naturaleza en áreas privadas de la Cordillera de la Costa en la Región de Los Ríos, Chile. *Investigaciones Geográficas (España)*, 75, 227–247.