

## ON SOME APPLICATIONS OF GRAPH THEORY III

BY

P. ERDŐS, A. MEIR, V. T. SÓS, and P. TURÁN

*In memory of Leo Moser, a friend and colleague*

1. In the first and second parts of this sequence we dealt with applications of graph theory to distance distribution in certain sets in euclidean spaces, to potential theory, to estimations of the transfinite diameter [1] and to value distribution of "triangle functionals" (e.g. perimeter, area of triangles) [2]. The basic tool is provided in all these applications by the result formulated as Lemma 2. This, an essentially pure logical result, proves to be a very flexible and versatile instrument in applications.

Here the same method is used in an abstract setting. First we deduce certain results for the density of a given family of subsets of an abstract set  $S$  in another family of subsets of the same  $S$ . Then we apply the results obtained to distance distribution in certain (e.g. totally bounded or compact) sets in metric spaces, in particular in a normed linear function space. Applications of this method to functionals on Hilbert spaces were given by Katona [3].

2. Let  $S$  denote an infinite set and  $F$  an infinite family of its finite subsets satisfying

(2.1)  $F$  contains arbitrary large subsets

(2.2)  $f \in F$  and  $f_1 \subset f$  imply  $f_1 \in F$ .

Let  $G$  be a given family of finite subsets of  $S$ . We shall be interested in the relative density of  $G$ -subsets in  $F$ -subsets.

For fixed  $f \in F$  and fixed integer  $k$  we denote by  $L_k(f, G)$  the number of sets  $g \in G$  with  $|g| = k$  such that  $g \subset f$ . Then for fixed  $n$ ,  $n \geq k$  we define

$$(2.3) \quad l_{n,k}(F, G) = \frac{1}{\binom{n}{k}} \inf_{\substack{f \in F \\ |f|=n}} L_k(f, G).$$

The quantities  $l_{n,k}(F, G)$  are lower bounds for the density of  $G$ -subsets of cardinality  $k$  in  $F$ -subsets of cardinality  $n$ . As we shall prove later, the following result holds:

LEMMA 1. For  $n \geq k$

$$(2.4) \quad l_{n+1,k}(F, G) \geq l_{n,k}(F, G).$$

---

Received by the editors July 7, 1971.

It follows thus from (2.4) that

$$(2.5) \quad \lambda_k(F, G) \stackrel{\text{def}}{=} \lim_{n \rightarrow \infty} l_{n,k}(F, G)$$

exists for every fixed  $k$ . By (2.3) clearly  $0 \leq \lambda_k(F, G) \leq 1$ .

As an immediate consequence of the definitions and (2.4) we obtain:

**COROLLARY.** *If every subset  $f \in F$  with  $|f|=N$  contains a subset  $g \in G$  (with  $|g|=k$ ), then for  $n \geq N$*

$$(2.6) \quad l_{n,k}(F, G) \geq \binom{N}{k}^{-1}.$$

*In particular,  $\lambda_k(F, G) \geq \binom{N}{k}^{-1}$ .*

Although (2.6), in general, is a weak lower bound for  $l_{n,k}$ , in some cases it yields nontrivial conclusions (see [2]).

3. In case  $k=2$  we can improve (2.6) substantially on using the graph theoretic Lemma 2. In fact we shall show that in certain cases we can determine the best possible lower bounds for  $l_{n,2}$  and the exact values of  $\lambda_2(F, G)$ .

Our main result for  $k=2$  is the following:

**THEOREM 1.** *Suppose every subset  $f$  in  $F$  with  $|f|=N+1$  contains a pair  $g \in G$ . Then for  $n \geq N+1$ ,*

$$(3.1) \quad l_{n,2}(F, G) \geq \frac{1}{N} - \frac{1}{n}.$$

*In particular,  $\lambda_2(F, G) \geq \frac{1}{N}$ .*

4. Applications. (i) Let  $S$  be a set,  $\phi$  and  $\psi$  functionals defined on all  $k$ -tuplets in  $S$ . Let  $F$  consist of all finite subsets  $\{x_1, \dots, x_n\}$  of  $S$  for which

$$\max_{1 \leq i_1 < \dots < i_k \leq n} \phi(x_{i_1}, \dots, x_{i_k}) \leq 1.$$

Let  $G$  consist of all those finite subsets  $\{y_1, \dots, y_m\}$  for which

$$\min_{1 \leq i_1 < \dots < i_k \leq m} \psi(y_{i_1}, \dots, y_{i_k}) \leq \theta$$

with some fixed  $0 < \theta < 1$ .

If among any set of  $N$  points in  $S$  there is a  $k$ -tuple such that the corresponding value of  $\psi$  is  $\leq \theta$ , then among any set of  $n (> N)$  points there are at least

$$\binom{n}{k} / \binom{N}{k}$$

$k$ -tuplets whose corresponding  $\psi$ -value is  $\leq \theta$ . Specializing this result we get Theorem 1 of [2].

(ii) Let  $K$  be a compact (and therefore totally bounded) set in a complete metric space  $(X, \rho)$ . Let the sequence of positive numbers  $d_i = d_i(K)$  (the “packing constants”) be defined by

$$(4.1) \quad d_i = \sup_{\substack{K \subset K \\ |K|=i}} \inf_{\substack{x, y \in K \\ x \neq y}} \rho(x, y), \quad i = 2, 3, \dots$$

and the “critical indices”  $i_j (j=2, 3, \dots)$  by

$$(4.2) \quad d_2 = \dots = d_{i_2} > d_{i_2+1} = \dots = d_{i_3} > \dots.$$

Observe that from the definition it follows that  $d_{i+1} \leq d_i$  for all  $i$  and  $d_{i_j+1} = d_{i_j+1}$ . Because of the total boundedness of  $K$ ,  $\lim_{i \rightarrow \infty} d_i = 0$ .

In order to apply our previous result, we choose all finite subsets of  $K$  as  $F$  and all pairs of points  $\{P_r, P_s\}$  in  $K$  for which  $\rho(P_r, P_s) \leq \theta$  ( $0 < \theta \leq d_2$ ) as  $G$ . Then we have

**THEOREM 2.** *For any finite subset  $f$  of  $K$  and arbitrary fixed  $\theta$  ( $0 < \theta \leq d_2$ ) let  $L(f, \theta)$  denote the number of pairs  $\{P_r, P_s\}$  in  $f$  satisfying  $\rho(P_r, P_s) \leq \theta$ . Let*

$$(4.3) \quad l_n(K, \theta) = \frac{1}{\binom{n}{2}} \inf_{\substack{f \subset K \\ |f|=n}} L(f, \theta)$$

and let  $i_j$  be the integer so that

$$(4.4) \quad d_{i_j+1} \leq \theta < d_{i_j}.$$

Then for  $n > i_j$

$$(4.5) \quad l_n(K, \theta) > \frac{1}{i_j} - \frac{1}{n}.$$

Moreover, inequality (4.5), in general, cannot be replaced by  $l_n(K, \theta) \geq (i_j)^{-1}$ . If we define the “lower distance distribution” of  $K$  for  $0 < \theta \leq d_2$  by

$$(4.6) \quad \lambda(K, \theta) \stackrel{\text{def}}{=} \lim_{n \rightarrow \infty} l_n(K, \theta),$$

we have with the above used notation

**THEOREM 3.** *If  $K$  is a perfect, compact set in a complete metric space  $(X, \rho)$ , then for  $j=2, 3, \dots$*

$$(4.7) \quad \lambda(K, \theta) = \frac{1}{i_j}, \quad d_{i_j+1} \leq \theta < d_{i_j}.$$

Thus,  $\lambda(K, \theta)$  is a right-continuous step function with jumps at  $\theta = d_{i_j}, j=3, 4, \dots$

A particular case of Theorem 2 deserves special attention.

**THEOREM 4.** *Let  $X$  denote the set of functions  $\{x(t)\}$  such that  $x(t) \in C[0, 1]$ ,  $x(0) = 0$  and  $|x(t_1) - x(t_2)| \leq |t_1 - t_2|$  whenever  $0 \leq t_1 < t_2 \leq 1$ . Let the distance of any two functions  $x, y$  in  $X$  be defined by the usual maximum norm  $\|x - y\|$ .*

Then for  $\nu=1, 2, \dots$  if  $n > 2^\nu$  and  $x_1, \dots, x_n$  are any functions in  $X$ , the number of distances  $\|x_i - x_j\|$  which are  $\leq 2/\nu$  is at least

$$\frac{n^2}{2^\nu} - \frac{n}{2}$$

This estimate, in general, is best possible.

An illuminating interpretation of Theorem 4 is that the probability that randomly chosen  $x, y \in X$  satisfy  $\|x - y\| \leq 2/\nu$  is at least  $1/(2^{\nu-1})$  ( $\nu=1, 2, \dots$ ).

**5. Proofs.** We need the following:

LEMMA 2 [6]: Let  $\Gamma$  be a graph (with simple edges and no loops) having  $n$  vertices and  $e$  edges. Let  $n = N \cdot m + \nu, 0 \leq \nu < N$  and suppose that

$$e > \frac{N-1}{2N} (n^2 - \nu^2) + \binom{\nu}{2}$$

Then  $\Gamma$  contains a complete subgraph of order  $N + 1$ .

In order to prove our Theorem 1, let  $f$  be any fixed set in  $F$  with  $|f| = n, n \geq N + 1$ . Denote the elements of  $f$  by  $x_1, \dots, x_n$ . Corresponding to  $f$  we define a graph on the vertices  $P_1, \dots, P_n$  as follows: The pair  $(P_i, P_j)$  should be an edge in  $\Gamma$  if and only if the pair  $(x_i, x_j)$  is *not* in  $G$ . Then, by the assumption of Theorem 1,  $\Gamma$  cannot contain a complete subgraph of order  $N + 1$ . Thus, by Lemma 2 the number of edges  $e$  in  $\Gamma$  satisfies

$$(5.1) \quad e \leq \frac{N-1}{2N} (n^2 - \nu^2) + \binom{\nu}{2}$$

where  $n = N \cdot m + \nu, 0 \leq \nu < N$ . Returning to  $f$ , inequality (5.1) implies that at least

$$\binom{n}{2} - \frac{N-1}{2N} (n^2 - \nu^2) - \binom{\nu}{2}$$

pairs  $\{x_i, x_j\}$  are members of  $G$ . In other words

$$(5.2) \quad L_2(f, G) \geq \binom{n}{2} - \frac{N-1}{2N} (n^2 - \nu^2) - \binom{\nu}{2}$$

Since as one easily calculates, the right-hand expression is

$$\geq \frac{n^2}{2} \left( \frac{1}{N} - \frac{1}{n} \right)$$

and  $f$  is an arbitrary set in  $F$ , (3.1) follows from (5.2).

Inequality (4.5) is a consequence of Theorem 1. Namely, if  $\theta$  satisfies (4.4), then by the definition (4.1), among any set of  $i_j + 1$  points in  $K$  there is a pair  $(P_r, P_s)$  with distance  $\rho(P_r, P_s) \leq \theta$ . Hence the conditions of Theorem 1 are satisfied with  $N = i_j$ .

Now let  $K$  be a perfect set and  $n = i_j \cdot m$  ( $m$  an integer). From the definition of the

packing constants it follows that there exist points, say  $Q_1, \dots, Q_{i_j}$  in  $K$  such that  $\rho(Q_r, Q_s) > \theta$  for  $1 \leq r < s \leq i_j$ . Since  $K$  is perfect, for each  $r$ ,  $1 \leq r \leq i_j$ , there exist  $m$  points in  $K$ , say  $Q_{r,1}, Q_{r,2}, \dots, Q_{r,m}$ , "near"  $Q_r$  so that  $\rho(Q_{r,p}, Q_{s,q}) > \theta$  whenever  $1 \leq r < s \leq i_j$ , for all values of  $p, q$ . Hence in the set of  $n$  elements  $\{Q_{r,p}\}$ , ( $1 \leq r \leq i_j, 1 \leq p \leq m$ ), the number of distances  $\leq \theta$  is not greater than  $i_j \binom{m}{2}$ . This implies that

$$l_n(K, \theta) \leq \frac{m-1}{n-1} < \frac{1}{i_j}.$$

This completes the proof of Theorem 2.

The proof of Theorem 3 follows from the above. Namely, since  $l_n(K, \theta)$  is an increasing sequence,  $l_n(K, \theta) < 1/i_j$  holds for all  $n > i_j$ . Hence (4.7) follows from (4.5) and (4.6).

Theorem 4 follows from (4.5) and a result of Newman and Raymon [5] (see also [4]). Namely, in our notation, it was shown in [5] that for the set  $X$  of Theorem 4,

$$d_{2^v+1} = d_{2^v+2} = \dots = d_{2^{v+1}} = \frac{2}{v+1}, v = 0, 1, \dots$$

and thus  $i_v = 2^{v-1}, v = 1, 2, \dots$

Finally, we prove Lemma 1. Suppose

$$l_{n+1,k}(F, G) = \binom{n+1}{k}^{-1} L_k(f^*, G).$$

Let  $f_1, f_2, \dots, f_{n+1}$  denote all subsets of  $f^*$  with cardinality  $n$ . By (2.2),  $f_i \in F$  for  $1 \leq i \leq n+1$ . Now, if for some  $g \in G$  we have  $g \subset f^*$ , then  $g \subset f_i$  will hold for  $n-k+1$  of the  $f_i$ 's. Hence

$$l_{n+1,k}(F, G) = \binom{n+1}{k}^{-1} \frac{1}{n-k+1} \sum_{i=1}^{n+1} L_k(f_i, G)$$

which by the definition (2.3) of the  $L_{n,k}$ 's is

$$\geq \binom{n+1}{k}^{-1} \frac{1}{n-k+1} \cdot \binom{n}{k} (n+1) l_{n,k}(F, G) = l_{n,k}(F, G).$$

REFERENCES

1. P. Erdős, A. Meir, V. T. Sós and P. Turán, *On some applications of graph theory I*, Discrete Mathematics (to appear).
2. P. Erdős, A. Meir, V. T. Sós and P. Turán, *On some applications of graph theory II*, *Studies in Pure Mathematics*, Academic Press, (1971), 89-100.
3. Gy. Katona, *Gráfok, vektorok és valószínűség-számítási egyenlőtlenségek* (in Hungarian, with English and Russian abstracts). *Mat. Lapok*. Vol. 20, Fasc. 1-2 (1969), 123-127.
4. A. N. Kolmogorov and V. M. Tihomirov,  $\epsilon$ -entropy and  $\epsilon$ -capacity of sets in function spaces, *Uspehi Mat. Nauk.* no. 2 (86), 14 (1959), 3-86; English transl., *Amer. Math. Soc. Transl.* (2) 17 (1961), 277-364.

5. D. J. Newman and L. Raymon, *Optimally separated contractions*, Amer. Math. Monthly 77 (1970), 58–59.

6. P. Turán, *Egy gráfelméleti szélsőértékfeladatról*. (Hungarian with German abstract.) Mat. Lapok. 49 (1941), 436–452. Reproduced in English in the Appendix to P. Turán, *On the theory of graphs*, Colloq. Math., 1954.

HUNGARIAN ACADEMY OF SCIENCES,  
BUDAPEST, HUNGARY

UNIVERSITY OF ALBERTA,  
EDMONTON, ALBERTA