

disordered eating behaviours between Muslim and non-Muslim females. Studies that did not report prevalence data on these outcomes or included male participants were excluded. Following PRISMA guidelines, a comprehensive search was conducted across Medline, Embase, PsycINFO, and Scopus up to August 18, 2024, supplemented by references and grey literature. Methodological quality was assessed using the ROBINS-E and JBI Prevalence tools. Relevant data were extracted and analysed through quantitative and narrative syntheses.

Results: Eleven studies met the eligibility criteria. Overall, findings indicated lower rates of body dissatisfaction and disordered eating behaviours among Muslim females, particularly among those practising veiling, compared with non-Muslim females. However, conflicting results were noted in several studies, possibly due to acculturation and sociocultural stressors. The heterogeneity in study methodologies and cultural contexts limited the generalizability of these findings. While religiosity appeared to be a protective factor, the variability in results underscores the need for more robust, longitudinal studies to better understand the complex interplay of religious, cultural, and psychosocial factors.

Conclusion: This review highlights the importance of integrating cultural and religious contexts into effective interventions and prevention strategies for body dissatisfaction and disordered eating in Muslim populations.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Pharmacogenomics-Guided Antidepressant Therapy as a Solution to Treatment-Resistant Depression: A Systematic Review

Ms Leah Rose Alexander

University of Buckingham Medical School, Buckingham, United Kingdom

doi: [10.1192/bjo.2025.10191](https://doi.org/10.1192/bjo.2025.10191)

Aims: Treatment-resistant depression (TRD), defined as failure to achieve remission despite adequate antidepressant trials, remains a pervasive clinical conundrum, linked to reduced quality of life, increased mortality, and higher healthcare costs. Pharmacogenomics-guided treatment (PGT) offers a potential solution to the ongoing challenge of TRD by using genomic testing to identify genetic factors influencing drug metabolism and response. This systematic review aimed to assess whether PGT improves symptom remission in adults with TRD compared with treatment as usual (TAU). Secondary objectives included evaluating the impact of PGT on treatment response and overall symptom improvement.

Methods: A systematic review of clinical studies published up to 8 July 2023 was conducted across PubMed and Cochrane Library databases based on Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. Search terms were defined using the ‘Population, Intervention, Comparison, Outcome’ (PICO) framework. Studies were selected using predefined inclusion and exclusion criteria. Clinical trials or observational studies evaluating the efficacy of PGT for TRD in at least 10 adult participants were included. The quality of the included studies was assessed using the Critical Appraisal Skills Program (CASP) checklist.

Results: The systematic literature search identified 307 records, 15 of which were included in a qualitative synthesis. Among 14 studies evaluating remission, 13 (92.9%) reported improvements with PGT

compared with TAU, with 8 (57.1%) showing significant gains. For treatment response, 12 out of 13 studies (92.3%) demonstrated enhancements with PGT, 8 (61.5%) of which were significant. All 8 studies assessing symptom improvement reported benefits with PGT, with 5 (62.5%) showing significant results. Notable findings included more consistent effects in patients with fewer prior antidepressant failures and improved response rates even in the absence of remission. However, variability in study designs, scoring systems, and definitions of TRD limited the generalisability of results.

Conclusion: PGT demonstrates promise in enhancing remission, treatment response, and symptom outcomes in TRD. By tailoring treatment to individual genetic profiles, this approach may reduce the trial-and-error process associated with antidepressant therapies, thereby improving patient well-being and alleviating healthcare burdens. Nevertheless, further studies, including meta-analyses and cost-benefit evaluations, are warranted to establish the clinical and economic viability of PGT for widespread implementation in healthcare systems like the NHS.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Understanding Subgenual Cingulate Functional Connectivity in Major Depressive Disorder Using 7T fMRI

Dr Laith Alexander^{1,2}, Ms Mu Li³, Professor James Murrough³ and Dr Laurel Morris^{3,4}

¹King’s College London, London, United Kingdom; ²South London and Maudsley NHS Foundation Trust, London, United Kingdom;

³Mount Sinai School of Medicine, New York, USA and ⁴Nuffield Department of Clinical Neurosciences, Oxford, United Kingdom

doi: [10.1192/bjo.2025.10192](https://doi.org/10.1192/bjo.2025.10192)

Aims: We aimed to determine differences in subgenual anterior cingulate cortex (sgACC) resting-state functional connectivity (rsFC) in Major Depressive Disorder (MDD) vs. healthy volunteers (HV) using 7-Tesla functional magnetic resonance imaging (fMRI). Abnormalities in the sgACC are linked to MDD, but the sgACC is anatomically and functionally diverse, including Brodmann area (BA) 25 (Cg25) and the subgenual portion of area 32 (Cg32). The differences in rsFC between Cg25 and Cg32 in MDD compared with HVs have not been directly examined. High-resolution 7T fMRI offers an unrivalled opportunity to measure differences in rsFC between these two subregions which otherwise suffer from signal dropout.

Methods: We used resting state 7T fMRI to compare rsFC between Cg25 and Cg32 in 40 patients with MDD, and 38 HVs. Within the MDD group, we correlated rsFC changes with anhedonia (SHAPS) and anxiety (STICSA) scores together with baseline high-sensitivity C-reactive protein (hsCRP) measures.

Results: Across all 78 participants, Cg25 and Cg32 showed regionally distinct rsFC patterns despite their proximity. Cg25 had increased rsFC to the orbitofrontal cortex, amygdala, hippocampus and dorsolateral (dl)PFC/BA46, while Cg32 showed increased rsFC to the perigenual (pg) and dorsal (d)ACC, dlPFC/BA9, posterior cingulate cortex (PCC), ventral striatum, and ventral tegmental area. When comparing MDD patients to HV, both Cg25 and Cg32 exhibited increased rsFC to the anterior (ant)PFC/BA10, amygdala and hypothalamus, together with key nodes of the default mode network (DMN), including pgACC, rostral ventromedial prefrontal

cortex (vmPFC) and the PCC. rsFC to nodes of the central executive and salience networks, such as the right dlPFC/BA46 and the bilateral insula, was decreased. Within the MDD group, Cg32-antPFC/BA10 and Cg32-dlPFC/BA9 rsFC was positively correlated with anhedonia scores; additionally, subthreshold clusters were identified in the ventral striatum, pgACC and hypothalamus. Cg25-antPFC/BA10 and Cg25-PCC rsFC was negatively correlated with anxiety scores. Cg32 rsFC to the insula, dlPFC/BA9 and dmPFC/BA10 showed negative correlations with hsCRP measures.

Conclusion: These findings suggest that sgACC subregions have distinct rsFC patterns which are altered in MDD. rsFC changes are differentially related to symptoms of anhedonia and anxiety, together with inflammatory status. This has important implications for the development of targeted neuromodulation treatment strategies.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Covid-19 Associated Psychosis in Children and Young Person – a Systematic Review of Case Reports

Dr Amey Angane and Dr Victoria Fernandez Garcia De Las Heras
South West London St George's Mental Health NHS Trust, London, United Kingdom

doi: [10.1192/bjo.2025.10193](https://doi.org/10.1192/bjo.2025.10193)

Aims: The primary objective of this descriptive systematic review of case reports is to describe the clinical commodities, presentations and outcomes in children and adolescents presenting with onset of non-delirious psychosis during or shortly after a SARS-CoV-2 infection and to find out statistically various other factors that might be linked to demographics of young people. The review also explores if the clinical presentation of the Covid-19 psychosis is different from early onset non-organic psychosis occurring in children and adolescents.

Methods: On 23 September 2023, the author searched six electronic databases including PubMed, Scopus, Web of Science, PsycInfo, Google Scholar, and CINAHL, using the following search terms: (COVID-19 OR SARS-CoV-2* OR Severe Acute Respiratory Syndrome Coronavirus 2* OR COVID*) AND (Psychosis) AND (Adolescent OR Children OR Teenager). An updated search was completed on 10 August 2024. Search results from six databases were manually checked to remove any duplication. The extracted data was then arranged in a standardised template. The extracted data included: demographic characteristics of the patients including age, gender, ethnicity, past personal and family psychiatric history, clinical features including neurological and psychotic symptoms and management including outcome.

Results: This descriptive systematic review identified 15 cases of incident psychosis in patients with antecedent or concurrent Covid-19. Out of 15 cases, 9 were males, 4 were females and 2 did not report any sex. The mean age of patients in our sample was 15.1 years with 2 cases not reporting the actual age. Delusions were present in all cases (100%) of patients, whereas hallucinations were reported in only 33% of the cases. Disorganised speech or behaviour was reported in 40% of the cases. Psychotic symptoms lasted from approximately 7–90 days. Family history was positive for 2/15 cases (7.5%) with psychosis and BPAD respectively. Only 2 cases had past personal history of mental illness (Depression and anxiety). All patients received antipsychotic medications as a part of the treatment, whereas 33% patients received intravenous immunoglobulins concurrently with antipsychotic medication. Full remission was obtained in nearly all cases after treatment.

Conclusion: Covid-19 related psychosis differs in various aspects and should be considered as a separate entity when considering the assessment and management. It differs in many ways from a typical early onset psychotic episode both in presentation and treatment response.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Societal Attitudes Towards Transgender Individuals: A Multicentric Survey Across Urban, Rural and Cosmopolitan Cities in India

Dr Arokia Antonysamy¹, Dr Amutha Govindaraju²,
Dr Helena Selvakodi² and Dr Shanmugapriya Samiyappan²

¹Cygnat Health Care, London, United Kingdom and ²Madurai Medical College Hospitals, Madurai, India

doi: [10.1192/bjo.2025.10194](https://doi.org/10.1192/bjo.2025.10194)

Aims: The aim of this study is to investigate public attitudes towards transgender individuals, seeking to understand the relationship between socio-demographic factors, gender beliefs and approach towards this minority group. The research acknowledges the widespread stigma faced by transgender people due to the incongruence between their gender identity and sex assigned at birth, which negatively impacts their access to resources and overall well-being.

Methods: This is a cross-sectional survey conducted across rural, suburban and cosmopolitan cities in India. A total of 500 participants were randomly selected using multistage sampling. Inclusion criteria for participants included the age range 18–60 years, no history of mental illness, and having no known family members with transgender characteristics.

This specific demographic targeting aimed to isolate general public perception, excluding potentially biased viewpoints from those with direct personal connections to transgender individuals or pre-existing mental health conditions.

Results: The study revealed a predominantly young adult sample, with 52.6% of respondents falling between the ages of 26 and 40. The sample was fairly distributed across both genders, male (56%), females (44%). Majority were married (62.4%), educated at graduate level (61%) and employed. This socio-demographic breakdown provides context for interpreting the attitudinal data.

Key findings indicated a concerning trend: younger, unmarried males with higher education demonstrated more negative attitudes towards transgender individuals. This finding contradicts some expectations, as higher education is often associated with more liberal and inclusive viewpoints. Furthermore, the study found that highly educated individuals, in general, held more negative attitudes, while married individuals tended to express more positive attitudes. Interestingly, there was no significant difference in attitudes towards transgender men compared with transgender women.

Conclusion: The study concludes by emphasizing the need for targeted educational interventions, particularly aimed at younger, unmarried, highly educated males. The authors argue that these interventions are crucial for promoting cultural competence and ensuring that transgender individuals are afforded their due rights. By focusing on education and awareness, the study suggests a potential pathway for mitigating the negative attitudes identified and fostering a more inclusive and accepting society for transgender individuals. The study highlights the importance of understanding the complex interplay between socio-demographic factors and