

the pandemic. On the other hand, despite these differences, the two groups were similar in their will to live.

Conclusions: Holocaust survivors seem to be more vulnerable to the COVID-19 pandemic, strengthening the vulnerability hypothesis. Policy makers and practitioners should pay special attention to this particularly vulnerable population during these difficult times.

Disclosure: No significant relationships.

Keywords: COVID-19; Holocaust survivors; mental health; loneliness

O085

Clinical profile of patients admitted in an acute psychiatric ward before and during the COVID-19 lockdown

A. Giménez Palomo*, N. Baldaquí, L. Colomer, F. Gutiérrez, E. Pujal, P. Barrio, M. Bioque, E. Vieta and I. Pacchiarotti

Department Of Psychiatry And Psychology, University Of Barcelona, Hospital Clínic, Barcelona, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.295

Introduction: The COVID-19 pandemic entailed several changes in health and medical assistance, economy, and lifestyle. In the Acute Psychiatric Ward of the Hospital Clínic of Barcelona, the implementation of restrictive measures was necessary in order to ensure patients' safety.

Objectives: To compare clinical profiles and course of hospitalization of patients admitted before and during the COVID-19 lockdown in our Acute Psychiatric Ward.

Methods: All patients admitted from January 7th to February 25th and from March 19th to May 7th of 2020 in the Acute Psychiatric Hospitalization Unit of Hospital Clínic of Barcelona, Spain, were retrospectively included for analysis and divided into two groups according to the period when they were admitted. Statistical analyses were performed using SPSS, 23.0 version.

Results: A total of 117 inpatients were included (73 admitted before lockdown and 44 during lockdown), being 50.4% male, with a mean age of 42.4 (SD 15.73). Patients from the first group presented a significantly higher proportion of antidepressants prescription at discharge ($p<0.05$) and more substance use disorders ($p<0.05$). Regarding the lockdown group, 51% of patients manifested COVID-19-related stress. Time of hospitalization was significantly lower in the lockdown group ($p<0.05$), even though a significantly higher proportion of patients were discharged at home ($p<0.05$) compared with the first group.

Conclusions: The situation of lockdown led to a series of changes in our unit and also in the profile of patients admitted, having shorter admissions, lower prescription of antidepressants, and often COVID-related stress. These differences should be considered in future situations in which restrictive measures may be necessary.

Disclosure: No significant relationships.

Keywords: coronavirus; lockdown; Hospitalization; acute psychiatric ward

O086

The perception of individuals in society about protection measures from COVID-19 infection: The example of turkey

F. Oflaz¹ and F. Atkan^{2*}

¹Nursing (psychiatric Nursing), Koç University School of Nursing, Istanbul, Turkey and ²Nursing (psychiatric Nursing), Koç University Graduate School of Health Sciences, Istanbul, Turkey

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.296

Introduction: The course of the epidemics such as COVID-19 and SARS has taught us that the management of the epidemic depends primarily on people's adherence to and implementation of the recommended measures.

Objectives: This study aimed to determine the knowledge and opinions of individuals about COVID-19 and transmission methods, sources of information, application status about protection measures and related factors.

Methods: 1444 people participated into the digital survey between March 22-April 6, 2020 for this descriptive study. For data collection, a 12-questions questionnaire consisting of questions about the socio-demographic characteristics, information sources they used about the COVID-19, their thoughts the practices to prevent the transmission of COVID-19 was conducted. Using descriptive statistics and comparison tests, individuals' perceptions about methods of protection from Covid-19 and related variables were investigated.

Results: The participants have had sufficient knowledge about Covid-19 and measures. They were using social media platforms, official web sites and TV news to get information about the COVID-19. The rates of believing and applying measures such as staying distant from people, washing hands, staying at home, avoiding from public transportation, using alcohol disinfectants were quite high. The women, people living in large cities, healthcare workers, regular commuters to work believed in measures more, however, their level of anxiety and seeing themselves and their environment at risk were higher.

Conclusions: Despite all the positive results regarding coronavirus infection and protection measures, the fact that the epidemic is spreading rapidly indicates the need for studies to continuously evaluate what has changed in the process and as time increases.

Disclosure: No significant relationships.

Keywords: prevention measures; COVID-19; Turkey; perceptions

O087

Neuropsychiatric manifestations of SARS-CoV-2 infection

V. Nogueira^{1*}, M. Marguilho², I. Pereira¹, J. Teixeira¹ and M. Mendes¹

¹Clínica 4 - Unidade De Alcoologia E Novas Dependências, Centro Hospitalar Psiquiátrico de Lisboa, Lisboa, Portugal and ²Clínica 5, Centro Hospitalar Psiquiátrico de Lisboa, Lisboa, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.297

Introduction: Starting in December 2019, the coronavirus SARS-CoV-2 emerged and soon acquired a pandemic dimension. The evidence that 1 in 3 patients presented neuropsychiatric symptoms highlighted SARS-CoV-2 neurotropic properties. The involvement of the Central Nervous System (CNS) seems to be associated with poor prognosis, and it can occur independently of the respiratory system.

Objectives: To assess neuropsychiatric symptoms in SARS-CoV-2 patients and possible mechanisms of CNS invasion; to reflect on what changes should be made in order to avoid short and long-term complications.

Methods: A non-systematic literature review was performed, including publications between January and August 2020.

Results: The most frequent CNS presentations included fatigue (38-75%), headache (6,5-34%), nausea or vomiting (1-13,7%). Regarding PNS involvement, three kinds of hypoesthesia (hyposmia, hypogeusia, and hypopsia) were commonly present. Additionally, cases of neurological syndromes associated with SARS-CoV2 were reported, being related to a poor prognosis in cases such as brainstem infiltration. Another major concern regarding CNS involvement is the possibility of permanent neurological disabilities. Importantly there are reports of patients who tested positive for SARS-CoV-2 in CFS, without samples from nasopharyngeal swabs. Different hypothesis are postulated to explain possible mechanisms through which SARS-CoV-2 affects CNS, including: direct invasion through the olfactory nerve, hematogenous route through ACE-2 (angiotensin-converting enzyme) receptor expressed in blood-brain-barrier; or indirect mechanisms.

Conclusions: Here we discuss the neuropsychiatric manifestations of SARS-CoV-2 infection and the potential mechanisms by which they occur at an early stage. Awareness, prevention and early treatment of potential neuropsychiatric symptoms of COVID-19 should not be overlooked, especially because they seem to predict a worse prognosis.

Disclosure: No significant relationships.

Keywords: SARS-CoV-2; nervous system; COVID-19

O088

Alcohol consumption during COVID-19 pandemic: What have we learnt so far?

V. Nogueira*, M. Mendes, I. Pereira and J. Teixeira

Clínica 4 - Unidade De Alcoologia E Novas Dependências, Centro Hospitalar Psiquiátrico de Lisboa, Lisboa, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.298

Introduction: The current SARS-CoV-2 pandemic has many implications, one of them being alcohol consumption. The impact of long-term distancing measures in terms of alcohol use and misuse is yet unknown. Any increase, would not only add to the usual disease burden associated with alcohol, but also add to the COVID-19 load, given that alcohol use may weaken the immune response.

Objectives: To characterize and compare the pattern of alcohol consumption throughout the pandemic in patients with the diagnosis of Alcohol Use Disorder; to identify factors considered as most relevant in the increase of alcohol consumption.

Methods: We conducted a observational study in an outpatient population in Centro Hospitalar Psiquiátrico de Lisboa (Portugal) with diagnosis of Alcohol Use Disorder, 6 months after the pandemic lockdown. We characterized our sample regarding social,

demographic and clinical categories. We applied auto-filled questionnaires, particularly: Mental Health Inventory (MHI), Positive Mental Health Scale (PMHS) and Severity of Alcohol Dependence Questionnaire (SADQ-C).

Results: A total of 65 patients were included. More than 30% changed their drinking habits because of the pandemic. Nearly half of these increased consumption, and half decreased (16% vs 14%). The increase affected particularly men, and was related with the severity of alcohol dependence, stress-related coping strategies and psychopathology; on the other hand, a lowered level of consumption based on the decrease of alcohol accessibility and affordability.

Conclusions: The current situation is unique in terms of mass physical distancing and may trigger different behaviours that should be monitored. Governments should give public health warning about excessive alcohol consumption to protect vulnerable individuals.

Disclosure: No significant relationships.

Keywords: COVID-19; pandemics; alcohol consumption; Psychological Distress

O090

The coronavirus pandemic impact on involuntary hospitalization

C. Oliveira*, F. Caldas, M. Gonçalves and J. Freitas

Internamento C, Hospital de Magalhães Lemos, Porto, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.299

Introduction: Compulsory admission is the last line of intervention in individuals who suffer from severe mental disorders, based on the principles of therapeutic need and social protection. In Portugal, the law configures this measure as a hospitalization by court order. The SARS-CoV-2 coronavirus is a global public health emergency, with multifaceted consequences for people's lives and health. Several studies are showing a great impact of the pandemic on the overall mental health.

Objectives: The aim is to assess the impact of the pandemic on the number of involuntary hospitalizations, their socio-demographic and clinical characteristics, and study the differences between 2019 and 2020.

Methods: Consultation of all patient's clinical files that were involuntarily admitted in Hospital Magalhães Lemos during 2019 and 2020. Statistical analysis of data.

Results: The authors claim to investigate the impact of the pandemic on involuntary hospitalizations, the factors of admission and decompensation and other clinical characteristics, by comparing the involuntary hospitalizations during 2019 and 2020. The authors believe that the number of compulsory admissions increased significantly with the pandemic. They also believe that factors such as increased or relapsed consumption of alcohol and drugs, suicide attempts, missed appointments and long-term injectable medication are at the root of this increase in 2020.

Conclusions: This study helps to analyze the impact of the new coronavirus on compulsory hospitalizations and allows to understand the main factors that aggravate the underlying pathologies. Thus, understanding the targets of greater attention from psychiatrists to avoid the decompensation of patients in times of pandemic in which we currently live.