


Original Research

Cite this article: Keskin G and Yurt E.
Evaluation of the situations of coping with
mental trauma and trauma in emergency
service personnel who medically intervened to
earthquake affected people in the 2020 Izmir
earthquake. *Disaster Med Public Health Prep.*
18(e17), 1–9. doi: [https://doi.org/10.1017/
dmp.2023.237](https://doi.org/10.1017/dmp.2023.237).

Keywords:
Izmir earthquake; trauma level; coping with
stress; health-care worker; affected
responder's stress

Corresponding author:
Esra Yurt; Email: esrayalcin2017@gmail.com.

Evaluation of the Situations of Coping With Mental Trauma and Trauma in Emergency Service Personnel Who Medically Intervened to Earthquake Affected People in the 2020 Izmir Earthquake

Gülseren Keskin PhD¹ and Esra Yurt MSc² 

¹Atatürk Medical Vocational Training School, İzmir, Turkey and ²Ege University Pre-Hospital Emergency Health Services and Disaster Management Program

Abstract

Objective: This study aimed to evaluate the mental trauma and coping status of emergency health care workers who responded to the earthquake affected people in the October 2020 Izmir earthquake.

Methods: This was a descriptive and cross-sectional study. State-Trait Anxiety Inventory (STAI), Posttraumatic Cognition Inventory, Strategies for Coping with Earthquake Stress Scale (SCESS), and Post-earthquake Trauma Level Determination Scale (PETLDS) were used. Independent t-test, analysis of variance, Tukey's test, and Pearson's correlation analysis were used for statistical analysis. Statistical significance was accepted as $P \leq 0.05$.

Results: A total of 113 health-care professionals participated in the study. There is a positive and high degree of correlation between the religious coping sub-dimension of the SCESS and the PETLDS ($P < 0.001$). There was a moderate positive correlation between the religious coping sub-dimension of the SCESS and the state anxiety ($P < 0.001$) and trait anxiety ($P < 0.001$) subscales of the STAI.

Conclusions: In our study, mental trauma and post-traumatic coping levels of health-care workers who were exposed to earthquakes differed according to age, gender, and educational status. We found that post-earthquake trauma levels were directly related to state and trait anxiety levels and religious coping levels. We found that, after the earthquake, social support seeking behavior, state and trait anxiety levels increased; negative cognitions about self, negative cognitions about the world, and self-blame increased.

Practice implications: This study draws attention to the evaluation of emergency health workers who work at the forefront of disasters in terms of subsyndromal post-traumatic stress disorder (PTSD). It is recommended to make plans for psychologically protective and preventive interventions for health-care workers especially after disasters.

© The Author(s), 2024. Published by Cambridge University Press on behalf of Society for Disaster Medicine and Public Health, Inc. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

SDMPH
SOCIETY FOR DISASTER MEDICINE & PUBLIC HEALTH

In recent years, many disasters have occurred around the world. One of the events that affect human life is earthquakes. Compared with other disasters, earthquakes are uncontrollable, but its risk to affect human health is controllable by disaster risk reduction, such as building with appropriate building code, and anti-seismic structure. It affects more people by causing unwarranted damage more quickly.¹ Earthquakes are without a doubt one of the most life-threatening, catastrophic, and uncontrollable disasters. Every year, half a million earthquakes are estimated to occur around the world, but most of them go unnoticed because they strike remote areas or are too small. People suffer a variety of losses following an earthquake, depending on its size.² Major disasters can have a significant impact on populations and health systems, particularly in low- and middle-income countries that are unprepared. Early intervention in major disasters, which is used for disasters that cause extensive destruction and damage, is important. Major disaster acute response activities often last only a few weeks. Disasters, on the other hand, have long-term consequences, and disaster management entails more than just quick fixes.³ Disasters remain as events that society remembers when they happen and in the future. Even years later, people who were exposed to disasters still experience the effects of that day. The stress factor is also defended as a reaction mechanism of the person. It should not be overlooked that disasters such as earthquakes can cause acute distress but also lead to chronic stress through long-term deterioration in social and economic conditions.⁴

The Izmir earthquake, which occurred on October 30, 2020, caused loss of life and property in Bornova Bayraklı districts.⁵ It was determined that the epicenter of the earthquake was Samos Island near Seferihisar, the earthquake magnitude was 6.9, and the earthquake lasted 15.68 s.

The Bayraklı district, which was heavily damaged in the earthquake, is approximately 70 kilometers away from the epicenter. As a result of this earthquake, 117 people died, 1034 people were injured, and 15,000 people were left homeless.⁶

Some survivors suffer from long-term distress and psychopathology, such as post-traumatic stress disorder (PTSD), depression, and other mental illnesses. The most prevalent of these negative psychological effects is PTSD. It has been shown that the consequences of being exposed to a disaster, such as the presence of physical injury, fear of death, loss of loved ones, and property damage, are stronger predictors of PTSD than the type of disaster.⁷ The factors that occur before, during, and after a disaster have an impact on psychological problems that arise because of the disaster. People are influenced by a variety of factors, including personal, social, and cultural factors.⁸

PTSD is a common psychiatric disorder caused by an event or series of events that are so painful or stressful that they pose a significant risk to a person's life. Re-experiencing aspects of the traumatic event, feelings of helplessness, intense fear, frightening dreams, or avoiding the source of the trauma are all possible symptoms. PTSD may also experience physical symptoms, such as increased blood pressure and heart rate, fatigue, muscle tension, nausea, joint pain, headaches, back pain, or other types of pain. These symptoms have a significant impact on the affected people's everyday life.⁹ More severe exposure to a traumatic event is associated with more pronounced PTSD symptoms.⁸

Exposure to trauma in PTSD can result from first-hand experience, witnessing traumatic events, or learning of traumatic events that have occurred to close family members or peers. Anyone who feels fear, helplessness, or fear after a natural disaster such as an earthquake can be injured or even killed. Diagnosis of PTSD requires continuous recovery of the event, avoidance symptoms, adverse changes in general responsiveness, arousal, and increased responsiveness.²

Although many people have been exposed to traumatic events throughout their lives, some may meet the criteria for PTSD. But the majority develop sub-PTSD symptoms. Subthreshold symptoms are also considered clinically significant and may require treatment. Individuals with sub-PTSD exhibit higher rates of functional impairment of acute stress symptoms than healthy groups, although not as severely as those who meet the criteria for full PTSD. Insomnia, irritability, anxiety, startle reactions, and social withdrawal symptoms are common.^{10,11} It has been determined that sub-PTSD poses a risk for developing major depression, agoraphobia, substance use disorder and PTSD. It can be detected more frequently in military personnel, disaster relief workers, and health personnel. Therefore, considering the possibility of sub-syndromal PTSD in the days and weeks following trauma, clinicians may provide an opportunity for early treatment and a better prognosis.^{12,13}

In line with this information, it is essential to determine the coping levels of emergency service workers working in intense conditions due to exposure to possible trauma. Obtaining this information will play a critical role in preventing mental disorders that may occur due to the double position of emergency service personnel, both living the disaster and responding to the disaster. This study aimed to determine the psychological trauma and coping levels of the emergency service workers who were exposed to the 2020 İzmir earthquake and lived.

Methods

Study Design and Sample

This descriptive study was conducted on the nurses and physicians working in the emergency service who provided medical intervention to earthquake affected people in the 2020 İzmir earthquake (EQ-2020-000215-TUR).

The research population consisted of 55 men and 70 women between the ages of 18 and 65, consisting of physicians and nurses working in the emergency department of Ege University Faculty of Medicine. In a previous similar study, the sample size was calculated with the G*Power 3 program, taking into account the mean and standard deviation values of trauma variables in health-care workers.¹⁴ The number of individuals to be sampled was 125, with a 95% confidence interval and a 5% margin of error. Initially, 128 people were included in the study and were interviewed. Then, 15 doctors and nurses were excluded from the study because they filled in the scales incompletely. The study was completed with a total of 113 doctors and nurses. After the purpose of the research was explained to the participants, they were informed that participation in the study was completely voluntary and that they could withdraw from participating if desired. It was reported that the data of those who withdraw from the study would not be used in the research.

Inclusion criteria for the study were volunteering to participate in the research and working as a physician or nurse in the emergency department, and being on duty at the hospital during the İzmir earthquake. Exclusion criteria from the study included working outside the emergency department, being on leave on the day of the earthquake, and having a chronic mental illness (serious mental disorder, eg, schizophrenia, bipolar affective disorder).

Data Collection

A Google form was used to create the study questions. Participants were notified of the event through WhatsApp. The data were taken between September 15 and October 15, 2021. The survey is set up so that each participant can only fill out 1 form with their Google account. Participants who agreed to participate in the study were able to continue by clicking the "I agree to participate in the study" button before filling out the questionnaires. Consent was sought in this manner in the digital world. Participants who agreed to participate in the survey completed the data collecting tools online. The questionnaire was completed while taking the necessary precautions to ensure that no more than 1 response was allowed.

Data Collection Tools

Survey data is an online survey using the Sociodemographic data form, Strategies for Coping with Earthquake Stress Scale (SCCESS), Post-earthquake Trauma Level Determination Scale (PETLDS), State-Trait Anxiety Inventory (STAI), and Post-Traumatic Cognition Inventory (PTCI).

Socio-Demographic Information Form: The form developed by the researchers included diagnostic questions such as age, gender, marital status, educational status, occupation, history of psychiatric illness, history of mental illness in the family, and years of professional experience.¹⁵

Strategies for Coping with Earthquake Stress Scale (SCCESS): The scale measures people's coping strategies with earthquake stress. The scale was developed by Yöndem and Altay.¹⁶ There are 16 questions in the content of the scale. The scale consists of

3 sub-dimensions: Religious Coping (RC) (items 1, 2, 3, 4, and 5), Positive Reappraisal (PR) (items 6, 7, 8, 9, 10, and 11), and Seeking Social Support (SSS) (12, 13, 14, 15, and 16) items). It is a 4-point Likert-type scale. (1 = Never; 4 = Always). There is no total score. The score range for the religious coping subscale is 5-20 points, 6-20 points from the positive reappraisal subscale, and 5-20 points from the social support seeking subscale. A high score indicates that the individual uses that coping strategy more, while a low score indicates that the individual uses it less. For the reliability study of the scale, the Cronbach alpha internal consistency reliability coefficient was calculated as 0.85.

Post-Earthquake Trauma Level Determination Scale (PETLDS): PETLDS is a tool developed by Tanhan and Kayri to measure the stress levels experienced by individuals after an earthquake.¹⁷ It consists of 20 questions in total. sub-PTSD symptoms were scaled with a 5-point Likert scale. Expressions in the Likert style are in the form of; “strongly disagree,” “little agree,” “moderately agree,” “strongly agree,” and “completely agree.” The lowest score that can be obtained from the scale is 20, and the highest score is 100. In this context, low, medium, and high levels of posttraumatic stress were examined with posttraumatic CHAID analysis. CHAID analysis builds a predictive model, or tree, to help determine how variables best merge to explain the outcome in the given dependent variable. A score of 45 and below indicates no stress, 46-62 indicates intermediate stress, and a score of 63 and above indicates high stress. The increase in the scores obtained from the scale shows that the individuals’ level of being affected by the earthquake also increases. The internal consistency coefficient (Cronbach’s alpha) of the PETLDS was found to be .87.

State-Trait Anxiety Inventory (STAI): STAI: Spielberger et al. (1970) developed this inventory, which consists of 2 subscales, a trait and a state, each with 20 questions.¹⁸ It is appropriate for those over the age of 14. The State Anxiety Scale (SAS) determines how a person feels at a given time and under specific circumstances. The Trait Anxiety Scale (TAS) assesses how a person feels regardless of the scenario or circumstances. Öner and Le Compte conducted a study on the STAI Turkish adaptation, validity, and reliability.¹⁹ The highest score of 3 is given when the presence of these feelings is reported as many, and the lowest score of 1 is given if it is reported that they are absent. A score of 36 or below for the STAI subscales indicates no anxiety, 37-42 indicates mild anxiety, and a score of 43 and above indicates high anxiety. The overall score is used in the inventory evaluation. A higher score suggests that the person more anxious.

Post-traumatic Cognition Inventory (PTCI): This is a Likert self-assessment scale consisting of 36 items from 1 to 7. Scores from the scale range from 36 to 252. Foa et al. administered the scale to a sample including clinically diagnosed PTSD patients. The total score of the scale is 133 as the cut-off point for PTSD.²⁰ A high-scale total score indicates the density of false cognition associated with the traumatic experience.²⁰ Translated into Turkish by Güleç et al., validity and reliability studies were carried out.²¹ The scale has 3 subscales: Negative Perception of Oneself (NPO), Negative Perception of the World (NPW), and Self-Blame Subscale (SB). The internal consistency coefficient of the scale is 0.95.

Data Analysis

For statistical analysis, SPSS (IBM SPSS Statistics version 22.0; SPSS Inc., Chicago, IL) was used. The continuous variables were given an arithmetic mean and standard deviation, while the

categorical variables were given numbers and percentages. The normal distribution of data was determined using the Kolmogorov-Smirnov test. The link between the independent and dependent variables was determined using T- and F-tests. Reliability analysis was used to establish the scales’ reliability coefficients (Cronbach’s alpha). A 95 percent confidence interval was used to assess the results. For all analyses, statistical significance was considered as a *P*-value of 0.05.

Ethical Considerations

All procedures performed in studies involving human participants followed the ethical standards of institutions and/or national research committees and the 1964 Declaration of Helsinki and subsequent amendments or equivalent ethical standards. Before starting the study, it was approved by the Health Science University and the Institutional Review Board for Non-Intervention Clinical Studies of the Ministry of Health of Turkey (Number 21-7.1T/22.). The people involved in the study were informed of the purpose, methodology, expected benefits, and predictable risks of the study and obtained written consent. Informed consent was obtained from all individual participants in the study.

Results

Socio-demographic Findings

In the research, 113 people were reached and data collection was made. Of the people whose data were collected in the study, 66 (58.4%) were female and 47 (41.6%) were male. The age range of most of the participants is in the 26-30 age group. Additionally, 75 nurses (66.4%) and 38 physicians (33.6%) working in the emergency department participated in the study, and 78 (69%) of the participants have been working for less than 5 y. The rate of participants having a history of mental illness is between 3.5% and 4 people. There were 8 people (7.1%) with a family history of psychiatric illness. In the study, when the mental health status of the participants before the earthquake was evaluated, it was determined that 4 people were treated for psychiatric illness (serious mental disorder-schizophrenia, bipolar affective disorder were excluded) before the earthquake. It was determined that 2 of them were treated for somatic symptom disorder and the others for bulimia nervosa, insomnia disorder (Table 1).

Evaluation of the MEAN SCORE of the Scales

PETLDS mean scores show that the participants have intermediate stress. Participants scored the highest number of points in PR from the subscales of the SCESS and used PR coping the most; they used SSS and RC coping strategies less frequently, respectively. The average scores of the participants on STAI’s TAS and SAS scales were high. Participants scored above the cutoff score on the PTCI. This showed that post-traumatic negative cognitions were high (Table 2).

The Cronbach alpha value for SCESS, PETLDS, STAI, PTCI scales were found to be 0.80, 0.95, 0.53, 0.96, respectively.

Evaluation of PETLDS, SCESS, STAI, PTCI Scales and Sub-scales Scores According to Sociodemographic Characteristics

A significant difference was found between the groups in the positive reappraisal subscale of the SCESS according to the gender of the participants (*P* < 0.05). It was determined that men got higher scores on the scale. There is no significant difference

Table 1. Socio-demographical characteristics of participants (*N*: 113)

	<i>N</i>	%
Gender		
Woman	66	58.4
Man	47	41.6
Marital status		
Married	47	41.6
Single	65	57.5
Divorces		0.9
Age		
20–25	22	19.5
26–30	62	54.9
31–40	27	23.9
40 and above	2	1.8
Profession		
Nurse	75	66.4
Physician	38	33.6
Years of professional experience		
0–5	78	69.0
6–10	26	23.0
11–20	8	7.1
20 and above	1	0.9
Educational status		
University	81	71.7
Graduate	27	23.9
Doctorate	5	4.4
Diagnosed psychiatric illness		
Yes	4	3.5
No	109	96.5
Family history of mental illness		
Yes	8	7.1
No	105	92.9
Total	113	100

between the RC and SSS subscales of SCESS, PETLDS, STAI and subscale, PTCI and subscale scores according to the gender of the participants ($P > 0.05$) (Table 2).

When the scores obtained from the scales according to age were evaluated, a significant difference was found between the groups in the SCESS positive reappraisal subscale. It was determined that this difference was due to the 26–30 age group getting the highest score with the Bonferroni further analysis ($P < 0.05$) There is no significant difference between the RC and SSS subscales of SCESS, PETLDS, STAI and subscale, PTCI and subscale scores according to the age group of the participants ($P > 0.05$) (Table 3).

When the scale analysis was examined according to educational status, a statistically significant difference was found in the undergraduate and graduate groups between academic level and the positive reappraisal sub-dimension of the SCESS ($P = 0.042$; $P < 0.05$). With the Bonferroni advanced analysis performed to determine which group caused the difference, it was determined that the difference was because the university group scores were higher than those of the graduate group ($P = 0.042$; $P < 0.05$). There is no significant difference between the RC and SSS subscales of SCESS, PETLDS, STAI and subscale, PTCI and subscale scores according to the educational status of the participants ($P > 0.05$) (Table 4).

SCESS, PETLDS, STAI, PTCI scores did not differ according to marital status, profession, years of professional experience, diagnosed psychiatric illness family history of mental illness of the participants ($P > 0.05$).

Evaluation of the Correlation Among the PETLDS and SCESS, PTCI, STAI Scales

There is a positive and high degree of correlation between the religious coping sub-dimension of the SCESS and the PETLDS ($P < 0.001$). As the level of impact from the earthquake increased, the use of religious coping increased. There was a moderate positive correlation between the religious coping sub-dimension of the SCESS and the state anxiety ($P < 0.001$) and trait anxiety ($P < 0.001$) subscales of the STAI. As the level of impact from the earthquake increased, the level of anxiety also increased. In other words, those most affected by the earthquake have higher anxiety and use more religious coping mechanism.

A moderate correlation was found between the religious coping sub-dimension of SCESS and the negative perception about oneself sub-scale of PTCI ($P < 0.001$). There is a high degree of positive correlation found between the social support-seeking sub-dimension of SCESS and the negative perception about oneself subscale ($P < 0.001$), negative perception about the world subscale ($P < 0.001$), and self-blame ($P < 0.001$) subscale of PTCI ($P < 0.001$).

A high degree of positive correlation was also found between the social support seeking subscale of SCESS and the state anxiety ($P < 0.001$) and trait anxiety ($P < 0.001$) subscales of STAI.

There is a positive correlation between PETLDS and the negative perception about oneself subscale ($P < 0.001$), negative perception about the world subscale ($P < 0.001$), and self-blame of the PTCI. There is a significant positive correlation between the PETLDS and trait anxiety ($P < 0.001$) sub-scales of the STAI. The state anxiety ($P < 0.001$) and trait anxiety ($P < 0.001$) sub-scales of the STAI and negative perception about oneself, negative perception about the world, self-blame subscales of PTCI, and also showed a high degree of positive correlation.

Effect of Variables on PETLDS and SCESS Scores

To determine the variables that predict the coping strategies of the participants in our study, variables such as STAI, PETLDS PTCI, age group, education, and gender were analyzed. PTCI, gender, and education significantly predict SCESS, together explaining 48% of the total variance ($R = .68$; $R^2 = .48$; $F = 52.15$; $P < 0.05$). According to the standardized regression coefficient (β), PTCI ($\beta = .60$) took precedence over gender ($\beta = .23$) and education ($\beta = .17$) in the relative importance of the predictor variables on SCESS. Looking at the significance tests of the regression coefficient, it is seen that the predictive variables PTCI ($P < 0.05$), gender ($P < 0.05$), and education ($P < 0.05$) are significant predictors on SCESS. Additionally, Cohen's f^2 value was calculated to determine the effect size and $f^2 = .83$ for the anxiety variable. In terms of Cohen (1988) criteria, it can be said that the effect size in the current study is high.

When evaluating the variables that predict the status of being affected by the earthquake; PTCI, STAI, religious coping sub-dimension of the SCESS significantly predict PETLDS, together explaining 52% of the total variance ($R = .57$; $R^2 = .52$; $F = 34.18$; $P < 0.05$).

Table 2. Average scores from the scales and their evaluation by gender

Scales	(X ± SD)	Scale	Gender	(X ± SD)	P-Value
SCESS		SCESS			
RC	11.67 ± 2.56	RC	Woman	11.67 ± 2.56	0.770
			Man	11.61 ± 2.59	
				11.76 ± 2.54	
PR	14.46 ± 3.42	PR	Woman	14.46 ± 3.42	0.040
			Man	13.90 ± 3.18	
				15.26 ± 3.64	
SSS	13.05 ± 2.82	SSS	Woman	13.05 ± 2.82	0.250
			Man	12.80 ± 2.68	
				13.44 ± 3.02	
PETLDS	49.69 ± 19.02	PETLDS	Woman	49.55 ± 18.98	0.925
			Man	49.90 ± 18.31	
STAI		STAI			
TAS	44.67 ± 11.87	TAS	Woman	45.67 ± 12.37	0.655
			Man	45.12 ± 10.41	
SAS	44.66 ± 15.17	SAS	Woman	44.14 ± 14.38	0.534
			Man	44.22 ± 13.75	
PTCI		PTCI			
NPO	59.41 ± 29.10	NPO	Woman	58.15 ± 3.52	0.871
			Man	57.82 ± 2.71	
NPW	35.33 ± 13.24	NPW	Woman	36.42 ± 8.22	0.947
			Man	36.12 ± 5.34	
SB	11.20 ± 6.77	SB	Woman	11.73 ± 4.74	0.714
			Man	12.01 ± 3.54	

Abbreviations: Mean (X), standard deviation (SD), religious coping (RC), positive reappraisal (PR), seeking social support (SSS), Trait Anxiety(TAS), State Anxiety (SAS), negative perception about oneself (NPO), negative perception about the world (NPW), self-blame (SB), analysis of the Post-Traumatic Cognitions Inventory (PTCI), Post-Earthquake Trauma Level Determination Scale (PETLDS).

Discussion

In this study, mental trauma and coping with trauma situations in emergency service workers, who gave medical intervention to earthquake victims in the 2020 İzmir earthquake, were evaluated. In the study, the participant was determined to be intermediate traumatic and anxiety levels of the participants were above the average.^{17,19} Studies have shown that emergency room personnel are at high risk for mental health problems, including anxiety, depression, and psychological distress.^{22,23} In particular, it has been determined that post-traumatic stress disorder is approximately 2 to 3 times more common among various groups of emergency service personnel (eg, ambulance specialists, rescue workers) than in the general population.²² We defined as traumatic according to the results of the scale, as a clinical this may not be sufficient to meet the symptoms of PTSD, more often it may be considered sub-PTSD. This study has implications for the nosology and symptoms of PTSD. We believe that the part of health-care workers in our study can be evaluated as subsyndromal PTSD.²³ Identification of subsyndromal PTSD provides the opportunity for early treatment and better prognosis. As McFarlane and colleagues argue, subsyndromal PTSD should not be viewed merely as a normal response to an abnormal event.²⁴ Post-traumatic stress symptoms, even at the subsyndromal level, have been found to be associated with increased anxiety symptoms with significant social and occupational impairment, as is full PTSD. Administrative steps may be necessary to ensure that health-care personnel identified as having subsyndromal PTSD receive trauma-focused treatment and long-term follow-up.^{25,26}

When exposed to more stress, they should be considered as “symptoms that are a harbinger of later mental disorders.” We found that the younger participants and those with the highest education level (doctoral) were more affected by the earthquake, which may make them more at risk in terms of trauma. We found that men, the youngest group (20–25 y old), those with a master’s degree, use “positive reappraisal” as a coping mechanism. The subsyndromal group is not defined as completely non-pathological, because clinically significant impairments are significant. According to Mylle and Maes (2004), taxonomy is important because the diagnosis also defines the medico-legal positions of the subjects and the position of individuals with subthreshold PTSD is uncertain.²⁷ However, in the literature, PTSD symptoms are higher in the subsyndromal group compared with the healthy ones. Although not a clear diagnostic category, subthreshold PTSD may not be part of a normative response to trauma but should be considered because it can cause significant impairment and be chronic. It was researched to determine the stress level left on the health workers who experienced and responded to the 30 October İzmir earthquake. Because Turkey is in an earthquake zone, it faces damages resulting from disasters. It is necessary to take preventive measures to minimize the losses and damages that may occur after the earthquake and to prevent mental health problems that may arise after the quake.²⁸

A relationship was found between postearthquake trauma and negative perceptions of the individual and the world, self-blame. According to the theoretical premise of cognitive models of psychopathology, dysfunctional cognitive processing in the form

Table 3. Evaluation of SCESS, PETLDS, STAI, and PTCI according to age

Scales	Age (y)	(X ± SD)	P-Value
SCESS			
RC	20-25	11.31 ± 2.14	0.241
	36-30	12.05 ± 2.67	
	31-40	11.17 ± 2.56	
PR	20-25	13.86 ± 3.22	0.030
	26-30	15.23 ± 3.57	
	31-40	13.34 ± 2.93	
SSS	20-25	12.81 ± 2.88	0.253
	26-30	12.77 ± 2.77	
	31-40	13.82 ± 2.84	
PETLDS	20-25	52.13 ± 20.16	0.664
	26-30	48.17 ± 19.13	
	31-40	50.18 ± 18.31	
STAI			
TAS	20-25	44.13 ± 11.34	0.451
	26-30	44.18 ± 12.31	
	31-40	44.25 ± 12.58	
SAS	20-25	46.16 ± 11.38	0.801
	26-30	45.85 ± 13.28	
	31-40	45.98 ± 14.87	
PTCI			
NPO	20-25	59.13 ± 2.43	0.871
	26-30	59.12 ± 3.74	
	31-40	58.88 ± 2.15	
NPW	20-25	36.42 ± 8.24	0.912
	26-30	36.12 ± 5.32	
	31-40	35.87 ± 4.78	
SB	20-25	11.13 ± 3.75	0.997
	26-30	11.51 ± 4.85	
	31-40	11.45 ± 3.77	

Abbreviations: Mean (X), standard deviation (SD), religious coping (RC), positive reappraisal (PR), seeking social support (SSS), Trait Anxiety(TAS), State Anxiety (SAS), negative perception about oneself (NPO), negative perception about the world (NPW), self-blame (SB), analysis of the Post-Traumatic Cognitions Inventory (PTCI), Post-Earthquake Trauma Level Determination Scale (PETLDS).

of distorted evaluations is very important in the emergence of psychological disorders. Negative post-traumatic cognitions are important in the development, persistence, and treatment of PTSD. It has been determined that these cognitions contribute to the individual's sense of "present threat," thereby increasing vulnerability to develop PTSD.²⁹⁻³¹ Levels of postearthquake trauma were found to be linked to state and trait anxiety, as well as religious coping. Religious coping was found to be more prevalent, especially when the individual's negative self-perceptions increased. When people are confronted with trauma, crisis, or stressful events, they often use a variety of coping strategies. Coping can be defined as the thoughts and behaviors that individuals use to manage the internal and external demands of situations that are considered stressful. Spirituality may be quite beneficial in the aftermath of a terrible event. When faced with distress or trauma, most people turn to religion or spirituality as a coping mechanism, and they often find it to be effective.^{32,33} It appears to be the primary mechanism through which spirituality and religion influence psychological outcomes.³² When it comes to dealing with stress and major life events, religion's ability to provide meaning in life

Table 4. Evaluation of SCESS, PETLDS, STAI, and PTCI according to educational status

Scale	Status	(X ± SD)	P-Value
SCESS			
RC	University	10.81 ± 3.17	0.945
	Master	11.05 ± 1.18	
	Doctorate	11.12 ± 3.77	
PR	University	14.00 ± 3.34	0.04
	Master	15.92 ± 3.54	
	Doctorate	14.80 ± 2.58	
SSS	University	13.71 ± 3.45	0.318
	Master	13.15 ± 1.88	
	Doctorate	12.99 ± 2.84	
PETLDS	University	46.20 ± 2.75	0.592
	Master	45.25 ± 3.73	
	Doctorate	50.32 ± 2.85	
STAI			
TAS	University	43.95 ± 11.34	0.548
	Master	44.18 ± 14.24	
	Doctorate	44.25 ± 12.54	
SAS	University	45.24 ± 25.14	0.769
	Master	44.19 ± 39.91	
	Doctorate	45.12 ± 13.77	
PTCI			
NPO	University	59.25 ± 3.77	0.958
	Master	58.19 ± 3.15	
	Doctorate	58.18 ± 3.77	
NPW	University	36.42 ± 8.24	0.922
	Master	36.12 ± 5.32	
	Doctorate	36.14 ± 3.22	
SB	University	11.87 ± 2.89	0.898
	Master	12.14 ± 3.55	
	Doctorate	11.92 ± 6.89	

Abbreviations: Mean (X), standard deviation (SD), religious coping (RC), positive reappraisal (PR), seeking social support (SSS), Trait Anxiety(TAS), State Anxiety (SAS), negative perception about oneself (NPO), negative perception about the world (NPW), self-blame (SB), analysis of the Post-Traumatic Cognitions Inventory (PTCI), Post-Earthquake Trauma Level Determination Scale (PETLDS).

is especially apparent. Religious coping can be defined as "the degree to which religion is a part of the process of understanding and coping with critical life events".³⁴

Social assistance has been shown in studies to help people recover after a disaster. One of the most commonly reported coping mechanisms among survivors is seeking help. Individuals' perceptions about their family, friends, and society are referred to as perceived social. Understanding and satisfying the needs of disaster-affected people requires a familial context. Social support is a resource that changes the negative perception of self and affects psychological and physical well-being, psychosocial adaptation, and personal gains.^{32,35} It has been observed that increased social support improves mental well-being, particularly in cases of increased anxiety. It was observed that there was an increase in state and trait anxiety levels, negative cognitions about oneself, negative cognitions about the world, and self-blame increased the behavior of seeking social support after the earthquake in the study participants.

Evaluating the scores obtained from the age scale showed that there was a significant difference between the groups in the positive reassessment subscales of SCESS, with the 26- to 30-y-old group receiving the highest score. According to Ericsson's Psychosocial Theory, people of different ages encounter different life events, go through a series of life stages of psychosocial development, and the major social emotions at each life stage. It solves the conflict well.³⁶ Folkman and Lazarus (1980) have a contextual theory of aging.³⁷ They argue that different types of stressors appear with age, and that these differences in stressors affect coping strategies and health outcomes. Adults' stress exposure differs depending on their life context, according to the contextual theory of Folkman and Lazarus (1980).³⁷ As a result, it exhibits a variety of stress levels. According to 1 study, as people get older, they are less affected by stress and anger.³⁸ According to Socioemotional Selectivity Theory, which explains the change in emotional functioning with advancing age, it has been found that, as perceptions of time alive decrease (a natural consequence of aging), people prioritize achieving emotion-focused goals over knowledge-based goals.³⁹ In another study investigating the level of stress during the COVID-19 (coronavirus disease 2019) period, it is found that the stress level decreases with increasing age; the adolescent population perceives the highest amount of stress, while middle-aged adults perceive the least amount of stress.⁴⁰ According to another study, post-traumatic growth and post-traumatic stress have a low-level significant positive relationship. This demonstrates that those who have growth because of a traumatic event also have stress symptoms.⁴¹

According to the gender of the participants, it was determined that men got higher scores in the positive PR subscale of SCESS. In addition to the effects of the COVID-19 pandemic on physical health, it can also have several adverse effects on people's mental health. However, these adverse effects may differ according to gender. In a previous study, coping with negative situations in life was evaluated in young adults, and it was found that young male adults exhibit faster-coping strategies.⁴² It has been determined that men can make decisions faster than women and have more tendency to take risks. It has been determined that women have difficulty coping with negative situations, experience more anxiety, and show depressive symptoms.⁴³ At the same time, studies in the literature have found a significant relationship between young age and the female gender with the probability of developing these problems.⁴⁴ The sample formed with the majority of the middle age group may have affected the result. Again, in this context, the fact that the male group was significant even though the female group in the sample group was high revealed results incompatible with the literature.

It was determined that postgraduate graduates received high scores in the positive re-evaluation sub-dimension of the SCESS according to their education level. In the literature, Baysak et al. (2019) also found that positive re-evaluation increased as the level of education increased; In this context, the result supports the literature.⁴⁵ After disasters with devastating effects such as earthquakes, people's trauma response levels and coping levels may differ.⁴⁶ According to Erdener (2019), it was concluded that those who graduated from university and graduate school experienced more secondary trauma.⁴⁶

At the end of the study, there was no significant difference between the occupational groups of physicians and nurses in terms of coping with stress. Many factors cause occupational stress. Occupational stress has become a growing problem in health-care workers in recent years.⁴⁷ The reason why there was no difference

between occupational groups in the study may be because the sample mainly consisted of nurses and that both occupational groups had the same intense working conditions. In addition, both occupational groups act with high professional responsibility when intervening with the patient. There may be similar increases in work intensities, especially during disasters, which may give similar reactions to being affected by the disaster. Different results were also found in the studies. According to a study on mental health, nurses with occupational group-specific differences were found to have the highest mental health disorders and burnout.⁴⁸ According to another study, it was found that, while the state anxiety scores of doctors and nurses are very close to each other, the trait anxiety scores of nurses are higher than doctors. Still, this difference was not statistically significant.⁴⁹ A disaster is a natural event that has irreversible consequences. The psychological dimension of health workers affected by all the damage and who are involved in the response is among the issues that need to be addressed in detail. Implementation of recovery programs for health workers after disasters have been seen among the methods that can be effective. According to a published study, it was emphasized that health systems should support the health and well-being of doctors and nurses with evidence-based interventions to improve community health and increase the quality and safety of care must be provided.⁵⁰

In the study, no significant difference was found in the SCESS in terms of past mental health problems and having a family member with a psychiatric illness. In fact, those with a history of mental illness are often expected to have high postdisaster stress levels. Understanding the traumatic response within a psychopathological framework is vital. It is rarely possible for PTSD to occur in isolation following a traumatic event. The perception of isolation or lack of social connection with others is associated with loneliness. May evoke psychological responses that often occur in situations such as depression, anxiety, or substance use.²⁹ In the literature, it was determined that people with a previous history of mental illness showed depressive symptoms after the earthquake. It is seen that those who have a history of mental illness in the past have high levels of help expectations and postearthquake stress.^{51,52} In another study, it was reported that there might be differences in the level of the psychological impact of individuals in coping with trauma after an earthquake. It has been determined that, after disasters such as earthquakes that occur suddenly and whose consequences are unpredictable, there may be psychological problems that improve in a few weeks.⁵¹ According to Baysak et al. (2019), it was seen that mental health problems experienced in the past were effective in coping with postearthquake stress.⁴⁵ The low rate of participants with a history of mental illness and a family history of mental illness in the study may have affected the result.

Limitations

This study has several limitations. When analyzing the study's findings, it is important to keep in mind the study's limitations. The most significant limitation is that data were collected by means of self-report questionnaires with rating scales. This methodology is based on the participants' personal statements, and it is possible that their responses are not always accurate or precise. Second, because the data were gathered through online surveys, not all workers could be reached. Participants who had no prior experience with the Internet and were unable to access the survey may have responded less or not at all. Another limitation

is related to the sample size of the study. Data were collected from health-care personnel working in the emergency department of a hospital. Future studies may be considered to be repeated in more hospitals, with different clinical staff and with a larger sample size. The final limitation is that the study design was cross-sectional, outcomes and predictors include the time and participants when the study data were collected.

Conclusions

In our study, psychological trauma and posttraumatic coping levels of health-care workers exposed to earthquakes differed according to age, gender, and educational status. Our study Participants scored above the cutoff score on the PTCI. This result suggests that the participants may have sub-syndromal PTSD. Sub-syndromal PTSD occurs when persons experience many of the symptoms of PTSD, with some impairment, but do not meet full diagnostic criteria.^{13,20} It is important to conduct a 1-to-1 psychological evaluation for a definitive clinical diagnosis. Anxiety levels have been found to be high. Postearthquake trauma levels were found to be directly related to state and trait anxiety levels and religious coping levels. Again, a relationship was found between religious coping and the individual's negative cognitions. In our study, we found that the participants showed moderate levels of stress symptoms in terms of being affected by the earthquake, and high levels of negative postearthquake cognitions and anxiety. Participants' negative posttraumatic cognitions, gender, and education levels were found to be predictive variables on coping strategies; religious coping strategies, posttraumatic cognitions, anxiety levels were found to be predictive variables on being affected by the earthquake. Considering the increasing frequency of natural disasters around the world, it is important to evaluate the psychological impact of disasters on emergency medical personnel working at the forefront of disasters. Future research should focus on interventions to improve emergency health-care workers' knowledge and competencies in dealing with disaster exposure and disaster situations.

Implications for Health-care Workers' Practice

It emphasizes how much the psychological problems that may arise after a natural disaster affect the health-care workers who work in intense and critical conditions. Undoubtedly, both are responding and experiencing the event after a natural disaster that occurs unexpectedly can cause severe psychological destruction for people. Health-care workers are also among the groups that are most affected by this situation psychologically. It is inevitable for health-care workers to receive both environmental and professional support after this severe trauma. Participating in the in-house training, playing an active role in the practices, and being informed about what can be done due to the event, will play a role in alleviating this situation. It is important to know what to do after the trauma, prepare psychologically for the conditions and conditions of the case, and manage the course of the process well. Our results show that years of experience effectively cope with posttraumatic stress. Accordingly, more training may be required in orientation training groups.

Studies show that health-care workers' anxiety levels are high due to intense working conditions. The addition of psychological stress that may occur after trauma to health-care workers who are already working under extreme conditions can cause severe losses in their lives and is an issue that needs to be rehabilitated.

Policies and plans need to be made to manage the mental problems of health-care personnel after a disaster. It is important that government officials organize systematic, coordinated disaster mental health support services within the scope of postdisaster rehabilitation services to improve the resilience skills of health response workers. Coordination and comprehensive programs to address post-trauma are key to the success of an effective and sustainable mental health program after disaster.

Competing interests. We have no conflict of interest to declare.

References

1. Bavafa A, Khazaie H, Khaledi-Paveh B, *et al.* The relationship of severity of symptoms of depression, anxiety, and stress with sleep quality in earthquake survivors in Kermanshah. *J Inj Violence Res.* 2019;11(2): 225-232. doi: [10.5249/jivr.v11i2.1203](https://doi.org/10.5249/jivr.v11i2.1203)
2. Farooqui M, Quadri SA, Suriya SS, *et al.* Posttraumatic stress disorder: a serious post-earthquake complication. *Trends Psychiatry Psychother.* 2017;39(2):135-143. doi: [10.1590/2237-6089-2016-0029](https://doi.org/10.1590/2237-6089-2016-0029)
3. Goyet S, Rayamajhi R, Gyawali BN, *et al.* Post-earthquake health-service support, Nepal. *Bull World Health Organ.* 2018;96(4):286-291. doi: [10.2471/BLT.17.205666](https://doi.org/10.2471/BLT.17.205666)
4. Khatri GK, Tran TD, Baral S, *et al.* Experiences of an earthquake during pregnancy, antenatal mental health and infants' birthweight in Bhaktapur District, Nepal, 2015: a population-based cohort study. *BMC Pregnancy Childbirth.* 2020;20(1):414. doi: [10.1186/s12884-020-03086-5](https://doi.org/10.1186/s12884-020-03086-5)
5. Çalişkan A, Kaya G. Sociological view of social solidarity practices in earthquake: the example of Izmir earthquake. *J Suleyman Demirel Univ Vis.* 2021;12(32):1052-1077. doi: [10.21076/vizyoner.878817](https://doi.org/10.21076/vizyoner.878817)
6. Karadaş A, Öner E. The effects of alluvial geomorphology on the damage of the Sisam Earthquake in the Bornova Plain. *J Geog.* 2021;(42):139-153. doi: [10.26650/JGEOG2021-872890](https://doi.org/10.26650/JGEOG2021-872890)
7. Yang W, Cui K, Sim T, *et al.* Health-related quality of life and post-traumatic stress disorder in inpatients injured in the Ludian earthquake: a longitudinal study. *Health Qual Life Outcomes.* 2020;18(1):229. doi: [10.1186/s12955-020-01470-5](https://doi.org/10.1186/s12955-020-01470-5)
8. Matsumoto K, Sakuma A, Ueda I, *et al.* Psychological trauma after the Great East Japan Earthquake. *Psychiatry Clin Neurosci.* 2016;70(8):318-331. doi: [10.1111/pcn.12403](https://doi.org/10.1111/pcn.12403)
9. Marthoenis M, Ilyas A, Sofyan H, *et al.* Prevalence, comorbidity and predictors of post-traumatic stress disorder, depression, and anxiety in adolescents following an earthquake. *Asian J Psychiatr.* 2019;43:154-159. doi: [10.1016/j.ajp.2019.05.030](https://doi.org/10.1016/j.ajp.2019.05.030)
10. Mota NP, Tsai J, Sareen J, *et al.* High burden of subthreshold DSM-5 post-traumatic stress disorder in U.S. military veterans. *World Psychiatry.* 2016;15(2):185-186. doi: [10.1002/wps.20313](https://doi.org/10.1002/wps.20313)
11. Highfill-McRoy RM, Levine JA, Larson GE, *et al.* Predictors of symptom increase in subsyndromal PTSD among previously deployed military personnel. *Mil Med.* 2022;187(5-6):e711-e717. doi: [10.1093/milmed/usa/b034](https://doi.org/10.1093/milmed/usa/b034)
12. McLaughlin KA, Koenen KC, Friedman MJ, *et al.* Subthreshold posttraumatic stress disorder in the world health organization world mental health surveys. *Biol Psychiatry.* 2015;77(4):375-384. doi: [10.1016/j.biopsych.2014.03.028](https://doi.org/10.1016/j.biopsych.2014.03.028)
13. Wallace D, Moss AS, Hodges S. Sub-syndromal PTSD: what is important to know in military personnel and veterans? *Australas Psychiatry.* 2020;28(3):254-256. doi: [10.1177/1039856219881965](https://doi.org/10.1177/1039856219881965)
14. Demirtürk Selçuk E, Demirağ BC. The effect of COVID-19 outbreak on secondary traumatic stress and sleep quality in healthcare workers. *Türkiye Klinikleri J Nurs Sci.* 2021;13(3):533-543.
15. Yiğitoğlu GT, Keskin G, Köktaş NÇ. The traumatic impact of the COVID-19 pandemic: the possible role of rumination and uncertainty [published online ahead of print, 2023 Feb 24]. *Curr Psychol.* 2023;1-10. doi: [10.1007/s12144-023-04379-6](https://doi.org/10.1007/s12144-023-04379-6)
16. Yöndem ZD, Eren A. Validity and reliability studies of earthquake stress and coping scale (ESCS). *Turkish Psychol Couns Guide J.* 2008;3(30):60-75.

17. **Tanhan F, Kayri M.** The validity and reliability study of the scale for determining the level of trauma after the earthquake. *Educ Sci Theory Pract.* 2013;13(2):1021-1025
18. **Spielberger CD, Gorsuch RL, Lushene RE.** *Manual for State and Anxiety Inventory.* Consulting Psychologists; 1970.
19. **Öner N, Le Compte WA.** *State-Trait Anxiety Inventory Handbook.* 20th ed. Boğaziçi University Press; 1985:3-5.
20. **Foa EB, Ehlers A, Clark DM, et al.** The posttraumatic cognitions inventory (PTCI): development and validation. *Psychol Assess.* 1999;11(3):303-314.
21. **Güleç M, Kalafat T, Boysan M, et al.** Psychometric properties of the Turkish version of the Posttraumatic Cognitions Inventory (PTCI) in a nonclinical sample. *Arch Neuropsychiatr.* 2013;50:147-153.
22. **Taylor AM, Cocks JH, Ashe MC, et al.** Mental health in young adult emergency services personnel: a rapid review of the evidence. *Int J Disaster Risk Reduct.* 2023;88:103605. doi: [10.1016/j.ijdrr.2023.103605](https://doi.org/10.1016/j.ijdrr.2023.103605)
23. **Lu W, Mueser KT, Yanos PT, et al.** Post-Traumatic Cognitions Inventory (PTCI): psychometric properties in clients with serious mental illness and co-occurring PTSD. *Behav Cogn Psychother.* 2023;51(5):459-474. doi: [10.1017/S1352465823000140](https://doi.org/10.1017/S1352465823000140)
24. **McFarlane AC, Lawrence-Wood E, Van Hooff M, et al.** The need to take a staging approach to the biological mechanisms of PTSD and its treatment. *Curr Psychiatry Rep.* 2017;19(2):10.
25. **Lawrence-Wood E, Baur J, Lawrence A, et al.** The role of inhibitory processes in the relationship between subsyndromal PTSD symptoms and aggressive behaviour *J Psychiatr Res.* 2021;143:357-363. doi: [10.1016/j.jpsychires.2021.09.037](https://doi.org/10.1016/j.jpsychires.2021.09.037)
26. **McLaughlin K, Demyttenaere K, Koenen K, et al.** Threshold versus Subthreshold Posttraumatic Stress Disorder. In Bromet E, Karam E, Koenen K, et al. eds. *Trauma and Posttraumatic Stress Disorder: Global Perspectives from the WHO World Mental Health Surveys.* Cambridge University Press; 2018:273-286.
27. **Mylle J, Maes M.** Partial posttraumatic stress disorder revisited. *J Affect Disord.* 2004;78(1):37-48. doi: [10.1016/s0165-0327\(02\)00218-5](https://doi.org/10.1016/s0165-0327(02)00218-5)
28. **Akalın S, Şakiroğlu M, Eren S.** Examining some variables that predict earthquake prevention behavior: Aydın province example. *Dokuz Eylul Univ J Soc Sci Ins.* 2020;22(3):977-993.
29. **Hyland P, Murphy J, Shevlin M, et al.** Psychometric properties of the Posttraumatic Cognition Inventory within a Northern Ireland adolescent sample. *Br J Clin Psychol.* 2015;54(4):435-449. doi: [10.1111/bjc.12089](https://doi.org/10.1111/bjc.12089)
30. **Mills KL, Ewer P, Dore G, et al.** The feasibility and acceptability of a brief intervention for clients of substance use services experiencing symptoms of post traumatic stress disorder. *Addict Behav.* 2014;39(6):1094-1099. doi: [10.1016/j.addbeh.2014.03.013](https://doi.org/10.1016/j.addbeh.2014.03.013)
31. **van Heemstra HE, Scholte WF, Ehring T, et al.** Contextualizing cognitions: the relation between negative post-traumatic cognitions and post-traumatic stress among Palestinian refugees. *J Cogn Ther.* 2020;13(2):159-172.
32. **Mesidor JK, Sly KF.** Religious coping, general coping strategies, perceived social support, PTSD symptoms, resilience, and post-traumatic growth among survivors of the 2010 earthquake in Haiti. *Ment Health Relig Cult.* 2019;22:130-143.
33. **Park CL, Sacco SJ, Mills MA.** Do religious habits and coping help in the immediate aftermath of a crisis? Relations with Hurricane Katrina evacuees' acute stress symptoms and functional impairment. *Psychol Trauma.* 2019;11(6):563-570. doi: [10.1037/tra0000426](https://doi.org/10.1037/tra0000426)
34. **Pargament KI, Ano GG, Wachholtz AB.** The religious dimensions of coping: advances in theory, research, and practice. In Paloutzian RF, Park CL, eds. *Handbook of the Psychology of Religion and Spirituality.* Guilford Press; 2005:479-495
35. **Samuel R, Burger K.** Negative life events, self-efficacy, and social support: risk and protective factors for school dropout intentions and dropout. *J Educ Psychol.* 2020;112(5):973-986.
36. **Erikson E.** *The Life Cycle Completed.* Norton & Company; 1982.
37. **Folkman S, Lazarus RS.** An analysis of coping in a middle-aged community sample. *J Health Soc Behav.* 1980;21(3):219-239.
38. **Stone AA, Schwartz JE, Broderick JE, et al.** A snapshot of the age distribution of psychological well-being in the United States. *Proc Natl Acad Sci U S A.* 2010;107(22):9985-9990. doi: [10.1073/pnas.1003744107](https://doi.org/10.1073/pnas.1003744107)
39. **Sliwinski MJ, Freed S, Scott SB, et al.** Does chronic stress moderate age differences in emotional well-being? Testing predictions of strength and vulnerability integration. *J Gerontol B Psychol Sci Soc Sci.* 2021;76(6):1104-1113. doi: [10.1093/geronb/gbaa174](https://doi.org/10.1093/geronb/gbaa174)
40. **Singh S, Mani Pandey N, Datta M, et al.** Stress, internet use, substance use and coping among adolescents, young-adults and middle-age adults amid the 'new normal' pandemic era. *Clin Epidemiol Glob Health.* 2021;12:100885. doi: [10.1016/j.cegh.2021.100885](https://doi.org/10.1016/j.cegh.2021.100885)
41. **Kardaş F, Tanhan F.** Investigation of post-traumatic stress, post-traumatic growth and hopelessness levels of university students who experienced the Van earthquake. *J Yüzüncü Yıl Univ Fac Educ.* 2018;15(1):1-36.
42. **Salas-Rodríguez J, Gómez-Jacinto L, Hombrados-Mendieta I.** Life history theory: evolutionary mechanisms and gender role on risk-taking behaviors in young adults. *Pers Individ Dif.* 2021;175:110752.
43. **Yang Z, Luo Y, Zhou Q, et al.** COVID-19-related stressors and depression in Chinese adolescents: the effects of life history strategies and gender. *J Affect Disord.* 2022;304:122-127. doi: [10.1016/j.jad.2022.02.060](https://doi.org/10.1016/j.jad.2022.02.060)
44. **Serrano-Ripoll MJ, Meneses-Echavez JF, Ricci-Cabello I, et al.** Impact of viral epidemic outbreaks on mental health of healthcare workers: a rapid systematic review and meta-analysis. *J Affect Disord.* 2020;277:347-357. doi: [10.1016/j.jad.2020.08.034](https://doi.org/10.1016/j.jad.2020.08.034)
45. **Baysak E, Eroğlu ZM, Utku Ç, et al.** Evaluation of post-traumatic stress disorder, burnout and coping styles in emergency and intensive care unit employees. *J Clin Psychiatry.* 2019;22(1):36-47.
46. **Erdener M.** Investigation of Psychological Resilience and Secondary Traumatic Stress Levels of Professionals Working in the Field of Disaster [master's thesis]. Ankara, Turkey: Hacettepe University Institute of Social Sciences, Department of Social Work; 2019.
47. **Trifunovic N, Jatic Z, Kulenovic AD.** Identification of causes of the occupational stress for health providers at different levels of health care. *Med Arch.* 2017;71(3):169-172. doi: [10.5455/medarch.2017.71.169-172](https://doi.org/10.5455/medarch.2017.71.169-172)
48. **Diekmann K, Böckelmann I, Karlsen HR, et al.** Effort-reward imbalance, mental health and burnout in occupational groups that face mental stress. *J Occup Environ Med.* 2020;62(10):847-852. doi: [10.1097/JOM.0000000000001978](https://doi.org/10.1097/JOM.0000000000001978)
49. **Sakaoglu HH, Orbatu D, Emiroglu M, et al.** Spielberger state and trait anxiety level in healthcare professionals during the Covid-19 outbreak: a case of Tepecik Hospital. *Anatol J Med.* 2020;30(2):1-9.
50. **Melnik BM, Kelly SA, Stephens J, et al.** Interventions to improve mental health, well-being, physical health, and lifestyle behaviors in physicians and nurses: a systematic review. *Am J Health Promot.* 2020;34(8):929-941. doi: [10.1177/0890117120920451](https://doi.org/10.1177/0890117120920451)
51. **Kılıç C.** Health services use for earthquake-related psychological problems: results from the 1999 earthquakes in Turkey. *Turk Psikiyatri Derg.* 2008; 19(2):113-123.
52. **Uğur K, Kartal F, Mete B, et al.** The relationship between peritraumatic dissociation and anxiety level, perceived stress, anxiety sensitivity, and coping with earthquake stress in those with post-earthquake acute stress disorder. *Turk Psikiyatri Derg.* 2021;32(4):253-260.