

Review Article

Cite this article: Pedersen T, Raunkiær M, Graven V (2024) Barriers and facilitators for place of death: A scoping review. *Palliative and Supportive Care* 22(6), 2207–2220. <https://doi.org/10.1017/S1478951524001500>

Received: 1 March 2024

Revised: 3 June 2024

Accepted: 30 June 2024

Keywords:

Death; palliative care; palliative medicine; end-of-life; scoping review

Corresponding author: Tina Pedersen;

Email: Tina.Pedersen3@rsyd.dk

Tina Pedersen, M.Sc.^{1,2} , Mette Raunkiær, R.N., M.Sc., Ph.D.^{1,2}  and Vibeke Graven, M.Phil., Ph.D.^{1,2} 

¹REHPA, The Danish Knowledge Centre for Rehabilitation and Palliative Care, Odense University Hospital, Nyborg, Denmark and ²Department of Clinical Research, University of Southern Denmark, Odense, Denmark

Abstract

Objectives. Many factors influence where people die, but most people prefer to die at home. Investigating the factors affecting death at different locations can enhance end-of-life care and enable more people to die at their preferred place. The aim was to investigate barriers and facilitators affecting place of death and compare facilitators and barriers across different places of death.

Methods. A scoping review registered on Open Science Framework was conducted in accordance with the guidelines for Scoping Reviews (PRISMA-ScR). An electronic search of literature was undertaken in MEDLINE, EMBASE, PUBMED, PsycINFO, and CINAHL covering the years January 2013–December 2023. Studies were included if they described barriers and/or facilitators for place of death among adults.

Results. This review identified 517 studies, and 95 of these were included in the review. The review identified the following themes. Illness factors: disease type, dying trajectory, treatment, symptoms, and safe environment. Individual factors: sex, age, ethnicity, preferences, and for environmental factors the following were identified: healthcare inputs, education and employment, social support, economy, and place of residence.

Significance of results. The factors influencing place of death are complex and some have a cumulative impact affecting where people die. These factors are mostly rooted in structural aspects and make hospital death more likely for vulnerable groups, who are also less likely to receive palliative care and advanced care planning. Disease type and social support further impact the location of death. Future research is needed regarding vulnerable groups and their preferences for place of death.

Introduction

Understanding the complexity of the factors that lead to place of death is crucial for the provision of good end-of-life care. For many years, home death has been seen as an indicator of high-quality palliative care (Stajduhar and Davies 2005), and the general picture also shows that the majority of people wish to die at home (Ali et al. 2019; Fereidouni et al. 2021; García-Sanjuán et al. 2022; Gomes et al. 2013; Hoare et al. 2015; van Doorne et al. 2021; World Health Organization 2011). However, hospital is the most common place of death in Europe (Jarlbæk 2017; Jiang and May 2021; Orlovic et al. 2017) and Canada (Wilson et al. 2018), but the least preferred place to die along with care homes (Calanzani et al. 2014). Despite preferences for place of death, the number of home deaths is projected to decrease to less than 1 in 10 in 2030 in England and Wales, and the number of institutional deaths, such as deaths at care home or hospital, is expected to increase by 20% (Gomes and Higginson 2008). In recent years, there has been a growing emphasis on end-of-life decision-making and advanced care planning, involving discussions among patients, family members, caregivers, and healthcare professionals to anticipate future healthcare choices, including the preferred place of death (Abel et al. 2013; Burghout et al. 2023). A conceptual model developed by Gomes and Higginson identifies several key factors, which are determinants for place of death (Gomes and Higginson 2006). This model emphasizes how place of death is influenced by an interplay between illness-related-, individual-, and environmental-factors (Gomes and Higginson 2006), and it has inspired other studies within the field (Billingham and Billingham 2013; Burge et al. 2015; García-Sanjuán et al. 2022; Gomes et al. 2015). However, the Quality of Death and Dying Index finds that dying in the preferred place is less important than other aspects, for example managing pain or discomfort and being treated in a clean and safe place (Sepulveda et al. 2022). This challenges the perception of home death as one of the important indicators of a good death (De Roo et al. 2014; Pollock 2015). With the projected increase in institutional deaths, there is a pressing need for a deeper understanding of the factors influencing place of death to develop sufficient end-of-life care options

© The Author(s), 2024. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

(Gomes and Higginson 2008). Existing literature on factors influencing place of death is often focused on one specific place of death, such as home death (Balasundram et al. 2023; Bannon et al. 2018). Accordingly, this review aims to identify factors influencing death at home, hospice, care home and hospital. Hence, this review aims to identify barriers and facilitators affecting place of death and compare the factors across different places of death. By “facilitator” and “barrier,” the review identifies both actor driven factors such as preferences plus contextual and structural factors of importance for place of death such as illness and socioeconomic status.

Methods

Study design and registration

This scoping review is conducted in accordance with PRISMA-ScR guidelines for scoping reviews (PRISMA 2023). It is registered in Open Science Framework, accessible via the link <https://doi.org/10.17605/OSF.IO/564ZA>. There has been no quality assessment of the included studies.

Eligibility criteria

Studies were eligible for inclusion if they addressed barriers and/or facilitators concerning place of death, included adult patients (≥ 18 years), and focused regions within Europe, United Kingdom, and Canada. The restriction to these countries is due to their similar welfare systems. Studies were included if they were published between 2013 and 2023, and were written in English, Danish, Norwegian, or Swedish language. There were no exclusion criteria besides not fulfilling the predetermined criteria of inclusion.

Searches and information sources

The first author (TP) selected search terms in collaboration with VG and MR. TP conducted the final search with guidance from a librarian between January 2023 and February 2023. The search terms were divided into 3 search blocks. Each search was tailored to the specific database, and MESH terms were used if possible. The first block includes the search terms “wish” or “prefer.” The second includes “place of death,” “end-of-life,” “EoL,” “place of care.” The third block uses proximity operators when possible and includes the following search terms “home death,” “hospice death,” “hospital death,” “nursing home death,” or “care home death.” Each block is combined by the Boolean operator AND. An electronic search was undertaken in PubMed, CINAHL, EMBASE, PsycInfo, and MEDLINE. The searches were initially run on January 9, 2023. The final database searches were conducted on February 9, 2023. An additional search was conducted on December 13, 2023. The search string tailored for EMBASE can be seen in Table 1. The other searches are available on request of the corresponding author.

Study selection

For inclusion in the title/abstract screening, consensus between 2 independent reviewers was required. One reviewer (TP) assessed all titles/abstracts, while 2 reviewers (MR and VG) each evaluated half of the titles/abstracts for eligibility. In the full-text phase, TP reviewed all studies and consulted MR or VG in cases of uncertainty. Covidence was utilized during the screening process to

Table 1. Key search terms and example of search string

EMBASE
#1 (wish* or prefer*).mp
#2 exp “place of death”/
#3 end-of-life*.mp
#4 EoL*.mp
#5 place of care*.mp
#6 home adj2 death*.mp
#7 hospice adj2 death*.mp
#8 hospital adj2 death*.mp
#9 nursing home adj2 death*.mp
#10 care home adj2 death*.mp
#11 2 or 3 or 4 or 5
#12 6 or 7 or 8 or 9 or 10
#13 1 and 11 and 12
#14 limit 13 to yr = “2013-Current”

remove duplicates, assist with title and abstract screening and register the reason for exclusion of studies. Quantitative, qualitative studies, and reviews were included.

Data charting process and data items

TP performed data extraction using a study specific data extraction form. The following data were extracted: first author and year, title, country of first author, countries discussed, study aim, and study design (Appendix 1). For the analysis data on barriers and facilitators for place of death was extracted.

Synthesis of results

The initial coding of the 95 included studies draws inspiration from Gomes and Higginson’s (2006) conceptual model, categorizing barriers and facilitators into illness-related-, individual-, and environmental factors. Subsequently, these factors for different places of death are further grouped using inductively inspired thematic coding, resulting in various subthemes. The coding of the illness factors yielded the following subthemes: disease type, dying trajectory, treatment, symptoms, and safe environment. Meanwhile, the coding of the individual factors led to the identification the following subthemes: sex, age, ethnicity, and preferences. Similarly, the coding of the environmental factors resulted in the following subthemes: healthcare input, education and employment, social support, economy, and place of residence. Tables 3–5 provide illustrations of the likelihood of each factor being associated with death at different places, along with the number of studies supporting the respective factor and place of death.

Results

Selection of source evidence

The database search revealed 1096 potential studies; 517 studies were initially screened for the appropriateness of title/abstract, with 97 studies subsequently included for full text screening resulting in 73 included studies. After assessing the reference lists of the included studies, 18 studies were included. The additional search in December resulted in inclusion of 3 studies. A total of 95 studies were included in the review. A PRISMA flow chart (Figure 1) outlines the selection process, including reasons for exclusion.




Figure 1. Flow chart screening process.

Characteristics of included studies

Appendix 1 presents the extracted data from the 95 included studies. Table 2 illustrates year published, number of studies, and the respective references for the included studies. Hence 30 studies were published between 2013 and 2015, 27 studies between 2016 and 2018, 29 studies between 2019 and 2021 and 9 studies between 2022 and 2023. Most of the included studies are quantitative ($n = 74$), followed by qualitative ($n = 11$), reviews ($n = 8$), and mixed methods design ($n = 2$).

Synthesis of results

Illness factors

Table 3 illustrates how different types of illness affects place of death. Cancer types, including lung, brain, prostate, and colorectal cancer, are grouped together due to their similar effects on

place of death. Overall, the review indicates that having cancer tends to facilitate death at home (Cabañero-Martínez et al. 2019; Costa et al. 2016; Houttekier et al. 2014; Hunt et al. 2014a; Kamisetty et al. 2015; Nilsson et al. 2020, 2021; Quinn et al. 2020; Sayma et al. 2020; Sleeman et al. 2014). However, hematological cancer is a barrier for home death (Gao et al. 2013; McCaughan et al. 2018; Öhlén et al. 2017; Raziee et al. 2017), but a facilitating factor for hospital death (Gao et al. 2013; Howell et al. 2017, 2013; McCaughan et al. 2018, 2019; Öhlén et al. 2017; Sheridan et al. 2021). The illness trajectory for hematological cancer often involves aggressive treatment up to death (McCaughan et al. 2019). Hematological patients who discuss preferred place of death are more likely not to die in hospital, whereas those who do not discuss preferred place of death, or those who receive hematological treatment close to death are more likely to die in hospital (Howell et al. 2017). Additionally, hematological patients are less likely to receive palliative care and advanced care planning, which facilitates

Table 2. Characteristics of included studies

Year published	Number of studies	References
2013–2015	30	(Abel et al. 2013; Ahearn et al. 2013; Brogaard et al. 2013; Burge et al. 2015; Dasch et al. 2015; De Roo et al. 2014; Domínguez-Berjón et al. 2015; Gage et al. 2015; Gao et al. 2013; Gomes et al. 2013; Gomes et al. 2015; Guerriere et al. 2015; Higginson et al. 2013; Houttekier et al. 2014; Howell et al. 2013; Hunt et al. 2014a; Hunt et al. 2014b; Håkanson et al. 2015; Jayaraman and Joseph 2013; Kamisetty et al. 2015; Ko et al. 2014; Koffman et al. 2014; Livingston et al. 2013; Pinzon et al. 2013; Purdy et al. 2015; Reyniers et al. 2014; Seal et al. 2015; Sharpe et al. 2015; Sleeman et al. 2014; Varani et al. 2015)
2016–2018	27	(Bannon et al. 2018; Black et al. 2016; Costa et al. 2016; de Graaf et al. 2016; Gisquet et al. 2016; Gomes et al. 2018; Howell et al. 2017; Johnson et al. 2018; Lovell et al. 2017; Luta et al. 2016; McCaughan et al. 2018; McEwen et al. 2018; Nieder et al. 2016; O'Sullivan and Higginson 2016; Oosterveld-Vlug et al. 2018; Pooler et al. 2018; Raziee et al. 2017; Reyniers et al. 2016; Schou-Andersen et al. 2016; Tanuseputro et al. 2018; Van Den Block et al. 2017; Wahid et al. 2018; Wales et al. 2018; Winthereik et al. 2018; Wye et al. 2016; Ziway et al. 2017; Öhlén et al. 2017),
2019–2021	29	(Alcorn et al. 2020; Archibald et al. 2021; Cabañero-Martínez et al. 2019; Cai et al. 2021; Davies et al. 2021; Dixon et al. 2019; Isenberg et al. 2020; Jiang and May 2021; Kalseth and Halvorsen 2020; Kern et al. 2020; Kjellstadli et al. 2020; Kjellstadli et al. 2019; Larsen et al. 2020; Martinsson et al. 2020; McCaughan et al. 2019; Mieras et al. 2019; Neergaard et al. 2019; Nilsson et al. 2020; Nilsson et al. 2021; Nolasco et al. 2020; Orlovic et al. 2020; Quinn et al. 2020; Rasch-Westin et al. 2019; Sayma et al. 2020; Sheridan et al. 2021; Skorstengaard et al. 2020; Stajduhar 2020; Van Spall et al. 2021; Wales et al. 2020; Wilson et al. 2020)
2022–2023	9	(Balasundram et al. 2023; Burghout et al. 2023; Campos et al. 2022; Driller et al. 2022; Ervik et al. 2023; García-Sanjuán et al. 2022; Jordan et al. 2023; Nysæter et al. 2022; Schnakenberg et al. 2022).

death in hospital (Howell et al. 2017; McCaughan et al. 2019). Moreover, diseases of the digestive system (Cabañero-Martínez et al. 2019), the respiratory system (Gomes et al. 2018; Kalseth and Halvorsen 2020; Orlovic et al. 2020; Sleeman et al. 2014), liver disease (Cabañero-Martínez et al. 2019; Gomes et al. 2018; Houttekier et al. 2014; Jordan et al. 2023), HIV/AIDS (Cabañero-Martínez et al. 2019; Gomes et al. 2018), kidney disease (Lovell et al. 2017), tumors, (Cabañero-Martínez et al. 2019) and ALS (Domínguez-Berjón et al. 2015) increase the likelihood of death in hospital. However, irrespective of disease type, close relationships with healthcare staff and feeling safe in hospital are factors leading the patient to prefer care and death in hospital (Howell et al. 2017, 2013; McCaughan et al. 2018, 2019; Sheridan et al. 2021). Table 3 also highlights differences in the dying trajectories. For example, having more hospital days prior to death (Gomes et al. 2015; Jiang and May 2021; Kern et al. 2020; Varani et al. 2015) or receiving life-prolonging treatment (Campos et al. 2022; Oosterveld-Vlug et al. 2018; Reyniers et al. 2016) are associated with hospital death. Additionally, patients with open awareness of dying are more likely to die at home (Hunt et al. 2014b; Kern et al. 2020; Nysæter et al. 2022; Pooler et al. 2018) as opposed to late recognition of dying (McCaughan et al. 2019), they are more likely to die at home rather than hospital.

Individual factors

Table 4 presents an inconclusive evidence regarding gender differences in place of death. Concerning age there appears to be a tendency indicating that lower age makes hospital death more likely (Cabañero-Martínez et al. 2019; Gomes et al. 2018; Houttekier et al. 2014; Luta et al. 2016; Martinsson et al. 2020; Nolasco et al. 2020), whereas older age increases the likelihood of death in a care home (Black et al. 2016; Dixon et al. 2019; Domínguez-Berjón et al. 2015; Houttekier et al. 2014; Kalseth and Halvorsen 2020; Kamisetty et al. 2015; Livingston et al. 2013; Luta et al. 2016; Sheridan et al. 2021; Sleeman et al. 2014; Wales et al. 2020). Being non-white or belonging to an ethnic minority decreases the likelihood of dying at home or in hospice and increases the chance of dying in hospital (Cabañero-Martínez et al. 2019; Higginson et al. 2013; Jiang and May 2021; Koffman et al. 2014). While variations in place of death

for different ethnicities are not extensively investigated, they may be attributed to differences in preferences influenced by culture or access to palliative care (Sharpe et al. 2015). Overall, expressed preference appears to play a significant role in determining place of death by increasing the likelihood of home death (Brogaard et al. 2013; Costa et al. 2016; Dixon et al. 2019; García-Sanjuán et al. 2022; Gomes et al. 2015; Hunt et al. 2014a; McCaughan et al. 2018; Neergaard et al. 2019; Nysæter et al. 2022; Rasch-Westin et al. 2019; Sayma et al. 2020; Schou-Andersen et al. 2016; Seal et al. 2015; Sheridan et al. 2021; Wales et al. 2018). Conversely, not having a preference or unknown preference are factors that increase the likelihood of hospital death (Abel et al. 2013; Ahearn et al. 2013; Burghout et al. 2023; Dixon et al. 2019; Howell et al. 2017; Kern et al. 2020; McCaughan et al. 2019; Orlovic et al. 2020).

Environmental factors

Table 5 identifies various environmental factors impacting the place of death. For instance, receiving palliative care (Archibald et al. 2021; Balasundram et al. 2023; Bannon et al. 2018; Brogaard et al. 2013; Burge et al. 2015; De Roo et al. 2014; Dixon et al. 2019; Gage et al. 2015; Gomes et al. 2015; Higginson et al. 2013; Johnson et al. 2018; Kern et al. 2020; Ko et al. 2014; Larsen et al. 2020; Pooler et al. 2018; Quinn et al. 2020; Tanuseputro et al. 2018; Varani et al. 2015; Wahid et al. 2018; Wye et al. 2016) and/or advanced care planning (Ahearn et al. 2013; Archibald et al. 2021; Burghout et al. 2023; Driller et al. 2022; Pooler et al. 2018; Sayma et al. 2020; Skorstengaard et al. 2020; Wahid et al. 2018) increases the chances of dying at home, whereas not receiving palliative care (Nieder et al. 2016) and/or advanced care planning (Ahearn et al. 2013; Burghout et al. 2023; Dixon et al. 2019; Howell et al. 2017; Kern et al. 2020; McCaughan et al. 2019; Orlovic et al. 2020) increases the chance of dying in hospital. Social factors such as living with others (Brogaard et al. 2013; Cai et al. 2021; Costa et al. 2016; Dixon et al. 2019; Gao et al. 2013; García-Sanjuán et al. 2022; Guerriere et al. 2015; Higginson et al. 2013; Houttekier et al. 2014; Neergaard et al. 2019; Pinzon et al. 2013), being married or having a partner (Cai et al. 2021; Houttekier et al. 2014; Öhlén et al. 2017), or having a family caregiver (Archibald et al. 2021; Costa et al. 2016; Ervik et al. 2023; Gomes et al. 2013; Kern et al. 2020; Ko et al. 2014;

Table 3. Illness factors associated with place of death

	Facilitating death in	Number of studies	Barrier for death in	Number of studies
Factors related to illness				
<i>Disease type</i>				
Cancer	Home (Cabañero-Martínez et al. 2019; Costa et al. 2016; Houttekier et al. 2014; Hunt et al. 2014a; Kamisetty et al. 2015; Nilsson et al. 2020; Nilsson et al. 2021; Quinn et al. 2020; Sayma et al. 2020; Sleeman et al. 2014)	43% 10/23	Hospital (Oosterveld-Vlug et al. 2018; Orlovic et al. 2020; Öhlén et al. 2017)	60% 3/5
Haematological cancer	Hospital (Gao et al. 2013; Howell et al. 2013; Howell et al. 2017; McCaughan et al. 2018; McCaughan et al. 2019; Sheridan et al. 2021; Öhlén et al. 2017)	100% 7/7	Home (Gao et al. 2013; McCaughan et al. 2018; Raziee et al. 2017; Öhlén et al. 2017)	80% 4/5
Neurodegenerative disease	Care home (Alcorn et al. 2020; Black et al. 2016; Cabañero-Martínez et al. 2019; Costa et al. 2016; Dixon et al. 2019; Houttekier et al. 2014; Jayaraman and Joseph 2013; Kalseth and Halvorsen 2020; Livingston et al. 2013; Sleeman et al. 2014)	43% 10/23	Hospital (Black et al. 2016; Cabañero-Martínez et al. 2019; Gomes et al. 2018; Houttekier et al. 2014; Kalseth and Halvorsen 2020; Orlovic et al. 2020; Schnakenberg et al. 2022)	88% 7/8
Liver disease	Hospital (Cabañero-Martínez et al. 2019; Gomes et al. 2018; Houttekier et al. 2014; Jordan et al. 2023)	100% 4/4	Home (Jiang and May 2021)	100% 1/1
Disease of the digestive system	Hospital (Cabañero-Martínez et al. 2019)	100% 1/1	–	–
Disease of the circulatory system	Home (Black et al. 2016; Cabañero-Martínez et al. 2019; Kalseth and Halvorsen 2020)	60% 3/5	Hospital (Cabañero-Martínez et al. 2019; Gomes et al. 2018; Kalseth and Halvorsen 2020) Care home (Jayaraman and Joseph 2013; Kalseth and Halvorsen 2020; Sleeman et al. 2014)	43% 3/7 43% 3/7
Disease of respiratory system	Hospital (Gomes et al. 2018; Kalseth and Halvorsen 2020; Orlovic et al. 2020; Sleeman et al. 2014)	80% 4/5	Home (Kalseth and Halvorsen 2020) Care home (Sleeman et al. 2014)	50% 1/2 50% 1/2
Cerebrovascular disease	–	–	Care home (Jayaraman and Joseph 2013; Sleeman et al. 2014)	100% 2/2
HIV/AIDS	Hospital (Cabañero-Martínez et al. 2019; Gomes et al. 2018)	100% 2/2	–	–
Kidney disease	Hospital (Lovell et al. 2017)	100% 1/1	–	–
Sepsis or bleeding	Hospital (McCaughan et al. 2019)	100% 1/1	–	–
Tumours	Hospital (Cabañero-Martínez et al. 2019)	100% 1/1	–	–
ALS	Hospital (Dominguez-Berjón et al. 2015)	100% 1/1	–	–
Mental illness	Care home (Wilson et al. 2020)	100% 1/1	–	–
Organ failure	Hospital (Houttekier et al. 2014)	100% 1/1	–	–
Comorbidity	–	–	Home (Raziee et al. 2017) Care home (Alcorn et al. 2020)	50% 1/2 50% 1/2
Stroke	Care home (Costa et al. 2016)	100% 1/1	–	–
<i>Dying trajectory</i>				
More hospital days prior to death	Hospital (Gomes et al. 2015; Jiang and May 2021; Kern et al. 2020; Varani et al. 2015)	80% 4/5	Home (Kern et al. 2020; Kjellstadli et al. 2020)	100% 2/2
Fewer hospital days prior to death	Home (Burge et al. 2015; Kern et al. 2020)	100% 2/2	–	–
Early conversation about prognosis	Home (McCaughan et al. 2018) Hospice (McCaughan et al. 2018)	50% 1/2 50% 1/2	–	–

(Continued)

Table 3. (Continued.)

	Facilitating death in	Number of studies	Barrier for death in	Number of studies
Death close to diagnosis	Home (Sheridan et al. 2021) Hospital (Howell et al. 2013)	50% 1/2 50% 1/2	Home (Sheridan et al. 2021) Hospice (Sheridan et al. 2021)	50% 1/2 50% 1/2
Open awareness of dying	Home (Hunt et al. 2014b; Kern et al. 2020; Nysæter et al. 2022; Pooler et al. 2018)	100% 4/4	–	–
Late recognition of dying	Hospital (McCaughan et al. 2019)	100% 1/1	Home (Kern et al. 2020; Sayma et al. 2020)	100% 2/2
Acute hospitalisation or care need	Hospital (Alcorn et al. 2020; Black et al. 2016; de Graaf et al. 2016; Reyniers et al. 2014; Seal et al. 2015; Van Spall et al. 2021; Wahid et al. 2018; Wales et al. 2020)	88% 8/9	Home (Cai et al. 2021; Reyniers et al. 2014) Care home (Alcorn et al. 2020; Van Spall et al. 2021)	50% 2/4 50% 2/4
Delayed discharge from hospital	Hospital (McCaughan et al. 2019; Sayma et al. 2020)	100% 2/2	–	–
Longer stay at care home	Care home (Costa et al. 2016)	100% 1/1	–	–
Late stage of disease	Care home (Costa et al. 2016) Hospital (Kamisetty et al. 2015)	50% 1/2 50% 1/2	–	–
<i>Treatment</i>				
In-hospital treatment	Hospital (McCaughan et al. 2019; Mieras et al. 2019; Orlovic et al. 2020)	100% 3/3	–	–
Life prolonging treatment	Hospital (Campos et al. 2022; Oosterveld-Vlug et al. 2018; Reyniers et al. 2016)	100% 3/3	Home (Campos et al. 2022; De Roo et al. 2014)	100% 2/2
<i>Symptoms</i>				
Lower care dependency	Hospital (Schnakenberg et al. 2022)	100% 1/1	–	–
Low symptom burden	Home (Kern et al. 2020)	100% 1/1	–	–
High symptom burden	–	–	Care home (Alcorn et al. 2020)	100% 1/1
Well managed symptoms/pain	Home (Hunt et al. 2014a; McEwen et al. 2018)	100% 2/2	–	–
Low functional status	Home (Higginson et al. 2013; Neergaard et al. 2019)	100% 2/2	–	–
<i>Safe environment</i>				
Safe environment at home	Home (Wales et al. 2018)	100% 1/1	–	–
Feeling safe or at home at hospital	Hospital (Howell et al. 2017; McCaughan et al. 2018; McCaughan et al. 2019; O'Sullivan and Higginson 2016; Reyniers et al. 2016)	100% 5/5	–	–

Pooler et al. 2018; Sayma et al. 2020; Wahid et al. 2018) increase the likelihood of home death. Whereas, living alone (Ahearn et al. 2013; Houttekier et al. 2014; Lovell et al. 2017), being single, widowed, or divorced (Domínguez-Berjón et al. 2015; Gao et al. 2013; Nilsson et al. 2021; Nolasco et al. 2020) increase the possibility of hospital death. The absence of a family caregiver or when family caregiving is experienced burdensome decrease the possibility of home death (Bannon et al. 2018; de Graaf et al. 2016; Kern et al. 2020; O'Sullivan and Higginson 2016; Sayma et al. 2020; Seal et al. 2015; Wahid et al. 2018). Residence in rural areas is associated with higher likelihood of dying at home (Houttekier et al. 2014; Jayaraman and Joseph 2013; Kern et al. 2020; Neergaard et al. 2019; Nilsson et al. 2020) as opposed to hospital (Cabañero-Martínez et al. 2019; Dasch et al. 2015; Gomes et al. 2018; Luta et al.

2016; Öhlén et al. 2017), while urban residents are more likely to die in hospital (Dasch et al. 2015; Gomes et al. 2018; Håkanson et al. 2015; Houttekier et al. 2014; Luta et al. 2016; Nilsson et al. 2020; Öhlén et al. 2017) rather than at home (Dasch et al. 2015; Håkanson et al. 2015; Kern et al. 2020; Neergaard et al. 2019). Some of these differences might be explained by urban residents living closer to hospital as opposed to rural residents (Kalseth and Halvorsen 2020; Ziway et al. 2017). Additionally, living in non-deprived or affluent areas increases the chance of dying at home (Bannon et al. 2018; Dixon et al. 2019; Gao et al. 2013; Neergaard et al. 2019; Raziee et al. 2017; Sleeman et al. 2014), whereas living in deprived or non-affluent areas increases the chance of a hospital death (Davies et al. 2021; Neergaard et al. 2019; Nolasco et al. 2020; Ziway et al. 2017), and lowers the chance of home death

Table 4. Individual factors associated with place of death

	Facilitating death in	Number of studies	Barrier for death in	Number of studies
Individual factors				
<i>Sex</i>				
Male	Home (Archibald et al. 2021; Black et al. 2016; Dasch et al. 2015; Dixon et al. 2019; Houttekier et al. 2014; Jayaraman and Joseph 2013; Kalseth and Halvorsen 2020; Schou-Andersen et al. 2016) Hospital (Cabañero-Martínez et al. 2019; Dasch et al. 2015; Houttekier et al. 2014; Jayaraman and Joseph 2013; Luta et al. 2016; Nolasco et al. 2020; Orlovic et al. 2020; Van Spall et al. 2021)	44% 8/18 44% 8/18	Home (Gao et al. 2013; Kjellstadli et al. 2020; Sleeman et al. 2014) Care home (Cabañero-Martínez et al. 2019; Dasch et al. 2015; Jayaraman and Joseph 2013)	43% 3/7 43% 3/7
Female	Care home (Cabañero-Martínez et al. 2019; Dasch et al. 2015; Houttekier et al. 2014; Jayaraman and Joseph 2013; Livingston et al. 2013; Luta et al. 2016; Sheridan et al. 2021)	50% 7/14	Home (Black et al. 2016; Dasch et al. 2015; Dixon et al. 2019; Jayaraman and Joseph 2013; Kalseth and Halvorsen 2020) Hospital (Cabañero-Martínez et al. 2019; Luta et al. 2016; Nolasco et al. 2020; Orlovic et al. 2020; Van Spall et al. 2021)	50% 5/10 50% 5/10
<i>Age</i>				
Younger age	Hospital (Cabañero-Martínez et al. 2019; Gomes et al. 2018; Houttekier et al. 2014; Luta et al. 2016; Martinsson et al. 2020; Nolasco et al. 2020)	60% 6/10	–	–
Increasing age	Care home (Black et al. 2016; Dixon et al. 2019; Domínguez-Berjón et al. 2015; Houttekier et al. 2014; Kalseth and Halvorsen 2020; Kamisetty et al. 2015; Livingston et al. 2013; Luta et al. 2016; Sheridan et al. 2021; Sleeman et al. 2014; Wales et al. 2020)	39% 11/28	Home (Bannon et al. 2018; Black et al. 2016; Dixon et al. 2019; Hunt et al. 2014b; Kalseth and Halvorsen 2020)	41% 5/12
<i>Ethnicity</i>				
Ethnic minority/non-white	Hospital (Cabañero-Martínez et al. 2019; Higginson et al. 2013; Jiang and May 2021; Koffman et al. 2014)	100% 4/4	Hospice (Higginson et al. 2013; Koffman et al. 2014) Home (Koffman et al. 2014; Sharpe et al. 2015)	50% 2/4 50% 2/4
<i>Preferences</i>				
Patient's preference	Home (Brogaard et al. 2013; Costa et al. 2016; Dixon et al. 2019; García-Sanjuán et al. 2022; Gomes et al. 2015; Hunt et al. 2014a; McCaughan et al. 2018; Neergaard et al. 2019; Nysæter et al. 2022; Rasch-Westin et al. 2019; Sayma et al. 2020; Schou-Andersen et al. 2016; Seal et al. 2015; Sheridan et al. 2021; Wales et al. 2018)	68% 15/22	–	–
No preference/unknown preference	Hospital (Abel et al. 2013; Ahearn et al. 2013; Burghout et al. 2023; Dixon et al. 2019; Howell et al. 2017; Kern et al. 2020; McCaughan et al. 2019; Orlovic et al. 2020)	100% 8/8	–	–
Discussion of preferred place of death with family	Home (Gomes et al. 2015; Pooler et al. 2018)	66% 2/3	–	–

(Gao et al. 2013; Higginson et al. 2013; Sleeman et al. 2014). Higher education (Gisquet et al. 2016; Houttekier et al. 2014; Nilsson et al. 2021), income (Neergaard et al. 2019; Schou-Andersen et al. 2016), and socioeconomic status (Sharpe et al. 2015) increases the likelihood of home death, and decreases the chance of hospital death

(Davies et al. 2021; Van Spall et al. 2021), whereas lower income (Wales et al. 2020) and socioeconomic status (Domínguez-Berjón et al. 2015) increases the likelihood of hospital death, and decreases the chance of home death (Higginson et al. 2013; Schou-Andersen et al. 2016; Wales et al. 2020).

Table 5. Environmental factors associated with place of death

	Facilitating death in	Number of studies	Barrier for death in	Number of studies
Environmental factors				
<i>Healthcare input</i>				
Home care provision (nursing and family physician visits, distinct or community nurse visits, home support, GP home visits)	Home (Bannon et al. 2018; Burge et al. 2015; Cai et al. 2021; Ervik et al. 2023; García-Sanjuán et al. 2022; Gomes et al. 2015; Isenberg et al. 2020; Kern et al. 2020; Kjellstadli et al. 2020; McEwen et al. 2018; Sayma et al. 2020; Tanuseputro et al. 2018; Varani et al. 2015; Wahid et al. 2018)	100% 14/14	–	–
Intensity of home care	Home (Cai et al. 2021; Ervik et al. 2023; Kalseth and Halvorsen 2020; Kjellstadli et al. 2019; Neergaard et al. 2019; Winthereik et al. 2018)	100% 6/6	–	–
Low availability of care home beds	Home (Houttekier et al. 2014)	100% 1/1	Care home (Kalseth and Halvorsen 2020; Sleeman et al. 2014)	100% 2/2
High availability of care home beds	Care home (Houttekier et al. 2014; Kalseth and Halvorsen 2020; Sleeman et al. 2014)	100% 3/3	–	–
Low availability of hospital beds	Home (Houttekier et al. 2014)	100% 1/1	–	–
High availability of hospital beds	Hospital (Gomes et al. 2015; Houttekier et al. 2014; Jiang and May 2021; Kalseth and Halvorsen 2020; Van Spall et al. 2021)	83% 5/6	–	–
Low availability of hospice beds	–	–	Hospice (McCaughan et al. 2018)	100% 1/1
Palliative care provision	Home (Archibald et al. 2021; Balasundram et al. 2023; Bannon et al. 2018; Brogaard et al. 2013; Burge et al. 2015; De Roo et al. 2014; Dixon et al. 2019; Gage et al. 2015; Gomes et al. 2015; Higginson et al. 2013; Johnson et al. 2018; Kern et al. 2020; Ko et al. 2014; Larsen et al. 2020; Pooler et al. 2018; Quinn et al. 2020; Tanuseputro et al. 2018; Varani et al. 2015; Wahid et al. 2018; Wye et al. 2016)	95% 20/21	Hospital (Jiang and May 2021; Johnson et al. 2018; Nieder et al. 2016; Oosterveld-Vlug et al. 2018; Purdy et al. 2015; Quinn et al. 2020; Van Spall et al. 2021)	100% 7/7
No palliative care provision	Hospital (Nieder et al. 2016)	100% 1/1	Home (Kern et al. 2020; Wahid et al. 2018)	100% 2/2
Discussion preferred place of death with healthcare professionals	Home (Bannon et al. 2018; De Roo et al. 2014; Dixon et al. 2019; Nysæter et al. 2022)	100% 4/4	–	–
Advanced care planning	Home (Ahearn et al. 2013; Archibald et al. 2021; Burghout et al. 2023; Driller et al. 2022; Pooler et al. 2018; Skorstengaard et al. 2020; Wahid et al. 2018)	63% 7/11	–	–
No advanced care planning	Hospital (Ahearn et al. 2013; Burghout et al. 2023; Dixon et al. 2019; Howell et al. 2017; Kern et al. 2020; McCaughan et al. 2019; Orlovic et al. 2020)	88% 7/8	–	–

(Continued)

Table 5. (Continued.)

	Facilitating death in	Number of studies	Barrier for death in	Number of studies
Good coordination of care	Home (Balasundram et al. 2023; Wahid et al. 2018)	100% 2/2	–	–
Lack of coordination of care	–	–	Home (Sayma et al. 2020; Wahid et al. 2018)	100% 2/2
Effective communication	Home (Wahid et al. 2018)	100% 1/1	–	–
Ineffective communication	Hospital (McCaughan et al. 2018; McCaughan et al. 2019)	100% 2/2	Home (Sayma et al. 2020; Wahid et al. 2018)	100% 2/2
Inadequate skills for healthcare staff	–	–	Home (Reyniers et al. 2014; Sayma et al. 2020; Wahid et al. 2018)	60% 3/5
Adequate skills for healthcare staff	Home (Burge et al. 2015; Nysæter et al. 2022)	100% 2/2	–	–
<i>Education and employment</i>				
No education or lower/medium level of education	Home (Neergaard et al. 2019) Care home (Houttekier et al. 2014) Hospital (Cabañero-Martínez et al. 2019)	33% 1/3 33% 1/3 33% 1/3	Home (Neergaard et al. 2019; Nilsson et al. 2021)	66% 2/3
High level of education	Home (Gisquet et al. 2016; Houttekier et al. 2014; Nilsson et al. 2021)	100% 3/3	–	–
Being employed	Home (Gisquet et al. 2016) Hospital (Neergaard et al. 2019)	50% 1/2 50% 1/2	–	–
Being unemployed	–	–	Hospital (Neergaard et al. 2019)	100% 1/1
<i>Social support</i>				
Living with relatives	Home (Brogaard et al. 2013; Cai et al. 2021; Costa et al. 2016; Dixon et al. 2019; Gao et al. 2013; García-Sanjuán et al. 2022; Guerriere et al. 2015; Higginson et al. 2013; Houttekier et al. 2014; Neergaard et al. 2019; Pinzon et al. 2013)	100% 11/11	–	–
Living alone	Hospital (Ahearn et al. 2013; Houttekier et al. 2014; Lovell et al. 2017)	75% 3/4	Home (Cai et al. 2021; Costa et al. 2016; Guerriere et al. 2015; O'Sullivan and Higginson 2016)	100% 4/4
Being married	Home (Cai et al. 2021; Houttekier et al. 2014; Öhlén et al. 2017)	60% 3/5	Hospital (Domínguez-Berjón et al. 2015; Nilsson et al. 2021) Care home (Jayaraman and Joseph 2013; Kalseth and Halvorsen 2020)	40% 2/5 40% 2/5
Being single, widowed or divorced	Hospital (Domínguez-Berjón et al. 2015; Gao et al. 2013; Nilsson et al. 2021; Nolasco et al. 2020)	57% 4/7	Home (Gao et al. 2013; Sleeman et al. 2014) Hospice (Gao et al. 2013; Sleeman et al. 2014) Hospital (Domínguez-Berjón et al. 2015; Gomes et al. 2018)	33% 2/6 33% 2/6 33% 2/6
Family's preference	Home (Bannon et al. 2018; Gomes et al. 2015; Kern et al. 2020; Pinzon et al. 2013; Sayma et al. 2020)	45% 5/11	–	–
Good relationship with healthcare staff	Home (Archibald et al. 2021; Dixon et al. 2019; Gomes et al. 2013; Hunt et al. 2014b; Kern et al. 2020; Nysæter et al. 2022)	83% 6/7	–	–

(Continued)

Table 5. (Continued.)

	Facilitating death in	Number of studies	Barrier for death in	Number of studies
Caregiver receiving support	Home (Archibald et al. 2021; Dixon et al. 2019; Gomes et al. 2013; Hunt et al. 2014b; Kern et al. 2020)	100% 5/5	–	–
Having a family caregiver	Home (Archibald et al. 2021; Costa et al. 2016; Ervik et al. 2023; Gomes et al. 2013; Kern et al. 2020; Ko et al. 2014; Pooler et al. 2018; Sayma et al. 2020; Wahid et al. 2018)	100% 9/9	–	–
Not having a family caregiver or care as a burden	Hospital (Gomes et al. 2015)	100% 1/1	Home (Bannon et al. 2018; de Graaf et al. 2016; Kern et al. 2020; O'Sullivan and Higginson 2016; Sayma et al. 2020; Seal et al. 2015; Wahid et al. 2018)	100% 7/7
<i>Economy</i>				
Higher income	Home (Neergaard et al. 2019; Schou-Andersen et al. 2016)	100% 2/2	Hospital (Davies et al. 2021; Van Spall et al. 2021)	100% 2/2
Lower income	Hospital (Wales et al. 2020)	100% 1/1	Home (Schou-Andersen et al. 2016; Wales et al. 2020)	100% 2/2
Higher socioeconomic status	Home (Sharpe et al. 2015) Hospice (Sharpe et al. 2015)	50% 1/2 50% 1/2	–	–
Lower socioeconomic status	Hospital (Dominguez-Berjón et al. 2015)	100% 1/1	Home (Higginson et al. 2013)	100% 1/1
High care cost	–	–	Home (Kern et al. 2020; Wahid et al. 2018)	66% 2/3
<i>Place of residence</i>				
Living in urban area	Hospital (Dasch et al. 2015; Gomes et al. 2018; Houttekier et al. 2014; Håkanson et al. 2015; Luta et al. 2016; Nilsson et al. 2020; Öhlén et al. 2017)	63% 7/11	Home (Dasch et al. 2015; Håkanson et al. 2015; Kern et al. 2020; Neergaard et al. 2019)	66% 4/6
Living in rural area	Home (Houttekier et al. 2014; Jayaraman and Joseph 2013; Kern et al. 2020; Neergaard et al. 2019; Nilsson et al. 2020)	83% 5/6	Hospital (Cabañero-Martínez et al. 2019; Dasch et al. 2015; Gomes et al. 2018; Luta et al. 2016; Öhlén et al. 2017)	50% 5/10
Living in deprived/low affluence area	Hospital (Davies et al. 2021; Neergaard et al. 2019; Nolasco et al. 2020; Ziway et al. 2017)	100% 4/4	Home (Gao et al. 2013; Higginson et al. 2013; Sleeman et al. 2014)	100% 3/3
Living in non-deprived/high affluence area	Home (Bannon et al. 2018; Dixon et al. 2019; Gao et al. 2013; Neergaard et al. 2019; Raziee et al. 2017; Sleeman et al. 2014)	60% 6/10	Home (Neergaard et al. 2019) Hospital (Ziway et al. 2017)	50% 1/2 50% 1/2
Living close to care home, hospice or hospital	Care home (Ziway et al. 2017) Hospital (Kalseth and Halvorsen 2020) Hospice (Ziway et al. 2017)	33% 1/3 33% 1/3 33% 1/3	–	–

Discussion

This scoping review found evidence for a complex network of factors that impact place of death. It reveals how various diseases, such as hematological cancer or undergoing active treatment, influence where individuals die (McCaughan et al. 2018). However not all the identified diseases were well investigated. The review shows a tendency toward hospital death for people that have not expressed a preference for place of death (Abel et al. 2013; Ahearn et al. 2013; Burghout et al. 2023; Dixon et al. 2019; Howell et al.

2017; Kern et al. 2020; McCaughan et al. 2019; Orlovic et al. 2020), have not engaged in advanced care planning (Ahearn et al. 2013; Burghout et al. 2023; Dixon et al. 2019; Howell et al. 2017; Kern et al. 2020; McCaughan et al. 2019; Orlovic et al. 2020), and have not received palliative care (Nieder et al. 2016). Conversely, hospital death is more likely for those with an expressed preference for hospital death, those having a good relationship with health-care staff, and feeling safe at hospital (Howell et al. 2017, 2013; McCaughan et al. 2018, 2019; Sheridan et al. 2021). In general

patients' expressed preferences seem to have a strong influence on place of death, especially for home death (Brogaard et al. 2013; Costa et al. 2016; Dixon et al. 2019; García-Sanjuán et al. 2022; Gomes et al. 2015; Hunt et al. 2014a; McCaughan et al. 2018; Neergaard et al. 2019; Nysæter et al. 2022; Rasch-Westin et al. 2019; Sayma et al. 2020; Schou-Andersen et al. 2016; Seal et al. 2015; Sheridan et al. 2021; Wales et al. 2018). Home death is positively influenced by family caregivers having the same preference as their loved one (Bannon et al. 2018; Gomes et al. 2015; Kern et al. 2020; Pinzon et al. 2013; Sayma et al. 2020), having a strong social support system and living with others (Brogaard et al. 2013; Cai et al. 2021; Costa et al. 2016; Dixon et al. 2019; Gao et al. 2013; García-Sanjuán et al. 2022; Guerriere et al. 2015; Higginson et al. 2013; Houttekier et al. 2014; Neergaard et al. 2019; Pinzon et al. 2013). Nonetheless, caring for a loved one at home can be experienced burdensome for family caregivers, causing physical, psychosocial, and financial stress (Stajduhar 2013; Wahid et al. 2018). Consequently, home death may not always be a possibility even though home death the preferred place of death.

This review most clearly indicates that being white, having higher income, higher socioeconomic status, higher education, and stronger social support increases the chances of dying at home (Neergaard et al. 2019; Schou-Andersen et al. 2016; Sharpe et al. 2015), and that not holding these privileges decreases the chance of a home death (Higginson et al. 2013; Schou-Andersen et al. 2016; Wales et al. 2020), and increases the possibility of a hospital death (Domínguez-Berjón et al. 2015; Wales et al. 2020). The factors linked to these social advantages are also linked with an increased possibility of dying in the preferred place of death. Hence studies find how higher socioeconomic status (Gao et al. 2013; Gisquet et al. 2016; Wales et al. 2020), living with others (Brogaard et al. 2013; Cai et al. 2021), being in a relationship (García-Sanjuán et al. 2022) is found positively associated with congruence between preferred and actual place of death. The influence of socioeconomic status can partly be explained by the ability to purchase additional home care services, which enables a preferred death at home (Wales et al. 2018). Moreover, people with higher socioeconomic status are more likely to have conversations about death and advance care planning directives (Hoare et al. 2015), which may influence the higher levels of congruence between preferred and actual place of death.

Another factor of importance is the impact of ethnicity in relation to place of death. The impact of ethnicity is discussed in a systematic review by Gomes et al., and raises questions about equity and equal access to both palliative care and advanced care planning for vulnerable groups and people from ethnic minority groups (Gomes and Higginson 2006). As pointed out by Stajduhar (2020) being privileged increases the chance of receiving palliative and end-of-life care, assistance with advanced care planning, support for family caregivers, and increases the likelihood of dying in the preferred place. Despite the positive association between privilege and preferred place of death, it also needs to be acknowledged that home death is not always the preferred or optimal place of death, even though there has been a tendency to regard home death as the gold standard (De Roo et al. 2014; Pollock 2015). As Pollock (2015) notes a death is not necessarily good just because it occurs at home. Home death can also be associated with feelings of loneliness, being inadequately supported, having poor symptom control, being distressed and fearful (Pollock 2015). By contrast, death in hospital can be peaceful and with sufficient pain and symptom management (Howell et al. 2017; McCaughan et al. 2019; Orlovic et al. 2020). The Quality of Death and Dying Index reveals that symptom

management is a factor of utmost importance near the end-of-life (Sepulveda et al. 2022). Therefore, we suggest that when advance care planning occurs, appropriate consideration should be given to the place in which symptoms can be best managed. As Pollock (2015) warns, assumptions that home is the best place of death risks denying the patient what might actually be best for them.

Limitations

The categorization of barriers and facilitators into illness-, individual-, and environmental factors is seen in other studies within the field (Burge et al. 2015; García-Sanjuán et al. 2022; Gomes et al. 2015; Neergaard et al. 2019). However, the strength of this review is the investigation of different places of death that sheds light on how vulnerable groups are in risk of experiencing inequities related to access to palliative care and advanced care planning. This potentially denies them the expression of preference and may determine their place of death. Some limitations need to be considered when interpreting the results. First, the review included studies across Europe, United Kingdom, and Canada due to their similar welfare systems, however they are not fully comparable which may cause some of the identified barriers and facilitators to be more or less relevant. Second, the consequence of grouping barriers and facilitators into the most or least likely place of death leaves limited space for showing detailed results of the included studies. However, this is considered helpful to provide a clear picture of this complex field.

Implication for future research

We suggest further research is urgently needed regarding the relationship between socioeconomic status, advanced care planning, palliative care, and place of death. This research should take into account how preference for place of death and actual place of death are determined. In addition, further attention is needed to discuss the ethical imperative behind the ideal of home death as the optimal place to die, and to investigate "good" deaths in other places.

Conclusion

In conclusion, this scoping review has demonstrated the complexity of factors influencing where people die. Several of these factors are rooted in structural conditions that work to restrict the access of underprivileged persons to end-of-life care and an opportunity to express their preferred place of death. Focusing on symptom management in addition to place of death may contribute to allow more people to receive sufficient end-of-life care and death in the best place possible.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S1478951524001500>.

Acknowledgments. The authors wish to thank PhD Jahan Shabnam for valuable comments during the process.

Funding. This research received no specific grant from any funding agency, commercial, or not-for-profit sectors.

Competing interests. The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this study.

Ethical approval. No ethical approval was required for this research as the study is a review of existing, published papers.

References

- Abel J, Pring A, Rich A, *et al.* (2013) The impact of advance care planning of place of death, a hospice retrospective cohort study. *BMJ Supportive & Palliative Care* 3(2), 168. doi:10.1136/bmjspcare-2012-000327
- Ahearn DJ, Nidh N, Kallat A, *et al.* (2013) Offering older hospitalised patients the choice to die in their preferred place. *Postgraduate Medical Journal* 89(1047), 20–24. doi:10.1136/postgradmedj-2012-131161
- Alcorn G, Murray SA and Hockley J (2020) Care home residents who die in hospital: Exploring factors, processes and experiences. *Age and Ageing* 49(3), 468–480. doi:10.1093/ageing/afz174
- Ali M, Capel M, Jones G, *et al.* (2019) The importance of identifying preferred place of death. *BMJ Support Palliat Care* 9(1), 84–91. doi:10.1136/bmjspcare-2015-000878
- Archibald N, Bakal JA, Richman-Eisenstat J, *et al.* (2021) Early integrated palliative care bundle impacts location of death in interstitial lung disease: A pilot retrospective study. *American Journal of Hospice & Palliative Medicine* 38(2), 104–113. doi:10.1177/1049909120924995
- Balasundram S, Holm A, Benthien KS, *et al.* (2023) Increasing the chance of dying at home: Roles, tasks and approaches of general practitioners enabling palliative care: A systematic review of qualitative literature. *BMC Primary Care* 24(1), 77. doi:10.1186/s12875-023-02038-0
- Bannon F, Cairnduff V, Fitzpatrick D, *et al.* (2018) Insights into the factors associated with achieving the preference of home death in terminal cancer: A national population-based study. *Palliative and Supportive Care* 16(6), 749–755. doi:10.1017/S1478951517000876
- Billingham MJ and Billingham SJ (2013) Congruence between preferred and actual place of death according to the presence of malignant or non-malignant disease: A systematic review and meta-analysis. *BMJ Support Palliat Care* 3(2), 144–154. doi:10.1136/bmjspcare-2012-000292
- Black H, Waugh C, Munoz-Arroyo R, *et al.* (2016) Predictors of place of death in South West Scotland 2000–2010: Retrospective cohort study. *Palliative Medicine* 30(8), 764–771. doi:10.1177/0269216315627122
- Brogaard T, Neergaard MA, Sokolowski I, *et al.* (2013) Congruence between preferred and actual place of care and death among Danish cancer patients. *Palliative Medicine* 27(2), 155–164. doi:10.1177/0269216312438468
- Burge F, Lawson B, Johnston G, *et al.* (2015) Preferred and actual location of death: What factors enable a preferred home death? *Journal of Palliative Medicine* 18(12), 1054–1059. doi:10.1089/jpm.2015.0177
- Burghout C, Nahar-van Venrooij LMW, Bolt SR, *et al.* (2023) Benefits of structured advance care plan in end-of-life care planning among older oncology patients: A retrospective pilot study. *Journal of Palliative Care* 38(1), 30–40. doi:10.1177/08258597221119660
- Cabañero-Martínez MJ, Nolasco A, Melchor I, *et al.* (2019) Place of death and associated factors: A population-based study using death certificate data. *European Journal of Public Health* 29(4), 608–615. doi:10.1093/eurpub/cky267
- Cai J, Zhang L, Guerriere D, *et al.* (2021) What variables contribute to the achievement of a preferred home death for cancer patients in receipt of home-based palliative care in Canada? *Cancer Nursing* 44(3), 214–222. doi:10.1097/NCC.0000000000000863
- Calanzani N, Moens K, Cohen J, *et al.* (2014) Choosing care homes as the least preferred place to die: A cross-national survey of public preferences in seven European countries. *BMC Palliative Care* 13(1), 48. doi:10.1186/1472-684X-13-48
- Campos E, Isenberg SR, Lovblom LE, *et al.* (2022) Supporting the heterogeneous and evolving treatment preferences of patients with heart failure through collaborative home-based palliative care. *Journal of the American Heart Association* 11(19), 1–10. doi:10.1161/JAHA.122.026319
- Costa V, Earle CC, Esplen MJ, *et al.* (2016) The determinants of home and nursing home death: A systematic review and meta-analysis. *BMC Palliative Care* 15(1), 8. doi:10.1186/s12904-016-0077-8
- Dasch B, Blum K, Gude P, *et al.* (2015) Place of death: Trends over the course of a decade: A population-based study of death certificates from the years 2001 and 2011. *Deutsches Ärzteblatt International* 112(29-30), 496–504. doi:10.3238/arztebl.2015.0496
- Davies JM, Maddocks M, Chua KC, *et al.* (2021) Socioeconomic position and use of hospital-based care towards the end of life: A mediation analysis using the English Longitudinal Study of Ageing. *The Lancet Public Health* 6(3), e155–e163. doi:10.1016/S2468-2667%2820%2930292-9
- de Graaf E, Zweers D, Valkenburg ACH, *et al.* (2016) Hospice assist at home: Does the integration of hospice care in primary healthcare support patients to die in their preferred location-A retrospective cross-sectional evaluation study. *Palliative Medicine* 30(6), 580–586. doi:10.1177/0269216315626353
- De Roo ML, Miccinesi G, Onwuteaka-Philipsen BD, *et al.* (2014) Actual and preferred place of death of home-dwelling patients in four European countries: Making sense of quality indicators. *PLoS ONE* 9(4), e93762. doi:10.1371/journal.pone.0093762
- Dixon J, King D and Knapp M (2019) Advance care planning in England: Is there an association with place of death? Secondary analysis of data from the National Survey of Bereaved People. *BMJ Supportive & Palliative Care* 9(3), 316–325. doi:10.1136/bmjspcare-2015-000971
- Dominguez-Berjón MF, Esteban-Vasallo MD, Zoni AC, *et al.* (2015) Place of death and associated factors among patients with amyotrophic lateral sclerosis in Madrid (Spain). *Amyotroph Lateral Scler Frontotemporal Degener* 17(1-2), 62–68. doi:10.3109/21678421.2015.1089908
- Driller B, Talseth-Palmer B, Hole T, *et al.* (2022) Cancer patients spend more time at home and more often die at home with advance care planning conversations in primary health care: A retrospective observational cohort study. *BMC Palliative Care* 21(1), 1–10. doi:10.1186/s12904-022-00952-1
- Ervik B, Donnem T and Johansen M-L (2023) Dying at “home” - a qualitative study of end-of-life care in rural Northern Norway from the perspective of health care professionals. *BMC Health Services Research* 23(1), 1359. doi:10.1186/s12913-023-10329-6
- Fereidouni A, Rassouli M, Salesi M, *et al.* (2021) Preferred place of death in adult cancer patients: A systematic review and meta-analysis. *Frontiers in Psychology* 12, 704590. doi:10.3389/fpsyg.2021.704590
- Gage H, Holdsworth LM, Flannery C, *et al.* (2015) Impact of a hospice rapid response service on preferred place of death, and costs. *BMC Palliative Care* 14(101088685), 75. doi:10.1186/s12904-015-0065-4
- Gao W, Ho YK, Verne J, *et al.* (2013) Changing patterns in place of cancer death in England: A population-based study. *PLoS Medicine* 10(3), e1001410. doi:10.1371/journal.pmed.1001410
- García-Sanjuán S, Fernández-Alcántara M, Clement-Carbonell V, *et al.* (2022) Levels and determinants of place-of-death congruence in palliative patients: A systematic review. *Frontiers in Psychology* 12, 80786. doi:10.3389/fpsyg.2021.807869
- Gisquet E, Julliard S and Geoffroy-Perez B (2016) Do social factors affect the place of death? Analysis of home versus institutional death over 20 years. *Journal of Public Health* 38(4), e472–e479. doi:10.1093/pubmed/fdv167
- Gomes B, Calanzani N, Gysels M, *et al.* (2013) Heterogeneity and changes in preferences for dying at home: A systematic review. *BMC Palliative Care* 12(1), 7. doi:10.1186/1472-684X-12-7
- Gomes B, Calanzani N, Koffman J, *et al.* (2015) Is dying in hospital better than home in incurable cancer and what factors influence this? A population-based study. *BMC Medicine* 13(101190723), 235. doi:10.1186/s12916-015-0466-5
- Gomes B and Higginson IJ (2006) Factors influencing death at home in terminally ill patients with cancer: Systematic review. *Bmj* 332(7540), 515–521. doi:10.1136/bmj.38740.614954.55
- Gomes B and Higginson IJ (2008) Where people die (1974–2030): Past trends, future projections and implications for care. *Palliative Medicine* 22(1), 33–41. doi:10.1177/0269216307084606
- Gomes B, Pinheiro MJ, Lopes S, *et al.* (2018) Risk factors for hospital death in conditions needing palliative care: Nationwide population-based death certificate study. *Palliative Medicine* 32(4), 891–901. doi:10.1177/0269216317743961
- Guerriere D, Husain A, Marshall D, *et al.* (2015) Predictors of place of death for those in receipt of home-based palliative care services in Ontario, Canada. *Journal of Palliative Care* 31(2), 76–88. doi:10.1177/082585971503100203
- Häkanson C, Öhlén J, Morin L, *et al.* (2015) A population-level study of place of death and associated factors in Sweden. *Scandinavian Journal of Public Health* 43(7), 744–751. doi:10.1177/1403494815595774

- Higginson IJ, Sarmiento VP, Calanzani N, *et al.* (2013) Dying at home—is it better: A narrative appraisal of the state of the science. *Palliative Medicine* 27(10), 918–924. doi:10.1177/0269216313487940
- Hoare S, Morris ZS, Kelly MP, *et al.* (2015) Do patients want to die at home? A systematic review of the UK literature, focused on missing preferences for place of death. *PLoS One* 10(11), e0142723. doi:10.1371/journal.pone.0142723
- Houttekier D, Cohen J, Pepersack T, *et al.* (2014) Dying in hospital: A study of incidence and factors related to hospital death using death certificate data. *European Journal of Public Health* 24(5), 751–756. doi:10.1093/eurpub/ckt209
- Howell DA, Wang HI, Roman E, *et al.* (2017) Preferred and actual place of death in haematological malignancy. *BMJ Supportive & Palliative Care* 7(2), 150–157. doi:10.1136/bmjspcare-2014-000793
- Howell DA, Wang H-I, Smith AG, *et al.* (2013) Place of death in haematological malignancy: Variations by disease sub-type and time from diagnosis to death. *BMC Palliative Care* 12(1), 42. doi:10.1186/1472-684X-12-42
- Hunt KJ, Shlomo N and Addington-Hall J (2014a) End-of-life care and achieving preferences for place of death in England: Results of a population-based survey using the VOICES-SF questionnaire. *Palliative Medicine* 28(5), 412–421. doi:10.1177/0269216313512012
- Hunt KJ, Shlomo N and Addington-Hall J (2014b) End-of-life care and preferences for place of death among the oldest old: results of a population-based survey using VOICES-short form. *Journal of Palliative Medicine* 17(2), 176–182. doi:10.1089/jpm.2013.0385
- Isenberg SR, Tanuseputro P, Spruin S, *et al.* (2020) Cost-effectiveness of investment in end-of-life home care to enable death in community settings. *Medical Care* 58(8), 665–673. doi:10.1097/mlr.0000000000001320
- Jarlbæk L (2017) *Dødsstedet I Danmark 2012-2014*. Nyborg: REHPA - Videncenter for Rehabilitering og Palliation.
- Jayaraman J and Joseph KS (2013) Determinants of place of death: A population-based retrospective cohort study. *BMC Palliative Care* 12(1), 19. doi:10.1186/1472-684X-12-19
- Jiang J and May P (2021) Proportion of deaths in hospital in European countries: Trends and associations from panel data (2005–2017). *European Journal of Public Health* 31(6), 1176–1183. doi:10.1093/eurpub/ckab169
- Johnson M, Allgar V, Chen H, *et al.* (2018) The complex relationship between household income of family caregivers, access to palliative care services and place of death: A national household population survey. *Palliative Medicine* 32(2), 357–365. doi:10.1177/0269216317711825
- Jordan RI, Elmokhallalati Y, Corless L, *et al.* (2023) Quality of end-of-life care with non-malignant liver disease: Analysis of the VOICES National Survey of Bereaved People. *Liver International* 43(2), 308–316. doi:10.1111/liv.15428
- Kalseth J and Halvorsen T (2020) Relationship of place of death with care capacity and accessibility: A multilevel population study of system effects on place of death in Norway. *BMC Health Services Research* 20(1), 1–12. doi:10.1186/s12913-020-05283-6
- Kamisetty A, Magennis P, Mayland C, *et al.* (2015) Where do patients treated for oral cancer die? A 20-year cohort study 1992–2011. *The British Journal of Oral & Maxillofacial Surgery* 53(10), 1015–1020. doi:10.1016/j.bjoms.2015.10.003
- Kern H, Corani G, Huber D, *et al.* (2020) Impact on place of death in cancer patients: A causal exploration in southern Switzerland. *BMC Palliative Care* 19(1), N.PAG–N.PAG. doi:10.1186/s12904-020-00664-4
- Kjellstadli C, Allore H, Husebo BS, *et al.* (2020) General practitioners' provision of end-of-life care and associations with dying at home: A registry-based longitudinal study. *Family Practice* 37(3), 340–347. doi:10.1093/fampra/cmz059
- Kjellstadli C, Han L, Allore H, *et al.* (2019) Associations between home deaths and end-of-life nursing care trajectories for community-dwelling people: A population-based registry study. *BMC Health Services Research* 19(1), 698. doi:10.1186/s12913-019-4536-9
- Ko W, Miccinesi G, Beccaro M, *et al.* (2014) Factors associated with fulfilling the preference for dying at home among cancer patients: The role of general practitioners. *Journal of Palliative Care* 30(3), 141–150. doi:10.1177/082585971403000303
- Koffman J, Ho YK, Davies J, *et al.* (2014) Does ethnicity affect where people with cancer die? A population-based 10 year study. *PLoS ONE* 9(4), e95052. doi:10.1371/journal.pone.0095052
- Larsen AMN, Neergaard MA, Andersen MF, *et al.* (2020) Increased rate of home-death among patients in a Danish general practice. *Danish Medical Journal* 67(10), 1–8.
- Livingston G, Lewis-Holmes E, Pitfield C, *et al.* (2013) Improving the end-of-life for people with dementia living in a care home: An intervention study. *International Psychogeriatrics* 25(11), 1849–1858. doi:10.1017/S1041610213001221
- Lovell N, Jones C, Baynes D, *et al.* (2017) Understanding patterns and factors associated with place of death in patients with end-stage kidney disease: A retrospective cohort study. *Palliative Medicine* 31(3), 283–288. doi:10.1177/0269216316655747
- Luta X, Panczak R, Maessen M, *et al.* (2016) Dying among older adults in Switzerland: Who dies in hospital, who dies in a nursing home? *BMC Palliative Care* 15, 1–11. doi:10.1186/s12904-016-0156-x
- Martinsson L, Lundström S and Sundelöf J (2020) Better quality of end-of-life care for persons with advanced dementia in nursing homes compared to hospitals: A Swedish national register study. *BMC Palliative Care* 19(1), 135. doi:10.1186/s12904-020-00639-5
- McCaughan D, Roman E, Smith AG, *et al.* (2018) Determinants of hospital death in haematological cancers: Findings from a qualitative study. *BMJ Supportive & Palliative Care* 8(1), 78–86. doi:10.1136/bmjspcare-2016-001289
- McCaughan D, Roman E, Smith AG, *et al.* (2019) Haematology nurses' perspectives of their patients' places of care and death: A UK qualitative interview study. *European Journal of Oncology Nursing* 39, 70–80. doi:10.1016/j.ejon.2019.02.003
- McEwen R, Asada Y, Burge F, *et al.* (2018) Associations between home death and the use and type of care at home. *Journal of Palliative Care* 33(1), 26–31. doi:10.1177/0825859717751933
- Mieras A, Pasman HRW, Onwuteaka-Philipsen BD, *et al.* (2019) Is in-hospital mortality higher in patients with metastatic lung cancer who received treatment in the last month of life? A retrospective cohort study. *Journal of Pain and Symptom Management* 58(5), 805–811. doi:10.1016/j.jpainsymman.2019.06.026
- Neergaard MA, Brunoe AH, Skorstengaard MH, *et al.* (2019) What socioeconomic factors determine place of death for people with life-limiting illness? A systematic review and appraisal of methodological rigour. *Palliative Medicine* 33(8), 900–925. doi:10.1177/0269216319847089
- Nieder C, Tollali T, Haukland E, *et al.* (2016) Impact of early palliative interventions on the outcomes of care for patients with non-small cell lung cancer. *Supportive Care in Cancer* 24(10), 4385–4391. doi:10.1007/s00520-016-3278-z
- Nilsson J, Axelsson B, Holgersson G, *et al.* (2020) Geographical differences in likelihood of home death among palliative cancer patients: A national population-based register study. *Anticancer Research* 40(7), 3897–3903. doi:10.21873/anticancer.14380
- Nilsson J, Holgersson G, Ullenhag G, *et al.* (2021) Socioeconomy as a prognostic factor for location of death in Swedish palliative cancer patients. *BMC Palliative Care* 20(1), 1–7. doi:10.1186/s12904-021-00736-z
- Nolasco A, Fernandez-Alcantara M, Pereyra-Zamora P, *et al.* (2020) Socioeconomic inequalities in the place of death in urban small areas of three Mediterranean cities. *International Journal for Equity in Health* 19(1), 214. doi:10.1186/s12939-020-01324-y
- Nysæter TM, Olsson C, Sandsdalen T, *et al.* (2022) Preferences for home care to enable home death among adult patients with cancer in late palliative phase – A grounded theory study. *BMC Palliative Care* 21(1), 1–10. doi:10.1186/s12904-022-00939-y
- Öhlén J, Cohen J and Håkanson C (2017) Determinants in the place of death for people with different cancer types: A national population-based study. *Acta Oncol* 56(3), 455–461. doi:10.1080/0284186x.2016.1250946
- Oosterveld-Vlug M, Donker G, Atsma F, *et al.* (2018) How do treatment aims in the last phase of life relate to hospitalizations and hospital mortality? A mortality follow-back study of Dutch patients with five types of cancer. *Supportive Care in Cancer* 26(3), 777–786. doi:10.1007/s00520-017-3889-z

- Orlovic M, Callender T, Riley J, *et al.* (2020) Impact of advance care planning on dying in hospital: Evidence from urgent care records. *PLoS ONE* **15**(12 December), e0242914. doi:10.1371/journal.pone.0242914
- Orlovic M, Marti J and Mossialos E (2017) Analysis of end-of-life care, out-of-pocket spending, and place of death in 16 European Countries And Israel. *Health Affairs* **36**(7), 1201–1210. doi:10.1377/hlthaff.2017.0166
- O'Sullivan EM and Higginson IJ (2016) 'I'll continue as long as I can, and die when I can't help it': A qualitative exploration of the views of end-of-life care by those affected by head and neck cancer (HNC). *BMJ Supportive & Palliative Care* **6**(1), 43–51. doi:10.1136/bmjspcare-2014-000664
- Pinzon LCE, Claus M, Perrar KM, *et al.* (2013) Dying with dementia: Symptom burden, quality of care, and place of death. *Deutsches Arzteblatt International* **110**(12), 195–202. doi:10.3238/arztebl.2013.0195
- Pollock K (2015) Is home always the best and preferred place of death? *BMJ: British Medical Journal* **351**, h4855. doi:10.1136/bmj.h4855
- Pooler C, Richman-Eisenstat J and Kalluri M (2018) Early integrated palliative approach for idiopathic pulmonary fibrosis: A narrative study of bereaved caregivers' experiences. *Palliative Medicine* **32**(9), 1455–1464. doi:10.1177/0269216318789025
- PRISMA (2023) PRISMA for scoping reviews. <http://www.prisma-statement.org/Extensions/ScopingReviews> (accessed 2023).
- Purdy S, Lasseter G, Griffin T, *et al.* (2015) Impact of the Marie Curie cancer care delivering choice programme in Somerset and North Somerset on place of death and hospital usage: A retrospective cohort study. *BMJ Supportive & Palliative Care* **5**(1), 34–39. doi:10.1136/bmjspcare-2013-000645
- Quinn KL, Hsu AT, Smith G, *et al.* (2020) Association between palliative care and death at home in adults with heart failure. *Journal of the American Heart Association* **9**(5), e013844. doi:10.1161/jaha.119.013844
- Rasch-Westin M, Helde-Frankling M and Bjorkhem-Bergman L (2019) Death at home: Predictive factors in a medical home care unit. *BMJ Supportive & Palliative Care* **14**, e350–e352. doi:10.1136/bmjspcare-2019-001932
- Raziee H, Saskin R and Barbera L (2017) Determinants of home death in patients with cancer: A population-based study in Ontario, Canada. *Journal of Palliative Care* **32**(1), 11–18. doi:10.1177/0825859717708518
- Reyniers T, Deliens L, Pasman HR, *et al.* (2016) Reasons for end-of-life hospital admissions: results of a survey among family physicians. *Journal of Pain and Symptom Management* **52**(4), 498–506. doi:10.1016/j.jpainsymman.2016.05.014
- Reyniers T, Houttekier D, Cohen J, *et al.* (2014) What justifies a hospital admission at the end of life? A focus group study on perspectives of family physicians and nurses. *Palliative Medicine* **28**(7), 941–948. doi:10.1177/0269216314522317
- Sayma M, Saleh D, Kerwat D, *et al.* (2020) A qualitative inquiry into the barriers and facilitators to achieving home death. *BMJ Supportive & Palliative Care* **10**(2), e18. doi:10.1136/bmjspcare-2016-001260
- Schnakenberg R, Fassmer AM, Allers K, *et al.* (2022) Characteristics and place of death in home care recipients in Germany – An analysis of nationwide health insurance claims data. *BMC Palliative Care* **21**(1), 1–8. doi:10.1186/s12904-022-01060-w
- Schou-Andersen M, Ullersted MP, Jensen AB, *et al.* (2016) Factors associated with preference for dying at home among terminally ill patients with cancer. *Scandinavian Journal of Caring Sciences* **30**(3), 466–476. doi:10.1111/scs.12265
- Seal K, Murray CD and Seddon L (2015) Family stories of end-of-life cancer care when unable to fulfill a loved one's wish to die at home. *Palliative and Supportive Care* **13**(3), 473–483. doi:10.1017/S1478951514000017
- Sepulveda JMG, Baid D, Johnson FR, *et al.* (2022) What is a good death? A choice experiment on care indicators for patients at end of life. *Journal of Pain and Symptom Management* **63**(4), 457–467. doi:10.1016/j.jpainsymman.2021.11.005
- Sharpe KH, Cezard G, Bansal N, *et al.* (2015) Policy for home or hospice as the preferred place of death from cancer: Scottish Health and Ethnicity Linkage Study population cohort shows challenges across all ethnic groups in Scotland. *BMJ Supportive & Palliative Care* **5**(4), 443–451. doi:10.1136/bmjspcare-2013-000485
- Sheridan R, Roman E, Smith AG, *et al.* (2021) Preferred and actual place of death in haematological malignancies: A report from the UK haematological malignancy research network. *BMJ Supportive and Palliative Care* **11**(1), 7–16. doi:10.1136/bmjspcare-2019-002097
- Skorstengaard MH, Jensen AB, Andreassen P, *et al.* (2020) Advance care planning and place of death, hospitalisation and actual place of death in lung, heart and cancer disease: A randomised controlled trial. *BMJ Supportive & Palliative Care* **10**(4), e37. doi:10.1136/bmjspcare-2018-001677
- Sleeman KE, Ho YK, Verne J, *et al.* (2014) Reversal of English trend towards hospital death in dementia: A population-based study of place of death and associated individual and regional factors, 2001–2010. *BMC Neurology* **14**(1), 59–59. doi:10.1186/1471-2377-14-59
- Stajduhar KI (2013) Burdens of family caregiving at the end of life. *Clinical and Investigative Medicine Medecine Clinique et Experimentale* **36**(3), E121–E126. doi:10.25011/cim.v36i3.19722
- Stajduhar KI (2020) Provocations on privilege in palliative care: Are we meeting our core mandate? *Progress in Palliative Care* **28**(2), 89–93. doi:10.1080/09699260.2019.1702334
- Stajduhar KI and Davies B (2005) Variations in and factors influencing family members' decisions for palliative home care. *Palliative Medicine* **19**(1), 21–32. doi:10.1191/0269216305pm963oa
- Tanuseputro P, Beach S, Chalifoux M, *et al.* (2018) Associations between physician home visits for the dying and place of death: A population-based retrospective cohort study. *PLoS ONE* **13**(2), e0191322. doi:10.1371/journal.pone.0191322
- Van Den Block L, Winne K, Miccinesi G, *et al.* (2017) Final transitions to place of death: Patients and families wishes. *Journal of Public Health* **39**(4), e302–e311. doi:10.1093/pubmed/fdw097
- van Doorne I, van Rijn M, Dofferhoff SM, *et al.* (2021) Patients' preferred place of death: Patients are willing to consider their preferences, but someone has to ask them. *Age Ageing* **50**(6), 2004–2011. doi:10.1093/ageing/afab176
- Van Spall HGC, Hill AD, Longdi F, *et al.* (2021) Temporal trends and sex differences in intensity of healthcare at the end of life in adults with heart failure. *Journal of the American Heart Association* **10**(1), 1–13. doi:10.1161/JAHA.120.018495
- Varani S, Dall'Olio FG, Messana R, *et al.* (2015) Clinical and demographic factors associated to the place of death in advanced cancer patients assisted at home in Italy. *Progress in Palliative Care* **23**(2), 61–67. doi:10.1179/1743291X14Y.00000000094
- Wahid AS, Sayma M, Jamshaid S, *et al.* (2018) Barriers and facilitators influencing death at home: A meta-ethnography. *Palliative Medicine* **32**(2), 314–328. doi:10.1177/0269216317713427
- Wales J, Kalia S, Moineddin R, *et al.* (2020) The impact of socioeconomic status on place of death among patients receiving home palliative care in Toronto, Canada: A retrospective cohort study. *Journal of Palliative Care* **35**(3), 167–173. doi:10.1177/0825859719855020
- Wales J, Kurahashi AM and Husain A (2018) The interaction of socioeconomic status with place of death: A qualitative analysis of physician experiences. *BMC Palliative Care* **17**(1), N.PAG–N.PAG. doi:10.1186/s12904-018-0341-1
- Wilson DM, Shen Y, Errasti-Ibarrondo B, *et al.* (2018) The location of death and dying across Canada: A study illustrating the socio-political context of death and dying. *Societies* **8**(4), 112.
- Wilson R, Hegglun N, Higginson IJ, *et al.* (2020) End-of-life care and place of death in adults with serious mental illness: A systematic review and narrative synthesis. *Palliative Medicine* **34**(1), 49–68. doi:10.1177/0269216319867847
- Winthereik AK, Hjertholm P, Neergaard MA, *et al.* (2018) Propensity for paying home visits among general practitioners and the associations with cancer patients' place of care and death: A register-based cohort study. *Palliative Medicine* **32**(2), 376–383. doi:10.1177/0269216317727387
- World Health Organization (2011) *Palliative Care for Older People: Better Practices*. Copenhagen: WHO Regional Office for Europe.
- Wye L, Lasseter G, Simmonds B, *et al.* (2016) Electronic palliative care coordinating systems (EPaCCS) may not facilitate home deaths: A mixed methods evaluation of end of life care in two English counties. *Journal of Research in Nursing* **21**(2), 96–107. doi:10.1177/1744987116628922
- Ziway SR, Samad D, Johnson CD, *et al.* (2017) Impact of place of residence on place of death in Wales: An observational study. *BMC Palliative Care* **16**, 1–6. doi:10.1186/s12904-017-0261-5