

# THE BLOCK STRUCTURE OF COMPLETE LATTICE ORDERED EFFECT ALGEBRAS

GEJZA JENČA

(Received 4 May 2005; revised 28 June 2006)

Communicated by M. Jackson

## Abstract

We prove that every for every complete lattice-ordered effect algebra  $E$  there exists an orthomodular lattice  $O(E)$  and a surjective full morphism  $\phi_E : O(E) \rightarrow E$  which preserves blocks in both directions: the (pre)image of a block is always a block. Moreover, there is a 0, 1-lattice embedding  $\phi_E^* : E \rightarrow O(E)$ .

2000 *Mathematics subject classification*: primary 06C15; secondary 03G12, 81P10.

## 1. Introduction

Effect algebras have recently been introduced by Foulis and Bennett in [9] for the study of the foundations of quantum mechanics. The class of effect algebras includes orthomodular lattices and a subclass equivalent to MV-algebras (see [4]).

In [30], Riečanová proved that every lattice-ordered effect algebra is a union of (essentially) MV-algebras. This result is a generalization of the well-known fact that every orthomodular lattice is a union of Boolean algebras. Generalizing the terminology from orthomodular lattices, a maximal sub-MV-effect algebra of a lattice-ordered effect algebra is called *a block*. Later, Riečanová and Jenča proved in [24] that the set of all sharp elements of a lattice-ordered effect algebra forms an orthomodular lattice. Both papers show that the class of lattice-ordered effect algebras generalizes the class of orthomodular lattices in a very natural way. In [18] a new class, called *homogeneous effect algebras* was introduced and most of the results from [30]

---

This research is supported by grant VEGA G-1/3025/06 of MŠ SR. This work was supported by the Slovak Research and Development Agency under the contract No. APVV-0071-06.

© 2007 Australian Mathematical Society 1446-7887/07 \$A2.00 + 0.00

and [24] were generalized for the homogeneous case. The main result of [18] is that every homogeneous effect algebra is a union of effect algebras satisfying the Riesz decomposition property.

Intuitively, one can consider the class of lattice-ordered effect algebras as an “unsharp” generalization of the class of orthomodular lattices, and the class of homogeneous effect algebras as an “unsharp” generalization of the class of orthoalgebras (see [10]). In these generalizations, the role of Boolean algebras is played by MV-effect algebras and by effect algebras with the Riesz decomposition property. The problems concerning this generalization were examined, for example, in [31] and [19]. The present paper continues this line of research.

The basic question we deal with in this paper is: “How are the blocks in a complete lattice-ordered effect algebra organized?”. The main result is that for every complete lattice-ordered effect algebra  $E$ , there exists an orthomodular lattice  $O(E)$  and a surjective full morphism of effect algebras  $\phi_E : O(E) \rightarrow E$  such that for every block  $B$  of  $O(E)$ ,  $\phi_E(B)$  is a block and for every block  $M$  of  $E$ ,  $\phi_E^{-1}(M)$  is a block of  $O(E)$ . This shows that the block structure of every complete lattice-ordered effect algebra is the same as the block structure of some orthomodular lattice. For the finite case, this result was proved in [19]. Moreover, we prove that the lattice  $E$  can be 0, 1-embedded into the lattice  $O(E)$ .

Our construction of  $O(E)$  is based on certain relations on the set of all quotients (that is, comparable pairs of elements) of  $E$ . We hope that it will be possible to adapt the techniques we have developed in this paper in order to deal with the more general orthocomplete non-lattice-ordered case. Most of the theorems were designed with this long-term goal in mind.

## 2. Definition and basic relationships

An *effect algebra* is a partial algebra  $(E; \oplus, 0, 1)$  with a binary partial operation  $\oplus$  and two nullary operations 0, 1 satisfying the following conditions.

- (E1) If  $a \oplus b$  is defined then  $b \oplus a$  is defined and  $a \oplus b = b \oplus a$ .
- (E2) If  $a \oplus b$  and  $(a \oplus b) \oplus c$  are defined then  $b \oplus c$  and  $a \oplus (b \oplus c)$  are defined and  $(a \oplus b) \oplus c = a \oplus (b \oplus c)$ .
- (E3) If  $a \oplus b = a \oplus c$  then  $b = c$ .
- (E4) If  $a \oplus b = 0$  then  $a = 0$ .
- (E5) For every  $a \in E$  there is an  $a' \in E$  such that  $a \oplus a' = 1$ .
- (E6) For every  $a \in E$ ,  $a \oplus 0 = a$ .

Effect algebras were introduced by Foulis and Bennett in their paper [9]. In the original paper, a different but equivalent set of axioms was used.

In their paper [26], Chovanec and Kôpka introduced an essentially equivalent structure called a *D-poset*. Their definition is an abstract algebraic version the *D-poset of fuzzy sets*, introduced by Kôpka in the paper [25]. Another equivalent structure was introduced by Giuntini and Greuling in [12]. We refer the reader to [7] for more information on effect algebras and related topics.

One can construct examples of effect algebras from an arbitrary partially ordered abelian group  $(G, \leq)$  in the following way: choose any positive  $u \in G$ ; then, for  $0 \leq a, b \leq u$ , define  $a \oplus b$  if and only if  $a + b \leq u$  and put  $a \oplus b = a + b$ . With such a partial operation  $\oplus$ , the interval  $[0, u]$  becomes an effect algebra  $([0, u], \oplus, 0, u)$ . Effect algebras which arise from partially ordered abelian groups in this way are called *interval effect algebras*, see [2].

In an effect algebra  $E$ , we write  $a \leq b$  if and only if there is a  $c \in E$  such that  $a \oplus c = b$ . It is easy to check that for every effect algebra the relation  $\leq$  is a partial order on  $E$ . Moreover, it is possible to introduce a new partial operation  $\ominus$ ;  $b \ominus a$  is defined if and only if  $a \leq b$  and then  $a \oplus (b \ominus a) = b$ . It can be proved that, in an effect algebra,  $a \oplus b$  is defined if and only if  $a \leq b'$ , if and only if  $b \leq a'$ . The partial operations  $\oplus$  and  $\ominus$  are connected by the rules

$$(2.1) \quad a \oplus b = (a' \ominus b)'$$

$$(2.2) \quad a \ominus b = (a' \oplus b)'$$

Let  $E_0 \subseteq E$  be such that  $1 \in E_0$  and, for all  $a, b \in E_0$  with  $a \geq b$ ,  $a \ominus b \in E_0$ . Since  $a' = 1 \ominus a$  and  $a \oplus b = (a' \ominus b)'$ ,  $E_0$  is closed with respect to  $\oplus$  and  $'$ . We then say that  $(E_0, \oplus, 0, 1)$  is a *subeffect algebra of  $E$* . Another way to construct a substructure of an effect algebra  $E$  is to restrict  $\oplus$  to an interval  $[0, a]$ , where  $a \in E$ , letting  $a$  act as the unit element. We denote such effect algebra by  $[0, a]_E$ .

EXAMPLE 1. Let  $G$  be the set of all real functions, partially ordered in the usual way. Let  $u$  be the constant function  $u(x) = 1$ . Then the restriction of  $+$  from  $G$  to the set  $[0, u]$  gives rise to an effect algebra, which we denote by  $[0, 1]^{[0,1]}$ . Note that  $[0, 1]^{[0,1]}$  is a complete distributive lattice.

Let  $E_1, E_2$  be effect algebras. A map  $\phi : E_1 \rightarrow E_2$  is called a *morphism of effect algebras* if and only if it satisfies the following condition.

(M1)  $\phi(1) = 1$  and, for all  $a, b \in E_1$ , if  $a \oplus b$  exists in  $E_1$  then  $\phi(a) \oplus \phi(b)$  exists in  $E_2$  and  $\phi(a \oplus b) = \phi(a) \oplus \phi(b)$ .

Every morphism preserves  $'$ ,  $0$ ,  $\leq$  and  $\ominus$ .

A morphism  $\phi : E_1 \rightarrow E_2$  of effect algebras is called *full* if and only if the following condition is satisfied.

(M2) If  $\phi(a) \oplus \phi(b)$  exists in  $E_2$  and  $\phi(a) \oplus \phi(b) \in \phi(E_1)$  then there exist  $a_1, b_1 \in E_1$  such that  $a_1 \oplus b_1$  exists in  $E_1$  and  $\phi(a) = \phi(a_1)$  and  $\phi(b) = \phi(b_1)$ .

A bijective and full morphism is called an *isomorphism of effect algebras*.

An ideal of an effect algebra  $E$  is a subset  $I$  of  $E$  satisfying the condition

$$a, b \in I \quad \text{and} \quad a \oplus b \text{ exists} \quad \iff \quad a \oplus b \in I.$$

The set of all ideals of an effect algebra  $E$  is denoted by  $I(E)$ .  $I(E)$  is a complete lattice with respect to inclusion.

An element  $c$  of an effect algebra is *central* (see [14]) if and only if  $[0, c]$  is an ideal and, for every  $x \in E$ , there is a decomposition  $x = x_1 \oplus x_2$  such that  $x_1 \leq c$  and  $x_2 \leq c'$ . It can be shown that this decomposition is unique. The set  $C(E)$  of all central elements of an effect algebra is called *the centre of  $E$* .  $C(E)$  is a Boolean algebra. For every central element  $c$  of  $E$ ,  $E$  is isomorphic to  $[0, c]_E \times [0, c']_E$ . For every central element  $c$  of  $E$  and every element  $a \in E$ ,  $a \wedge c$  exists and the mapping  $a \mapsto a \wedge c$  is a full morphism from  $E$  onto  $[0, c]_E$ ; in other words,  $(a_1 \oplus a_2) \wedge c = (a_1 \wedge c) \oplus (a_2 \wedge c)$  and  $a = (a \wedge c) \oplus (a \wedge c')$ .

If  $E$  is an effect algebra such that  $(E, \leq)$  is a lattice, then we say that  $E$  is a *lattice ordered effect algebra*. If  $(E, \leq)$  is a complete lattice, then we say that  $E$  is a *complete lattice ordered effect algebra*. An *orthoalgebra*  $E$  (see [11]) is an effect algebra such that  $a \leq a'$  implies  $a = 0$ . It is easy to check that an effect algebra  $E$  is an orthoalgebra if and only if  $a \wedge a' = 0$  for all  $a \in E$ .

EXAMPLE 2. Recall, that an *orthomodular lattice* is an algebra  $(O; \vee, \wedge, ', 0, 1)$  such that  $(O; \vee, \wedge, 0, 1)$  is a bounded lattice and  $a \leq b$  if and only if  $b' \leq a'$ ,  $a'' = a$ ,  $a \wedge a' = 0$ ,  $(a \vee b)' = a' \wedge b'$ , and the orthomodular law

$$(2.3) \quad a \leq b \quad \implies \quad b = a \vee (b \wedge a')$$

is satisfied. Equip  $O$  with a partial operation as follows:  $a \oplus b$  is defined if and only if  $a \leq b'$  and then  $a \oplus b = a \vee b$ . Then  $(O; \oplus, 0, 1)$  is an orthoalgebra. On the other hand, for every lattice-ordered orthoalgebra  $(O; \oplus, 0, 1)$ ,  $(O; \vee, \wedge, ', 0, 1)$  is an orthomodular lattice. However, there exist non-lattice-ordered orthoalgebras.

EXAMPLE 3. An *MV-algebra* (see [4, 28]) is a commutative semigroup  $(M; \oplus, \neg, 0)$  satisfying the identities  $x \oplus 0 = x$ ,  $\neg\neg x = x$ ,  $x \oplus \neg 0 = \neg 0$  and

$$x \oplus \neg(x \oplus \neg y) = y \oplus \neg(y \oplus \neg x).$$

There is a natural partial order in an MV-algebra, given by  $y \leq x$  if and only if  $x = x \oplus \neg(x \oplus \neg y)$ . Every MV-algebra  $(M; \oplus, \neg, 0)$  can be considered as an effect algebra  $(M; \oplus, 0, \neg 0)$ , when we restrict the operation  $\oplus$  to the domain  $\{(x, y) : x \leq \neg y\}$ .

Every lattice-ordered effect algebra satisfies the de Morgan laws. More generally, for an interval  $[0, a]$  and  $x, y \in [0, a]$ , we have

$$(2.4) \quad a \ominus (x \vee y) = (a \ominus x) \wedge (a \ominus y)$$

$$(2.5) \quad a \ominus (x \wedge y) = (a \ominus x) \vee (a \ominus y).$$

A substitution  $a = b'$  and an application of (2.2) now yields

$$(2.6) \quad b \oplus (x \vee y) = (b \oplus x) \vee (b \oplus y)$$

$$(2.7) \quad b \oplus (x \wedge y) = (b \oplus x) \wedge (b \oplus y),$$

for all  $x, y \leq b'$ .

Let  $E$  be a lattice-ordered effect algebra. For a pair of elements  $a, b \in E$ , the following conditions are equivalent:

- $a \ominus (a \wedge b) = (a \vee b) \ominus b$ .
- $a \ominus (a \wedge b) \leq b'$ .
- There are  $a_1, a_2$  such that  $a = a_1 \leq a_2, a_1 \leq b, a_2 \leq b'$ .
- There are  $a_1, c, b_1$  such that  $a_1 \oplus c \oplus b_1$  exists and  $a = a_1 \oplus c$  and  $b = b_1 \oplus c$ .

If  $a, b$  satisfy any (or, equivalently, all) of these conditions then we say that  $a, b$  are *compatible* (in symbols  $a \leftrightarrow b$ ). It is easy to check that  $a \leq b$  or  $a \leq b'$  implies that  $a \leftrightarrow b$ . Moreover,  $a \leftrightarrow b$  if and only if  $a \leftrightarrow b'$ . We say that a subset  $A \subseteq E$  is compatible if and only if we have  $a \leftrightarrow b$  for all  $a, b \in A$ . If  $M$  is a lattice ordered effect algebra such that  $M$  is a compatible subset of  $M$  then we say that  $M$  is an *MV-effect algebra*. It was proved in [5] that there is a natural, one-to-one correspondence between MV-effect algebras and MV-algebras, as outlined in Example 3. Every MV-effect algebra is a distributive lattice. A lattice-ordered effect algebra is an MV-effect algebra if and only if, for all elements  $a, b$ ,

$$a \wedge b = 0 \implies a \leq b',$$

that means, the sum of every disjoint pair exists (see [1]). An orthoalgebra that is an MV-effect algebra is a *Boolean algebra*.

Let  $L$  be a lattice. We say that  $L_0 \subseteq L$  is a *full sublattice* of  $L$  if and only if

- for all  $A \subseteq L_0$  such that  $\bigvee A$  exists in  $L, \bigvee A \in L_0$ , and
- for all  $A \subseteq L_0$  such that  $\bigwedge A$  exists in  $L, \bigwedge A \in L_0$ .

Note that a full sublattice of a complete lattice is complete.

Let  $E$  be a lattice-ordered effect algebra. A subeffect algebra of  $E$  that is maximal with respect to the property of being an MV-effect algebra is called a *block*. According to [30], blocks coincide with maximal compatible subsets of  $E$ . Moreover, every block is a full sublattice of  $E$ , (see [24]). Since every singleton is a compatible set, a lattice-ordered effect algebra is a union of its blocks.

If  $L$  is a compatible sublattice of a lattice-ordered effect algebra, then there is a block  $M \supseteq L$ . Since  $M$  is a distributive lattice,  $L$  is distributive as well.

If  $E$  is an orthomodular lattice then every block is a Boolean algebra. Thus, the fact that every lattice-ordered effect algebra is a union of its blocks is a generalization of the well-known fact that every orthomodular lattice is a union of Boolean algebras.

We say that an element  $a$  of an effect algebra is *sharp* if and only if  $a \wedge a' = 0$ . We write  $S(E)$  for the set of all sharp elements of an effect algebra  $E$ . An orthoalgebra can be characterized by  $E = S(E)$ . Every central element is sharp, hence  $C(E) \subseteq S(E)$ . In general,  $S(E)$  is not closed with respect to  $\oplus$ , (see [15]). However, by [24], if  $E$  is lattice ordered, then  $S(E)$  is a subeffect algebra and a full sublattice of  $E$ . For a block  $M$  of a lattice-ordered effect algebra, we have  $S(E) \cap M = S(M) = C(M)$ , (see [24]).

An effect algebra  $E$  is called *homogeneous* if and only if, for all  $u, v_1, v_2 \in E$  such that  $u \leq v_1 \oplus v_2 \leq u'$ , there are  $u_1, u_2$  such that  $u_1 \leq v_1, u_2 \leq v_2$ , and  $u = u_1 \oplus u_2$ . Homogeneous effect algebras were introduced in [18]. Every orthoalgebra and every lattice-ordered effect algebra is homogeneous. The set of all sharp elements of a homogeneous effect algebra is closed with respect to  $\oplus$ , hence it forms an orthoalgebra.

In [18], most of the results (concerning compatibility, blocks, and sharp elements) from [30] and [24] were generalized for the homogeneous case. The situation is more complicated here. In a homogeneous effect algebra, the blocks need not be lattice-ordered anymore (they only satisfy the *Riesz decomposition property*) and the notion of compatibility has to be generalized as well.

EXAMPLE 4. Let  $B$  be a Boolean algebra with at least two elements. Let us equip  $B \times B$  with a partial  $\oplus$  operation as follows:  $\langle x_1, x_2 \rangle \oplus \langle y_1, y_2 \rangle$  is defined if and only if  $x_1 \wedge y_2 = x_2 \wedge y_1 = 0$  and then

$$\langle x_1, x_2 \rangle \oplus \langle y_1, y_2 \rangle = \langle x_1 \vee y_1 \vee (x_2 \wedge y_2), x_2 \vee y_2 \vee (x_1 \wedge y_1) \rangle.$$

Then  $(B \times B; \oplus, \langle 0, 0 \rangle, \langle 1, 1 \rangle)$  is an effect algebra denoted by  $D^B$ . In  $D^B$ , we have  $\langle x_1, x_2 \rangle \leq \langle y_1, y_2 \rangle$  if and only if  $x_1 \leq y_1$  and  $x_2 \leq y_2$ . Thus,  $D^B$  is the same lattice as the “ordinary” Boolean lattice  $B \times B$ . However, if  $B$  has more than one element then  $D^B$  is not an MV-effect algebra: we have  $\langle 1, 0 \rangle \wedge \langle 0, 1 \rangle = 0$  but  $\langle 1, 0 \rangle \oplus \langle 0, 1 \rangle$  does not exist.

It is easy to check that

$$\langle x_1, x_2 \rangle' = \langle x_2', x_1' \rangle$$

and that

$$S(D^B) = \{ \langle x, x \rangle : x \in B \}.$$

Note that  $S(D^B)$  is a Boolean algebra. This implies that  $S(D^B) = C(D^B)$ . Since, for  $|B| > 1$ , there are unsharp elements in  $D^B$ ,  $D^B$  is not an orthomodular lattice.

### 3. Generalized test spaces

In this section we present a slightly generalized version of the notion of *test space*, originally introduced by Foulis and Randall in [11] and [29]. Despite of its relative simplicity, the notion of test space (and its generalizations) is a very useful tool for construction of orthoalgebras and effect algebras. See for example [6], [8], or [19] for constructions that use test spaces.

Let  $X$  be a nonempty set and let  $\mathcal{N}, \mathcal{T}$  be subsets of  $2^X$ . We say that a triple  $(X, \mathcal{T}, \mathcal{N})$  is a *generalized test space* if and only if the following conditions are satisfied.

(GTS1)  $X = \cup_{t \in \mathcal{T}} t$ .

(GTS2)  $\mathcal{N}$  is an ideal of  $2^X$ , that is,  $\mathcal{N}$  is nonempty and for all  $\mathbf{o}_1, \mathbf{o}_2 \subseteq X$  we have  $\mathbf{o}_1 \cup \mathbf{o}_2 \in \mathcal{N}$  if and only if  $\mathbf{o}_1, \mathbf{o}_2 \in \mathcal{N}$ .

(GTS3) For all  $t_1 \subseteq t_2 \subseteq X$  such that  $t_1 \in \mathcal{T}$ , we have  $t_2 \in \mathcal{T}$  if and only if  $t_2 \setminus t_1 \in \mathcal{N}$ .

(GTS4) For all  $t_1 \subseteq t_2 \subseteq X$  such that  $t_2 \setminus t_1 \in \mathcal{N}$ , we have  $t_1 \in \mathcal{T}$  if and only if  $t_2 \in \mathcal{T}$ .

A generalized test space is a *test space* if and only if  $\mathcal{N} = \{\emptyset\}$ . For a test space, the Axioms (GTS2) and (GTS4) collapse to tautologies and (GTS3) transforms to

(TS) If  $t_1, t_2 \in \mathcal{T}$  and  $t_1 \subseteq t_2$ , then  $t_1 = t_2$ .

(GTS1) and (TS) are the original axioms of a test space.

EXAMPLE 5. Let  $X$  be the system of all measurable subsets of the real interval  $[0, 1]_{\mathbb{R}}$  and let  $\mathcal{T}$  be the set of all finite systems  $t \subseteq X$  such that the elements of  $t$  are measurable, pairwise disjoint and  $\mu(\dot{\cup}_{A \in t} A) = 1$ . Let  $\mathcal{N}$  be the set of all finite pairwise disjoint systems of sets with zero measure. Then  $(X, \mathcal{T}, \mathcal{N})$  is a generalized test space.

Throughout this section, we assume that  $(X, \mathcal{T}, \mathcal{N})$  is a generalized test space.

An element of  $\mathcal{T}$  is called a *test*. Since  $X$  is nonempty, (GTS1) implies that there is at least one test. We say that a subset  $f$  of  $X$  is an *event* if and only if there is a test  $t \in \mathcal{T}$  such that  $f \subseteq t$ .

LEMMA 3.1. *Every element of  $\mathcal{N}$  is an event.*

PROOF. Let  $t_1$  be a test and let  $\mathbf{o} \in \mathcal{N}$ . Put  $t_2 = t_1 \cup \mathbf{o}$ . We have  $t_2 \setminus t_1 \subseteq \mathbf{o} \in \mathcal{N}$ . Since  $\mathcal{N}$  is an ideal of sets,  $t_2 \setminus t_1 \in \mathcal{N}$ . By (GTS3),  $t_1 \in \mathcal{T}$  and  $t_2 \setminus t_1 \in \mathcal{N}$  imply that  $t_2$  is a test. Thus,  $\mathbf{o} \subseteq t_2$  is an event. □

The elements of  $\mathcal{N}$  are called *null events*.

We say that two events  $f, g$  of a generalized test space  $(X, \mathcal{T}, \mathcal{N})$  are

- (i) *orthogonal* (in symbols  $f \perp g$ ) if and only if  $f \cap g \in \mathcal{N}$  and  $f \cup g$  is an event.
- (ii) *local complements* (in symbols  $f \text{ loc } g$ ) if and only if  $f \cap g \in \mathcal{N}$  and  $f \cup g$  is a test.
- (iii) *perspective* (in symbols  $f \sim g$ ) if and only if they share a local complement.

Note that every pair of tests is perspective, since  $\emptyset$  is a local complement of every test.

LEMMA 3.2. *For an event  $f$  we have  $f \sim \emptyset$  if and only if  $f \in \mathcal{N}$ .*

PROOF. Suppose that  $f \sim \emptyset$ . There is a test  $t$  such that  $f \text{ loc } t$ . Since  $f \cup t$  is a test, (GTS3) implies that  $f \cup t \setminus t = f \setminus f \cap t$  is a null event. Since  $f \text{ loc } t$ ,  $f \cap t$  is a null event. Therefore,  $f = (f \setminus f \cap t) \cup (f \cap t)$  is a null event.

Suppose that  $f \in \mathcal{N}$ . Since  $f$  is an event, there is a test  $t \supseteq f$ . Since  $t \cap f = f \in \mathcal{N}$  and  $t \cup f = t \in \mathcal{T}$ ,  $f \text{ loc } t$ . Since  $t$  is a test,  $\emptyset \text{ loc } t$ . Therefore,  $f \sim \emptyset$ . □

LEMMA 3.3. *Let  $t$  be a test and let  $f$  be an event such that  $f \sim t$ . Then  $f$  is a test.*

PROOF. Let  $h$  be a local complement shared by  $f$  and  $t$ . Both  $t \cup h$  and  $t$  are tests. By (GTS3),  $t \cup h \setminus t \in \mathcal{N}$ . Since  $t \text{ loc } h$ ,  $t \cap h \in \mathcal{N}$ . Therefore,  $h = (t \cup h \setminus t) \cup (t \cap h) \in \mathcal{N}$  and, since  $f \cup h \setminus f \subseteq h$ ,  $f \cup h \setminus f \in \mathcal{N}$ . By (GTS4),  $f \cup h \in \mathcal{T}$  and  $f \cup h \setminus f \in \mathcal{N}$  imply that  $f \in \mathcal{T}$ . □

We say that a generalized test space is *algebraic* if and only if for all events  $f, g, h$  we have

$$f \sim g \quad \text{and} \quad f \text{ loc } h \quad \implies \quad g \text{ loc } h.$$

LEMMA 3.4. *In an algebraic generalized test space, perspectivity is transitive.*

PROOF. Suppose that  $f \sim g \sim h$ . There are events  $u_1, u_2$  such that  $f \text{ loc } u_1 \text{ loc } g$  and  $g \text{ loc } u_2 \text{ loc } h$ . Since  $u_2 \text{ loc } h$  and  $u_1 \sim u_2$ ,  $u_1 \text{ loc } h$ . Therefore,  $f \sim h$ . □

Note that, in an algebraic generalized test space, both  $\mathcal{N}$  and  $\mathcal{T}$  are equivalence classes of  $\sim$ .

LEMMA 3.5. *In an algebraic generalized test space, if  $f_1 \sim f_2$  and  $f_1 \perp g$  then  $f_2 \perp g$  and  $f_1 \cup g \sim f_2 \cup g$ .*

PROOF. Let  $h$  be a local complement of  $f_1 \cup g$ . Since  $f_1 \text{ loc } h \cup g$  and  $f_1 \sim f_2$ ,  $f_2 \text{ loc } h \cup g$ . This implies that  $f_2 \perp g$ . Moreover, as  $f_1 \cup g \text{ loc } h \text{ loc } f_2 \cup g$ ,  $f_1 \cup g \sim f_2 \cup g$ . □

**THEOREM 3.6.** *Let  $(X, \mathcal{T}, \mathcal{N})$  be an algebraic generalized test space. Let  $\mathcal{E}$  be the set of all events of  $(X, \mathcal{T}, \mathcal{N})$ . Define on  $\mathcal{E}/\sim$  a relation  $\perp$  and a partial operation  $\oplus$  with domain  $\perp$  in the following way:  $[\mathbf{f}]_{\sim} \perp [\mathbf{g}]_{\sim}$  if and only if  $\mathbf{f} \perp \mathbf{g}$  and then  $[\mathbf{f}]_{\sim} \oplus [\mathbf{g}]_{\sim} = [\mathbf{f} \cup \mathbf{g}]_{\sim}$ . Then  $(\mathcal{E}/\sim, \oplus, \mathcal{N}, \mathcal{T})$  is an orthoalgebra.*

**PROOF.** Let us prove that  $\perp$  and  $\oplus$  are well-defined. Suppose that  $\mathbf{f}_1 \sim \mathbf{f}_2$  and  $\mathbf{g}_1 \sim \mathbf{g}_2$  and  $\mathbf{f}_1 \perp \mathbf{g}_1$ . By Lemma 3.5,  $\mathbf{f}_1 \perp \mathbf{g}_2$  and  $\mathbf{f}_1 \cup \mathbf{g}_1 \sim \mathbf{f}_1 \cup \mathbf{g}_2$ . Again, by Lemma 3.5, this implies that  $\mathbf{f}_2 \perp \mathbf{g}_2$  and  $\mathbf{f}_1 \cup \mathbf{g}_2 \sim \mathbf{f}_2 \cup \mathbf{g}_2$ .

(E1) is trivially true, so let us prove (E2). If both sides of the associative equality exist, they are (obviously) equal. Suppose that  $([\mathbf{f}]_{\sim} \oplus [\mathbf{g}]_{\sim}) \oplus [\mathbf{h}]_{\sim}$  exists. Then  $\mathbf{f} \perp \mathbf{g}$  and  $([\mathbf{f}]_{\sim} \oplus [\mathbf{g}]_{\sim}) = [\mathbf{f} \cup \mathbf{g}]_{\sim}$ . Therefore  $\mathbf{f} \cup \mathbf{g} \perp \mathbf{h}$  and we see that  $[\mathbf{f}]_{\sim} \oplus ([\mathbf{g}]_{\sim} \oplus [\mathbf{h}]_{\sim})$  exists.

Let us prove (E3). Suppose that  $[\mathbf{f}]_{\sim} \oplus [\mathbf{h}]_{\sim} = [\mathbf{g}]_{\sim} \oplus [\mathbf{h}]_{\sim}$ , that means,  $\mathbf{f}, \mathbf{g} \perp \mathbf{h}$  and  $\mathbf{f} \cup \mathbf{h} \sim \mathbf{g} \cup \mathbf{h}$ ; let  $\mathbf{u}$  be a common local complement of  $\mathbf{f} \cup \mathbf{h}$  and  $\mathbf{g} \cup \mathbf{h}$ . We see that  $\mathbf{h} \cup \mathbf{u}$  is a common local complement of  $\mathbf{f}$  and  $\mathbf{g}$ , therefore  $\mathbf{f} \sim \mathbf{g}$  and  $[\mathbf{f}]_{\sim} = [\mathbf{g}]_{\sim}$ .

(E4) is trivial, so let us prove (E5). For an event  $\mathbf{f}$ ,  $[\mathbf{f}]_{\sim}$  is just the set of all local complements of  $\mathbf{f}$ . Obviously,  $[\mathbf{f}]_{\sim}$  can be characterized by the property  $[\mathbf{f}]_{\sim} \oplus [\mathbf{f}]'_{\sim} = \mathcal{T}$ .

The proof of (E6) is trivial.

Finally, let us prove that  $[\mathbf{f}]_{\sim} \leq [\mathbf{f}]'_{\sim}$  implies  $[\mathbf{f}]_{\sim} = \mathcal{N}$ . Since  $\mathbf{f} \perp \mathbf{f}$ , we have  $\mathbf{f} \cap \mathbf{f} = \mathbf{f} \in \mathcal{N}$ . Therefore  $[\mathbf{f}]_{\sim} = \mathcal{N}$  and we see that  $(\mathcal{E}/\sim, \oplus, \mathcal{N}, \mathcal{T})$  is an orthoalgebra. □

The orthoalgebra  $(\mathcal{E}/\sim, \oplus, \mathcal{N}, \mathcal{T})$  is called *the orthoalgebra of the generalized test space  $(X, \mathcal{T}, \mathcal{N})$ .*

### 4. The generalized test space of quotients

In this section we introduce our main tool. For every homogeneous effect algebra  $E$  we shall construct a generalized test space  $\Omega(E)$ , where tests are finite sets of comparable pairs (called *quotients*) with certain properties.

The origins of the notion of a quotient and the relations  $\nearrow$  and  $\searrow$  lie in lattice theory, see for example [13, Section III.1]. However, the definitions of  $\nearrow$  and  $\searrow$  introduced here do not coincide with their lattice-theoretical versions, even in the case of a lattice-ordered effect algebra (see the remark following Proposition 4.3). In the case of an MV-effect algebra, our definitions coincide with their lattice-theoretical counterparts.

Let  $E$  be an effect algebra and let  $P$  be a subposet of  $E$ . Let  $a/b$  denote an ordered pair of elements of  $P$  satisfying  $a \geq b$ . We say that  $a/b$  is a *quotient of  $P$* . The set

of all quotients of  $P$  is denoted by  $Q(P)$ . We say that  $c/d$  is a subquotient of  $a/b$  (in symbols  $c/d \sqsubseteq a/b$  or  $a/b \supseteq c/d$ ) if and only if  $b \leq d \leq c \leq a$ .

If  $a > b$ , we say that  $a/b$  is *proper*, otherwise we say that  $a/b$  is *null*.

We write  $a/b \nearrow c/d$  if and only if  $a \leq c, b \leq d, c \ominus a = d \ominus b$  and  $(c \ominus a) \wedge (a \ominus b) = 0$ . We write  $a/b \searrow c/d$  if and only if  $c \leq a, d \leq b, a \ominus c = b \ominus d$  and  $(a \ominus c) \wedge (a \ominus b) = 0$ . It is easy to check that  $a/b \searrow c/d$  or  $a/b \nearrow c/d$  implies that  $a \ominus b = c \ominus d$ .

**PROPOSITION 4.1.** *Let  $E$  be an effect algebra and let  $a/b, c/d \in Q(E)$ . The following are equivalent:*

- (i)  $a/b \searrow c/d$ ;
- (ii)  $c/d \nearrow a/b$ ;
- (iii)  $b'/a' \nearrow d'/c'$ ;
- (iv)  $d'/c' \searrow b'/a'$ .

**PROOF.** The proofs of the equivalence of (i) and (ii) and of (iii) and (iv) are trivial.

Let us prove that (i) implies (iii). Since  $c \leq a, a' \leq c'$ . Since  $d \leq b, b' \leq d'$ . We have

$$d' \ominus b' = (d' \ominus b')'' = (d \oplus b')' = (b' \oplus d)' = b \ominus d.$$

Similarly,  $c' \ominus a' = a \ominus c$  and  $b' \ominus a' = a \ominus b$ . Therefore,

$$d' \ominus b' = b \ominus d = a \ominus c = c' \ominus a'$$

and

$$(d' \ominus b') \wedge (b' \ominus a') = (b \ominus d) \wedge (a \ominus b) = (a \ominus c) \wedge (a \ominus b) = 0.$$

Let us prove that (iii) implies (i). By the previous parts of the proof,

$$b'/a' \nearrow d'/c' \implies d'/c' \searrow b'/a' \implies c/d \nearrow a/b \implies a/b \searrow c/d. \quad \square$$

**EXAMPLE 6.** Let  $[0, 1]^{[0,1]}$  be the effect algebra of all real functions of a real variable  $[0, 1] \rightarrow [0, 1]$ . For  $a/b, c/d \in Q([0, 1]^{[0,1]})$  we have  $a/b \searrow c/d$  if and only if, for all  $x \in [0, 1]$ ,  $a(x) \neq c(x)$  or  $b(x) \neq d(x)$  imply  $a(x) = b(x) \geq c(x) = d(x)$ .

For example, we may take

$$\begin{aligned} a(x) &= |x - 0.5| + 0.5, & c(x) &= 1 - x, \\ b(x) &= x, & d(x) &= 0.5 - |x - 0.5| \end{aligned}$$

(see Figure 1).

**EXAMPLE 7.** Let  $E$  be a 6-element effect algebra with two atoms  $a, b$ , satisfying  $a \oplus a \oplus a = a \oplus b \oplus b = 1$ . On  $Q(E)$ , the  $\searrow$  relation is not transitive: we have  $1/a' \searrow a \oplus b/b$  and  $a \oplus b/b \searrow a/0$ , but  $1/a' \not\searrow a/0$ , because  $a \wedge a' = a \neq 0$ .




FIGURE 1.  $a/b \searrow c/d$  in  $[0, 1]^{[0,1]}$

PROPOSITION 4.2. For every homogeneous effect algebra,  $\nearrow$  and  $\searrow$  are transitive.

PROOF. Let  $E$  be a homogeneous effect algebra and let  $a/b, c/d, e/f \in Q(E)$ . Assume that  $a/b \nearrow c/d$  and  $c/d \nearrow e/f$ . Obviously  $a \leq e$  and  $b \leq f$ . We see that

$$e \ominus a = (e \ominus c) \oplus (c \ominus a) = (f \ominus d) \oplus (d \ominus b) = f \ominus b.$$

Suppose that  $x \leq e \ominus a, a \ominus b$ . Since  $e \ominus a \leq (a \ominus b)' \leq x'$  and  $e \ominus a = (e \ominus c) \oplus (c \ominus a)$ , we obtain

$$x \leq (e \ominus c) \oplus (c \ominus a) \leq x'.$$

Since  $E$  is homogeneous, there exist  $x_1 \leq e \ominus c$  and  $x_2 \leq c \ominus a$  such that  $x = x_1 \oplus x_2$ . However, as  $c \ominus d = a \ominus b$ , we have  $(e \ominus c) \wedge (c \ominus d) = (e \ominus c) \wedge (a \ominus b) = 0$  and thus  $x_1 \leq e \ominus c, a \ominus b$  implies that  $x_1 = 0$ . As  $x_2 \leq c \ominus a, a \ominus b$  and  $(c \ominus a) \wedge (a \ominus b) = 0$ ,  $x_2 = 0$ . Therefore,  $(e \ominus a) \wedge (a \ominus b) = 0$  and  $a/b \nearrow e/f$ .

Assume that  $a/b \searrow c/d$  and  $c/d \searrow e/f$ . By Proposition 4.1, this is equivalent to  $b'/a' \nearrow d'/c'$  and  $d'/c' \nearrow f'/e'$ . Since  $\nearrow$  is transitive,  $b'/a' \nearrow f'/e'$  and hence  $a/b \searrow e/f$ . □

PROPOSITION 4.3. Let  $E$  be a lattice-ordered effect algebra and let  $a/b, c/d \in Q(E)$ . Then

- (i)  $a/b \nearrow c/d$  if and only if  $a \leftrightarrow d, a \vee d = c, a \wedge d = b$ ,
- (ii)  $a/b \searrow c/d$  if and only if  $c \leftrightarrow b, c \vee b = a, c \wedge b = d$ .

PROOF. (i) Suppose that  $a/b \nearrow c/d$ . Since  $(c \ominus a) \oplus (a \ominus b) \oplus b$  exists, the set  $\{c \ominus a, a \ominus b, b\}$  is compatible and can be embedded into a block  $M$ . As  $a = (a \ominus b) \oplus b$ ,  $a \in M$ . Since  $c \ominus a = d \ominus b$ , we see that  $d = (d \ominus b) \oplus b = (c \ominus a) \oplus b \in M$ . Therefore,  $a \leftrightarrow d$ . We see that

$$c \ominus (a \vee d) = (c \ominus a) \wedge (c \ominus d) = (d \ominus b) \wedge (a \ominus b) = 0.$$

Therefore  $c = a \vee d$ .

Since  $a \leftrightarrow d$  and  $c \ominus d = a \ominus b$ ,

$$c \ominus d = (a \vee d) \ominus d = a \ominus (a \wedge d) = a \ominus b.$$

Therefore  $a \wedge d = b$ .

Suppose that  $a \leftrightarrow d$ ,  $a \vee d = c$  and  $a \wedge d = b$ . Obviously,  $a \wedge d \leq a$  and  $d \leq a \vee d$ . Since  $a \leftrightarrow d$ ,  $(a \vee d) \ominus a = d \ominus (a \wedge b)$  and it is easy to check that  $[a \ominus (a \wedge d)] \wedge [d \ominus (a \wedge d)] = 0$ .

(ii) This follows from (i) by a permutation of  $\{a, b, c, d\}$ . □

REMARK. In lattice theory, the relation  $\nearrow$  is defined by the rule

$$a/b \nearrow c/d \iff a \vee d = c \quad \text{and} \quad a \wedge d = b$$

and  $\searrow$  is defined in a dual way. By Proposition 4.3, our “effect-algebraic  $\nearrow$ ” is more restrictive than the original lattice-theoretical  $\nearrow$ . Both definitions coincide for MV-effect algebras, because in this case the additional condition  $a \leftrightarrow d$  is clearly satisfied.

In what follows, the symbol  $\equiv$  denotes the transitive closure of  $(\searrow \cup \nearrow)$ . Obviously,  $\equiv$  is an equivalence relation.

EXAMPLE 8. Let  $a/b, c/d \in Q([0, 1]^{[0,1]})$ . We have  $a/b \equiv c/d$  if and only if for all  $x \in [0, 1]$

$$a(x) \neq c(x) \quad \text{or} \quad b(x) \neq d(x) \implies a(x) = b(x) \quad \text{and} \quad c(x) = d(x).$$

We say that quotients  $a/b$  and  $c/d$  are *disjoint* if and only if for all  $x/y$  and  $z/w$ ,

$$a/b \supseteq x/y \equiv z/w \subseteq c/d \implies x = y.$$

We say that  $a/b$  and  $c/d$  are *orthogonal* (in symbols  $a/b \perp c/d$ ) if and only if  $a/b$  and  $c/d$  are disjoint and  $(a \ominus b) \oplus (c \ominus d)$  exists in  $E$ . We say that a finite set  $\mathbf{f}$  of quotients is *pairwise orthogonal* if and only if any two distinct elements of  $\mathbf{f}$  are orthogonal. We say that a finite set of quotients  $\mathbf{f} = \{a_1/b_1, \dots, a_n/b_n\}$  is *orthogonal* if and only if  $\mathbf{f}$  is pairwise orthogonal and the sum  $|\mathbf{f}|$  defined by

$$|\mathbf{f}| = (a_1 \ominus b_1) \oplus \dots \oplus (a_n \ominus b_n)$$

exists in  $E$ .

EXAMPLE 9. In  $[0, 1]^{[0,1]}$ , we have  $a/b \perp c/d$  if and only if, for all  $x \in [0, 1]$ , the intervals  $(b(x), a(x)]$  and  $(d(x), c(x)]$  are disjoint.




FIGURE 2.  $a/b \perp c/d$  in  $[0, 1]^{[0,1]}$

EXAMPLE 10. In an orthoalgebra, we have  $a/b \perp c/d$  if and only if  $(a \ominus b) \oplus (c \ominus d)$  exists.

Note that, for  $a/b \nearrow c/d$  and  $x/y \sqsubseteq a/b$ , there is  $x_0/y_0 \sqsubseteq c/d$  with  $x/y \nearrow x_0/y_0$ . Indeed, we may put  $x_0 = x \oplus (c \ominus a)$  and  $y_0 = y \oplus (c \ominus a)$ . There is an analogous relationship between  $\searrow$  and  $\sqsubseteq$ .

PROPOSITION 4.4. Let  $E$  be a homogeneous effect algebra and let  $a/b, c/d, e/f \in Q(E)$ . If  $a/b \equiv c/d$  and  $c/d$  is disjoint with  $e/f$ , then  $a/b$  is disjoint with  $e/f$ .

PROOF. Suppose that  $a/b \nearrow c/d$ . Let  $x/y$  and  $z/w$  be such that

$$a/b \sqsupseteq x/y \equiv z/w \sqsubseteq e/f.$$

There is  $x_0/y_0 \sqsubseteq c/d$  with  $x/y \nearrow x_0/y_0$ . However, since  $c/d \sqsupseteq x_0/y_0 \equiv z/w \sqsubseteq e/f$  and  $c/d, e/f$  are disjoint,  $x_0 = y_0$  and hence  $x = y$ .

Similarly,  $a/b \searrow c/d$  implies that  $a/b$  is disjoint with  $e/f$ . The rest of the proof is a simple induction. □

Let  $E$  be a homogeneous effect algebra. Let us extend the relation  $\equiv$  to the set of all finite subsets of  $Q(E)$ : for two finite sets of quotients  $\mathbf{f}$  and  $\mathbf{g}$  we write  $\mathbf{f} \equiv \mathbf{g}$  if and only if the following symmetric pair of conditions is satisfied.

- For every proper  $a/b \in \mathbf{f}$  there is exactly one  $c/d \in \mathbf{g}$  such that  $a/b \equiv c/d$ .
- For every proper  $a/b \in \mathbf{g}$  there is exactly one  $c/d \in \mathbf{f}$  such that  $a/b \equiv c/d$ .

It is obvious that  $\equiv$  is an equivalence relation on the set of all finite sets of quotients. Note that  $\mathbf{f} \equiv \emptyset$  if and only if  $\mathbf{f}$  contains only null quotients.

LEMMA 4.5. Let  $E$  be a homogeneous effect algebra and let  $\mathbf{f}, \mathbf{g}$  be finite sets of quotients. If  $\mathbf{f} \equiv \mathbf{g}$  and  $\mathbf{f}$  is (pairwise) orthogonal, then  $\mathbf{g}$  is (pairwise) orthogonal.

PROOF. Suppose that  $\mathbf{f}$  is pairwise orthogonal. Let  $a_1/b_1, a_2/b_2 \in \mathbf{g}$  such that  $a_1/b_1 \neq a_2/b_2$ . If one of  $a_1/b_1, a_2/b_2$  is null then  $a_1/b_1 \perp a_2/b_2$ , so let us assume

that both  $a_1/b_1, a_2/b_2$  are proper. Since  $\mathbf{f} \equiv \mathbf{g}$ , there are  $c_1/d_1, c_2/d_2 \in \mathbf{f}$  such that  $c_1/d_1 \equiv a_1/b_1, c_2/d_2 \equiv a_2/b_2$  and  $c_1/d_1 \neq c_2/d_2$ . Since  $\mathbf{f}$  is pairwise orthogonal,  $c_1/d_1 \perp c_2/d_2$ . Therefore,  $a_1/b_1 \perp a_2/b_2$ .

Suppose that  $\mathbf{f}$  is orthogonal. Then  $\mathbf{g}$  is pairwise orthogonal and it remains to observe that the elements occurring in the sum  $|\mathbf{f}|$  are (up to some zeros) the same as the elements occurring in the sum  $|\mathbf{g}|$ . Therefore,  $\mathbf{g}$  is orthogonal.  $\square$

LEMMA 4.6. *Let  $E$  be a homogeneous effect algebra and let*

$$\mathbf{f} = \{a_1/b_1, \dots, a_n/b_n\} \subseteq Q(E)$$

*be a pairwise orthogonal  $n$ -element set. Let*

$$\mathbf{g} = \{c_1/d_1, \dots, c_n/d_n\} \subseteq Q(E)$$

*be a finite set such that, for all  $i \in \{1, \dots, n\}$ ,  $c_i/d_i \equiv a_i/b_i$ . Then  $\mathbf{f} \equiv \mathbf{g}$ .*

PROOF. We have to prove that, for proper  $a_j/b_j$  and  $c_i/d_i$ ,  $a_j/b_j \equiv c_i/d_i$  implies that  $i = j$ . Suppose that  $i \neq j$ . Since  $\mathbf{f}$  is pairwise orthogonal,  $a_i/b_i \perp a_j/b_j$ . However,  $a_i/b_i \equiv c_i/d_i \equiv a_j/b_j$ . Therefore,  $a_i/b_i \equiv a_j/b_j$  and this is a contradiction with  $a_i/b_i \perp a_j/b_j$ .  $\square$

Note that we cannot omit the assumption that  $\mathbf{f}$  is pairwise orthogonal from Lemma 4.6. To see this, let  $\mathbf{f} = \{a_1/b_1, a_2/b_2\}$  be such that  $a_1/b_1 \equiv a_2/b_2$  and  $a_1/b_1$  is proper. Then for  $\mathbf{g} = \{a_1/b_1, a_1/b_1\}$  we have  $\mathbf{g} \neq \mathbf{f}$ .

Let  $E$  be a homogeneous effect algebra. Let  $\mathcal{T}$  be the set of all finite orthogonal sets  $\mathbf{t} \subseteq Q(E)$  with  $|\mathbf{t}| = 1$  and let  $\mathcal{N}$  be the set of all finite sets of null quotients. We define a triple  $\Omega(E)$  by  $\Omega(E) = (Q(E), \mathcal{T}, \mathcal{N})$ . It is evident that  $\Omega(E)$  forms a generalized test space. Note that, for all events  $\mathbf{f}, \mathbf{g}$  of  $\Omega(E)$ ,  $\mathbf{f} \equiv \mathbf{g}$  implies that  $\mathbf{f} \sim \mathbf{g}$ .

The main aim of the following two sections of this paper is to prove the following theorem.

THEOREM 4.7. *Let  $E$  be a complete lattice-ordered effect algebra. Let  $\mathbf{f}$  be a finite set of quotients of  $E$ . Then the following are equivalent:*

- (a)  $\mathbf{f}$  is an event of  $\Omega(E)$ ;
- (b)  $\mathbf{f}$  is an orthogonal set of quotients;
- (c)  $\mathbf{f}$  is a pairwise orthogonal set of quotients.

It will then turn out that  $\Omega(E)$  is an algebraic generalized test space. Later we shall prove that the orthoalgebra  $O(E)$  of  $\Omega(E)$  is actually an orthomodular lattice with the same block structure as  $E$ .

### 5. Reduced quotients

Let  $E$  be an effect algebra and let  $a/b \in Q(E)$ . We say that  $a/b$  is a *reduced quotient* if and only if

$$a/b \searrow c/d \implies a/b = c/d.$$

Note that  $a/b$  is reduced if and only if  $x \leq b$  and  $x \wedge (a \oplus b) = 0$  imply  $x = 0$ .

A null quotient  $a/a$  is reduced if and only if  $a = 0$ . In an orthoalgebra, a proper quotient  $a/b$  is reduced if and only if  $b = 0$ . On the other hand, in a totally ordered effect algebra every proper quotient is reduced.

We say that a finite set  $\{a_1/b_1, \dots, a_n/b_n\}$  of quotients is *compatible* if and only if  $\{a_1, b_1, \dots, a_n, b_n\}$  is a compatible set.

The aim of this section is to show that for every pairwise orthogonal finite set of quotients in a complete lattice-ordered effect algebra there exists a compatible pairwise orthogonal finite set  $\mathbf{f}_R$  of reduced quotients with  $\mathbf{f} \equiv \mathbf{f}_R$ .

EXAMPLE 11. A quotient  $a/b \in Q([0, 1]^{[0,1]})$  is reduced if and only if

$$a(x) = b(x) \implies a(x) = b(x) = 0.$$

An effect algebra  $E$  is *sharply dominating* if and only if, for every  $x \in E$ , the element  $x^\uparrow$  defined by

$$x^\uparrow = \bigwedge \{t : t \in [x, 1] \cap S(E)\}$$

exists and is sharp. It is easy to see that in a sharply dominating effect algebra  $E$ , the element  $x^\downarrow$  defined by

$$x^\downarrow = \bigvee \{t : t \in [0, x] \cap S(E)\}$$

exists and is sharp for all  $x \in E$ . Moreover, we have  $x^{\uparrow'} = x'^\downarrow$  and  $x^{\downarrow'} = x'^\uparrow$ . We say that  $x^\uparrow$  is the *sharp cover* of  $x$  and that  $x^\downarrow$  is the *sharp kernel* of  $x$ . In his paper [3], Cattaneo proved that for every sharply dominating effect algebra the set of all sharp elements forms a subeffect algebra which is an orthoalgebra. See [16] for another version of the proof.

EXAMPLE 12. The lattice-ordered effect algebra  $D^B$  from Example 4 is sharply dominating, even if  $B$  is incomplete. We have

$$\begin{aligned} \langle x_1, x_2 \rangle^\uparrow &= \langle x_1 \vee x_2, x_1 \vee x_2 \rangle \quad \text{and} \\ \langle x_1, x_2 \rangle^\downarrow &= \langle x_1 \wedge x_2, x_1 \wedge x_2 \rangle. \end{aligned}$$

We say that an effect algebra  $E$  is *orthocomplete* if and only if every chain has a supremum in  $E$ . See [23] and [22] for results on orthocomplete effect algebras. A lattice-ordered effect algebra is orthocomplete if and only if it is a complete lattice.

**PROPOSITION 5.1.** ([21, Corollary 5]) *Every orthocomplete homogeneous effect algebra is sharply dominating. Moreover, for every block  $M$ ,  $x \in M$  implies that  $[x^\downarrow, x], [x, x^\uparrow] \subseteq M$ .*

Since all complete lattice-ordered effect algebras are orthocomplete and homogeneous, we may apply Proposition 5.1 for them. Note that Proposition 5.1 implies that every subset of a complete lattice-ordered effect algebra of the form  $[x^\downarrow, x] \cup [x, x^\uparrow]$  is compatible.

**LEMMA 5.2.** *Let  $E$  be a complete lattice-ordered effect algebra. For all  $y \in E$ ,  $y^\downarrow = y^{\uparrow'}$  is the greatest element of the set*

$$(5.1) \quad \{x \in E : x \leq y' \text{ and } y \wedge x = 0\}.$$

**PROOF.** Let us prove that  $y^\downarrow$  is the upper bound of the set (5.1). For every  $x$ ,  $x \leq y'$ . Therefore,  $x \leftrightarrow y$  and there is a block  $M \supseteq \{y, x\}$ . By [21, Lemma 1],  $y \wedge x = 0$  implies that  $y^\uparrow \wedge x = 0$ . Since  $M$  is an MV-effect algebra, this implies that  $x \leq y^{\uparrow'} = y^\downarrow$ .

Since  $y^\downarrow \leq y'$  and

$$y \wedge y^\downarrow = y \wedge y^{\uparrow'} \leq y^\uparrow \wedge y^{\uparrow'} = 0,$$

$y^\downarrow$  belongs to (5.1). □

**PROPOSITION 5.3.** *Let  $E$  be a complete lattice-ordered effect algebra and let  $a/b \in Q(E)$ . The following are equivalent:*

- (i)  $a/b$  is reduced;
- (ii)  $b \wedge (a \ominus b)^{\uparrow'} = 0$ ;
- (iii)  $a \leq (a \ominus b)^\uparrow$ ;
- (iv)  $b \leq (a \ominus b)^\uparrow \ominus (a \ominus b)$ .

**PROOF.** (i)  $\implies$  (ii): Suppose that  $x \leq b$ ,  $x \leq (a \ominus b)^{\uparrow'}$ . Since  $x \leq b$ ,  $x \leq (a \ominus b)'$ . By Lemma 5.2,  $x \leq (a \ominus b)^{\uparrow'}$  implies that  $x \wedge (a \ominus b) = 0$ . Since  $a/b$  is reduced, this implies that  $x = 0$ .

(ii)  $\implies$  (iii): Let  $M$  be a block of  $E$  with  $a, b \in M$ . By Proposition 5.1,  $(a \ominus b)^\uparrow \in M$  and hence  $(a \ominus b)^{\uparrow'} \in M$ . Since  $(a \ominus b)^{\uparrow'}$  is sharp,  $(a \ominus b)^{\uparrow'}$  is central in  $M$ . Thus, we may compute

$$a \wedge (a \ominus b)^{\uparrow'} = ((a \ominus b) \oplus b) \wedge (a \ominus b)^{\uparrow'} = ((a \ominus b) \wedge (a \ominus b)^{\uparrow'}) \oplus (b \wedge (a \ominus b)^{\uparrow'}).$$

By Lemma 5.2,  $(a \ominus b) \wedge (a \ominus b)^{\uparrow'} = 0$ . By assumption,  $b \wedge (a \ominus b)^{\uparrow'} = 0$ . Since  $a \wedge (a \ominus b)^{\uparrow'} = 0$  and  $M$  is an MV-effect algebra,  $a \leq (a \ominus b)^{\uparrow''} = (a \ominus b)^{\uparrow}$ .

(iii)  $\implies$  (iv): We see that  $a = (a \ominus b) \oplus b \leq (a \ominus b)^{\uparrow}$ , hence  $b \leq (a \ominus b)^{\uparrow} \ominus (a \ominus b)$ .

(iv)  $\implies$  (i): Suppose that  $x \leq b, x \wedge (a \ominus b) = 0$ . Since  $x \leq b \leq (a \ominus b)^{\uparrow} \ominus (a \ominus b)$ , we have  $x \leq (a \ominus b)'$ . By Lemma 5.2,  $x \leq (a \ominus b)'$  and  $x \wedge (a \ominus b) = 0$  imply  $x \leq (a \ominus b)^{\downarrow} = (a \ominus b)^{\uparrow'}$ . Since  $(a \ominus b)^{\uparrow} \wedge (a \ominus b)^{\uparrow'} = 0, x = 0$ .  $\square$

The following lemma is crucial.

LEMMA 5.4. *Let  $E$  be a complete lattice-ordered effect algebra. Let  $a/b$  be a reduced quotient of  $E$  and let  $M$  be a block of  $E$  with  $a \ominus b \in M$ . Then  $a, b \in M$ .*

PROOF. By Proposition 5.3,  $a \leq (a \ominus b)^{\uparrow}$ . This implies that  $a \in [a \ominus b, (a \ominus b)^{\uparrow}]$ . Therefore, by Proposition 5.1,  $a \in M$ . Since  $a, (a \ominus b) \in M, b = a \ominus (a \ominus b) \in M$ .  $\square$

COROLLARY 5.5. *Let  $E$  be a complete lattice-ordered effect algebra, let  $\mathbf{f}$  be a finite set of reduced quotients of  $E$  such that  $\{a \ominus b : a/b \in \mathbf{f}\}$  is compatible. Then  $\mathbf{f}$  is a compatible set of quotients.*

PROOF. Let  $a/b, c/d \in \mathbf{f}$ . Since  $a \ominus b \leftrightarrow c \ominus d$ , there exists a block  $M$  with  $a \ominus b, c \ominus d \in M$ . By Lemma 5.4,  $a, b, c, d \in M$ . Therefore,  $\{a/b, c/d\}$  is a compatible set of quotients. Thus,  $\mathbf{f}$  is a compatible set of quotients.  $\square$

COROLLARY 5.6. *Every reduced pairwise orthogonal finite set of quotients of a complete lattice-ordered effect algebra is compatible.*

PROOF. This follows from Corollary 5.5.  $\square$

EXAMPLE 13. If  $a/b$  is a quotient of  $[0, 1]^{[0,1]}$  then  $a^R/b^R$  is given by

$$a^R(x) = \begin{cases} a(x) & \text{if } a(x) > b(x), \\ 0 & \text{if } a(x) = b(x), \end{cases} \quad b^R(x) = \begin{cases} b(x) & \text{if } a(x) > b(x), \\ 0 & \text{if } a(x) = b(x). \end{cases}$$

(See Figure 3.)

Let  $E$  be a complete lattice-ordered effect algebra. Let us introduce a mapping  $R : Q(E) \rightarrow Q(E)$ , given by  $a/b \mapsto a^R/b^R$ , where  $a^R = a \wedge (a \ominus b)^{\uparrow}$  and  $b^R = b \wedge (a \ominus b)^{\uparrow}$ . We say that  $a^R/b^R$  is the reduct of  $a/b$ .

PROPOSITION 5.7. *Let  $E$  be a complete lattice-ordered effect algebra. For every  $a/b \in Q(E)$ ,  $a/b \searrow a^R/b^R$  and  $a^R/b^R$  is reduced.*




FIGURE 3.  $a/b$  and  $a^R/b^R$  in  $[0, 1]^{0,11}$

PROOF. Obviously,  $a^R \leq a$  and  $b^R \leq b$ . Let  $M$  be a block with  $a, b \in M$ . By Proposition 5.1,  $(a \ominus b)^\dagger \in M$ . Since  $(a \ominus b)^\dagger$  is sharp,  $(a \ominus b)^\dagger$  is central in  $M$ . Therefore,

$$\begin{aligned} a \ominus a^R &= a \ominus (a \wedge (a \ominus b)^\dagger) = a \wedge (a \ominus b)^{\dagger'} = (b \oplus (a \ominus b)) \wedge (a \ominus b)^{\dagger'} \\ &= (b \wedge (a \ominus b)^{\dagger'}) \oplus ((a \ominus b) \wedge (a \ominus b)^{\dagger'}) = b \wedge (a \ominus b)^{\dagger'} \\ &= b \ominus (b \wedge (a \ominus b)^\dagger) = b \ominus b^R. \end{aligned}$$

Moreover,

$$(a \ominus b) \wedge (a \ominus a^R) = (a \ominus b) \wedge a \wedge (a \ominus b)^{\dagger'} = 0.$$

Let us prove that  $a^R/b^R$  is reduced. By Proposition 5.3, this is equivalent to  $b^R \wedge (a^R \ominus b^R)^{\dagger'} = 0$ . Since  $a/b \searrow a^R/b^R$ ,  $a \ominus b = a^R \ominus b^R$ . Thus,

$$b^R \wedge (a^R \ominus b^R)^{\dagger'} = b \wedge (a \ominus b)^\dagger \wedge (a \ominus b)^{\dagger'} = 0. \quad \square$$

PROPOSITION 5.8. Let  $E$  be a complete lattice ordered effect algebra and let  $a/b, c/d \in Q(E)$ . Then  $a/b \equiv c/d$  if and only if  $a^R/b^R = c^R/d^R$ .

PROOF. If  $a^R/b^R = c^R/d^R$  then  $a/b \searrow a^R/b^R = c^R/d^R \nearrow c/d$ .

Suppose that  $a/b \nearrow c/d$ . By Proposition 4.3,  $a \leftrightarrow d$  so there is a block  $M$  with  $\{a, b, c, d\} \subseteq M$ . By Proposition 5.1,  $(c \ominus d)^\dagger \in M \cap S(E) = C(M)$ . Therefore,

$$\begin{aligned} c^R &= c \wedge (c \ominus d)^\dagger = ((c \ominus a) \oplus a) \wedge (c \ominus d)^\dagger \\ &= ((c \ominus a) \wedge (c \ominus d)^\dagger) \oplus (a \wedge (c \ominus d)^\dagger). \end{aligned}$$

By Lemma 5.2,  $c \ominus a \leq (c \ominus d)'$  and  $(c \ominus a) \wedge (c \ominus d) = 0$  imply that

$$c \ominus a \leq (c \ominus d)^{\dagger'} = (c \ominus d)^{\dagger'}.$$

Therefore,  $(c \ominus a) \wedge (c \ominus d)^\dagger = 0$  and

$$c^R = a \wedge (c \ominus d)^\dagger = a \wedge (a \ominus b)^\dagger = a^R.$$

As a consequence,  $d^R = c^R \ominus (c \ominus d) = a^R \ominus (a \ominus b) = b^R. \quad \square$

Let  $E$  be a complete lattice-ordered effect algebra. For a finite  $n$ -element set  $\mathbf{f} = \{a_1/b_1, \dots, a_n/b_n\}$  we write  $\mathbf{f}^R = \{a_1^R/b_1^R, \dots, a_n^R/b_n^R\}$ .

**PROPOSITION 5.9.** *Let  $E$  be a complete lattice-ordered effect algebra. Let  $\mathbf{f}$  be a finite pairwise orthogonal set of quotients. Then  $\mathbf{f} \equiv \mathbf{f}^R$  and  $\mathbf{f}^R$  is a pairwise orthogonal compatible set of quotients.*

**PROOF.** By Lemma 4.6,  $\mathbf{f} \equiv \mathbf{f}^R$ . By Lemma 4.5,  $\mathbf{f}^R$  is pairwise orthogonal. By Corollary 5.5,  $\mathbf{f}^R$  is compatible. □

### 6. Compatible sets of quotients

In this section we are going to prove a restriction of Theorem 4.7 for compatible events, (see Proposition 6.6.) Using Proposition 5.9, it is then possible to extend the result to the general case.

Let  $D$  be a bounded distributive lattice. Up to isomorphism, there exists a unique Boolean algebra  $B(D)$  such that  $D$  is a 0, 1-sublattice of  $B(D)$  and  $D$  generates  $B(D)$  as a Boolean algebra. This Boolean algebra is called an  $R$ -generated Boolean algebra. We refer to [13, Section II.4] for an overview of results concerning  $R$ -generated Boolean algebras. See also [17] and [27]. For every element  $x$  of  $B(D)$ , there exists a finite chain  $x_1 \leq \dots \leq x_n$  in  $D$  such that  $x = x_1 + \dots + x_n$ , where  $+$  denotes the symmetric difference, as in Boolean rings. We then say that  $\{x_i\}_{i=1}^n$  is a  $D$ -chain representation of  $x$ . It is easy to see that every element of  $B(D)$  has a  $D$ -chain representation of even length.

Note that if  $D_1$  is a 0, 1-sublattice of a distributive lattice  $D_2$  then  $B(D_1)$  is a subalgebra of  $B(D_2)$ .

**LEMMA 6.1.** ([20, Lemma 7]). *Let  $L$  be a finite 0, 1-sublattice of an MV-effect algebra  $M$ . The mapping  $\phi_L : B(L) \rightarrow M$  given by*

$$(6.1) \quad \phi_L(x) = \bigoplus_{i=1}^n (x_{2i} \ominus x_{2i-1}),$$

where  $\{x_i\}_{i=1}^{2n}$  is a  $L$ -chain representation of  $x$ , is a faithful surjective homomorphism of effect algebras. The value of  $\phi_L(x)$ , as given by (6.1), does not depend on the choice of  $\{x_i\}_{i=1}^{2n}$ .

Note that, since every compatible 0, 1-sublattice  $L$  of a lattice-ordered effect algebra is a sublattice of some block  $M$ , Lemma 6.1 is true even if we merely suppose that  $L$  is a compatible 0, 1-sublattice of a lattice-ordered effect algebra.

Let  $L$  be a lattice. An element  $a$  of  $L$  is *join-irreducible* if and only if  $a = b \vee c$  implies that  $a = b$  or  $a = c$ ; it is *meet-irreducible* if and only if  $a = b \wedge c$  implies that  $a = b$  or  $a = c$ . The set of all nonzero join-irreducible elements of a lattice  $L$  is denoted by  $J(L)$  and the set of all non-unit meet-irreducible elements of a lattice  $L$  is denoted by  $M(L)$ .

Let  $L$  be a finite distributive lattice. Then the mapping  $r : L \rightarrow 2^{J(L)}$  given by  $r(x) = \{a \in J(L) : a \leq x\}$  is a 0, 1-embedding of  $L$  into  $2^{J(L)}$ . Since, for every finite distributive lattice  $L$ ,  $r(L)$  R-generates  $2^{J(L)}$ , the injective mapping  $r : L \rightarrow 2^{J(L)}$  uniquely extends to an isomorphism of Boolean algebras  $\hat{r} : B(L) \rightarrow 2^{J(L)}$ .

In what follows,  $\succ_P$  denotes the usual covering relation on a poset  $P$ , so that  $a \succ_P b$  if and only if  $b$  is a maximal element of the set  $\{x \in P : x < a\}$ . In a finite distributive lattice  $L$ , we have  $a \succ_L b$  if and only if  $\hat{r}(a) \setminus \hat{r}(b)$  is a singleton.

Let  $L$  be a finite distributive lattice. We have  $a \in J(L)$  if and only if there is a unique  $b$  such that  $a \succ_L b$ . Therefore  $\{a + b : a \succ_L b, a \in J(L)\}$  is the set of all atoms of  $B(L)$ .

Let  $L$  be a finite 0, 1-sublattice of a lattice-ordered effect algebra  $E$ , and let  $\mathbf{f} = \{a_1/b_1, \dots, a_n/b_n\}$  be a compatible set of quotients such that  $\mathbf{f} \subseteq Q(L)$ . We write

$$+\mathbf{f} = a_1 + b_1 + \dots + a_n + b_n,$$

where the  $+$  on the right-hand side is taken in  $B(L)$ .

**LEMMA 6.2.** *Let  $E$  be a complete lattice-ordered effect algebra and let  $\{a/b, c/d\}$  be compatible. Let  $L \supseteq \{a, b, c, d\}$  be a finite compatible 0, 1-sublattice of  $E$ . Then  $a/b \equiv c/d$  implies that  $a + b = c + d$  in  $B(L)$ .*

**PROOF.** By Proposition 5.8,  $a/b \equiv c/d$  implies that  $a^R/b^R = c^R/d^R$ . Let  $M$  be a block of  $E$  such that  $L \subseteq M$ . Since  $a \oplus b \in M$ , Lemma 5.4 implies  $a^R/b^R \in Q(M)$ . Therefore  $\{a/b, c/d, a^R/b^R = c^R/d^R\}$  is a compatible set of quotients. Let  $L_1$  be a finite compatible lattice such that  $L \subseteq L_1 \subseteq M$  and  $\{a^R, b^R\} \subseteq L_1$ .

By Proposition 5.7,  $a/b \searrow a^R/b^R$ . By Proposition 4.3,  $b \vee a^R = a$  and  $b \wedge a^R = b^R$ . Therefore, we may calculate in  $B(L_1)$  that

$$a + b = (b \vee a^R) + b = a^R + (b \wedge a^R) = a^R + b^R.$$

Similarly,  $c/d \searrow c^R/d^R$  implies that  $c + d = c^R + d^R$ . Therefore

$$a + b = a^R + b^R = c^R + d^R = c + d$$

in  $B(L_1)$  and, since  $B(L)$  is a subalgebra of  $B(L_1)$ ,  $a + b = c + d$  in  $B(L)$ . □

PROPOSITION 6.3. *Let  $E$  be a complete lattice-ordered effect algebra and let  $\{a/b, c/d\}$  be compatible. Let  $L \supseteq \{a, b, c, d\}$  be a finite compatible 0, 1-sublattice of  $E$ . Suppose that  $a \succ_L b$ . Then  $a/b \equiv c/d$  if and only if  $a + b = c + d$  in  $B(L)$ .*

PROOF. Suppose that  $a + b = c + d$ . Since  $a \succ_L b$ ,  $\hat{r}(a) \setminus \hat{r}(b)$  is a singleton. Let  $e \in \hat{r}(a) \setminus \hat{r}(b)$ . Since  $e$  is join-irreducible and nonzero, there is a single element  $f \in L$  such that  $e \succ_L f$ . We claim that  $a/b \searrow e/f$ . Indeed,

$$a \wedge (b \vee e) = (a \wedge b) \vee (a \wedge e) = b \vee e,$$

hence  $a \geq b \vee e \geq b$ . Since  $a \succ_L b$ , we have either  $b \vee e = a$  or  $b \vee e = b$ . However,  $b \vee e = b$  implies that  $e \in \hat{r}(b)$ , which contradicts  $e \in \hat{r}(a) \setminus \hat{r}(b)$ . Therefore  $b \vee e = a$ .

Since  $L$  is distributive, the intervals  $[b, b \vee e]$  and  $[b \wedge e, e]$  are isomorphic. As  $b \vee e = a \succ_L b$ ,  $e \succ_L b \wedge e$ . Since  $e$  is join-irreducible and nonzero,  $e$  covers exactly one element, hence  $b \wedge e = f$ . We have proved that  $b \vee e = a$  and  $a \wedge e = f$ . By Proposition 4.3,  $a/b \searrow e/f$ .

Since  $a + b = c + d$ ,  $\hat{r}(a) \setminus \hat{r}(b) = \hat{r}(c) \setminus \hat{r}(d)$  and  $\hat{r}(c) \setminus \hat{r}(d)$  is a singleton. This implies that  $c \succ_L d$  and, as for  $a/b$ , we deduce  $c/d \searrow e/f$ . Therefore,  $a/b \searrow e/f \nearrow c/d$  and  $a/b \equiv c/d$ .

The converse implication follows by Lemma 6.2. □

LEMMA 6.4. *Let  $E$  be a complete lattice-ordered effect algebra and let  $\{a/b, c/d\}$  be compatible. Let  $L \supseteq \{a, b, c, d\}$  be a finite compatible 0, 1-sublattice of  $E$ . Then  $a/b$  and  $c/d$  are orthogonal if and only if  $a + b$  and  $c + d$  are disjoint in  $B(L)$ .*

PROOF. Suppose that  $a/b$  and  $c/d$  are orthogonal and that  $a + b$  and  $c + d$  are not disjoint in  $B(L)$ . Then there exists  $e \in (\hat{r}(a) \setminus \hat{r}(b)) \cap (\hat{r}(c) \setminus \hat{r}(d))$ .

Let  $a_0/b_0, \dots, a_n/b_n$  and  $c_0/d_0, \dots, c_k/d_k$  be sequences of quotients of  $L$  such that

$$a = a_0 \succ_L b_0 = a_1 \succ_L b_1 = a_2 \succ_L \dots \succ_L b_{n-1} = a_n \succ_L b_n = b$$

and

$$c = c_0 \succ_L d_0 = c_1 \succ_L d_1 = c_2 \succ_L \dots \succ_L d_{k-1} = c_k \succ_L d_k = d.$$

In  $2^{J(L)}$ , we have

$$\hat{r}(a) \setminus \hat{r}(b) = \dot{\cup}_{i=0}^n (\hat{r}(a_i) \setminus \hat{r}(b_i))$$

and similarly for  $\hat{r}(c) \setminus \hat{r}(d)$ . Therefore  $e \in (\hat{r}(a_i) \setminus \hat{r}(b_i)) \cap (\hat{r}(c_j) \setminus \hat{r}(d_j))$  for some  $i, j$ . Since  $a_i \succ_L b_i$  and  $c_j \succ_L d_j$ , this implies that  $\hat{r}(a_i) \setminus \hat{r}(b_i) = \hat{r}(c_j) \setminus \hat{r}(d_j)$ , which means  $a_i + b_i = c_j + d_j$ . By Proposition 6.3,  $a_i/b_i \equiv c_j/d_j$ . This contradicts  $a/b \perp c/d$ .

Suppose that  $a + b$  and  $c + d$  are disjoint in  $B(L)$ . Let  $x/y, z/w \in Q(E)$  be such that

$$a/b \supseteq x/y \equiv z/w \subseteq c/d.$$

Obviously  $\{a, b, x, y\}$  and  $\{c, d, z, w\}$  are compatible sets. Since  $x \ominus y = z \ominus w \leq a, c$  and  $x \ominus y = z \ominus w \leq b', d'$ , it follows that  $\{a, b, c, d, x \ominus y = z \ominus w\}$  is compatible as well. By Lemma 5.4, this implies that the sets of quotients

$$\begin{aligned} \mathbf{f}_1 &:= \{a/b, x/y, a^R/b^R, x^R/y^R\}, & \mathbf{f}_2 &:= \{c/d, z/w, c^R/d^R, z^R/w^R\}, & \text{and} \\ \mathbf{g} &:= \{a/b, c/d, a^R/b^R, c^R/d^R, x^R/y^R\} \end{aligned}$$

are compatible. Moreover, by Proposition 5.8,  $x^R/y^R = z^R/w^R$ . Let  $L_1, L_2$  and  $K$  be finite compatible 0, 1-sublattices of  $E$  such that  $\mathbf{f}_1 \subseteq Q(L_1), \mathbf{f}_2 \subseteq Q(L_2)$  and  $\mathbf{g} \subseteq Q(K)$ .

Obviously,  $a + b \geq_{B(L_1)} x + y$ . By Lemma 6.2,  $a + b =_{B(L_1)} a^R + b^R$  and  $x + y =_{B(L_1)} x^R + y^R$ . Therefore  $a^R + b^R \geq_{B(L_1)} x^R + y^R$ . Since  $B(L_1 \cap K)$  is a subalgebra of  $B(L_1)$ , we have  $a^R + b^R \geq_{B(L_1 \cap K)} x^R + y^R$ . Since  $B(L_1 \cap K)$  is a subalgebra of  $B(K)$ , this implies that  $a^R + b^R \geq_{B(K)} x^R + y^R$ . Similarly,  $c^R + d^R \geq_{B(K)} z^R + w^R = x^R + y^R$ . Since  $a + b$  and  $c + d$  are disjoint in  $B(L)$ , they are disjoint in  $B(L \cap K)$  and hence also in  $B(K)$ . By Lemma 6.2,  $a^R + b^R =_{B(K)} a + b$  and  $c^R + d^R =_{B(K)} c + d$ . Thus  $a^R + b^R$  and  $c^R + d^R$  are disjoint elements of  $B(K)$ . This implies that  $x^R + y^R = 0$ , so  $x^R = y^R$  and hence  $x = y$ . □

**PROPOSITION 6.5.** *Let  $E$  be a complete lattice-ordered effect algebra. Let  $\mathbf{f} = \{a_1/b_1, \dots, a_n/b_n\} \subseteq Q(E)$  be a compatible set of quotients. Let  $L$  be a finite compatible 0, 1-sublattice of  $E$  with  $\{a_1, b_1, \dots, a_n, b_n\} \in L$ . Then  $\mathbf{f}$  is orthogonal if and only if, for all  $i \neq j, a_i + b_i$  and  $a_j + b_j$  are disjoint in  $B(L)$ .*

**PROOF.** ( $\Rightarrow$ ): This follows from Lemma 6.4.

( $\Leftarrow$ ): By Lemma 6.4, the elements of  $\mathbf{f}$  are pairwise disjoint. It remains to prove that

$$|\mathbf{f}| = (a_1 \ominus b_1) \oplus \dots \oplus (a_n \ominus b_n)$$

exists. By assumption, the sum

$$(a_1 + b_1) \oplus \dots \oplus (a_n + b_n)$$

exists in the effect algebra  $B(L)$ . Since  $\phi_L$  is a morphism of effect algebras, the sum

$$\phi_L(a_1 + b_1) \oplus \dots \oplus \phi_L(a_n + b_n)$$

exists in  $E$ . It remains to observe that, for all  $i, \phi_L(a_i + b_i) = a_i \ominus b_i$ . □

It is now clear that, for every finite compatible 0, 1-sublattice  $L$  of a complete lattice ordered effect algebra  $E$ ,

$$\{a/b : a \succ_L b, a \in J(L)\}$$

is a compatible test of  $\Omega(E)$ . On the other hand, for a finite compatible and orthogonal set of quotients  $\mathbf{f} = \{a_1/b_1, \dots, a_n/b_n\}$  we have

$$+\mathbf{f} = (a_1 + b_1) \dot{\vee} \dots \dot{\vee} (a_n + b_n)$$

in every  $B(L)$  with  $\mathbf{f} \subseteq Q(L)$ , where  $L$  is a finite compatible 0, 1-sublattice of  $E$ .

**PROPOSITION 6.6.** *Let  $E$  be a complete lattice-ordered effect algebra. Let  $\mathbf{f}$  be a finite compatible set of quotients of  $E$ . Then the following are equivalent:*

- (a)  $\mathbf{f}$  is an event of  $\Omega(E)$ ;
- (b)  $\mathbf{f}$  is an orthogonal set of quotients;
- (c)  $\mathbf{f}$  is pairwise orthogonal.

**PROOF.** (a) implies (b) and (b) implies (c) by definition.

To show that (c) implies (a), we shall prove that there exists a compatible and orthogonal finite set  $\mathbf{t} \supseteq \mathbf{f}$  with  $|\mathbf{t}| = 1$ . Let  $\mathbf{f} = \{a_1/b_1, \dots, a_n/b_n\}$ . Let  $L$  be a finite compatible 0, 1-sublattice of  $E$  with  $\{a_1, b_1, \dots, a_n, b_n\} \subseteq L$ . Let  $(c_i)_{i=1}^{2k}$  be an  $L$ -chain representation of the complement of  $a_1 + b_1 + \dots + a_n + b_n$  in  $B(L)$ . By Proposition 6.5,

$$\mathbf{t} = \{a_1/b_1, \dots, a_n/b_n, c_2/c_1, \dots, c_{2k}/c_{2k-1}\}$$

is orthogonal. By Lemma 6.1, we have  $|\mathbf{t}| = 1$ . □

**COROLLARY 6.7.** *Let  $E$  be a complete lattice-ordered effect algebra. Let  $\mathbf{f} \subseteq Q(E)$  and  $\mathbf{g} \subseteq Q(E)$  be events of  $\Omega(E)$  such that  $\mathbf{f} \cup \mathbf{g}$  is compatible. Let  $L$  be a finite compatible 0, 1-sublattice of  $E$  with  $\mathbf{f}, \mathbf{g} \subseteq Q(L)$ . Then  $\mathbf{f} \perp \mathbf{g}$  if and only if  $+\mathbf{f} \perp_{B(L)} +\mathbf{g}$  and  $\mathbf{f} \text{ loc } \mathbf{g}$  if and only if  $+\mathbf{f} \perp_{B(L)} +\mathbf{g}$  and  $\phi_L((+\mathbf{f}) \dot{\vee} (+\mathbf{g})) = 1$ .*

**PROOF.** This follows from Propositions 6.5 and 6.6. □

**PROOF (Proof of Theorem 4.7).** Let  $\mathbf{f}$  be a finite pairwise orthogonal set of quotients. Since  $\mathbf{f} \equiv \mathbf{f}^R$ ,  $\mathbf{f}^R$  is pairwise orthogonal. By Corollary 5.6,  $\mathbf{f}^R$  is compatible. By Proposition 6.6,  $\mathbf{f}^R$  is an event of  $\Omega(E)$ , therefore there exists a test  $\mathbf{t}_0 \supseteq \mathbf{f}^R$ . Put  $\mathbf{t} = \mathbf{f} \cup (\mathbf{t}_0 \setminus \mathbf{f}^R)$ . By Lemma 4.6,  $\mathbf{t} \equiv \mathbf{t}_0$ . By Lemma 4.5,  $\mathbf{t}$  is a test. □

**PROPOSITION 6.8.** *Let  $E$  be a complete lattice-ordered effect algebra and let  $\mathbf{f}, \mathbf{g}$  be events of  $\Omega(E)$ . Then  $\mathbf{f} \perp \mathbf{g}$  if and only if, for all  $x/y \in \mathbf{f}$  and  $z/w \in \mathbf{g}$ , we have  $x/y \perp z/w$ .*

**PROOF.**  $\mathbf{f} \perp \mathbf{g}$  if and only if  $\mathbf{f} \cap \mathbf{g} \in \mathcal{N}$  and  $\mathbf{f} \cup \mathbf{g}$  is an event of  $\Omega(E)$ . The rest follows by Theorem 4.7. □

### 7. $\Omega(E)$ is algebraic

Let  $\mathbf{f}$  be a finite orthogonal set of quotients and let  $z/w \in Q(E)$ . We say that  $z/w$  is covered by  $\mathbf{f}$  if and only if there are  $z_1/w_1, \dots, z_n/w_n$  such that

- $z = z_1$ ,
- $w_n = w$ ,
- for all  $1 \leq i < n$ ,  $w_i = z_{i+1}$ ,
- for all  $1 \leq i \leq n$ , there are  $c/d$  and  $e/f$  such that  $z_i/w_i \equiv c/d \sqsubseteq e/f \in \mathbf{f}$ .

**PROPOSITION 7.1.** *Let  $E$  be a complete lattice-ordered effect algebra and let  $\mathbf{t}$  be a test of  $\Omega(E)$ . Let  $z/w$  be such that, for all  $e/f \in \mathbf{t}$ ,  $z \ominus w \leftrightarrow e \ominus f$ . Then  $z/w$  is covered by  $\mathbf{t}$ .*

**PROOF.** Let us write  $\mathbf{t} = \{e_1/f_1, \dots, e_m/f_m\}$ . By Corollary 5.5,  $\mathbf{t}^R \cup \{z^R/w^R\}$  is a compatible set of quotients. Let  $L$  be a finite compatible 0, 1-sublattice of  $E$  with  $\mathbf{t}^R \cup \{z^R/w^R\} \subseteq Q(L)$ . By Lemma 6.1,

$$\phi_L\left( (e_1^R + f_1^R) \dot{\vee} \dots \dot{\vee} (e_m^R + f_m^R) \right) = (e_1^R \ominus f_1^R) \oplus \dots \oplus (e_m^R \ominus f_m^R) = |\mathbf{t}| = 1.$$

Since  $\phi_L$  is faithful, this implies that  $(e_1^R + f_1^R) \dot{\vee} \dots \dot{\vee} (e_m^R + f_m^R) = 1$ .

Let  $z_1/w_1, \dots, z_n/w_n \subseteq Q(L)$  be such that

- $z^R = z_1$ ,
- $w_n = w^R$ ,
- for all  $1 \leq i < n$ ,  $z_i = w_{i+1}$ ,
- for all  $1 \leq i \leq n$ ,  $z_i \succ_L w_i$ .

In  $B(L)$ , we have

$$z^R + w^R = (z_1 + w_1) \dot{\vee} \dots \dot{\vee} (z_n + w_n).$$

Since each  $z_i + w_i$  is an atom of  $B(L)$ , we see that for every  $1 \leq i \leq n$  there exists some  $1 \leq j \leq m$  such that  $z_i + w_i \leq e_j^R + f_j^R$ . Therefore there exist  $c, d \in L$  such that  $z_i + w_i = c + d$  and  $e_j^R \geq c \succ d \geq f_j^R$ . By Proposition 6.3,  $z_i/w_i \equiv c/d$ .

Since  $z^R/w^R$  is covered by  $\mathbf{t}^R$ ,  $z/w$  is covered by  $\mathbf{t}$ . □

**THEOREM 7.2.** *For every complete lattice-ordered effect algebra  $E$ ,  $\Omega(E)$  is an algebraic generalized test space.*

**PROOF.** Let  $\mathbf{f}, \mathbf{g}, \mathbf{h}$  be such that  $\mathbf{f} \sim \mathbf{g}, \mathbf{g} \text{ loc } \mathbf{h}$ . We shall prove that  $\mathbf{f} \text{ loc } \mathbf{h}$ .

There is an event  $\mathbf{u}$  such that  $\mathbf{f} \text{ loc } \mathbf{u}$  and  $\mathbf{g} \text{ loc } \mathbf{u}$ . Since  $|\mathbf{f}| \oplus |\mathbf{h}| = 1$ , it suffices to prove that every pair of quotients  $a/b \in \mathbf{f}$  and  $c/d \in \mathbf{h}$  is disjoint. Assume the contrary and let  $x/y, z/w$  be proper quotients such that

$$a/b \supseteq x/y \equiv z/w \sqsubseteq c/d.$$

Since  $(x \ominus y) \oplus |\mathbf{u}|$  and  $(z \ominus w) \oplus |\mathbf{g}|$  exist and  $x \ominus u = z \ominus w$ , we see that, for all  $e/f \in \mathbf{u} \cup \mathbf{g}$ ,  $x \ominus y = z \ominus w \leq (e \ominus f)'$ . Therefore  $x \ominus y = z \ominus w \leftrightarrow e \ominus f$ . By Proposition 7.1, this implies that  $x/y$  is covered by the test  $\mathbf{u} \cup \mathbf{g}$ . However, since  $x/y \sqsubseteq a/b \in \mathbf{f} \perp \mathbf{u}$ ,  $x/y$  is disjoint with every element of  $\mathbf{u}$ . Therefore  $x/y$  is covered by  $\mathbf{g}$ . In particular, there exists a proper quotient  $x_1/y_1 \sqsubseteq x/y$  such that  $x_1/y_1 \sqsubseteq p/q$  for some  $p/q \in \mathbf{g}$ . As  $x_1/y_1 \sqsubseteq x/y$  and  $x/y \equiv z/w$ , there exists a proper quotient  $z_1/w_1 \sqsubseteq z/w$  such that  $z_1/w_1 \equiv x_1/y_1$ . Obviously  $z_1 \ominus w_1 \leq z \ominus w$  implies that, for all  $e/f \in \mathbf{u} \cup \mathbf{g}$ ,  $x_1 \ominus y_1 = z_1 \ominus w_1 \leq (e \ominus f)'$  and hence  $x_1 \ominus y_1 = z_1 \ominus w_1 \leftrightarrow e \ominus f$ . By Proposition 7.1, this implies that  $z_1/w_1$  is covered by the test  $\mathbf{u} \cup \mathbf{g}$ . Since  $z_1/w_1 \sqsubseteq z/w \sqsubseteq c/d \in \mathbf{h} \perp \mathbf{g}$ ,  $z_1/w_1$  is covered by  $\mathbf{u}$ . In particular, there is  $r/s \in \mathbf{u}$  such that there is a proper quotient  $z_2/w_2 \sqsubseteq z_1/w_1, r/s$ . As  $z_1/w_1 \equiv x_1/y_1$ , there is a proper quotient  $x_2/y_2 \sqsubseteq x_1/y_1$  such that  $x_2/y_2 \equiv z_2/w_2$ . We see that

$$p/q \supseteq x_1/y_1 \supseteq x_2/y_2 \equiv z_2/w_2 \sqsubseteq z_1/w_1 \sqsubseteq r/s$$

and  $x_2/y_2$  is proper. This is a contradiction with  $r/s \perp p/q$ . □

For a complete lattice-ordered effect algebra, we denote the orthoalgebra of  $\Omega(E)$  by  $O(E)$ .

**COROLLARY 7.3.** *For every complete lattice-ordered effect algebra  $E$ , the mapping  $\phi_E : O(E) \rightarrow E$  given by  $\phi_E([\mathbf{f}]_{\sim}) = |\mathbf{f}|$  is a surjective full morphism of effect algebras.*

**PROOF.** It is easy to check that  $\phi_E$  is a morphism of effect algebras. Let  $s, t \in E$  and suppose that  $s \oplus t$  exists. Then, in  $\Omega(E)$ ,  $\{s/0\} \perp \{s \oplus t/s\}$  and hence, in  $O(E)$ , the sum  $[\{s/0\}]_{\sim} \oplus [\{s \oplus t/s\}]_{\sim}$  exists. Since  $\phi_E([\{s/0\}]_{\sim}) = s$  for all  $s \in E$ ,  $\phi_E$  is surjective. □

To abbreviate our notations, let us write

- $\mathbf{f} \lesssim \mathbf{g}$  instead of  $[\mathbf{f}]_{\sim} \leq [\mathbf{g}]_{\sim}$ ,
- $a/b \perp \mathbf{f}$  instead of  $\{a/b\} \perp \mathbf{f}$ ,

- $a/b \lesssim \mathbf{f}$  instead of  $\{a/b\} \lesssim \mathbf{f}$ .

PROPOSITION 7.4. *Let  $E$  be a complete lattice-ordered effect algebra and let  $\mathbf{f}, \mathbf{g}$  be events of  $\Omega(E)$ . Then  $\mathbf{f} \lesssim \mathbf{g}$  if and only if, for all  $a/b \in Q(E)$ ,  $a/b \perp \mathbf{g}$  implies  $a/b \perp \mathbf{f}$ .*

PROOF. Suppose that  $\mathbf{f} \lesssim \mathbf{g}$ . There is an event  $\mathbf{v}$  such that  $\mathbf{v} \perp \mathbf{f}$  and  $\mathbf{v} \cup \mathbf{f} \sim \mathbf{g}$ . If  $a/b \perp \mathbf{g}$  then  $a/b \perp \mathbf{v} \cup \mathbf{f}$  and  $a/b \perp \mathbf{f}$ .

Suppose that, for all  $a/b \in Q(E)$ ,  $a/b \perp \mathbf{g}$  implies that  $a/b \perp \mathbf{f}$ . Let  $\mathbf{u}$  be a local complement of  $\mathbf{g}$ . By assumption, every quotient in  $\mathbf{u}$  is orthogonal to  $\mathbf{f}$ . By Proposition 6.8, this implies that  $\mathbf{f} \cup \mathbf{u}$  is an event. Let  $\mathbf{v}$  be a local complement of  $\mathbf{f} \cup \mathbf{u}$ . Then  $\mathbf{u}$  is a local complement of  $\mathbf{f} \cup \mathbf{v}$ . Consequently,  $\mathbf{f} \cup \mathbf{v} \sim \mathbf{g}$  and  $\mathbf{f} \lesssim \mathbf{g}$ .  $\square$

COROLLARY 7.5. *Let  $E$  be a complete lattice-ordered effect algebra and let  $\mathbf{f}, \mathbf{g}$  be events of  $\Omega(E)$ . Then  $\mathbf{f} \sim \mathbf{g}$  if and only if, for all  $a/b \in Q(E)$ ,  $a/b \perp \mathbf{f}$  if and only if  $a/b \perp \mathbf{g}$ .*

PROOF. This follows from Proposition 7.4.  $\square$

PROPOSITION 7.6. *Let  $E$  be a complete lattice-ordered effect algebra and let  $\mathbf{f}, \mathbf{g}$  be events of  $\Omega(E)$ . Then  $\mathbf{f} \lesssim \mathbf{g}$  if and only if, for all  $x/y \in \mathbf{f}$ ,  $x/y \lesssim \mathbf{g}$ .*

PROOF. Suppose that for all  $x/y \in \mathbf{f}$  we have  $x/y \lesssim \mathbf{g}$ . Let  $\mathbf{h}$  be a local complement of  $\mathbf{g}$ . Then  $\mathbf{f} \lesssim \mathbf{g}$  if and only if  $\mathbf{f} \perp \mathbf{h}$ . Let  $x/y \in \mathbf{f}$  and  $z/w \in \mathbf{h}$ . Since  $x/y \lesssim \mathbf{g} \perp \mathbf{h} \gtrsim z/w$ , we see that  $x/y \perp z/w$ . By Proposition 6.8,  $\mathbf{f} \perp \mathbf{h}$ .  $\square$

### 8. Perspectivity of sharp and compatible events

We say that an event  $\mathbf{f}$  of  $\Omega(E)$  *sharp* if and only if  $|\mathbf{f}|$  is sharp.

PROPOSITION 8.1. *Let  $E$  be a complete lattice-ordered effect algebra. Let  $\mathbf{g}$  be a sharp event of  $\Omega(E)$  and let  $\mathbf{f}$  be an event of  $\Omega(E)$ . Then  $\mathbf{f} \lesssim \mathbf{g}$  if and only if  $|\mathbf{f}| \leq |\mathbf{g}|$ .*

PROOF. Obviously,  $\mathbf{f} \lesssim \mathbf{g}$  implies that  $|\mathbf{f}| \leq |\mathbf{g}|$ .

Suppose that  $|\mathbf{f}| \leq |\mathbf{g}|$  and that  $a/b \perp \mathbf{g}$ . By Proposition 7.4, it suffices to prove that  $a/b \perp \mathbf{f}$ . Suppose that  $a/b \not\perp \mathbf{f}$ . By Proposition 6.8, this implies that  $a/b \not\perp c/d$  for some  $c/d \in \mathbf{f}$ . As  $(a \ominus b) \oplus |\mathbf{g}|$  exists,  $(a \ominus b) \oplus (c \ominus d)$  exists. Therefore  $a/b$  is not disjoint with  $c/d$ . In particular,  $(a \ominus b) \wedge (c \ominus d) > 0$ . However, we then have

$$0 < (a \ominus b) \wedge (c \ominus d) \leq a \ominus b \leq |\mathbf{g}'| \quad \text{and}$$

$$0 < (a \ominus b) \wedge (c \ominus d) \leq c \ominus d \leq |\mathbf{f}| \leq |\mathbf{g}|.$$

This is a contradiction with  $|\mathbf{g}| \in S(E)$ .  $\square$

**COROLLARY 8.2.** *Let  $E$  be a complete orthomodular lattice. Then the mapping  $\psi : O(E) \rightarrow E$  given by  $\psi([f]_{\sim}) = |f|$  is an isomorphism.*

**PROOF.** The proof is a trivial application of Proposition 8.1 and is omitted. □

**PROPOSITION 8.3.** *Let  $E$  be a complete lattice-ordered effect algebra. Let  $f, g$  be compatible events of  $\Omega(E)$  such that  $f \cup g$  is compatible. Let  $L$  be a finite compatible 0, 1-sublattice of  $E$  with  $f \cup g \subseteq Q(L)$ . Then  $f \lesssim g$  if and only if  $+f \leq +g$  in  $B(L)$ .*

**PROOF.** Let  $(c_i)_{i=1}^{2k}$  be an  $L$ -chain representation of the complement of  $+g$  in  $B(L)$  and write  $h = \{c_2/c_1, \dots, c_{2k}/c_{2k-1}\}$ . By Corollary 6.7,  $g \text{ loc } h$ .

Since  $\Omega(E)$  is algebraic,  $f \lesssim g$  is equivalent to  $f \perp h$ . By Corollary 6.7,  $f \perp h$  if and only if  $+f \perp +h$ . Obviously  $+f \perp +h$  if and only if  $+f \leq +g$ . □

**COROLLARY 8.4.** *Let  $M$  be a complete MV-effect algebra. Then the mapping  $\psi : O(M) \rightarrow B(M)$  given by  $\psi([f]_{\sim}) = +f$  is an isomorphism of effect algebras.*

**PROOF.** Let us prove that  $\psi$  is well-defined: suppose that  $f \sim g$ . Since  $M$  is an MV-effect algebra,  $f \cup g$  is compatible. By Proposition 8.3,  $+f = +g$ . Obviously  $\psi$  is surjective. Suppose that  $\psi(f) = \psi(g)$ , which means  $+f = +g$ . By Proposition 8.3,  $f \sim g$ .

It remains to prove that  $\psi$  is a homomorphism. Suppose that  $[f]_{\sim} \perp [g]_{\sim}$ . This implies that  $f \perp g$ . Let  $L$  be a finite 0, 1-sublattice of  $E$  with  $f \cup g \subseteq Q(L)$ . By Corollary 6.7,  $\psi(f) \perp \psi(g)$  and, obviously,

$$\psi([f]_{\sim} \oplus [g]_{\sim}) = \psi([f \cup g]_{\sim}) = (+f) \oplus (+g) = \psi([f]_{\sim}) \oplus \psi([g]_{\sim}). \quad \square$$

### 9. $O(E)$ is a lattice

Let  $f = \{a_1/b_2, \dots, a_n/b_n\}$  be a compatible event of  $E$ . In what follows,  $\langle f \rangle$  denotes the (finite distributive) 0, 1-sublattice of  $E$  generated by the set  $\{a_1, b_1, \dots, a_n, b_n\}$ .

Let  $f$  be an event of  $E$ . We denote the test

$$\{a/b : a \in J(\langle f^R \rangle), a \succ_L b\}$$

by  $t_f$ . We write

$$f^* = \{a/b \in t_f : a/b \perp f\}.$$

We have  $f \sim f^R$  and  $f^R \text{ loc } f^*$ . Since  $\Omega(E)$  is algebraic,  $f \text{ loc } f^*$ .

**PROPOSITION 9.1.** *Let  $E$  be a complete lattice-ordered effect algebra. Let  $f$  be an event of  $\Omega(E)$  and let  $a/b \in Q(E)$ . Then  $a/b \perp f$  if and only if  $a/b$  is covered by  $f^*$ .*

PROOF. Suppose that  $a/b \perp f$ . We shall prove that  $a/b$  is covered by  $f^*$ . For all  $c/d \in f$ ,  $a \ominus b \leftrightarrow c \ominus d$ . Therefore, by Corollary 5.5,  $\{a^R/b^R\} \cup f^R$  is a compatible set of quotients. Obviously  $\langle f^R \rangle \subseteq \langle f^R \cup \{a^R/b^R\} \rangle$ . As  $f^* \subseteq Q(\langle f^R \rangle)$ , this implies that  $\{a^R/b^R\} \cup f^*$  is compatible. Therefore, for all  $c/d \in f^R \cup f^*$ ,  $a \ominus b \leftrightarrow c \ominus d$ . By Proposition 7.1,  $a/b$  is covered by the test  $f^R \cup f^*$ . Since  $a/b \perp f^R$ ,  $a/b$  is covered by  $f^*$ .

Suppose that  $a/b$  is covered by  $f^*$ . As  $f \text{ loc } f^*$ , this implies that  $a/b \lesssim f^*$  and hence  $a/b \perp f$ . □

COROLLARY 9.2. *Let  $E$  be a complete lattice-ordered effect algebra. Let  $f, g$  be events of  $\Omega(E)$ . Then  $g \lesssim f$  if and only if every  $a/b \in g$  is covered by  $f^*$ .*

PROOF. Since  $f \text{ loc } f^*$ ,  $g \lesssim f$  if and only if  $g \perp f^*$ . By Proposition 6.8,  $g \perp f^*$  if and only if every  $a/b \in g$  is orthogonal to  $f^*$ . By Proposition 9.1,  $a/b \perp f^*$  if and only if  $a/b$  is covered by  $f^*$ . □

Let us write, for  $a/b \in Q(E)$  and  $p \in S(E)$ ,  $(a/b) \sqcap p = a \wedge p/b \wedge p$ . Note that the reduction map is a special case of  $\sqcap$  since  $a^R/b^R = (a/b) \sqcap (a \ominus b)^\dagger$ .

LEMMA 9.3. *Let  $E$  be a complete lattice-ordered effect algebra and let  $p, a, b \in E$ . If  $p \leftrightarrow a, b$  then  $\{a/b\} \sim \{(a/b) \sqcap p, (a/b) \sqcap p'\}$ .*

PROOF. Let  $L$  be a finite compatible 0, 1-sublattice of  $E$  with  $a, b, p \in L$ . An easy computation in  $B(L)$  yields  $(a \wedge p + b \wedge p) \dot{\vee} (a \wedge p' + b \wedge p') = a + b$ . By Proposition 8.3,  $\{a/b\} \sim \{(a/b) \sqcap p, (a/b) \sqcap p'\}$ . □

LEMMA 9.4. *If  $E$  is a complete lattice-ordered effect algebra then*

$$s \wedge (s \wedge t)^\dagger \leftrightarrow t \wedge (s \wedge t)^\dagger \quad \text{for all } s, t \in E.$$

PROOF. We have

$$s \wedge t = s \wedge (s \wedge t) \leq s \wedge (s \wedge t)^\dagger.$$

Similarly,  $s \wedge t \leq t \wedge (s \wedge t)^\dagger$ . Thus, we have

$$s \wedge (s \wedge t)^\dagger, t \wedge (s \wedge t)^\dagger \in [s \wedge t, (s \wedge t)^\dagger].$$

By Proposition 5.1,  $[s \wedge t, (s \wedge t)^\dagger]$  is a compatible set, hence

$$s \wedge (s \wedge t)^\dagger \leftrightarrow t \wedge (s \wedge t)^\dagger. \quad \square$$

LEMMA 9.5. *Let  $E$  be a complete lattice ordered effect algebra. Let  $a/b, c/d \in Q(E)$  be reduced. Then  $x/y \lesssim (a/b), (c/d)$  if and only if*

$$\begin{aligned} x/y &\lesssim (a/b) \sqcap ((a \ominus b) \wedge (c \ominus d))^\dagger \quad \text{and} \\ x/y &\lesssim (c/d) \sqcap ((a \ominus b) \wedge (c \ominus d))^\dagger. \end{aligned}$$

PROOF. We may assume that  $x/y$  is reduced. Suppose that  $x/y \lesssim (a/b), (c/d)$ . Then  $x \ominus y \leq (a \ominus b) \wedge (c \ominus d) \leq ((a \ominus b) \wedge (c \ominus d))^\dagger$ . By Proposition 5.1, since  $\{x \ominus y, a \ominus b, (a \ominus b) \wedge (c \ominus d)\}$  is a compatible set,  $\{x \ominus y, a \ominus b, ((a \ominus b) \wedge (c \ominus d))^\dagger\}$  is a compatible set. As  $x/y$  and  $a/b$  are reduced quotients, Corollary 5.5 implies that  $\{x, y, a, b, ((a \ominus b) \wedge (c \ominus d))^\dagger\}$  is a compatible set. By Lemma 9.3, we have

$$\{x/y\} \lesssim \{a/b\} \sim \left\{ (a/b) \sqcap ((a \ominus b) \wedge (c \ominus d))^\dagger, (a/b) \sqcap ((a \ominus b) \wedge (c \ominus d))^{\dagger'} \right\}.$$

Let  $u$  be a local complement of

$$\left\{ (a/b) \sqcap ((a \ominus b) \wedge (c \ominus d))^\dagger, (a/b) \sqcap ((a \ominus b) \wedge (c \ominus d))^{\dagger'} \right\}.$$

Since  $\{x, y, a, b, ((a \ominus b) \wedge (c \ominus d))^\dagger\}$  is a compatible set and  $x/y \perp u$ , we have  $x \ominus y \leftrightarrow e \ominus f$ , for all

$$e/f \in \left\{ (a/b) \sqcap ((a \ominus b) \wedge (c \ominus d))^\dagger, (a/b) \sqcap ((a \ominus b) \wedge (c \ominus d))^{\dagger'} \right\} \cup u.$$

Therefore, by Proposition 7.1,  $x/y$  is covered by

$$\left\{ (a/b) \sqcap ((a \ominus b) \wedge (c \ominus d))^\dagger, (a/b) \sqcap ((a \ominus b) \wedge (c \ominus d))^{\dagger'} \right\} \cup u.$$

However,  $x/y \perp u$  and, since  $x \ominus y \leq ((a \ominus b) \wedge (c \ominus d))^\dagger \in S(E)$ ,  $x/y$  and  $(a/b) \sqcap ((a \ominus b) \wedge (c \ominus d))^{\dagger'}$  are disjoint. Therefore,  $x/y \lesssim (a/b) \sqcap ((a \ominus b) \wedge (c \ominus d))^\dagger$ . Symmetrically, one can prove  $x/y \lesssim (c/d) \sqcap ((a \ominus b) \wedge (c \ominus d))^\dagger$ .

The converse implication follows by Lemma 9.3. □

LEMMA 9.6. *Let  $E$  be a complete lattice-ordered effect algebra. Suppose that  $a/b, c/d \in Q(E)$  are such that  $a \ominus b \leftrightarrow c \ominus d$ . Then  $[(a/b)]_{\sim} \wedge [(c/d)]_{\sim}$  exists in  $O(E)$  and equals  $[(a^R \wedge c^R / (b^R \vee d^R) \wedge (a^R \vee c^R))]_{\sim}$ .*

PROOF. Suppose that  $x/y \lesssim a/b, c/d$ . Since  $\{a \ominus b, c \ominus d, x \ominus y\}$  is mutually compatible, Corollary 5.5 implies that  $\{x^R/y^R, a^R/b^R, c^R/d^R\}$  is a compatible set of quotients. Thus, there is a finite compatible sublattice  $L$  of  $E$  with  $\{x^R/y^R, a^R/b^R, c^R/d^R\} \subseteq Q(L)$ . By Proposition 8.3,  $x^R + y^R \leq a^R + b^R, c^R + d^R$  in  $B(L)$ . A simple calculation in  $B(L)$  then yields

$$(a^R + b^R) \wedge (c^R + d^R) = (a^R \wedge c^R) + ((b^R \vee d^R) \wedge (a^R \vee c^R)),$$

hence we obtain

$$x^R + y^R \leq (a^R \wedge c^R) + ((b^R \vee d^R) \wedge (a^R \vee c^R)) \leq a^R + b^R, c^R + d^R.$$

Again, by Proposition 8.3, we obtain

$$x^R/y^R \lesssim a^R \wedge c^R / (b^R \vee d^R) \wedge (a^R \vee c^R) \lesssim a^R/b^R, c^R/d^R. \quad \square$$

LEMMA 9.7. *Let  $E$  be a complete lattice-ordered effect algebra. For all  $a/b, c/d \in Q(E)$ ,  $[(a/b)]_{\sim} \wedge [(c/d)]_{\sim}$  exists in  $O(E)$ .*

PROOF. We may assume that  $a/b$  and  $c/d$  are reduced. Following Lemma 9.5,  $[(a/b)]_{\sim} \wedge [(c/d)]_{\sim}$  exists if and only if

$$\left[ \left\{ (a/b) \sqcap ((a \ominus b) \wedge (c \ominus d))^{\uparrow} \right\} \right]_{\sim} \wedge \left[ \left\{ (c/d) \sqcap ((a \ominus b) \wedge (c \ominus d))^{\uparrow} \right\} \right]_{\sim}$$

exists, and if one exists, and hence the other does too, then they are equal. Let  $M$  be a block of  $E$  with  $a \ominus b, (a \ominus b) \wedge (c \ominus d) \in M$ . By Proposition 5.1,  $((a \ominus b) \wedge (c \ominus d))^{\uparrow} \in M$ . Since  $M \cap S(M) = C(M)$ ,  $((a \ominus b) \wedge (c \ominus d))^{\uparrow}$  is central in  $M$ . Therefore,

$$\begin{aligned} (9.1) \quad & \left| (a/b) \sqcap ((a \ominus b) \wedge (c \ominus d))^{\uparrow} \right| \\ &= \left( a \wedge ((a \ominus b) \wedge (c \ominus d))^{\uparrow} \right) \ominus \left( b \wedge ((a \ominus b) \wedge (c \ominus d))^{\uparrow} \right) \\ &= (a \ominus b) \wedge ((a \ominus b) \wedge (c \ominus d))^{\uparrow}. \end{aligned}$$

Similarly, we obtain

$$\left| (c/d) \sqcap ((a \ominus b) \wedge (c \ominus d))^{\uparrow} \right| = (c \ominus d) \wedge ((a \ominus b) \wedge (c \ominus d))^{\uparrow}.$$

By Lemma 9.4 (putting  $s = a \ominus b$  and  $t = c \ominus d$ ),

$$(a \ominus b) \wedge ((a \ominus b) \wedge (c \ominus d))^{\uparrow} \leftrightarrow (c \ominus d) \wedge ((a \ominus b) \wedge (c \ominus d))^{\uparrow}.$$

By Lemma 9.6,

$$\left[ \left\{ (a/b) \sqcap ((a \ominus b) \wedge (c \ominus d))^{\uparrow} \right\} \right]_{\sim} \wedge \left[ \left\{ (c/d) \sqcap ((a \ominus b) \wedge (c \ominus d))^{\uparrow} \right\} \right]_{\sim}$$

exists in  $O(E)$ . □

THEOREM 9.8. *Let  $E$  be a complete lattice-ordered effect algebra. Then  $O(E)$  is an orthomodular lattice.*

PROOF. It is well known that an orthoalgebra is a lattice if and only if it is a (lower or upper) semilattice. Therefore, it suffices to prove that for every pair  $\mathbf{f}, \mathbf{g}$  of events of  $\Omega(E)$ ,  $[\mathbf{f}]_{\sim} \wedge [\mathbf{g}]_{\sim}$  exists in the orthoalgebra  $O(E)$ . Let us write

$$\mathbf{f}^{**} = \{a_1/b_1, \dots, a_n/b_n\} \quad \text{and} \quad \mathbf{g}^{**} = \{c_1/d_1, \dots, c_m/d_m\}.$$

For  $i \in \{1, \dots, n\}$  and  $j \in \{1, \dots, m\}$  let  $e_{ij}/f_{ij}$  be such that

$$[e_{ij}/f_{ij}]_{\sim} = [a_i/b_i]_{\sim} \wedge [c_j/d_j]_{\sim}$$

and let

$$\mathbf{h} = \{e_{ij}/f_{ij} : i \in \{1, \dots, n\} \text{ and } j \in \{1, \dots, m\}\}.$$

By Theorem 4.7,  $\mathbf{h}$  is an event of  $\Omega(E)$  and, by Proposition 7.6,  $\mathbf{h} \lesssim \mathbf{f}^{**}, \mathbf{g}^{**}$ . It remains to prove that, for every  $\mathbf{u} \lesssim \mathbf{f}^{**}, \mathbf{g}^{**}$ , we have  $\mathbf{u} \lesssim \mathbf{h}$ . Let  $x/y \in \mathbf{u}$ . By Corollary 9.2,  $x/y$  is covered by  $\mathbf{f}^{**}$  and  $\mathbf{g}^{**}$ . By a simple induction with respect to  $m$  and  $n$ , it is easy to prove that

$$\{x/y\} \sim \{x_{ij}/y_{ij} : i \in \{1, \dots, n\} \text{ and } j \in \{1, \dots, m\}\},$$

where for each  $x_{ij}/y_{ij}$  we have  $x_{ij}/y_{ij} \sqsubseteq a_i/b_i, c_j/d_j$ . This implies that, for all  $i \in \{1, \dots, n\}$  and  $j \in \{1, \dots, m\}$ ,

$$[x_{ij}/y_{ij}]_{\sim} \leq [a_i/b_i]_{\sim} \wedge [c_j/d_j]_{\sim}.$$

Consequently, by Proposition 7.6,  $x/y \lesssim \mathbf{h}$  and, again by Proposition 7.6,  $\mathbf{u} \lesssim \mathbf{h}$ .  $\square$

### 10. $\phi_E, \phi_E^*$ , compatibility and blocks

In this section, we shall show that there is one to one correspondence between blocks of a complete lattice-ordered effect algebra  $E$  and blocks of  $O(E)$ . Under  $\phi_E$ , the (pre)image of a block is always a block. Moreover, we prove that  $E$ , as a lattice, embeds into  $O(E)$ .

LEMMA 10.1. *Let  $E$  be a complete lattice-ordered effect algebra and let  $\mathbf{f}$  be a reduced event of  $\Omega(E)$ . Then*

$$\mathbf{f} \sim \{|\mathbf{f}|^\downarrow/0\} \cup \{(a/b) \cap |\mathbf{f}|^\downarrow : a/b \in \mathbf{f}\}$$

PROOF. Since  $\mathbf{f}$  is reduced,  $\mathbf{f}$  is compatible. Let  $M$  be a block with  $\mathbf{f} \subseteq Q(M)$ . Since  $|\mathbf{f}| \in M$ ,  $|\mathbf{f}|^\downarrow \in M$ . Therefore, for all  $a/b \in \mathbf{f}$  we have  $|\mathbf{f}|^\downarrow \leftrightarrow a, b$ . By Lemma 9.3,

$$\{a/b\} \sim \{(a/b) \cap |\mathbf{f}|^\downarrow, (a/b) \cap |\mathbf{f}|^\downarrow\},$$

hence

$$(10.1) \quad \mathbf{f} \sim \bigcup_{a/b \in \mathbf{f}} \{(a/b) \cap |\mathbf{f}|^\downarrow, (a/b) \cap |\mathbf{f}|^\downarrow\}.$$

Since  $|\mathbf{f}|^\uparrow \in C(M)$ , we have

$$|\mathbf{f}| = \bigoplus_{a/b \in \mathbf{f}} a \ominus b = \bigoplus_{a/b \in \mathbf{f}} (a \ominus b) \wedge |\mathbf{f}|^\downarrow \bigoplus_{a/b \in \mathbf{f}} (a \ominus b) \wedge |\mathbf{f}|^\downarrow.$$

Since  $|f|^\downarrow \leq |f|$  and  $|f|^\downarrow \in S(E)$ , we see that

$$|f|^\downarrow = \bigoplus_{a/b \in f} (a \ominus b) \wedge |f|^\downarrow.$$

Since, for all  $a/b \in f$ ,

$$|(a/b) \cap |f|^\downarrow| = (a \wedge |f|^\downarrow) \ominus (b \wedge |f|^\downarrow),$$

we obtain

$$|f|^\downarrow = \bigoplus_{a/b \in f} \{|(a/b) \cap |f|^\downarrow|\} = \{|f|^\downarrow/0\}.$$

By Proposition 8.1, this implies that

$$\bigcup_{a/b \in f} \{(a/b) \cap |f|^\downarrow\} \sim \{|f|^\downarrow/0\}.$$

The rest follows from (10.1). □

**LEMMA 10.2.** *Let  $E$  be a complete lattice-ordered effect algebra and let  $[f]_\sim, [g]_\sim \in O(E)$ . Then  $[f]_\sim \leftrightarrow_{O(E)} [g]_\sim$  if and only if  $\phi_E([f]_\sim) \leftrightarrow_E \phi_E([g]_\sim)$ .*

**PROOF.** We may assume that  $f$  and  $g$  are reduced. By Lemma 10.1, we have

$$\begin{aligned} f &\sim \{|f|^\downarrow/0\} \cup \{(a/b) \cap |f|^\downarrow : a/b \in f\}^R, \\ g &\sim \{|g|^\downarrow/0\} \cup \{(a/b) \cap |g|^\downarrow : a/b \in g\}^R. \end{aligned}$$

Let  $M$  be a block with  $|f|, |g| \in M$ . Then  $\{|f|^\downarrow, |f|\} \cup \{|g|^\downarrow, |g|\} \in M$ . Since, for all  $c/d \in \{|f|^\downarrow/0\} \cup \{(a/b) \cap |f|^\downarrow : a/b \in f\}^R \cup \{|g|^\downarrow/0\} \cup \{(a/b) \cap |g|^\downarrow : a/b \in g\}^R$ ,  $c \ominus d \in M$ , we have  $c, d \in M$  for all such  $c/d$ . Let  $L$  be a finite compatible 0, 1-sublattice of  $E$  such that

$$\{|f|^\downarrow/0\} \cup \{(a/b) \cap |f|^\downarrow : a/b \in f\}^R \cup \{|g|^\downarrow/0\} \cup \{(a/b) \cap |g|^\downarrow : a/b \in g\}^R \subseteq Q(L).$$

Let  $t_L$  be the test  $\{e/f : e \in J(L) \text{ and } e \succ_L f\}$ .

By Proposition 8.3 and Corollary 6.7, it is easy to check that for every  $c/d \in Q(L)$  there exists  $h \subseteq t_L$  such that  $h \sim \{c/d\}$ . Therefore  $[f]_\sim$  and  $[g]_\sim$  are covered by the word  $(\{(a/b) \cap |f|^\downarrow : a/b \in f\})_\sim$ . Thus  $[f]_\sim \leftrightarrow [g]_\sim$ . □

**THEOREM 10.3.** *Let  $E$  be a complete lattice-ordered effect algebra.*

- (a) *If  $M$  is a block of  $E$  then  $\phi_E^{-1}(M)$  is a block of  $O(E)$ .*
- (b) *If  $B$  is a block of  $O(E)$  then  $\phi_E(B)$  is a block of  $E$ .*

PROOF. (a) By Lemma 10.2,  $\phi_E^{-1}(M)$  is a compatible subset of  $O(E)$ . We shall prove that  $\phi_E^{-1}(M)$  is a maximal compatible subset of  $O(E)$ . Let  $y \in O(E)$  and suppose that, for all  $x \in \phi_E^{-1}(M)$ ,  $x \leftrightarrow y$ . Then,  $\phi_E(\{y\} \cup \phi_E^{-1}(M)) \supseteq M$  is compatible in  $E$  by Lemma 10.2. Since  $M$  is a maximal compatible subset of  $E$ ,  $\phi_E(\{y\} \cup \phi_E^{-1}(M)) = M$ . Therefore  $\phi_E(y) \in M$  and  $y \in \phi_E^{-1}(M)$ .

(b)  $\phi_E(B)$  is compatible. Let  $M \supseteq \phi_E(B)$  be a block of  $E$ . By part (a),  $\phi_E^{-1}(M)$  is a block of  $O(E)$ . By the maximality of  $B$ ,  $B = \phi_E^{-1}(M)$  and we see that  $\phi_E(B) = \phi_E(\phi_E^{-1}(M)) = M$ . □

For a complete lattice-ordered effect algebra  $E$ , a mapping  $\phi_E^* : E \rightarrow O(E)$  is defined by  $\phi_E^*(x) = [\{x/0\}]_{\sim}$ . Note that  $\phi_E(\phi_E^*(x)) = x$ .

LEMMA 10.4. *Let  $E$  be a complete lattice-ordered effect algebra, let  $a/b \in Q(E)$  be reduced and let  $p \in S(E)$  be such that  $p \leftrightarrow a \ominus b$ . Then  $(a/b) \sqcap p$  is reduced.*

PROOF. We shall prove that  $x \leq b \wedge p$  and  $x \wedge ((a \wedge p) \ominus (b \wedge p)) = 0$  imply that  $x = 0$ .

Note that, since  $a/b$  is reduced and  $x, p \leftrightarrow a \ominus b$ ,  $\{x, a, b, p\}$  is a compatible set; let  $M \supseteq \{x, a, b, p\}$  be a block of  $E$ . We have  $(a \wedge p) \ominus (b \wedge p) = (a \ominus b) \wedge p$  since  $p$  is central in  $M$ . Moreover,

$$(10.2) \quad \begin{aligned} x \wedge (a \ominus b) &= x \wedge \left( ((a \ominus b) \wedge p) \vee ((a \ominus b) \wedge p') \right) \\ &= (x \wedge (a \ominus b) \wedge p) \vee (x \wedge (a \ominus b) \wedge p') = x \wedge (a \ominus b) \wedge p' \end{aligned}$$

and, since  $x \leq p$ ,  $x \wedge (a \ominus b) \wedge p' = 0$ . Since  $x \leq b$ ,  $x \wedge (a \ominus b) = 0$  and  $a/b$  is reduced,  $x = 0$ . □

THEOREM 10.5. *Let  $E$  be a complete lattice ordered effect algebra. Then  $\phi_E^*$  is a injective 0, 1-lattice homomorphism.*

PROOF. It is obvious that  $\phi_E^*(0) = 0_{O(E)}$  and that  $\phi_E^*(1) = 1_{O(E)}$  and that  $\phi_E^*$  is injective. Let  $a, c \in E$ . By Lemma 9.5,

$$\begin{aligned} \phi_E^*(a) \wedge \phi_E^*(c) &= [\{a/0\}]_{\sim} \wedge [\{c/0\}]_{\sim} = [\{a/0\} \sqcap (a \wedge c)^\dagger]_{\sim} \wedge [\{c/0\} \sqcap (a \wedge c)^\dagger/0]_{\sim} \\ &= [\{a \wedge (a \wedge c)^\dagger/0\}]_{\sim} \wedge [\{c \wedge (a \wedge c)^\dagger/0\}]_{\sim}. \end{aligned}$$

By Lemma 9.4,  $a \wedge (a \wedge c)^\dagger \leftrightarrow a \wedge (a \wedge c)^\dagger$ , hence we may apply Lemma 9.6 to obtain

$$[\{a \wedge (a \wedge c)^\dagger/0\}]_{\sim} \wedge [\{c \wedge (a \wedge c)^\dagger\}]_{\sim} = [\{a \wedge c \wedge (a \wedge c)^\dagger/0\}]_{\sim} = [\{a \wedge c/0\}]_{\sim}.$$

It remains to prove that  $\phi_E^*$  preserves joins; in other words, that

$$[\{a/0\}]_{\sim} \vee [\{c/0\}]_{\sim} = [\{a \vee c/0\}]_{\sim}.$$

This is equivalent to  $[\{1/a\}]_{\sim} \wedge [\{1/c\}]_{\sim} = [\{1/a \vee c\}]_{\sim}$ . We have

$$(1/a)^R = a'^{\uparrow}/a \wedge a'^{\uparrow}, \quad (1/c)^R = c'^{\uparrow}/c \wedge c'^{\uparrow}.$$

By Lemma 9.5,

$$\begin{aligned} & [\{a'^{\uparrow}/a \wedge a'^{\uparrow}\}]_{\sim} \wedge [\{c'^{\uparrow}/c \wedge c'^{\uparrow}\}]_{\sim} \\ &= \left[ \left\{ (a'^{\uparrow}/a \wedge a'^{\uparrow}) \sqcap (a' \wedge c')^{\uparrow} \right\} \right]_{\sim} \wedge \left[ \left\{ (c'^{\uparrow}/c \wedge c'^{\uparrow}) \sqcap (a' \wedge c')^{\uparrow} \right\} \right]_{\sim}. \end{aligned}$$

We see that

$$\begin{aligned} (a'^{\uparrow}/a \wedge a'^{\uparrow}) \sqcap (a' \wedge c')^{\uparrow} &= (a' \wedge c')^{\uparrow} / a \wedge (a' \wedge c')^{\uparrow} \\ (c'^{\uparrow}/c \wedge c'^{\uparrow}) \sqcap (a' \wedge c')^{\uparrow} &= (a' \wedge c')^{\uparrow} / c \wedge (a' \wedge c')^{\uparrow} \end{aligned}$$

and that, by Lemma 10.4, both quotients are reduced. Moreover, since

$$(a' \wedge c')^{\uparrow} \ominus (a \wedge (a' \wedge c')^{\uparrow}) = (1 \ominus a) \wedge (a' \wedge c')^{\uparrow} = a' \wedge (a' \wedge c')^{\uparrow}$$

and, similarly,

$$(a' \wedge c')^{\uparrow} \ominus (c \wedge (a' \wedge c')^{\uparrow}) = c' \wedge (a' \wedge c')^{\uparrow},$$

Lemma 9.4 implies that they are compatible. Therefore, we may apply Lemma 9.6 to compute the meet of their perspectivity classes. After an easy computation we obtain

$$\begin{aligned} & \left[ (a' \wedge c')^{\uparrow} / a \wedge (a' \wedge c')^{\uparrow} \right]_{\sim} \wedge \left[ (a' \wedge c')^{\uparrow} / c \wedge (a' \wedge c')^{\uparrow} \right]_{\sim} \\ &= \left[ (a' \wedge c')^{\uparrow} / (a \vee c) \wedge (a' \wedge c')^{\uparrow} \right]_{\sim}. \end{aligned}$$

Finally, it remains to observe that

$$(1/a \vee c)^R = (a' \wedge c')^{\uparrow} / (a \vee c) \wedge (a' \wedge c')^{\uparrow}. \quad \square$$

**COROLLARY 10.6.** *Let  $E$  be a complete lattice-ordered effect algebra. Then  $\phi_E^*(S(E))$  is a sub-orthomodular lattice of  $O(E)$ .*

**PROOF.** By Theorem 10.5,  $\phi_E^*(S(E))$  is closed with respect to  $0$ ,  $1$ ,  $\vee$ , and  $\wedge$ . It remains to prove that  $\phi_E^*(S(E))$  is closed with respect to  $'$ . Let  $a \in S(E)$ . Then  $\phi_E^*(a) = [\{a/0\}]_{\sim}$ . In  $O(E)$ , we have  $[\{a/0\}]_{\sim} = [\{1/a\}]_{\sim}$ . Since  $a, a' \in S(E)$ ,  $\{1/a\}$  is a sharp event of  $\Omega(E)$ . By Proposition 8.1,  $|\{1/a\}| = |\{a'/0\}|$  implies that  $\{1/a\} \sim \{a'/0\}$  and we see that

$$[\{1/a\}]_{\sim} = [\{a'/0\}]_{\sim} = \phi_E^*(a') \in \phi_E^*(S(E)). \quad \square$$

## References

- [1] M. K. Bennett and D. J. Foulis, 'Phi-symmetric effect algebras', *Found. Phys.* **25** (1995), 1699–1722.
- [2] ———, 'Interval and scale effect algebras', *Adv. in Appl. Math.* **19** (1997), 200–215.
- [3] G. Cattaneo, 'A unified framework for the algebra of unsharp quantum mechanics', *Internat. J. Theoret. Phys.* **36** (1997), 3085–3117.
- [4] C. C. Chang, 'Algebraic analysis of many-valued logics', *Trans. Amer. Math. Soc.* **89** (1959), 74–80.
- [5] F. Chovanec and F. Kôpka, 'Boolean D-posets', *Tatra Mt. Math. Publ.* **10** (1997), 1–15.
- [6] A. Dvurečenskij and S. Pulmannová, 'Tensor product of D-posets and D-test spaces', *Rep. Math. Phys.* **34** (1994), 251–275.
- [7] A. Dvurečenskij and S. Pulmannová, *New Trends in Quantum Structures* (Kluwer, Dordrecht and Ister Science, Bratislava, 2000).
- [8] D. Foulis and M. K. Bennett, 'Tensor products of orthoalgebras', *Order* **10** (1993), 271–282.
- [9] D. J. Foulis and M. K. Bennett, 'Effect algebras and unsharp quantum logics', *Found. Phys.* **24** (1994), 1325–1346.
- [10] D. J. Foulis, R. Greechie and G. Rüttimann, 'Filters and supports in orthoalgebras', *Internat. J. Theoret. Phys.* **35** (1995), 789–802.
- [11] D. J. Foulis and C. H. Randall, 'Operational quantum statistics. I. Basic concepts', *J. Math. Phys.* **13** (1972), 1667–1675.
- [12] R. Giuntini and H. Greuling, 'Toward a formal language for unsharp properties', *Found. Phys.* **19** (1989), 931–945.
- [13] G. Grätzer, *General Lattice Theory*, second edition (Birkhäuser, 1998).
- [14] R. Greechie, D. Foulis and S. Pulmannová, 'The center of an effect algebra', *Order* **12** (1995), 91–106.
- [15] S. Gudder, 'S-dominating effect algebras', *Internat. J. Theoret. Phys.* **37** (1998), 915–923.
- [16] ———, 'Sharply dominating effect algebras', *Tatra Mt. Math. Publ.* **15** (1998), 23–30.
- [17] J. Hashimoto, 'Ideal theory for lattices', *Math. Japon.* **2** (1952), 149–186.
- [18] G. Jenča, 'Blocks of homogeneous effect algebras', *Bull. Austral. Math. Soc.* **64** (2001), 81–98.
- [19] ———, 'Finite homogeneous and lattice ordered effect algebras', *Discrete Math.* **272** (2003), 197–214.
- [20] ———, 'Boolean algebras R-generated by MV-effect algebras', *Fuzzy sets and systems* **145** (2004), 279–285.
- [21] ———, 'Sharp and meager elements in orthocomplete homogeneous effect algebras', Technical report, (FEI STU Bratislava, 2004).
- [22] G. Jenča and S. Pulmannová, 'Quotients of partial abelian monoids and the Riesz decomposition property', *Algebra Universalis* **47** (2002), 443–477.
- [23] ———, 'Orthocomplete effect algebras', *Proc. Amer. Math. Soc.* **131** (2003), 2663–2671.
- [24] G. Jenča and Z. Riečanová, 'On sharp elements in lattice ordered effect algebras', *BUSEFAL* **80** (1999), 24–29.
- [25] F. Kôpka, 'D-posets of fuzzy sets', *Tatra Mt. Math. Publ.* **1** (1992), 83–87.
- [26] F. Kôpka and F. Chovanec, 'D-posets', *Math. Slovaca* **44** (1994), 21–34.
- [27] H. M. MacNeille, 'Extension of a distributive lattice to a Boolean ring', *Bull. Amer. Math. Soc.* **45** (1939), 452–455.
- [28] D. Mundici, 'Interpretation of AF  $C^*$ -algebras in Lukasiewicz sentential calculus', *J. Funct. Anal.* **65** (1986), 15–63.
- [29] C. H. Randall and D. J. Foulis, 'Operational quantum statistics. II. Manual of operations and their logics', *J. Math. Phys.* **13** (1972), 1667–1675.

- [30] Z. Riečanová, 'A generalization of blocks for lattice effect algebras', *Internat. J. Theoret. Phys.* **39** (2000), 855–865.
- [31] ———, 'Continuous lattice effect algebras admitting order continuous states', *Fuzzy sets and systems* **136** (2003), 41–54.

Department of Mathematics  
Faculty of Electrical Engineering and Information Technology  
Ilkovičova 3  
812 19 Bratislava  
Slovakia  
e-mail: [gejza.jenca@stuba.sk](mailto:gejza.jenca@stuba.sk)